RECEIVED: February 27, 2019

REVISED: July 17, 2019

ACCEPTED: July 26, 2019

PUBLISHED: August 12, 2019

Sensitivity of the intensity frontier experiments for neutrino and scalar portals: analytic estimates

Kyrylo Bondarenko,^a Alexey Boyarsky,^a Maksym Ovchinnikov^a and Oleg Ruchayskiy^b

^a*Intituut-Lorentz, Leiden University,
Niels Bohrweg 2, 2333 CA Leiden, The Netherlands*

^b*Discovery Center, Niels Bohr Institute, Copenhagen University,
Blegdamsvej 17, DK-2100 Copenhagen, Denmark*

E-mail: bondarenko@lorentz.leidenuniv.nl,
boyarsky@lorentz.leidenuniv.nl, ovchinnikov@lorentz.leidenuniv.nl,
oleg.ruchayskiy@nbi.ku.dk

ABSTRACT: In recent years, a number of intensity frontier experiments have been proposed to search for feebly interacting particles with masses in the GeV range. We discuss how the characteristic shape of the experimental sensitivity regions — upper and lower boundaries of the probed region, the maximal mass reach — depends on the parameters of the experiments. We use the SHiP and the MATHUSLA experiments as examples. We find a good agreement of our estimates with the results of the Monte Carlo simulations. This simple approach allows to cross-check and debug Monte Carlo results, to scan quickly over the parameter space of feebly interacting particle models, and to explore how sensitivity depends on the geometry of experiments.

KEYWORDS: Beyond Standard Model, Fixed target experiments, Electroweak interaction, Higgs physics

ARXIV EPRINT: [1902.06240](https://arxiv.org/abs/1902.06240)

Contents

1	Introduction: searching for feebly coupled particles	1
2	Lower boundary of the sensitivity region: main factors	5
3	Upper boundary of the sensitivity curve	6
4	Maximal mass probed	8
5	Number and momentum distribution of mesons and W's at SHiP and MATHUSLA	9
5.1	B and D mesons	9
5.2	Mesons at MATHUSLA	10
5.3	W bosons	11
6	Calculation of sensitivities	14
6.1	Efficiencies	14
6.2	Lower bound	14
6.3	Upper bound	16
6.4	Maximal mass probed	18
7	Comparison with simulations	18
7.1	HNLs	19
7.2	Scalars	20
8	Conclusions	20
A	Portals	23
B	Production and detection of portal particles	24
B.1	Production in proton-proton collisions	24
B.1.1	HNL production	24
B.1.2	Quarkonia and heavy flavour baryons	25
B.1.3	Scalar production	26
B.2	Main decay channels	28
B.2.1	HNL	28
B.2.2	Scalar	28
B.2.3	Visible branching ratio	28
B.3	Comparison with scalar models used by SHiP and MATHUSLA collaborations	29
C	Relation between momentum of HNL and meson momentum	30

D Geometry of the experiments	30
D.1 SHiP	30
D.2 MATHUSLA	30
E Analytic estimation of the upper bound: details	31
E.1 Fits of the spectra	31
E.2 Upper bound estimation	32
F Details of the sensitivity curve drawing	33
G Analytic estimation of the lower bound for particular masses	34
H HNLs at MATHUSLA for small mass	34

1 Introduction: searching for feebly coupled particles

The construction of the Standard Model has culminated with the confirmation of one of its most important predictions — the discovery of the Higgs boson. The quest for new particles has not ended, however. The observed but unexplained phenomena in particle physics and cosmology (such as neutrino masses and oscillations, dark matter, baryon asymmetry of the Universe) indicate that other particles exist in the Universe. It is possible that these particles evaded detection so far because they are too heavy to be created at accelerators. Alternatively, some of the hypothetical particles can be sufficiently light (lighter than the Higgs or W boson), but interact very weakly with the Standard Model sector (we will use the term *feeble interaction* to distinguish this from the weak interaction of the Standard Model). In order to explore this latter possibility, the particle physics community is turning its attention to the so-called *Intensity Frontier experiments*, see e.g. [1] for an overview. Such experiments aim to create high-intensity particle beams and use large detectors to search for rare interactions of feebly interacting hypothetical particles.

New particles with masses much lighter than the electroweak scale may be directly responsible for some of the BSM phenomena, or can serve as mediators (or “*portals*”), coupling to states in the “hidden sectors” and at the same time interacting with the Standard Model particles. Such portals can be renormalizable (mass dimension ≤ 4) or be realized as higher-dimensional operators suppressed by the dimensional couplings Λ^{-n} , with Λ being the new energy scale of the hidden sector. In the Standard Model there can only be three *renormalizable* portals:

- a scalar portal that couples gauge singlet scalar to the $H^\dagger H$ term constructed from a Higgs doublet field H_a , $a = 1, 2$;
- a neutrino portal that couples new gauge singlet fermion to the $\epsilon_{ab} \bar{L}_a H_b$ where L_a is the SU(2) lepton doublet and ϵ_{ab} is completely antisymmetric tensor in two dimensions;
- a vector portal that couples the field strength of a new U(1) field to the U(1) hypercharge field strength.

Let us denote a new particle by X . The interaction of X with the SM is controlled by the *mixing angle* θ_X — a dimensionless parameter that specifies the mixing between X and the corresponding SM particle: the SM neutrinos for the neutrino portal, the Higgs boson for the scalar portal and the hyperfield for the vector portal. The searches for such particles are included in the scientific programs of many existing experiments [2–16]. Although the LHC is a flagship of the *Energy Frontier* exploration, its high luminosity (especially in the Run 3 and beyond) means that huge numbers of heavy flavored mesons and vector bosons are created. This opens the possibility of supplementing the High Luminosity phase of the LHC with *Intensity Frontier* experiments associated with the existing interaction points. Several such experiments have been proposed: CODEX-b [17], MATHUSLA [18, 19], FASER [20, 21], and AL3X [22]. *Given that all these experiments can probe similar parameter spaces, it is important to be able to assess their scientific reach in a consistent way, under clearly specified identical assumptions.*

Detailed Monte Carlo (MC) simulations of both production and decays, complemented with background studies and detector simulations, offer ultimate sensitivity curves for each of the experiments.

Such simulations are however difficult to reproduce and modify. The modifications are nevertheless routinely needed because

- (a) Geometrical configurations of most experiments are not fully fixed yet and it is important to explore changes of the science reach with the modification of experimental designs;
- (b) Production or decays of GeV-mass feeble interacting particles involving quarks and mesons often requires the description outside of the validity range of both perturbative QCD and low-energy meson physics and is, therefore, subject to large uncertainties. This is the case for example for both scalar and neutrino portals (see e.g. [23–28] as well as the discussion in section 7). In particular,
 - Different groups use different prescription for scalar production [1, 19, 27–29]
 - the decay width and hadronic branching fractions for scalars with masses from ~ 0.5 GeV to few GeV are subject to large uncertainties, see [25, 26];
 - multi-hadronic HNL decays are not accounted for by any of the existing simulation tools. Yet they account for the largest part of the HNLs with masses around few GeV [24, 30].
- (c) Monte Carlo simulations are done for a limited set of model parameters and it is difficult to explore the overall parameter space and/or modify the sensitivity estimates for extended models (see e.g. the discussion and approach in [31])

With this in mind we gathered in one place a sufficiently simple and fully controlled (semi)analytic estimates. Such estimates emphasize the main factors that influence the sensitivity: *(i)* dependence on the model (parameters, physical assumptions); *(ii)* dependence on the geometry of the experiment; *(iii)* factors, related to the beam energy, etc. We present the final number as a convolution of these factors, which allows to modify any of them at will. As a result one can efficiently compare between several experimental designs; to identify the main factors that influence the sensitivity reach of a particular experiment/model; to *reuse* existing Monte Carlo sensitivities by separating them into

the experimental efficiencies and physical input (model, production/decay phenomenology) with the subsequent modification of one of these factors; to scan over the parameter space of different models as compared to those used in the MC simulations.

It turns out that the ratio between the sensitivities of the experiments to a great extent does not depend on the specific model of new physics, and is determined mainly by the geometry and collision energies of the experiments, which allow a comparison of the sensitivities in a largely model-independent way. To illustrate this point, we compare the potentials of two proposed experiments: the LHC-based MATHUSLA experiment [18, 19, 32–34] and a proton fixed target experiment using the proton beam of the Super Proton Synchrotron (SPS) at CERN — SHiP [35–37]. We analyze their sensitivity to the neutrino [38–43] and scalar [44–51] portals. For particle masses $M_X \lesssim m_{B_c}$ ¹ the main production channels are decays of heavy flavored mesons and W bosons [33, 36] (see also appendix B.1 for a brief overview). We concentrate on the mass range $M_X \gtrsim m_K$, since the domain of lower masses for the HNL and Higgs-like scalar is expected to be probed by the currently running NA62 experiment [15, 52].

The sensitivity of the experiments is determined by the number of events that one expects to detect for a set of given parameters. In realistic experiments such events should be disentangled from the “background” signals.

For SHiP, detailed simulations have shown that the number of background events is expected to be very low, so that the experiment is “background free” [35, 53–55]. For MATHUSLA, the background is also expected to be low [18, 19], although no simulation studies of background have been performed. Even in the most favorable case of $N_{\text{bg}} \ll 1$ one needs *on average* $\bar{N}_{\text{events}} = 2.3$ expected signal events to observe at least one event with the probability higher than 90%.² However, due to the lack of spectrometer, mass reconstruction and particle identification at MATHUSLA, the meaning of the discovery of 2.3 events in the two experiments is very different as there is no way to associate the signal with a model in MATHUSLA and further consolidate the discovery.

For both experiments considered here the production point (“target”) is separated from the detector decay volume (of length l_{det}) by some macroscopic distance $l_{\text{target-det}}$ (see appendix D). For such experiments the sensitivity curve has a typical “cigar-like shape” in the plane “mass vs. interaction strength”, see figure 1.

The number of decay events in the decay volume factorizes into

$$N_{\text{events}} = \sum_M N_{\text{prod},M} \times P_{\text{decay},M}, \quad (1.1)$$

where $N_{\text{prod},M}$ is the number of particles X that are produced from a mother particle M and $P_{\text{decay},M}$ is the decay probability. For $N_{\text{prod},M}$ we have

$$N_{\text{prod},M} \approx N_M \times \text{BR}_{M \rightarrow X} \times \epsilon_{\text{decay},M} \quad (1.2)$$

¹By $m_{\dots}$ we denote the masses of lightest flavour mesons, for example, kaons (m_K), D^+ (m_D), B^+ (m_B), etc.

²To obtain 95% confidence limit one should assume $\bar{N}_{\text{events}} = 3$, as the Poisson probability to see *at least one* event, while expecting 3 “on average” is 0.9502.

Figure 1. A typical cigar-like shape of the sensitivity region of Intensity Frontier experiments. The *upper boundary* is determined by the condition $l_{\text{decay}} \sim l_{\text{target-det}}$, i.e. particles do not reach the detector. The *lower boundary* of the sensitivity region is determined by the parameters at which decays become too rare.

Here, N_M is the number of parent particles produced at the experiment; in the case of mesons $N_M = N_{\text{meson}} = 2N_{\bar{q}q} \times f_{\text{meson}}$, where f_{meson} is the fragmentation fraction of a quark q into a given hadron, and $N_M = N_W$ in the case of the W bosons. $\text{BR}_{M \rightarrow X}$ is the total branching ratio of decay of the parent particle into X (see appendix B.1). Finally, ϵ_{decay} is the *decay acceptance* — the fraction of particles X whose trajectory intersects the decay volume, so that they *could* decay inside it.

The probability of decay into a state that can be detected is given by³

$$P_{\text{decay},M} = \left[\exp\left(-\frac{l_{\text{target-det}}}{l_{\text{decay}}}\right) - \exp\left(-\frac{l_{\text{target-det}} + l_{\text{det}}}{l_{\text{decay}}}\right) \right] \times \epsilon_{\text{det}} \times \text{BR}_{\text{vis}}, \quad (1.3)$$

where the branching ratio BR_{vis} is the fraction of all decays producing final states that can be registered. Finally, $\epsilon_{\text{det}} \leq 1$ is the *detection efficiency* — a fraction of all decays inside the decay volume for which the decay products could be detected. In the absence of detector simulations we optimistically assume a detector efficiency of MATHUSLA of $\epsilon_{\text{det}} = 1$. The decay length l_{decay} in eq. (1.3) is defined as

$$l_{\text{decay}} = c\tau_X\beta_X\gamma_X, \quad (1.4)$$

where τ_X is the lifetime of the particle X (see appendix B.2), β_X is its velocity and γ_X is the γ factor (which depends on the mother particle that produces X).

³Here we ignored that particles travel slightly different distances depending on their off-axis angle. Eq. (1.3) also neglects the energy distribution of the produced particles, assuming that all of them travel with the same average energy. This is essential for proper determining of the upper boundary and we will return to this in section 3.

The production branching ratio and the lifetime behave with the mixing angle as

$$\text{BR}_{\text{meson} \rightarrow X} \propto \theta_X^2, \quad \tau_X \propto \theta_X^{-2} \quad (1.5)$$

At the lower bound of the sensitivity the decay probability behaves as $P_{\text{decay}} \propto l_{\text{det}}/l_{\text{decay}}$, and as a consequence of (1.5) the number of events scales as

$$N_{\text{events,lower}} \propto \theta_X^4/\gamma_X \quad (1.6)$$

At the upper bound $P_{\text{decay}} \approx e^{-l_{\text{target-det}}/l_{\text{decay}}}$, and

$$N_{\text{events,upper}} \propto \theta_X^2 e^{-C\theta_X^2/\gamma_X}, \quad (1.7)$$

where C is some numerical factor (that depends on properties of X).

Larger γ factor suppresses the exponents in the expression for the decay probability (1.3). From (1.6), (1.7) we see that this affects the upper and lower bounds of the cigar-like sensitivity plots in the opposite ways. For the lower bound, an experiment with the smaller average γ factor is sensitive to small coupling constants. For sufficiently large couplings, larger γ factor ensures that particles do not decay before reaching the detector, thus increasing the sensitivity to the upper range of the sensitivity curve.

The paper is organized as follows. In sections 2–4 we discuss the lower and upper boundaries of the sensitivity region, the maximal mass that can be probed and experimental parameters that affect them. In section 5 we discuss the total amount and energy distribution of charm- and beauty mesons at both SHiP and MATHUSLA experiments, as well as the contribution from the W bosons. In section 6 we summarize and discuss our results, while in section 7 we compare our approach with results of official simulations. Finally, in section 8 we make conclusions. Appendices A–H provide details of computations and relevant supplementary information.

2 Lower boundary of the sensitivity region: main factors

As we will see later (see section 6), the production from the W bosons does not give a contribution to the lower bound of the sensitivity curve for neither of the two experiments, and for neither of the two models discussed. So, in this section we will consider only the production from the mesons.

Let us first estimate the *lower boundary* of the sensitivity region, where $l_{\text{decay}} \gg l_{\text{det}}, l_{\text{target-det}}$. For the number of events (1.1) we have

$$N_{\text{events,lower}} \approx N_{\text{meson}} \times \text{BR}_{M \rightarrow X} \times \frac{\langle l_{\text{det}} \rangle}{c\tau_X \langle \gamma_X \rangle} \times \epsilon_X, \quad (2.1)$$

where $\epsilon_X \equiv \epsilon_{\text{prod}} \times \epsilon_{\text{decay}} \times \text{BR}_{\text{vis}}$ is the overall efficiency and τ_X is the lifetime of the particle X (see the discussion below eq. (1.4)). The particles are assumed to be relativistic (we will see below when this assumption is justified), so that $\beta_X \approx 1$. We estimate the γ factor γ_X from that of the parent meson:

$$\gamma_X \approx \gamma_{\text{meson}} \frac{\langle E_X^{\text{rest}} \rangle}{M_X}, \quad (2.2)$$

Figure 2. The number of decay events for HNL with mass $M_N = 3$ GeV as a function of U_ϵ^2 . The number of mesons is taken $N_{\text{meson}} = 10^{14}$, the γ factor is $\langle \gamma_N \rangle = 15$, the efficiency $\epsilon = 1$, and the distances $l_{\text{target-det}} = l_{\text{det}} = 50$ m. The decay width can be found from eq. (B.6). The dashed blue line corresponds to U_{max}^2 (Equation (3.1)), while the dashed red line corresponds to the estimate of the upper bound based on eq. (3.3). Small discrepancy between the position of the upper bound and the estimate is caused by logarithmic errors in (3.3).

The average formula (2.2) does not take into account the distribution of HNLs (scalars) in the meson rest frame — some of the new particles fly in the direction of the parent meson and have γ_X larger than (2.2), while the other fly in the opposite direction. We show below that this *does not play a role* for the lower boundary of the sensitivity curve while the *upper* boundary is exponentially sensitive to the high γ -factor tail of the distribution and therefore cannot be determined from eq. (2.2). For the experiments like FASER this difference plays an essential role, see [56].

Since at the lower bound $N_{\text{events}} \propto \theta_X^4$ (see eq. (1.6)), for the ratio of the mixing angles at the lower bound, we have

$$\frac{(\theta_{X,\text{lower}}^{\text{SHiP}})^2}{(\theta_{X,\text{lower}}^{\text{MAT}})^2} = \sqrt{\frac{N_{\text{events}}^{\text{MAT}}}{N_{\text{events}}^{\text{SHiP}}}} \simeq \sqrt{\frac{N_{\text{meson}}^{\text{MAT}}}{N_{\text{meson}}^{\text{SHiP}}} \times \frac{l_{\text{det}}^{\text{MAT}}}{l_{\text{det}}^{\text{SHiP}}} \times \frac{\langle \gamma_{\text{meson}}^{\text{SHiP}} \rangle}{\langle \gamma_{\text{meson}}^{\text{MAT}} \rangle} \times \frac{\epsilon_{\text{MAT}}}{\epsilon_{\text{SHiP}}}}, \quad (2.3)$$

where we assumed that the same meson is the main production channel at both the SHiP and MATHUSLA experiments for the given mass M_X of the new particle, so the branching ratio $\text{BR}_{\text{meson} \rightarrow X}$ from eq. (2.1) disappears. Therefore, to make a comparison between the experiments we only need to know the total number of mesons, their average γ factor, the decay volume length and the overall efficiency.

3 Upper boundary of the sensitivity curve

If particles have sufficiently large interaction strength (i.e., the mixing angles), they decay before reaching the decay volume. This determines the upper bound of the sensitivity curve, that we call $\theta_{X,\text{upper}}^2$.

A useful quantity to consider is a mixing angle for which the amount of decays inside the decay volume is maximal, $\theta_{X,\text{max}}$. It can be found using the asymptotic behavior for the number of events N_{events} from the estimations (1.6), (1.7). In the domain $l_{\text{decay}} \gg l_{\text{target-det}}$

for a fixed mass M_X it follows that N_{events} monotonically grows as θ_X^4 with the increase of θ_X , while in the domain $l_{\text{decay}} \ll l_{\text{target-det}}$ it falls exponentially. The position of the maximum θ_{max} can be found from

$$l_{\text{decay}}(M_X, \theta_{\text{max}}^2) \simeq \begin{cases} 1.5l_{\text{target-det}}, & \text{if } l_{\text{target-det}} \simeq l_{\text{det}} \\ 0.5l_{\text{target-det}}, & \text{if } l_{\text{target-det}} \gg l_{\text{det}} \end{cases} \quad (3.1)$$

Using θ_{max} , we can estimate the value of θ_{upper} assuming that all the particles X have the same (average) energy $\langle E_X \rangle$. If we neglect the second exponent in the expression for the decay probability (1.3), then the formula for the number of events (1.1) becomes

$$N_{\text{events}} \simeq N_{\text{prod}} \times \epsilon_{\text{det}} \times \text{BR}_{\text{vis}} \times e^{-l_{\text{target-det}}/l_{\text{decay}}} \quad (3.2)$$

We can estimate the exponent in (3.2) as $l_{\text{target-det}}/l_{\text{decay}} \approx \theta_X^2/\theta_{\text{max}}^2$, see eq. (3.1). So imposing the condition $N_{\text{events}} \simeq 1$ in eq. (3.2) with the logarithmic precision we get

$$\theta_{\text{upper}}^2 \simeq \theta_{\text{max}}^2 \times \log \left[N_{\text{prod}}(\theta_{\text{max}}^2) \epsilon_{\text{det}} \text{BR}_{\text{vis}} \right]. \quad (3.3)$$

An example of the dependence of the number of events on θ_X^2 for the fixed mass M_X , together with the estimation of the θ_X for the maximal number of events given by (3.1) and the upper bound predicted by (3.3), is shown in figure 2.

Of course, it is not sufficient to use only the average energy $\langle E_X \rangle$ to estimate the position of the upper boundary. Indeed, the decrease of $c\tau_X$ with the growth of θ_X^2 can be compensated by the increase of the energy E_X and, therefore, of the γ -factor. As a result the particles with $E_X > \langle E_X \rangle$ can reach the detector even if the mixing angle θ_X is larger than the estimate (3.3).

The expression (3.3) helps to estimate how the sensitivity curve depends on the parameters of the experiment and on various assumptions. In particular, we can now estimate how large is a mistake from using $\langle E_X^{\text{rest}} \rangle$ in eq. (2.2) rather than the actual E_X distribution. In order to do that we replaced $\langle E_X^{\text{rest}} \rangle \rightarrow m_{\text{meson}}$ — the maximal energy of the particle X in the meson's rest frame. This substitution increases the γ_X by a factor of 2. The estimates (3.1)–(3.3) show that θ_{max}^2 and as a result θ_{upper}^2 will shift by the same factor of 2. This number indicates an upper bound on the possible error, introduced by the approximate treatment.

Next, we turn to the exact treatment. To this end we consider the energy distribution of the X particles,

$$f_X(E_X) = \frac{1}{N_X} \frac{dN_X}{dE_X}. \quad (3.4)$$

Taking into account this distribution, the formula for the decay probability (1.3) at the upper bound should be modified as

$$P_{\text{decay}} = \epsilon_{\text{det}} \times \text{BR}_{\text{vis}} \times \int_0^\infty dE_X f_X(E_X) \times \pi \left(\frac{\tau_X p_X}{l_{\text{target-det}} M_X} \right), \quad (3.5)$$

Figure 3. The function $\pi(y)$ that determines the position of the upper boundary (see eq. (3.6)). We assumed $l_{\text{target-det}} = l_{\text{det}}$.

where an argument of π function is $l_{\text{decay}}/l_{\text{target-det}}$ and we used the expression for the decay length (1.4). The function $\pi(y)$, defined via

$$\pi(y) \equiv \exp\left(-\frac{1}{y}\right) - \exp\left(-\frac{l_{\text{target-det}} + l_{\text{det}}}{l_{\text{target-det}}} \frac{1}{y}\right), \quad (3.6)$$

determines a “window” of energies in which the shape of $f_X(E_X)$ distribution (rather than the average number of particles) contributes to the overall probability. $\pi(y)$ is shown in figure 3. For small energies (small y) $\pi(y)$ is exponentially small, while for large energies (large y) $\pi(y)$ is inversely proportional to energy and decreases slowly. Therefore, a sufficiently long “tail” of high-energy mesons can contribute to the integral in (3.5), but this range cannot be estimated without knowledge of the distribution function f_X . We will discuss f_X for mesons and W bosons in section 5.

4 Maximal mass probed

The maximal mass probed by the experiment is defined as the mass at which the lower sensitivity bound meets the upper sensitivity bound. It can be estimated from the condition that the decay length, calculated at the lower bound θ_{lower} (see section 2), is equal to the distance from the target to the decay volume of the given experiment:

$$l_{\text{decay}}(M_{X,\text{max}}, \theta_{\text{lower}}^2(M_{X,\text{max}})) \simeq l_{\text{target-det}}. \quad (4.1)$$

The decay length (1.4) depends on the mass as $l_{\text{decay}} \propto M_X^{-\alpha-1}$, where the term α in the exponent approximates the behaviour of the lifetime with the mass, and the term 1 comes from the γ factor.

Using the condition (4.1), the maximal mass probed can be estimated as

$$M_{X,\text{max}} \propto \left(\frac{\langle E_X \rangle}{|\theta_{\text{lower}}|^2 l_{\text{target-det}}} \right)^{\frac{1}{\alpha+1}}, \quad (4.2)$$

which results in the following ratio of the maximal mass probed at the SHiP and MATHUSLA experiments:

$$\frac{M_{X,\max}^{\text{SHIP}}}{M_{X,\max}^{\text{MAT}}} \simeq \left(\frac{\langle E_X \rangle^{\text{SHIP}}}{\langle E_X \rangle^{\text{MAT}}} \times \frac{|\theta_{X,\text{lower}}^{\text{MAT}}|^2}{|\theta_{X,\text{lower}}^{\text{SHIP}}|^2} \times \frac{l_{\text{target-det}}^{\text{MAT}}}{l_{\text{target-det}}^{\text{SHIP}}} \right)^{\frac{1}{\alpha+1}}. \quad (4.3)$$

For Higgs-like scalars we have $\alpha \approx 2$, while for HNLs it is $\alpha \approx 5$, see appendix B.2.

The estimate of the maximal mass probed (4.2) is applicable only if the result does not exceed the kinematic threshold; for the production from B mesons for the HNLs it is $m_{B_c} - m_l$ or $m_B - m_l$ depending on whether amount of produced B_c mesons is large enough to be relevant for the production (see the discussion in section 5.1), and for the scalars it is $m_B - m_\pi$.

5 Number and momentum distribution of mesons and W 's at SHiP and MATHUSLA

In this section, we discuss the number and distribution of charm and beauty mesons and of W bosons at SHiP and MATHUSLA experiments. As we have seen, to estimate the lower boundary we need only the number of parent particles and their *average* γ factors (see eqs. (2.1), (4.3)). On the other hand, for the estimation of the upper boundary we also need the energy distribution of the mesons and W (see section 3).

5.1 B and D mesons

The main production channel of HNLs in the mass range $M_N \lesssim m_{D_s}$ is the two-body leptonic decay of D_s mesons. For masses $m_{D_s} \lesssim M_N \lesssim m_{B_c}$ the main contribution comes from decays of B mesons, see, e.g., [24].⁴ For masses $M_N \gtrsim 3$ GeV the main HNL production channel is determined by the value of the fragmentation fraction of B_c mesons, f_{B_c} : in the case $f_{B_c} \gtrsim 10^{-4}$ it is the two-body decay of the B_c meson, while for smaller values it is the two-body decay of the B^+ meson [31]. For the scalars the production from D mesons is negligible as compared to the $B^{+/\text{0}}$ mesons decays even for masses $m_K \lesssim M_S \lesssim m_D$. The B_c mesons are not relevant for their production (see, e.g., [23, 28]). The branching ratios of the production of the HNLs and the scalars used for our estimations are given in appendix B.1.

For the LHC energies the fragmentation fraction f_{B_c} was measured at the LHCb [57] and found to be $f_{B_c} \approx (2.6 \pm 1.3) \times 10^{-3}$. Earlier measurements at the Tevatron give a similar value $f_{B_c} \approx (2 \pm 1) \times 10^{-3}$ [58–60], which is in good agreement with [57]. Therefore, at the LHC the B_c decay is the main production channel for heavy HNLs. However, at the energies of the SHiP experiment, $\sqrt{s} \simeq 30$ GeV, currently there is no experimental data on f_{B_c} . Additionally, the theoretical predictions of f_{B_c} (see, e.g., [61–63]) disagree with the LHC and Tevatron measurements at least by an order of magnitude, which also makes them untrustable at SHiP's energies. As a result, the value of f_{B_c} at SHiP experiment is unknown. In order to estimate the effect of this uncertainty, we perform our analysis of

⁴This statement is true for HNLs with dominant mixing with $\nu_{e/\mu}$. For dominant mixing with ν_τ the main production channel is from τ leptons for $M_N \lesssim m_\tau$ and from B mesons for larger masses [24].

the sensitivity of the SHiP experiment for two extreme cases: (i) SHiP's f_{B_c} at the same level as at the LHC, and (ii) $f_{B_c} = 0$, “no B_c mesons”.

Let us now discuss the available data. For the SHiP experiment, the amounts of produced charmed and beauty mesons (except the B_c mesons) were obtained in detailed PYTHIA simulations; the corresponding numbers can be found in [64] and are reproduced in table 1. We estimate the spectrum of the B_c mesons from the spectrum of the B^+ mesons by rescaling the energy $E_{B_c} = (m_{B_c}/m_B)E_B$ for the events with B^+ mesons. For MATHUSLA experiment, the situation is different: there is no available data with detailed simulations that give us the relevant properties of the mesons, so we discuss them below.

5.2 Mesons at MATHUSLA

In order to estimate the number of mesons and their γ factors for the MATHUSLA experiment, one needs to know their p_T distribution at ATLAS/CMS in the MATHUSLA pseudorapidity range $0.9 < \eta < 1.6$ (see appendix D). The relevant distributions were measured for B^+ mesons by the CMS collaboration [65] (13 TeV) with the p_T cut $p_T^B \geq 10$ GeV, and for D^+/D^0 mesons by the ATLAS collaboration [66] (7 TeV) for $p_T^D \geq 3.5$ GeV. We show the spectra obtained in these papers in figure 4.

The low p_T mesons, unaccounted for these studies, are the most relevant for the MATHUSLA sensitivity estimate because of two reasons. Firstly, the p_T spectrum of the hadrons produced in pp collisions has a maximum at $p_T \sim \text{few GeV}$ (see, e.g., experimental papers [67, 68], theoretical paper [69] and references therein), and therefore we expect that most of the D or B mesons have p_T s below the LHC cuts. Secondly, low p_T mesons produce decay products with the smallest γ factor, and therefore with the shortest decay length (1.4) and the largest probability to decay inside the decay volume (here we consider the case $l_{\text{decay}} \gg l_{\text{target-det}}$). Therefore, by shifting the position of the peak to smaller p_T s, we increase the number of mesons and decrease their average γ factor, and both of these effects enhance the number of events at the lower bound (2.1). Therefore an accurate prediction of the distribution $d\sigma/dp_T$ in the domain of low p_T s is very important.

In order to evaluate the distribution of heavy flavored mesons at low p_T and also to estimate D meson production cross-section at $\sqrt{s} = 13$ TeV we use FONLL (Fixed Order + Next-to-Leading Logarithms) — a model for calculating the single inclusive heavy quark production cross section which convolutes perturbative cross section with non-perturbative fragmentation function, see [69–72] for details.

Predictions of FONLL have been calibrated against the accelerator data and were found to be in very good agreement, see e.g. [65, 66, 73–75]. In particular, comparison of the FONLL simulations of the production of the B^+ s with the measurements at the Tevatron and at the LHC showed that FONLL predicts the low p_T distribution accurately. We show the central values of the FONLL predictions down to $p_T = 0$, confronted with the measurements of the CMS [65] and ATLAS [66] collaborations in figure 4. As expected, the distributions have maxima, after which they fall. We see, however, that the central predictions of FONLL for the differential cross-sections typically lie below the uncertainty range of the experimental cross-section, which results in a somewhat lower total cross-sections. Namely, integrating the central predictions over the experimentally measured p_T s, we have

Figure 4. Comparison of the p_T spectra of B^+ , D^+ mesons predicted by the FONLL simulations (red points) with the measurements of the ATLAS and CMS collaborations [65, 66] (blue points with uncertainties bars). Only the central values of the FONLL predictions are shown. See text for details.

$\sigma_{D,\text{FONLL}}/\sigma_{D,\text{exp}} \approx 0.4$ and $\sigma_{B,\text{FONLL}}/\sigma_{B,\text{exp}} \approx 0.7$. However, as is demonstrated in the same papers [65, 66], the agreement between the FONLL predictions and the experimental data is much better if one uses the upper bound of the FONLL predictions defined by the theoretical uncertainties.

Using the results of the FONLL simulations, we find the amounts of low p_T mesons traveling in the MATHUSLA direction:

$$\frac{N_D|_{p_T < 3.5 \text{ GeV}}}{N_D|_{p_T > 3.5 \text{ GeV}} = 3.8, \quad \frac{N_B|_{p_T < 10 \text{ GeV}}}{N_B|_{p_T > 10 \text{ GeV}} = 5.7 \quad (5.1)$$

This justify our statement that most of the B and D mesons have the p_T below the cuts in the currently available experimental papers [65, 66].

FONLL does not provide the distributions of the D_s and the B_c mesons. We approximate their distributions by those the D^+ and B^+ distributions. In the case of the B_c mesons we justify this approximation by comparing the distributions provided by BCVEGPY 2.0 package [76] (that simulates the distribution of the B_c mesons and was tested at the LHC energies) for the B_c meson with that of FONLL for the B^+ meson. We conclude that the p_T and η distributions of B_c and B^+ have similar shapes.

The relevant parameters — the total number of mesons, the average γ factor of the mesons that are produced in the direction of the decay volume of the experiments and the geometric acceptances $\epsilon_{\text{geom,meson}}$ for the mesons — are given in table 1.

5.3 W bosons

The production channel from the decays of W bosons is only relevant for the MATHUSLA experiment since the center of mass energy at SHiP experiment is not enough to produce on-shell W bosons.

The total W boson production cross-section at the LHC energies $\sqrt{s} = 13 \text{ TeV}$ was measured in [77] as $\sigma_{W \rightarrow N+l} \approx 20.5 \text{ nb}$. The corresponding number of W bosons produced during the high luminosity phase of the LHC is

$$N_{W,\text{total}} \approx 6 \cdot 10^{11} \quad (5.2)$$

Experiment	N_D	N_B	$\langle\gamma_D\rangle$	$\langle\gamma_B\rangle$	$\epsilon_{\text{geom},D}$	$\epsilon_{\text{geom},B}$
MATHUSLA	4.4×10^{16}	3×10^{15}	2.6	2.3	1.3×10^{-2}	1.8×10^{-2}
SHiP	1.6×10^{18}	1.1×10^{14}	19.2	16.6	—	—

Table 1. Parameters of the SHiP and MATHUSLA experiments: the total number of all charmed/beauty hadrons; the average γ factor of mesons flying in the direction of the decay volumes of the experiment; the geometric acceptances for these hadrons. We take B_c meson distribution to be proportional to that of B^+ mesons, scaled by f_{B_c} . As a result, B_c gamma factor is the same as for B^+ mesons for SHiP and scaled by m_B/m_{B_c} for MATHUSLA, see discussion in section 5.1 and 5.2 For SHiP we assumed 5 years of operation (2×10^{20} protons on target) and for MATHUSLA we took the luminosity of the HL phase, $\mathcal{L}_h = 3000 \text{ fb}^{-1}$. Predictions are based on the FairSHiP simulations (SHiP) and on the FONLL simulations (MATHUSLA). See text for details.

The p_T distribution of the W bosons at the LHC in the pseudorapidity range $|\eta| < 2.5$ and for energies $\sqrt{s} = 7 - 8 \text{ TeV}$ was measured by the ATLAS and CMS collaborations [78, 79]. Their results show that most of the vector bosons are produced with low p_T (below 10 GeV or so). However, these results do not give us the magnitude of the W 's average momentum $\langle p_W \rangle$, needed to estimate the decay acceptance and the average momentum of HNLs.

In order to obtain $\langle p_W \rangle$ we have simulated the process $p + p \rightarrow W^\pm$ in MadGraph5 [80]. In the leading order we have obtained $\sigma_{W \rightarrow \nu + l} \approx 15.7 \text{ nb}$, which is in reasonable agreement with the prediction [77]. The resulting momentum distribution of W bosons is shown in figure 5 (left). A remark is in order here: at the leading order MadGraph5 does not predict the p_T distribution of W s, since the production process is $2 \rightarrow 1$ process and the colliding partons have $p_T = 0$; therefore, all of the W bosons in the simulations fly along the beam line, and the magnitude of their momentum is given by the longitudinal momentum p_L . The realistic p_T spectrum can only be obtained after implementation of the parton showering. However, based on the above-mentioned measurements [78, 79], the typical p_T 's of W bosons are significantly smaller than their typical p_L and therefore we chose to neglect the p_T momentum of the W bosons in what follows.

Having the W boson distribution dN_W/dp_W , we can obtain the $\epsilon_{\text{decay},W}$ and the average HNL momentum $\langle p_X \rangle$ by calculating the distribution of the particles in the energy E_X and the angle θ_X between the direction of motion of the X and the beam:

$$\frac{d^2 N_X^W}{dE_X d\cos(\theta_X)} = \int dp_W \frac{dN_W}{dp_W} \times \frac{d^2 \text{BR}_{W \rightarrow X}}{d\theta_X dE_X} \times P(\theta_X) \quad (5.3)$$

Here $d^2 \text{BR}_{W \rightarrow X} / d\theta_X dE_X$ is the differential production branching ratio, and $P(\theta_X)$ is a projector which takes the unit value if θ_X lies inside MATHUSLA's polar angle range and zero otherwise.

Let us compare the amounts of the X particles produced from the W bosons and from B mesons and flying in the direction of the decay volume. We have

$$N_{\text{prod},W}/N_{\text{prod},B} \approx \frac{N_W}{N_B} \times \frac{\text{BR}_{W \rightarrow X}}{\text{BR}_{B \rightarrow X}} \times \frac{\epsilon_{\text{decay},W}}{\epsilon_{\text{decay},B}} \approx \begin{cases} 10^{-3} \epsilon_{\text{decay},W}, & \text{scalars} \\ 10 \epsilon_{\text{decay},W}, & \text{HNLs,} \end{cases} \quad (5.4)$$

Figure 5. Left: momentum spectrum of W bosons produced in the pp collisions at $\sqrt{s} = 13$ TeV that is predicted by MadGraph5. Right: the energy spectrum of the HNLs produced in the decay of the W bosons and flying in the direction of the decay volume of the MATHUSLA experiment. The solid line corresponds to the spectrum obtained for the pseudorapidity range of the MATHUSLA experiment $\eta \in (0.9, 1.6)$, while the dashed line — to the spectrum for the HNLs flying in the direction $\eta \approx 1.3$.

where we used the amount of B mesons at the LHC and the decay acceptance from the table 2, the number of W s at the LHC (5.2) and the branching ratios of the scalar and HNL production from appendix B.1. Therefore we conclude that for scalars the production from the W s is not relevant, while for HNLs careful estimation is needed.

In the case of HNL, the differential branching ratio in the eq. (5.3) is

$$\frac{d^2 \text{BR}_{W \rightarrow N}}{d\theta_N dE_N} = \frac{1}{\Gamma_W} \frac{|\mathcal{M}_{W \rightarrow e+N}|^2}{8\pi} \frac{p_N}{E_W} \delta(M_N^2 + m_W^2 - 2E_N E_W + 2|\mathbf{p}_N||\mathbf{p}_W| \cos(\theta_N)) \quad (5.5)$$

The energy and angular distributions of the HNLs from the W bosons at MATHUSLA are almost independent of the HNL mass in the mass range of interest, $M_N \ll m_W$. It is an expected result because the kinematic in this limit should not depend on small HNL masses. The energy distribution for $M_N = 1$ GeV is shown in figure 5. The decay acceptance was found to be $\epsilon_{\text{decay}, W} \simeq 2\%$, while the average momentum of the produced HNLs is $\langle p_N \rangle \approx 62$ GeV.

The shape of the energy spectrum of the HNLs can be qualitatively understood in the following way. For a given value of the angle θ_N of the HNL, the energy distribution has a maximum at $E_{N, \text{max}}(\theta_N) = m_W/2 \sin(\theta_N)$,⁵ which corresponds to HNLs produced from the W bosons with some momentum $p_{W, \text{max}}(\theta)$. As a consequence, the largest amount of HNLs flying in the direction θ_N has an energy close to their maximum, see the dashed line at the right panel of figure 5. The total energy spectrum is a superposition of different angles and has a peak at $E_{N, \text{peak}} \approx 58$ GeV corresponding to the maximal angle possible at MATHUSLA, $\theta_{\text{MAT}}^{\text{max}} \approx 44^\circ$. From the other side, the maximal energy possible for HNLs at MATHUSLA is defined by the minimal angle $\theta_{\text{MAT}}^{\text{min}} \approx 22^\circ$, which explains why the spectrum tends to zero near the energy $E_{N, \text{max}} \approx 106$ GeV.

⁵This formula is valid for 2-body decay into massless particles.

6 Calculation of sensitivities

6.1 Efficiencies

Using the results of section 5, we have almost all ingredients needed to estimate the lower bound, the upper bound, the maximal mass probed and the total sensitivity curve. The only questions remaining are the following. The first one is the relation between the mesons spectra and the X particles spectra. The second one is the value of the overall efficiency

$$\epsilon = \epsilon_{\text{decay}} \times \epsilon_{\text{det}} \times \text{BR}_{\text{vis}}, \quad (6.1)$$

where the quantities $\epsilon_{\text{decay}}, \epsilon_{\text{det}}, \text{BR}_{\text{vis}}$ are the decay acceptance, detection efficiency and the visible branching correspondingly; they are defined by eqs. (1.2), (1.3).

We approximate the spectra of the X particles originating from the mesons and flying to the decay volume by the distributions of the mesons flying in the direction of the decay volume. To take into account the kinematics of the meson decays, we use the relation (2.2) between γ factors of the X particle and the meson in the expressions (1.3), (3.5) for the decay probability.

Let us discuss the efficiencies. For the HNLs at SHiP experiment, we used the values of ϵ_{decay} and ϵ_{det} provided by detailed FairSHiP simulations [81]. The results of the SHiP collaboration on the sensitivity to the scalars are not currently available, and for the product of $\epsilon_{\text{decay}} \cdot \epsilon_{\text{det}}$ we used the value for the HNL averaged over its mass, $\overline{\epsilon_{\text{decay}} \cdot \epsilon_{\text{det}}} \approx 0.2$.

For the MATHUSLA experiment there currently is no such detailed analysis of the efficiencies and background. In [19, 32] it is claimed that all the SM background can be rejected with high efficiency, but detailed simulations are needed for the justification of this statement. Here we optimistically use $\epsilon_{\text{det}} = 1$. For the decay acceptance of the particles produced from the mesons we use the geometric acceptance of the mesons at MATHUSLA, which we obtained using FONLL.⁶ For the decay acceptance of the HNLs produced in the decays of the W bosons we used the value $\epsilon_{\text{decay}, W} \approx 0.02$ obtained in section 5.3. All the parameters above, together with geometrical properties of the experiments are summarized in table 2. We estimate $\langle l_{\text{det}} \rangle$ and $\langle l_{\text{target-det}} \rangle$ using an assumption that the angular distribution of the X particles in the angular range of the decay volume is isotropic, see appendix D for details.

The last needed parameter is the visible decay branching fraction. Following [19, 31], for the visible decay branching fractions for both MATHUSLA and SHiP experiments we include only the decay channels of the X particle that contain at least two charged tracks. Our estimation of BR_{vis} is described in appendix B.2.3. The plots of the visible branching ratios for the HNLs and for the scalars are shown in figure 6.

6.2 Lower bound

Let us first compare the relevant parameters of the experiments summarized in tables 1, 2. One sees that the effective number of D mesons is approximately two orders of magnitude

⁶For the geometric acceptance as MATHUSLA we use the definition $\epsilon_{\text{geom}} = N^{\eta \in \eta_{\text{MAT}}} / N_{\text{meson}} \times \Delta\varphi / (2\pi)$, where $\eta_{\text{MAT}} \in (0.9; 1.6)$ and $\Delta\varphi = \pi/2$ are correspondingly pseudorapidity range of the MATHUSLA experiment and azimuthal size, see section D.

Figure 6. The branching ratio of decays of the HNLs (left) and the scalars (right) in visible states. The drop of the branching ratio for the HNLs mixing with ν_τ in the domain of HNL masses $\lesssim 1$ GeV is caused by the dominant invisible decay $N \rightarrow \pi^0 \nu_\tau$, while for the scalars of the same masses — by the decay $S \rightarrow \pi^0 \pi^0$.

Exp.	$\langle l_{\text{target-det}} \rangle$	$\langle l_{\text{det}} \rangle$	$\bar{\epsilon}_{X,D}$	$\bar{\epsilon}_{X,B}$	$\bar{\epsilon}_{X,W}$	$N_{D,\text{eff}}$	$N_{B,\text{eff}}$	$N_{W,\text{eff}}$
MAT	192 m	38 m	0.013	0.018	0.02	$5.7 \cdot 10^{14}$	$5.4 \cdot 10^{13}$	$1.2 \cdot 10^{10}$
SHiP	50 m	50 m	0.09	0.12	—	$1.4 \cdot 10^{17}$	$1.3 \cdot 10^{13}$	—

Table 2. Parameters of the SHiP and MATHUSLA experiments: the average length from the interaction point to the decay volume $\langle l_{\text{target-det}} \rangle$, the average length of the decay volume $\langle l_{\text{det}} \rangle$ (see appendix D for details), values of the overall efficiencies (6.1) averaged over the probed mass range of X for the particles X produced from D and B mesons, the effective number of the D and B mesons and W bosons defined by $N_{M,\text{eff}} = N_M \times \bar{\epsilon}_{X,M}$.

larger at SHiP,⁷ the effective numbers of B mesons are comparable between the experiments, and the average momenta (and therefore the γ factors) of the mesons produced in the direction of the decay volume are $\simeq 7 - 8$ times smaller at MATHUSLA. The latter is caused by (i) different beam configurations (colliding beams for MATHUSLA, fixed target for SHiP) (ii) their different geometric orientation relative to the proton beam direction (the decay volume of the SHiP experiment is located in the forward direction, while the one of MATHUSLA's is about 20° off-axis.)

Using the numbers from the tables 1, 2, for the ratio of the mixing angles at the lower bound (2.3) we have

$$\left. \frac{U_{\text{lower,SHIP}}^2}{U_{\text{lower,MAT}}^2} \right|_{M_N \lesssim m_D} \approx \frac{1}{5}, \quad \left. \frac{U_{\text{lower,SHIP}}^2}{U_{\text{lower,MAT}}^2} \right|_{M_N \gtrsim m_D} \simeq \left. \frac{\theta_{\text{lower,SHIP}}^2}{\theta_{\text{lower,MAT}}^2} \right|_{M_S \gtrsim m_K} \approx 5 \quad (6.2)$$

Qualitatively, for particles produced in the decays of the B mesons (HNLs with masses $M_N > m_D$ and scalars with masses $M_S > m_K$) MATHUSLA can probe mixing angles a

⁷By the effective number of the mesons we call the production of the number of mesons multiplied by the overall efficiency, $N_{\text{meson}} \cdot \epsilon_{X,\text{meson}}$.

Figure 7. The dependence of the number of events at SHiP and MATHUSLA evaluated at $U^2 = \theta_{\max}^2$ for the HNLs mixing with ν_e (left) and for scalars (right). Dashed lines denote the values for $U_{\max}^2$ for which the sensitivity of SHiP and MATHUSLA intersects the domain that has been closed by previous experiments (see, e.g., [36]).

factor $\simeq 5$ smaller than SHiP due to the smaller γ factor of the B mesons and larger effective number of B mesons (i.e. the total number of B mesons times the overall efficiency (6.1)). For the HNLs in the mass range $m_K \lesssim M_N \lesssim m_D$ the smallness of γ factor of the D mesons at MATHUSLA and the suppression of the number of events at SHiP by the overall efficiency cannot compensate the difference of two orders of magnitude in the effective numbers of the D mesons, and therefore the SHiP reaches a sensitivity which is about half an order of magnitude lower in U^2 . We note again that the result (6.2) was obtained under the optimistic condition $\epsilon_{\text{det}} = 1$ for MATHUSLA; after using a realistic efficiency the lower bound of the sensitivity at MATHUSLA will be changed by a factor $1/\sqrt{\epsilon_{\text{det}}}$, which will affect the ratio (6.2).

6.3 Upper bound

We show the dependence of the number of events at $\theta_X^2 = \theta_{\max}^2$ as a function of the mass for the HNLs mixing with ν_e and the scalars in figure 7. We see that by the maximal number of events the SHiP experiment is much better than the MATHUSLA experiment, which is explained by the shorter length to the decay volume and higher value of the average gamma factor.

With the energy distributions of the mesons and the W bosons obtained in section 5, let us now estimate their effect on the upper bound of the sensitivity. To do this, we introduce the width of the upper bound defined by

$$R = \theta_{\text{upper}}^2 / \theta_{\max}^2 \quad (6.3)$$

We take the HNLs as an example, commenting later on the difference with the scalar. We will be interested in the HNLs with $M_N \gtrsim 2 \text{ GeV}$ (for smaller masses θ_{upper}^2 lies deep inside the region excluded by the previous searches, see e.g. [24]). The HNLs in question are produced from the decays of B mesons and W bosons.

Our procedure of the estimation of the upper bound width is based on (3.5). As we already mentioned at the beginning of this section, in the case of the production from B

Figure 8. Ratios $R = \theta_{\text{upper}}^2 / \theta_{\text{max}}^2$ for the HNLs mixing with ν_e (left) and for scalars (right) at SHiP and MATHUSLA experiments. Solid lines are obtained by taking the account the energy distribution of the mother particles (B mesons and W bosons). Dashed lines are obtained under an assumption that all particles have the same average energy.

mesons we approximate the spectra of the HNLs by the spectra of the B mesons (so that the HNLs fly in the same direction as the B mesons) and take into the account the relation (2.2) between the B meson and the energies of HNLs. In the case of the production from the W bosons, we use the energy spectrum of the HNLs from figure 5. We approximate the shapes of the high-energy tails of these spectra by simple analytic functions. For the B mesons at SHiP, the fit is an exponential function, for the B mesons at MATHUSLA the fit is a power law function, while for the HNLs from the W bosons the fit is a linear function, see appendix E.1. Using the fits, we calculate the upper bound θ_{upper}^2 using the steepest descent method for the evaluation of the integral (3.5). The derivation of θ_{upper}^2 is given in appendix E.2.

Using θ_{upper}^2 , we present the upper bound width (6.3) in figure 8. We also show there the prediction of the estimations of the upper bound width which assume that all of the produced particles have the same energy, see eq. (3.3).

We see that for the particles from B mesons at SHiP and for the HNLs from the W bosons at MATHUSLA the broadening of the width due to the distribution is small, while for the particles from B mesons the distribution contributes significantly. This is a direct consequence of the behavior of the shape of the high-energy tails of the distributions. Namely, for the B mesons at SHiP, the number of high-energy HNLs is exponentially suppressed. For the HNLs originating from the W bosons the tail falls linearly, and naively the upper bound would be significantly improved. However, the distribution becomes zero not very far from $\langle p_N \rangle$, and the effect of the contribution is insignificant. Only for the B mesons at MATHUSLA the tail causes significant improvement of the width of the upper bound.

Finally, let us comment on the difference between the shapes of the width between the HNL and scalar cases. The lifetime τ_S is changed with the mass slower than τ_N , see the discussion in section B.2. In addition, $\text{Br}_{B \rightarrow S}$ behaves with the mass monotonically, while for HNLs new production channels appear at different masses. Therefore the upper bound of the sensitivity region for the scalars changes less steeply and more smoothly with their mass, see figure 8 (right panel).

Figure 9. Comparison of the sensitivity of the MATHUSLA experiment to the HNLs that are produced in decays of D and B mesons (including B_c) and in decays of W for the mixing with ν_e .

The comparison of the upper bound of the sensitivity for the HNLs originating from W bosons and B mesons is shown in figure 9. Our method of obtaining the sensitivity is summarized in appendix F. We see that the W s determine the upper bound. The reason for this is that the HNLs from W s have sufficiently larger average momentum, which compensates the production suppression (see eq. (5.4)).

6.4 Maximal mass probed

The smaller γ factor of the mesons at MATHUSLA adversely affects the upper bound of the sensitivity curve and thus the maximal mass probed. In particular, for the HNLs mixing with $\nu_{e/\mu}$, the maximal mass probed ratio (4.3) becomes

$$M_{\max}^{N,\text{SHIP}}/M_{\max}^{N,\text{MAT}} \approx 1.3, \quad (6.4)$$

which agrees well with the sensitivity plot from figure 12. For the other cases — the HNLs mixing with ν_τ and the scalars — the estimation of the maximal mass for the SHiP experiment based on the definition above exceeds the kinematic threshold, and therefore the result (4.3) is not valid. However, for the scalars the maximal mass for the MATHUSLA experiment is smaller than the kinematic threshold, which is still a consequence of smaller γ factor.

7 Comparison with simulations

Next we compare our sensitivity estimates with the results of the SHiP and MATHUSLA collaborations. Different groups used different phenomenology for HNL and, especially, for scalars. Therefore, we will use different prescriptions for production and/or decay in

Figure 10. Comparison of the sensitivities to the HNLs mixing with the electron flavor obtained in this paper (solid lines) with the results of the SHiP [31] and MATHUSLA [19] collaborations (dotted lines). For the MATHUSLA experiment, the contributions from both B, D mesons and from W bosons are shown separately. For the SHiP experiment, we consider the case of maximally possible contribution of B_c mesons, given by the fragmentation fraction $f_{B_c} = 2.6 \cdot 10^{-3}$ measured at LHC energies $\sqrt{s} = 13$ TeV [57]. Orange points, based on analytic estimates of the lower boundary, allow for simple cross-check of our results, see appendix G for details. Possible origins of the discrepancy at low masses at the left panel are discussed in appendix H.

different sections below, in order to facilitate the comparison of our approach with the Monte Carlo results of other groups. Our current view of the HNL phenomenology is summarized in [24] and for scalar in [28]. Our method of obtaining the sensitivity curves is summarized in appendix F.

7.1 HNLs

The results for the HNLs are shown in figure 10. To facilitate the cross-check of our results, we also provide simple analytic estimates of the lower boundary for several HNL masses (see appendix G). Small discrepancies between the simple estimation of the lower bound and numeric result are caused by the difference in the values of $1/\langle p_{\text{meson}} \rangle$ and $\langle 1/p_{\text{meson}} \rangle$, which actually defines the lower bound.

For the sensitivity of the SHiP experiment, there is good agreement of the sensitivity curves, with a slight difference in the maximal mass probed. We think that this is due to the difference in the average γ factors used in our estimation and those obtained in Monte Carlo simulations by the SHiP collaboration. Indeed, using the SHiP simulations results available in [31, 81], we have found that for the masses $M_N \simeq m_{B_c}$ the ratio of average γ factors is $\langle \gamma_N^{\text{analytic}} \rangle / \langle \gamma_N^{\text{simulations}} \rangle \simeq 0.8$, which seems to explain the difference.

For the sensitivity of MATHUSLA [19] to the HNLs produced in W decays there is good agreement for the entire mass range probed. For the sensitivity to the HNLs from B and D mesons, the situation is somewhat different. In the mass range $M_N \gtrsim m_{D_s}$, where the main production channel is the decay of the B mesons, there is reasonable agreement with our estimate. The discrepancy can be caused by higher average energy of

Figure 11. Sensitivity to the scalar portal particles for the SHiP (*left panel*) and MATHUSLA (*right panel*) experiments. Solid lines — results obtained in this work. Dashed lines — simulations of SHiP [1] (*left panel*) and MATHUSLA [19] (*right panel*). In order to facilitate the comparison with collaboration results *we have used different scalar production and decay models* in left and right panels: for comparison with MATHUSLA results we took the model from [19], while for comparison with SHiP we used the model from [29], see section 7.2. Orange points, based on analytic estimates of the lower boundary, allow for simple cross-check of our results, see appendix G.

the HNLs in the simulations, which simultaneously lifts up the lower and upper bounds of the sensitivity. The reason for the difference at masses $M_N < 1$ GeV is not known, see a discussion in appendix H.

7.2 Scalars

The comparison of our sensitivity estimates with the results of the SHiP and MATHUSLA experiments is presented in figure B.3. We also show the results of a simple analytic estimate of the lower bound for particular masses from appendix G. For the comparison with the sensitivity provided by the MATHUSLA collaboration we used the model of scalar production and decay given in [19], while comparing with the results of the SHiP collaboration — from [1]. A description of the models is given in appendix B.3.

The sensitivity curves are in good agreement. Small differences in the position of the maximal mass probed can be explained by different energy distributions of the scalars used in our estimate and in [19] and in [1].

8 Conclusions

In this work, we investigated the sensitivity of Intensity Frontier experiments to two models of new super-weakly interacting physics: *heavy neutral leptons* and *dark scalars*. We explored analytically the characteristic features of the experiment’s sensitivity regions: upper and lower boundaries and the maximal mass of new particles that can be probed. Our analytic analysis allows identifying the parameters responsible for the positions of the main “features” of these curves and to cross-check/validate the results of the Monte Carlo simulations. We analyse a number of experimental factors that contribute to the sensitivity

Figure 12. Comparison of the sensitivity of SHiP and MATHUSLA for the HNL. The production fraction of B_c mesons at SHiP energies $\sqrt{s} \approx 28$ GeV is not known, and the largest possible contribution is based on the production fraction measured at the LHC, $f(b \rightarrow B_c) = 2.6 \times 10^{-3}$. In the case of the SHiP experiment we used the overall efficiency calibrated against the Monte Carlo simulations [31] and also selected only those channels where at least two charged tracks from the HNL decay appear. In the case of the MATHUSLA experiment we optimistically used $\epsilon_{\text{det}} = 1$ for the detection efficiency.

Figure 13. Comparison of sensitivities of the SHiP and MATHUSLA experiments for the scalar portal model. In the case of the SHiP experiment we used the overall efficiency $\bar{\epsilon} = 0.2$, see the text for details. In the case of the MATHUSLA experiment we optimistically used $\epsilon_{\text{det}} = 1$ for the detection efficiency. We used the scalar phenomenology described in [28].

estimates: (i) the number of heavy flavour mesons traveling in the direction of the detector; (ii) their average momentum and the high-energy tail of the momentum distribution; (iii) geometry of the experiment; (iv) efficiency. We use SHiP and MATHUSLA as examples of the fixed target and LHC-accompanying Intensity Frontier experiments, respectively. Our analytic estimates agree well with the Monte Carlo-based sensitivities provided by the SHiP [31] and MATHUSLA [19] collaborations under similar assumptions about the overall efficiencies of the experiments.

Our main results are as follows. Our estimates of the sensitivities of the SHiP and MATHUSLA experiments to the HNLs are shown in figures 12 and to the dark scalars in figure 13.

Qualitatively both experiments can probe similar ranges of parameters. The SHiP has higher average γ factors of the mesons ($\langle\gamma_{\text{meson}}^{\text{SHiP}}\rangle/\langle\gamma_{\text{meson}}^{\text{MAT}}\rangle \simeq \mathcal{O}(10)$) and, as a result, significantly higher upper boundary of the sensitivity region than MATHUSLA (as the upper boundary is exponentially sensitive to the γ factor). As the consequence, the SHiP can probe higher masses for both HNLs and scalars than MATHUSLA (except of HNLs with dominant mixing with tau flavor). However, the W boson decays at the LHC would produce some highly boosted HNLs traveling to the MATHUSLA decay volume, partly mitigating this difference.

The SHiP experiment is able to probe lower mixing angles for HNLs with $M_N \lesssim m_{D_s}$ owing to the larger number of D mesons. MATHUSLA can probe lower mixing angles for the HNLs with $M_N \gtrsim m_{D_s}$ and the scalars for all masses, owing to the larger number of the B^{+0} mesons at the LHC (as charmed mesons contribute negligibly to the scalar production).

Uncertainties. According to the theoretical predictions the $d\sigma/dp_T$ distribution of B mesons at the LHC has a maximum at $p_T \sim \text{GeV}$, see figure 4. The region of low p_T is complicated for the theoretical predictions because of limitations of the applicability of the perturbative QCD. At the same time, these cross-sections have not been measured by neither the ATLAS, nor the CMS collaborations in the required kinematic range. The increase in the overall amount of low-momentum mesons shifts leftwards the position of the peak of the $d\sigma/dp_T$ distribution, thus decreasing their average momentum. Both factors lead to the increase of the number of events at the lower boundary. Therefore the uncertainty in the position of the lower boundary of the sensitivity region depends on both of these numbers such that the uncertainty in the position of the peak enters into the sensitivity estimate *squared*.

Another uncertainty comes from the background estimates. For the SHiP experiment, comprehensive background studies have proven that the yield of background events passing the online and offline event selections is negligible [35, 37]. For MATHUSLA such an analysis is not available at the time of writing. The Standard Model background at MATHUSLA is non-zero (due to neutrinos from LHC and atmosphere, cosmic rays, muons, etc), however, it is claimed to be rejected with high efficiency based on the topology of the events [19, 32]. It is not known how much this rejection affects the detection efficiency, ϵ_{det} . In this work, we conservatively assumed $\epsilon_{\text{det}} = 1$ for MATHUSLA, while for SHiP it was taken from the actual Monte Carlo simulations [31]. More detailed analysis of the MATHUSLA background should be performed, which could influence the sensitivities.

In case of the SHiP experiment, the main uncertainty for HNLs is the unknown production fraction of the B_c mesons at $\sqrt{s} \approx 28 \text{ GeV}$. It changes the position of the lower bound and consequently the maximal mass probed in a significant way, see figure 12.

Comparison with other works. We have compared our sensitivity estimates with the results of the Monte Carlo simulations presented by the SHiP collaboration [1, 31, 82] and with the estimates of the MATHUSLA physics case paper [19] (figures 10–11). For the

HNLs, the estimates are in good agreement with the results of the SHiP collaboration. In the case of MATHUSLA, there is a difference for HNLs with mass smaller than 1 GeV. It can be attributed to different branching for the HNL production used in our estimates and in the Monte Carlo simulations of [19], see discussion in appendix H. For the scalars, our estimates are in good agreement with the results from the SHiP and MATHUSLA collaboration. Small discrepancies between the sensitivities at the upper bound can be explained mainly by the difference in the meson energy spectrum used in our estimation and obtained in the Monte Carlo simulations.

Acknowledgments

We thank D. Curtin, J. Evans, R. Jacobsson and W. Valkenburg for fruitful discussions and comments on the manuscript. This project has received funding from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (GA 694896) and from the Netherlands Science Foundation (NWO/OCW).

A Portals

New particles with masses much lighter than the electroweak scale can couple to the Standard Model fields via renormalizable interactions with small dimensionless coupling constants (sometimes called “*portals*” as they can mediate interactions between the Standard Model and “hidden sectors”). In this work, we considered two *renormalizable* portals: scalar (or Higgs) portal and neutrino portal.

The scalar portal couples a gauge-singlet scalar S to the gauge invariant combination $H^\dagger H$ made of the Higgs doublet:

$$\mathcal{L}_{\text{scalar}} = \mathcal{L}_{\text{SM}} + \frac{1}{2}(\partial_\mu S)^2 - \frac{M_S^2}{2}S^2 + gSH^\dagger H + \mathcal{L}_{\text{int}} \quad (\text{A.1})$$

where g is the coupling constant and $\mathcal{L}_{\text{int}}$ are interaction terms that play no role in our analysis. After the spontaneous symmetry breaking the cubic term in (A.1) gives rise to the Higgs-like interaction of the scalar S with all massive particles with their mass times a small *mixing parameter*

$$\mathcal{L}_{S,\text{int}} = \theta S \left[\sum_f m_f \bar{f} f + M_W W_\mu^+ W_\mu^- + \dots \right] \quad \theta \equiv \frac{gv}{m_H} \ll 1 \quad (\text{A.2})$$

where g is the coupling in (A.1); v is the Higgs VEV; m_H is the Higgs mass; sum in (A.2) goes over all massive fermions (leptons and quarks); $W_\mu^\pm$ is the W boson and $\dots$ denote other interaction terms, not relevant for this work. The details of the phenomenology of the scalar portal are provided in [28] (see also [50, 83–85]). The computation of hadronic decay width of S is subject to large uncertainties at masses $M_S \sim \text{few GeV}$, where neither chiral perturbation theory nor perturbative QCD can provide reliable results (see a discussion in [25]).

We have also considered the neutrino portal where one adds to the Standard Model new gauge-singlet fermion — heavy neutral lepton N — that couples to the $\epsilon_{ab} \bar{L}_a H_b$ where

Meson M	B^+	B^0	B_s	B_c^+	D^+	D^0	D_s^+
MATHUSLA	0.324	0.324	0.088	$2.6 \cdot 10^{-3}$	0.225	0.553	0.105
SHiP	0.417	0.418	0.09	?	0.207	0.632	0.088

Table 3. The fragmentation fractions for heavy mesons at the LHC energies [57, 86, 87] and of the SHiP experiment [24, 88]. For SHiP the contribution of flavoured baryons or quarkonia states can be neglected, see [24]. For the LHC energies, the remaining 20-25% come of all heavy flavour quarks hadronize into baryons, mostly Λ_b states [86].

L_a is the SU(2) lepton doublet and ϵ_{ab} is absolutely antisymmetric tensor in 2 dimensions. Phenomenologically, HNL is massive Majorana particle that possesses “neutrino-like” interactions with W and Z bosons (the interaction with the Higgs boson does not play a role in our analysis and will be ignored). The interaction strength is suppressed as compared to that of ordinary neutrinos by a flavour-dependent factors (known as *mixing angles*) $U_\alpha \ll 1$ ($\alpha = \{e, \mu, \tau\}$).

B Production and detection of portal particles

B.1 Production in proton-proton collisions

The number of mesons is determined by the number of produced $q\bar{q}$ pairs and fragmentation fractions f_{meson} , that can be extracted from the experimental data [57, 86, 87]. We summarize the fragmentation fractions that we use for MATHUSLA in the table 3. For the SHiP experiment, all fragmentation fractions except for B_c meson are known to be close to the MATHUSLA’s ones [88]. The B_c meson fragmentation fraction at the energy of the SHiP experiment is unknown. In our estimations, we take it the same as for the MATHUSLA experiment.

B.1.1 HNL production

The production of the HNL in the decay of charmed and beauty mesons have been considered in [89, 90], see [24] for the recent review and summary of the results. The branching ratios multiplied by fragmentation fractions of D and B for the most relevant channels and the values of the fragmentation fractions from the table 3 are presented at the figure 14. We see that for the HNL mass range $m_N \gtrsim 3.5$ GeV the main production channel is B_c meson decay $B_c \rightarrow N + l$. This is a quite surprising fact, taking into account that B_c fragmentation fraction is of order 10^{-3} . To understand this result let us compare HNL production from B_c with production from the two-body B^+ decay. The decay widths for both cases are given by

$$\text{BR}(h \rightarrow \ell_\alpha N) \approx \frac{G_F^2 f_h^2 m_h m_N^2}{8\pi\Gamma_h} |V_h^{\text{CKM}}|^2 |U_\alpha|^2 K(m_N/m_h), \quad (\text{B.1})$$

Figure 14. Branching ratios multiplied by fragmentation fractions of D and B mesons decaying into HNL through e -type mixing (upper panel) and into the HNL through τ -type mixing (lower panel) for $U^2 = 1$. The values of fragmentation fractions are taken at LHC energies $\sqrt{s} = 13$ TeV, see table 3.

where we take $m_N \gg m_\ell$, and K is a kinematic suppression. Neglecting them, for the ratio for the numbers of HNLs produced by B_c and B^+ we obtain

$$\frac{N_{\text{HNL}}(B_c \rightarrow \ell N)}{N_{\text{HNL}}(B^+ \rightarrow \ell N)} \approx \underbrace{\frac{f_{B_c}}{f_{B^+}}}_{\approx 0.008} \times \underbrace{\frac{\Gamma_{B^+}}{\Gamma_{B_c}}}_{\approx 0.3} \times \underbrace{\left(\frac{f_{B_c}}{f_{B^+}}\right)^2}_{\approx 5} \times \underbrace{\frac{m_{B_c}}{m_{B^+}}}_{\approx 1.2} \times \underbrace{\left(\frac{V_{cb}^{\text{CKM}}}{V_{ub}^{\text{CKM}}}\right)^2}_{\approx 100} \approx 1.44. \quad (\text{B.2})$$

We see that the small fragmentation fraction of B_c meson is compensated by the ratio of the CKM matrix elements and meson decay constants.

HNLs can also be produced in the decays of the W bosons, $W \rightarrow N + l$. The corresponding branching ratio is

$$\text{BR}(W \rightarrow N + \ell_\alpha) \approx \frac{1}{\Gamma_W} \frac{G_F m_W^3}{6\sqrt{2}\pi} \approx 0.1 U_\alpha^2, \quad (\text{B.3})$$

where we have neglected the HNL and the lepton masses.

B.1.2 Quarkonia and heavy flavour baryons

Quarkonia states (especially Υ meson) can produce HNLs reaching 10 GeV in mass above the beauty meson threshold. The contribution of quarkonia decays to the production were found negligible at SHiP energies, see [24].

The LHC experiments have measured Υ production at both ATLAS and CMS [91, 92]. In the rapidity range $|y| < 2$, relevant for MATHUSLA, the cross-section is given by [93]

$$\sigma(pp \rightarrow \Upsilon(nS)) \times \text{BR}(\Upsilon \rightarrow \mu^+ \mu^-) \sim 10 \text{ nb} \quad (\text{B.4})$$

(as this is an order of magnitude estimate, we combine production of $1S$, $2S$ and $3S$ bottomonium states and neglected both statistical and systematic uncertainties of the cross-section measurement). Using $\text{BR}(\Upsilon \rightarrow \mu^+ \mu^-) \simeq 2.4 \times 10^{-2}$ [94] we find that during the high luminosity phase one can expect $N_\Upsilon \sim 10^{12}$. Large fraction of this mesons are traveling into the direction of the fiducial decay volume of MATHUSLA, as their distribution is sufficiently flat in the $|y| < 2$ rapidity range. This number should be multiplied by the branching ration $\text{BR}(\Upsilon \rightarrow N\nu)$, estimated in [24] to be at the level $\text{BR}(\Upsilon \rightarrow N\nu) \sim 10^{-5} U^2$, so that overall one expects in MATHUSLA detector about $10^7 U^2$ HNLs from Υ decays.

This number should be compared with those, produced from W -bosons (as we are above the B -meson threshold): $N_W \times \text{BR}(W \rightarrow N + l) \times \epsilon_N$, where N_W is given by (5.2), $\epsilon_N \approx 0.02$ is the geometric acceptance for the HNLs produced from W and flying into the MATHUSLA fiducial volume and the branching fraction is given by (B.3). The resulting number is $\sim 6 \times 10^8 U^2$ — exceeding the number of HNLs from Υ -mesons by about 2 orders of magnitude.

As table 3 demonstrates, about 25% of b -quarks at the LHC hadronize into the Λ_b^0 baryons. These baryons produce HNLs in the 3-body semi-leptonic decay $\Lambda_b^0 \rightarrow B + \ell + N$ where B is a baryon. The mass of the Λ_b^0 is $m_{\Lambda_b^0} \simeq 5.62 \text{ GeV}$. The decays $\Lambda_b^0 \rightarrow p + \ell^- + N$ are suppressed by the CKM matrix element V_{bu} , while the decays $\Lambda_b^0 \rightarrow \Lambda_c^+ + \ell^- + N$ can only produce HNLs with $M_N < m_{\Lambda_b^0} - m_{\Lambda_c^+} \simeq 3.35 \text{ GeV}$. HNLs of this mass are produced from more copious B -mesons and therefore Λ baryons can be neglected.

The contribution of heavy flavour baryon decays to the production were found negligible at SHiP energies, see [24].

B.1.3 Scalar production

The main difference in the phenomenology of the Higgs-like scalar S in comparison to HNLs is that the interaction of S with fermions is proportional to their mass, see section A. Therefore, its production at the mass range $M_S > M_K$ is dominated by the decay of the B^+ , B^0 , while the contribution from D mesons is negligible [23, 28]. The main production process is the 2-body decay

$$B \rightarrow X_{s/d} + S, \quad (\text{B.5})$$

where X_q is a hadron that contains the quark q . The branching ratios for these states are discussed in [28]. Here we only state the main points. For $B \rightarrow X_s + S$ we choose two lightest resonances X_s for each given spin and parity. Exceptions are pseudo-scalar and tensor mesons (there is only one known meson that has these properties, see [94]). We have found that each heavier meson from the “family” gives a smaller contribution to the branching ratio than the lighter one. For $B \rightarrow X_d + S$ we take only one meson $X_d = \pi$ since this channel has the largest kinematic threshold $m_B - m_\pi$. We summarize the list of the final states below:

Figure 15. Branching ratios of the scalar production in the process $B \rightarrow S + X$, where X denotes one of the mesons from the caption (B.5).

- Spin 0, odd parity: $X_q = \pi, K$;
- Spin 0, even parity: $X_q = K_0^*(700), K_0^*(1430)$;
- Spin 1, odd parity: $K_1(1270), K_1(1400)$;
- Spin 1, even parity: $K^*(892), K^*(1410)$;
- Spin 2, even parity: $K_2^*(1430)$.

The main source of the uncertainty is unknown quark squad of the $K_0^*(700)$ meson: it can be either a di-quark or a tetra-quark (see e.g. [95]). In the second case, the $K_0^*(700)$ contribution to the scalar production is unknown, which causes an uncertainty up to 30%. We consider it as the di-quark state.

The dependence of the branching ratios of the process (B.5) on the scalar mass is shown in figure 15.

We estimate the production of the scalars from the W bosons by the decay $W \rightarrow S + f + \bar{f}'$, where the summation over all the SM fermions species $f = l, q$ is taken. We obtained $\text{BR}_{W \rightarrow S} / \theta^2 \simeq 4 \cdot 10^{-3}$.

We mention in passing that the production of scalars from Υ (due to $b \rightarrow s$ transition) is not playing essential role, as the mass difference $m_\Upsilon - m_B < m_B$ and therefore one should compare the number of scalars produced from the bottomonium decays with the number of scalars from B -meson decays. The latter of B -mesons is several orders of magnitude higher (see table 1). In addition to that the branching ratio of $\Upsilon \rightarrow B + S$ is much smaller than $B \rightarrow K + S$ because the width of Υ is dominated by electromagnetic decays.

The $b \rightarrow s$ transitions also generate decays $\Lambda_b^0 \rightarrow \Lambda^0 + S$. However, the mass of thus produced scalar, $M_S < m_{\Lambda_b^0} - m_{\Lambda^0} \simeq 4.5 \text{ GeV}$ and thus is subdominant to the production from B mesons.

B.2 Main decay channels

B.2.1 HNL

The HNL has 3-body leptonic decays and different semileptonic modes. Following the paper [24], we estimate the decay width of HNL into hadronic states as a sum of decay widths of specific channels for the HNL with a mass lower as 1 GeV and use the decay into the quarks with QCD corrections for larger masses. In the latter mass region, the decay width of HNLs mixing with the flavor α can be approximated by the formula

$$\Gamma_N \approx g_{\text{eff}} |U_\alpha|^2 \frac{G_F^2 m_N^5}{192\pi^3}, \quad (\text{B.6})$$

where g_{eff} is a dimensionless factor that depends on the mass of HNL and changes from 1 to ~ 10 , see e.g. [24] for details.

The dependence of the proper lifetime $c\tau_N$ on the HNL mass at $U^2 = 1$ is given at the left panel of figure 16.

B.2.2 Scalar

The decay width of the scalar particle has large uncertainty in the scalar mass region $0.5 \text{ GeV} < M_S < 2 \text{ GeV}$ because of resonant nature of $S \rightarrow 2\pi$ decay, see [25] for the recent overview. At higher masses the decay width is determined by perturbative QCD calculations [96]. We omit the problem of pion resonance in this work using continuous interpolation between the sum of the decay channel at low masses and perturbative QCD at high ones.

For scalar mass region above 2 GeV one can naively estimate S decay width as $\Gamma_S \propto \sum_f \theta^2 y_f^2 M_S$. This estimation does not take into account three effects:

1. For the decay into quarks parameter y_q depends on scalar mass as $y_q \equiv M_q(M_S^2)/v$, where $M_q(M_S^2)$ is quark running mass, which gives logarithmic correction;
2. The decay into gluons has different M_S dependence, $\Gamma_S \propto \theta^2 M_S^3/v^2$, and dominates in the region $2 \text{ GeV} < M_S < 3.5 \text{ GeV}$ [23];
3. In the region M_S near 3.5 GeV new decay channels appear (into τ and c quark), and the kinematical factor is important.

Taking them into account, for the mass domain $3.5 \text{ GeV} < M_S < 5 \text{ GeV}$, near the threshold of production from B mesons, we made a fit to the total Γ_S and found that its behavior is $\Gamma_S \propto M_S^2$.

The dependence of the proper lifetime $c\tau_S$ on the scalar mass for $\theta^2 = 1$ is shown in figure 16 (right panel).

B.2.3 Visible branching ratio

We define the “visible” decay channels as those that contain at least two charged particles α in the final state. The corresponding decays are

$$X \rightarrow \alpha\alpha'Y, \quad X \rightarrow F\tilde{Y} \quad (\text{B.7})$$

Figure 16. The dependence of the proper lifetime $c\tau$ on the mass for the HNL (left panel, based on [24]) and the scalar (right panel, based on [28]).

where Y is arbitrary state, F is uncharged state that decays to n charged particles and $\tilde{Y}$ is a state with at least $2 - n$ charged particles (assuming $n < 2$). Using this definition, the decay $N \rightarrow 3\nu$ is identified as invisible decay, the decay $N \rightarrow \bar{\mu}\nu_\mu e$ — as visible decay, while the decay $N \rightarrow \eta\nu$ — as the visible decay if η meson decays into two charged particles, and as invisible decay otherwise. To take into account only visible decays of F , we include the factor $\text{BR}_{F \rightarrow \text{vis}}$ to the partial decay width $\Gamma_{X \rightarrow F\tilde{Y}}$. We take the values of $\text{BR}_{F \rightarrow \text{vis}}$ from [94]. For HNL/scalar masses $M > 2 \text{ GeV}$ we describe hadronic decays as having quarks and gluons in the final states. In this case we assume that any such decay will contain at least 2 charged tracks and therefore the whole hadronic width is visible.

B.3 Comparison with scalar models used by SHiP and MATHUSLA collaborations

Our model of the scalar production and decay described above differs from those used in [1, 19] for estimating the sensitivity. Namely, for scalar production in ref. [28] has summed over main exclusive channels $B \rightarrow X_{s/d} + S$.⁸ In ref. [19] the production from B mesons is estimated using the free quark model, while [1] considers only the channel $B \rightarrow KS$. This causes differences in the magnitude of the branching ratio and kinematic production thresholds. In particular, we note that the free quark model breaks down for large scalar masses $M_S \simeq 3 \text{ GeV}$ since the QCD enters the non-perturbative regime, and therefore it gives meaningless predictions for the production rate of heavy scalars.

As for the scalar decay width, because of theoretical uncertainty for the mass range $2m_\pi \lesssim M_S \lesssim 2m_D$ there is no agreement in the literature how to describe the scalar decays in this domain, see [25]. Our decay width differs significantly from the decay width used in [1, 19].

⁸There is 30% level uncertainty in the total production rate because of the $B \rightarrow K_0^*(700) + S$ channel. The meson $K_0^*(700)$ is not observed experimentally and it could be either a di-quark or a tetra-quark state (see, e.g. [95] and references therein). We did an estimation assuming that $K_0^*(700)$ is the di-quark state, in the other case this production channel is absent.

C Relation between momentum of HNL and meson momentum

The energy of the particle X at lab frame, E_X , is related to energy of X at meson's rest frame, E_X^{rest} , and meson energy E_{meson} at lab frame as

$$E_X(\theta, E_X^{\text{rest}}) = \frac{E_{\text{meson}}}{m_{\text{meson}}} \left(E_X^{\text{rest}} + |\mathbf{p}_X^{\text{rest}}| \frac{|\mathbf{p}_{\text{meson}}|}{E_{\text{meson}}} \cos(\theta) \right), \quad (\text{C.1})$$

where θ is the angle between the direction of motion of meson in lab frame and the direction of motion of the particle X in the meson's rest frame. At meson frame the angle distribution is isotropic, so for the average energy we obtain

$$\langle E_X \rangle = \gamma_{\text{meson}} \langle E_X^{\text{rest}} \rangle, \quad (\text{C.2})$$

where $\gamma_{\text{meson}} \equiv E_{\text{meson}}/m_{\text{meson}}$.

D Geometry of the experiments

D.1 SHiP

The SHiP experiment [37] is a fixed-target experiment using the proton beam of the Super Proton Synchrotron (SPS) at CERN. SPS can deliver $N_{\text{p.o.t.}} = 2 \times 10^{20}$ protons with the energy 400 GeV over a 5 year term. The SHiP will be searching for new physics in the largely unexplored domain of very weakly interacting particles with masses below $\mathcal{O}(10)$ GeV and $c\tau$ exceeding tens of meters. The overview of the experiment is as follows.

The proton beam hits a target [35, 36]. The target will be followed by a 5 m hadron stopper, intended to stop all $\pi^\pm$ and K mesons before they decay, and by a system of shielding magnets called active muon shield, constructed to sweep muons away from the fiducial decay volume. The whole active muon shield system is 34 m long.

The decay volume is a long pyramidal frustum vacuum chamber with the length

$$l_{\text{det}} = 50 \text{ m} \quad (\text{D.1})$$

and the cross-section $5 \text{ m} \times 10 \text{ m}$. It begins at

$$l_{\text{target-det}} = 50 \text{ m} \quad (\text{D.2})$$

downstream of the primary target respectively. The SHIP spectrometer downstream of the decay volume consists of a four-station tracker, timing detector, and an electromagnetic calorimeter and muon detector for particle identification. The detectors are seen from the interaction point at an angle $\theta \approx 25 \text{ mrad}$.

D.2 MATHUSLA

MATHUSLA (MAssive Timing Hodoscope for Ultra Stable neutraL pARticles) is a proposed experiment [32, 33] that consists of a $20 \text{ m} \times 200 \text{ m} \times 200 \text{ m}$ surface detector, installed above ATLAS or CMS detectors (see figure 17). The long-lived particles, created at the LHC collisions, travel 100+ meters of rock and decay within a large decay volume ($8 \times 10^5 \text{ m}^3$).

Figure 17. MATHUSLA experiment geometry. Adapted from [18].

Parameter	θ_1	θ_2	η_1	η_2	$\bar{l}_{\text{target-det, m}}$	$\bar{l}_{\text{det, m}}$	$\Delta\phi$
Value	44.3°	22.9°	0.9	1.6	192.5	38.5	$\pi/2$

Table 4. Parameters of MATHUSLA experiment [19]. For the definition of angles $\theta_{1,2}$ see figure 17, and $\Delta\phi$ is the azimuthal size of MATHUSLA.

Multi-layer tracker at the roof of the detector would catch charged tracks, originating from the particle decays. The ground between the ATLAS/CMS and MATHUSLA detector would serve as a passive shield, significantly reducing the Standard Model background (with the exception of neutrinos, muons and K_L^0 created near the surface). Assuming isotropic angular distribution of a given particle traveling to MATHUSLA, the average distance that it should travel to reach the MATHUSLA decay volume is equal to

$$\bar{l}_{\text{target-det}} \equiv \left\langle \frac{L_{\text{ground}}}{\sin \theta} \right\rangle = 192.5 \text{ m} \quad (\text{D.3})$$

where $L_{\text{ground}} = 100 \text{ m}$. The average distance a particle travels inside the decay volume, $\bar{l}_{\text{det}}$, is given by

$$\bar{l}_{\text{det}} \equiv \left\langle \frac{20 \text{ m}}{\sin \theta} \right\rangle = 38.5 \text{ m} \quad (\text{D.4})$$

Geometrical parameters of MATHUSLA experiment are summarized in table 4.

E Analytic estimation of the upper bound: details

In this section we estimate the ratio between θ_{max} and θ_{upper} — the quantities defined in section 3.

E.1 Fits of the spectra

The high-energy tail of the B mesons distribution function at SHiP is well described by the exponential distribution, see the left figure 18:

$$\frac{dN_B^{\text{SHIP}}}{dE} = f_0 e^{-E\delta}, \quad \delta \approx 3 \cdot 10^{-2} \text{ GeV}^{-1} \quad \text{and} \quad f_0 \approx 0.3 \text{ GeV}^{-1} \quad (\text{E.1})$$

Figure 18. Fits of the high energy tail of the distributions of the B mesons. *Left panel:* SHiP data is taken from [64]. *Right panel:* the FONLL simulations are performed for $\sqrt{s} = 13$ TeV and $|\eta| \sim 1$.

The distribution of the high energy B mesons at the LHC for energies $E_B \lesssim 300$ GeV can be approximated by the power law function, see the right panel in figure 18:

$$\frac{dN_B^{\text{MAT}}}{dE} \approx \tilde{f}_0 E^{-\alpha}, \quad \tilde{f}_0 \approx 1.6 \times 10^4 \text{ GeV}^{\alpha-1} \quad \text{and} \quad \alpha \simeq 4.6 \quad (\text{E.2})$$

Finally, the distribution of the HNLs originating from the W bosons can be approximated by the expression

$$\frac{dN_{N,W}}{dE_N} \approx F_0 \left(\frac{E_1 - E_N}{E_2} \right), \quad F_0 \approx 0.3 \text{ GeV}^{-1}, \quad E_1 = 60 \text{ GeV}, \quad E_2 \approx 107 \text{ GeV} \quad (\text{E.3})$$

E.2 Upper bound estimation

We start from the number of the events

$$N_{\text{events}}(M_X, \theta_X^2) = \tilde{N}_{\text{prod}}(M_X, \theta_X^2) \times \tilde{P}_{\text{decay}} \quad (\text{E.4})$$

Here for simplicity we defined quantities $\tilde{P}_{\text{decay}}$ and $\tilde{N}_{\text{decay}}$ defined by

$$P_{\text{decay}} = \epsilon_{\text{det}} \times \text{BR}_{\text{vis}} \tilde{P}_{\text{decay}}, \quad \tilde{N}_{\text{prod}} = N_{\text{prod}} \times \epsilon_{\text{det}} \times \text{BR}_{\text{vis}} \quad (\text{E.5})$$

The decay probability (3.5) can be rewritten in the form

$$\tilde{P}_{\text{decay}} = \int dE e^{-g(E)}, \quad (\text{E.6})$$

where $g(E) = l_{\text{target-det}} \Gamma_X M_X / E - \log \left(\frac{dN_X}{dE} \right)$ (for clarity we assumed $\frac{dN_X}{dE}$ to be dimensionless). The integral (3.5) can be evaluated as

$$\tilde{P}_{\text{decay}} \approx \sqrt{\frac{2\pi}{-g''(E_{\text{peak}})}} e^{-g(E_{\text{peak}})} \quad (\text{E.7})$$

where E_{peak} is the minimum of $g(E) = 0$, here we used the steepest descent approximation.

For the exponential spectrum (E.1) the peak energy and the probability are, correspondingly,

$$E_{\text{peak}} = \sqrt{\frac{l_{\text{target-det}} \Gamma_X M_X}{\delta}}, \quad P_{\text{decay}} \approx \sqrt{\pi} f_0 e^{-2E_{\text{peak}} \delta} \sqrt{\frac{E_{\text{peak}}}{\delta}}, \quad (\text{E.8})$$

while for the power law spectrum $f_E = f_0 E^{-\alpha}$ they are

$$E_{\text{peak}} = \frac{l_{\text{target-det}} \Gamma_X M_X}{\alpha}, \quad P_{\text{decay}} \approx \sqrt{\frac{2\pi}{\alpha}} f_0 e^{-\alpha} (E_{\text{peak}})^{-\alpha} \quad (\text{E.9})$$

Expressing then $\Gamma_X \propto U^2$ and $\theta_X^2 \equiv \theta_{\text{max}}^2(M_X) \times R$, for the upper bound given by the particles produced from B mesons one immediately arrives to

$$\theta_{X,\text{upper}}^{\text{SHiP}}(M_X) \approx \theta_{X,\text{max}}^{\text{SHiP}}(M_X) \frac{\log^2 \left(f_0 \sqrt{\pi} (\delta^{-3} \langle p_B \rangle)^{\frac{1}{4}} \tilde{N}_{\text{prod}}(M_X, \theta_{X,\text{max}}^{\text{SHiP}}(M_X)) \right)}{4 \langle p_B \rangle \delta}, \quad (\text{E.10})$$

$$\theta_{X,\text{upper}}^{\text{MAT}}(M_X) \approx \theta_{X,\text{max}}^{\text{MAT}}(M_X) \frac{\alpha}{\langle p_B \rangle} \left(\sqrt{\frac{2\pi}{\alpha^3}} \tilde{N}_{\text{prod}}(M_X, \theta_{X,\text{max}}^{\text{MAT}}(M_X)) \right)^{\frac{1}{\alpha-2}}. \quad (\text{E.11})$$

Similarly we can estimate the upper bound for the W bosons.

We estimate θ_{max} for the production from B mesons using the HNLs momenta given by eq. (2.2) with $\langle p_B \rangle = 12$ GeV for MATHUSLA and $\langle p_B \rangle = 88$ GeV for SHiP (see table 2), while for the production for W we use $\langle p_N \rangle \approx 62$ GeV, see section 5.3.

F Details of the sensitivity curve drawing

We draw the sensitivity curve for HNLs and scalars requiring

$$N_{\text{events}}(\theta_X, M_X) = \sum_{\text{meson}} N_{\text{events,meson}} + N_{\text{events},W} \geq 2.3, \quad (\text{F.1})$$

where numbers of decay events of particles produced from B, D mesons and W bosons are estimated as

$$N_{\text{events,meson}} = N_{q\bar{q}} \times f_{q \rightarrow \text{meson}} \text{Br}_{\text{meson} \rightarrow X} \times \epsilon_{\text{meson}} \int dp_{\text{meson}} f_{p_{\text{meson}}} \times \tilde{P}_{\text{decay}}(p_{\text{meson}}), \quad (\text{F.2})$$

$$N_{\text{events},W} = N_{W,\text{LHC}} \times \text{Br}_{W \rightarrow X} \times \epsilon_W \times \int dp_X f_{p_X,W} \times \tilde{P}_{\text{decay}}(p_X) \quad (\text{F.3})$$

Here, $N_{q\bar{q}}$ is the total number of the $q\bar{q}$ pairs that are produced in pp collisions at the high luminosity LHC or in p – target collisions at SHiP, $f_{p_{\text{meson}}}$ is the momentum distribution of the mesons that fly to the decay volume of the experiment (see section 5.1), and ϵ is the overall efficiency (see section 6.1). In the expression for the decay probability $\tilde{P}_{\text{decay}}$ the γ factor of the X particle is related to the meson momentum by the relation (C.2). $N_{W,\text{LHC}}$ is the number of the W bosons produced at the high luminosity LHC, and $f_{p_X,W}$ is the momentum distribution function of the X particles (see section 5.3).

X/M	M_X , GeV	$f_M \times \text{BR}_{M \rightarrow X}$	$\langle \gamma_X \rangle$	$c\tau_X$, m	ϵ	$U_{e,\text{lower}}^2$
$N, \nu_e/D$	0.5	$4.5 \cdot 10^{-2}$	30.1	$1.1 \cdot 10^{-2}$	$6.5 \cdot 10^{-2}$	$1.9 \cdot 10^{-9}$
$N, \nu_e/B$	3	$4 \cdot 10^{-4}$	20.2	$2 \cdot 10^{-6}$	0.13	$1.7 \cdot 10^{-8}$
S/B	0.5	88.2	30.1	$4.3 \cdot 10^{-9}$	0.14	$2.5 \cdot 10^{-11}$
S/B	2.5	21.4	20.2	$1 \cdot 10^{-9}$	0.2	$4.8 \cdot 10^{-12}$

Table 5. Table of parameters used in simple analytic estimation of the lower bound (2.1) for the SHiP experiment for particular masses of the HNLs with the pure electron mixing and the scalars. The columns are as follows: the type of the particle X and the mother particle, the branching ratio of the production of X at $\theta_X^2 = 1$, average γ factor, proper decay length $c\tau_X$ at $\theta_X^2 = 1$, overall efficiency (6.1), the mixing angle at the lower bound estimated as $N_{\text{events,lower}} = 2.3$, where $N_{\text{events,lower}}$ is given by (2.1).

X/M	M_X , GeV	$f_M \times \text{BR}_{M \rightarrow X}$	$\langle \gamma_X \rangle$	$c\tau_X$, m	ϵ	θ_{lower}^2
$N, \nu_e/D$	0.5	$4.5 \cdot 10^{-2}$	2.4	$1.1 \cdot 10^{-2}$	$6.5 \cdot 10^{-2}$	$1.2 \cdot 10^{-8}$
$N, \nu_e/B$	2.5	$2.6 \cdot 10^{-3}$	2.2	$5.4 \cdot 10^{-6}$	$1.8 \cdot 10^{-2}$	$2.4 \cdot 10^{-9}$
$N, \nu_e/W$	1	0.11	62	$6.5 \cdot 10^{-2}$	$1.9 \cdot 10^{-2}$	$1.7 \cdot 10^{-6}$
$N, \nu_e/W$	2.5	0.11	24.8	$5.4 \cdot 10^{-6}$	$1.9 \cdot 10^{-2}$	$1.1 \cdot 10^{-7}$
S/B	0.5	5.3	9	$4.3 \cdot 10^{-9}$	$1.8 \cdot 10^{-2}$	$4.2 \cdot 10^{-12}$
S/B	2.5	2.7	2	$1 \cdot 10^{-9}$	0.1	$4.9 \cdot 10^{-13}$

Table 6. Table of parameters used in simple analytic estimation (2.1) for the MATHUSLA experiment for particular masses of the HNLs and scalars. We use the description of the scalar phenomenology from [19]. The columns are as follows: the type of the particle X and the mother particle, the branching ratio of the production of X at $\theta_X^2 = 1$, average γ factor, proper decay length $c\tau_X$ at $\theta_X^2 = 1$, overall efficiency (6.1), the mixing angle at the lower bound estimated as $N_{\text{events,lower}} = 4$, where $N_{\text{events,lower}}$ is given by (2.1).

G Analytic estimation of the lower bound for particular masses

Here we make an analytic estimation of the lower bound for particular masses using the formula (2.1). The relevant parameters and the value θ_{lower}^2 for the SHiP and MATHUSLA experiments are given in tables 5, 6.

H HNLs at MATHUSLA for small mass

For HNLs with $M_N \lesssim m_{D_s}$, where the sensitivity curve is determined by the production from D mesons our sensitivity curve reproduces that of the MATHUSLA collaboration [19] in the range up to 1 GeV (figure 10). For smaller masses our estimates of the lower boundary differ by a factor ~ 3 (which would corresponds to the order-of-magnitude difference between the number of decay events). Moreover, the shapes of the sensitivity curves also differ.

Below we list several possible reasons for this discrepancy:

- a) different estimate of the number of the parent D mesons produced
- b) HNLs that are produced from the mesons that *do not fly* into the fiducial decay volume of MATHUSLA (they were not taken into account in our estimate)
- c) different HNL phenomenology (production and decay) used in comparison.
- d) Production from K -mesons was not taken into account in our estimates

Since the positions of the lower bounds in the mass range $1\text{ GeV} \lesssim M_N \lesssim m_{D_s}$ are in good agreement, we conclude that case a) with different amounts of D mesons is less probable, as it should shift the lower bound for the whole mass range $M_N \lesssim m_D$.

In order to estimate the amount of light HNLs produced from D mesons, we performed a MadGraph 5 simulation of a process $pp \rightarrow \bar{c}e^+\nu_e s$, whose kinematics corresponds to the main process for a production of light HNLs $M_N \lesssim 0.5\text{ GeV} - D \rightarrow Ne^+K$ [24]. We computed the ratio

$$\chi = \left(\frac{\sigma_{\bar{c}e^+\nu_e s, \text{ MAT}}}{\sigma_{\bar{c}e^+\nu_e s, \text{ tot}}} \right) / \left(\frac{\sigma_{c\bar{c}, \text{ MAT}}}{\sigma_{c\bar{c}, \text{ tot}}} \right), \quad (\text{H.1})$$

where the first fraction is the amount of the HNLs that fly in the decay volume of MATHUSLA, while the second one is the amount of $c\bar{c}$ pairs that fly in the same direction. We found $\chi \approx 1.7$, which is not enough to explain the discrepancy.

For c), we compared the decay widths of the HNLs used in our analysis with those, used in [19] (based on [97]). We found them to be different by 20 – 40% (for $U_e : U_\mu : U_\tau = 1$) with the decay width from [97] being smaller than that from [24]. The difference can reach up to 80% at small masses (below $\mathcal{O}(500)\text{ MeV}$).

Finally, production from K mesons would not explain why the discrepancy starts close to 1 GeV, much higher than production threshold from kaons.

We did not find the information about the HNL production ratios used in [19]. As we see, the cases b)–d) are not enough to explain a factor 10 in the number of events, and we assume that the main reason for the discrepancy is different production branching ratios adopted in [19].

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] J. Beacham et al., *Physics beyond colliders at CERN: beyond the Standard Model working group report*, [arXiv:1901.09966](https://arxiv.org/abs/1901.09966) [[INSPIRE](#)].
- [2] BELLE collaboration, *Search for a low mass particle decaying into $\mu^+\mu^-$ in $B^0 \rightarrow K^{*0}X$ and $B^0 \rightarrow \rho^0 X$ at Belle*, *Phys. Rev. Lett.* **105** (2010) 091801 [[arXiv:1005.1450](https://arxiv.org/abs/1005.1450)] [[INSPIRE](#)].
- [3] BABAR collaboration, *Search for low-mass dark-sector Higgs bosons*, *Phys. Rev. Lett.* **108** (2012) 211801 [[arXiv:1202.1313](https://arxiv.org/abs/1202.1313)] [[INSPIRE](#)].
- [4] LBNE collaboration, *The Long-Baseline Neutrino Experiment: exploring fundamental symmetries of the universe*, in *Snowmass 2013: Workshop on Energy Frontier*, Seattle, WA, U.S.A., 30 June–3 July 2013 [[arXiv:1307.7335](https://arxiv.org/abs/1307.7335)] [[INSPIRE](#)].

- [5] BABAR collaboration, *Search for a dark photon in e^+e^- collisions at BaBar*, *Phys. Rev. Lett.* **113** (2014) 201801 [[arXiv:1406.2980](#)] [[INSPIRE](#)].
- [6] LHCb collaboration, *Search for Majorana neutrinos in $B^- \rightarrow \pi^+ \mu^- \mu^-$ decays*, *Phys. Rev. Lett.* **112** (2014) 131802 [[arXiv:1401.5361](#)] [[INSPIRE](#)].
- [7] ATLAS collaboration, *Search for heavy Majorana neutrinos with the ATLAS detector in pp collisions at $\sqrt{s} = 8$ TeV*, *JHEP* **07** (2015) 162 [[arXiv:1506.06020](#)] [[INSPIRE](#)].
- [8] CMS collaboration, *Search for heavy Majorana neutrinos in $\mu^\pm \mu^\pm +$ jets events in proton-proton collisions at $\sqrt{s} = 8$ TeV*, *Phys. Lett. B* **748** (2015) 144 [[arXiv:1501.05566](#)] [[INSPIRE](#)].
- [9] BELLE collaboration, *Search for the dark photon and the dark Higgs boson at Belle*, *Phys. Rev. Lett.* **114** (2015) 211801 [[arXiv:1502.00084](#)] [[INSPIRE](#)].
- [10] NA64 collaboration, *Search for invisible decays of sub-GeV dark photons in missing-energy events at the CERN SPS*, *Phys. Rev. Lett.* **118** (2017) 011802 [[arXiv:1610.02988](#)] [[INSPIRE](#)].
- [11] BABAR collaboration, *Search for invisible decays of a dark photon produced in e^+e^- collisions at BaBar*, *Phys. Rev. Lett.* **119** (2017) 131804 [[arXiv:1702.03327](#)] [[INSPIRE](#)].
- [12] SHiP collaboration, *Prospects of the SHiP and NA62 experiments at CERN for hidden sector searches*, *PoS(NuFact2017)139* (2017) [[arXiv:1712.01768](#)] [[INSPIRE](#)].
- [13] A. Izmaylov and S. Suvorov, *Search for heavy neutrinos in the ND280 near detector of the T2K experiment*, *Phys. Part. Nucl.* **48** (2017) 984 [[INSPIRE](#)].
- [14] LHCb collaboration, *Search for dark photons produced in 13 TeV pp collisions*, *Phys. Rev. Lett.* **120** (2018) 061801 [[arXiv:1710.02867](#)] [[INSPIRE](#)].
- [15] NA62 collaboration, *Searches for very weakly-coupled particles beyond the Standard Model with NA62*, in *Proceedings, 13th Patras Workshop on Axions, WIMPs and WISPs, (PATRAS 2017)*, Thessaloniki, Greece, 15–19 May 2017, *DESY-PROC-2017-02*, (2018), pg. 145 [[arXiv:1711.08967](#)] [[INSPIRE](#)].
- [16] CMS collaboration, *Search for heavy neutral leptons in events with three charged leptons in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *Phys. Rev. Lett.* **120** (2018) 221801 [[arXiv:1802.02965](#)] [[INSPIRE](#)].
- [17] V.V. Gligorov, S. Knapen, M. Papucci and D.J. Robinson, *Searching for long-lived particles: a compact detector for exotics at LHCb*, *Phys. Rev. D* **97** (2018) 015023 [[arXiv:1708.09395](#)] [[INSPIRE](#)].
- [18] J.P. Chou, D. Curtin and H.J. Lubatti, *New detectors to explore the lifetime frontier*, *Phys. Lett. B* **767** (2017) 29 [[arXiv:1606.06298](#)] [[INSPIRE](#)].
- [19] D. Curtin et al., *Long-lived particles at the energy frontier: the MATHUSLA physics case*, [[arXiv:1806.07396](#)] [[INSPIRE](#)].
- [20] J.L. Feng, I. Galon, F. Kling and S. Trojanowski, *ForwArd Search ExpeRiment at the LHC*, *Phys. Rev. D* **97** (2018) 035001 [[arXiv:1708.09389](#)] [[INSPIRE](#)].
- [21] FASER collaboration, *FASER: ForwArd Search ExpeRiment at the LHC*, [[arXiv:1901.04468](#)] [[INSPIRE](#)].
- [22] V.V. Gligorov, S. Knapen, B. Nachman, M. Papucci and D.J. Robinson, *Leveraging the ALICE/L3 cavern for long-lived particle searches*, *Phys. Rev. D* **99** (2019) 015023 [[arXiv:1810.03636](#)] [[INSPIRE](#)].

- [23] F. Bezrukov and D. Gorbunov, *Light inflaton hunter's guide*, *JHEP* **05** (2010) 010 [[arXiv:0912.0390](#)] [[INSPIRE](#)].
- [24] K. Bondarenko, A. Boyarsky, D. Gorbunov and O. Ruchayskiy, *Phenomenology of GeV-scale heavy neutral leptons*, *JHEP* **11** (2018) 032 [[arXiv:1805.08567](#)] [[INSPIRE](#)].
- [25] A. Monin, A. Boyarsky and O. Ruchayskiy, *Hadronic decays of a light Higgs-like scalar*, *Phys. Rev. D* **99** (2019) 015019 [[arXiv:1806.07759](#)] [[INSPIRE](#)].
- [26] F. Bezrukov, D. Gorbunov and I. Timiryasov, *Uncertainties of hadronic scalar decay calculations*, [arXiv:1812.08088](#) [[INSPIRE](#)].
- [27] M.W. Winkler, *Decay and detection of a light scalar boson mixing with the Higgs boson*, *Phys. Rev. D* **99** (2019) 015018 [[arXiv:1809.01876](#)] [[INSPIRE](#)].
- [28] I. Boiarska, K. Bondarenko, A. Boyarsky, V. Gorkavenko, M. Ovchinnikov and A. Sokolenko, *Phenomenology of GeV-scale scalar portal*, [arXiv:1904.10447](#) [[INSPIRE](#)].
- [29] G. Krnjaic, *Probing light thermal dark-matter with a Higgs portal mediator*, *Phys. Rev. D* **94** (2016) 073009 [[arXiv:1512.04119](#)] [[INSPIRE](#)].
- [30] C.O. Dib, C.S. Kim and S. Tapia Araya, *Search for light sterile neutrinos from $W^\pm$ decays at the LHC*, [arXiv:1903.04905](#) [[INSPIRE](#)].
- [31] SHiP collaboration, *Sensitivity of the SHiP experiment to heavy neutral leptons*, *JHEP* **04** (2019) 077 [[arXiv:1811.00930](#)] [[INSPIRE](#)].
- [32] D. Curtin and M.E. Peskin, *Analysis of long lived particle decays with the MATHUSLA detector*, *Phys. Rev. D* **97** (2018) 015006 [[arXiv:1705.06327](#)] [[INSPIRE](#)].
- [33] J.A. Evans, *Detecting hidden particles with MATHUSLA*, *Phys. Rev. D* **97** (2018) 055046 [[arXiv:1708.08503](#)] [[INSPIRE](#)].
- [34] J.C. Helo, M. Hirsch and Z.S. Wang, *Heavy neutral fermions at the high-luminosity LHC*, *JHEP* **07** (2018) 056 [[arXiv:1803.02212](#)] [[INSPIRE](#)].
- [35] SHiP collaboration, *A facility to Search for Hidden Particles (SHiP) at the CERN SPS*, [arXiv:1504.04956](#) [[INSPIRE](#)].
- [36] S. Alekhin et al., *A facility to Search for Hidden Particles at the CERN SPS: the SHiP physics case*, *Rept. Prog. Phys.* **79** (2016) 124201 [[arXiv:1504.04855](#)] [[INSPIRE](#)].
- [37] SHiP collaboration, *The experimental facility for the Search for Hidden Particles at the CERN SPS*, *2019 JINST* **14** P03025 [[arXiv:1810.06880](#)] [[INSPIRE](#)].
- [38] P. Minkowski, $\mu \rightarrow e\gamma$ at a rate of one out of 10^9 muon decays?, *Phys. Lett. B* **67** (1977) 421 [[INSPIRE](#)].
- [39] T. Yanagida, *Horizontal gauge symmetry and masses of neutrinos*, *Conf. Proc. C* **7902131** (1979) 95 [[INSPIRE](#)].
- [40] S.L. Glashow, *The future of elementary particle physics*, *NATO Sci. Ser. B* **61** (1980) 687 [[INSPIRE](#)].
- [41] M. Gell-Mann, P. Ramond and R. Slansky, *Complex spinors and unified theories*, *Conf. Proc. C* **790927** (1979) 315 [[arXiv:1306.4669](#)] [[INSPIRE](#)].
- [42] R.N. Mohapatra and G. Senjanović, *Neutrino mass and spontaneous parity nonconservation*, *Phys. Rev. Lett.* **44** (1980) 912 [[INSPIRE](#)].

- [43] R.N. Mohapatra and G. Senjanović, *Neutrino masses and mixings in gauge models with spontaneous parity violation*, *Phys. Rev. D* **23** (1981) 165 [INSPIRE].
- [44] V. Silveira and A. Zee, *Scalar phantoms*, *Phys. Lett. B* **161** (1985) 136 [INSPIRE].
- [45] J. McDonald, *Gauge singlet scalars as cold dark matter*, *Phys. Rev. D* **50** (1994) 3637 [[hep-ph/0702143](#)] [INSPIRE].
- [46] T. Binoth and J.J. van der Bij, *Influence of strongly coupled, hidden scalars on Higgs signals*, *Z. Phys. C* **75** (1997) 17 [[hep-ph/9608245](#)] [INSPIRE].
- [47] C.P. Burgess, M. Pospelov and T. ter Veldhuis, *The minimal model of nonbaryonic dark matter: a singlet scalar*, *Nucl. Phys. B* **619** (2001) 709 [[hep-ph/0011335](#)] [INSPIRE].
- [48] R.M. Schabinger and J.D. Wells, *A minimal spontaneously broken hidden sector and its impact on Higgs boson physics at the Large Hadron Collider*, *Phys. Rev. D* **72** (2005) 093007 [[hep-ph/0509209](#)] [INSPIRE].
- [49] M.J. Strassler and K.M. Zurek, *Echoes of a hidden valley at hadron colliders*, *Phys. Lett. B* **651** (2007) 374 [[hep-ph/0604261](#)] [INSPIRE].
- [50] B. Patt and F. Wilczek, *Higgs-field portal into hidden sectors*, [[hep-ph/0605188](#)] [INSPIRE].
- [51] D. O’Connell, M.J. Ramsey-Musolf and M.B. Wise, *Minimal extension of the Standard Model scalar sector*, *Phys. Rev. D* **75** (2007) 037701 [[hep-ph/0611014](#)] [INSPIRE].
- [52] M. Drewes, J. Hajer, J. Klaric and G. Lanfranchi, *NA62 sensitivity to heavy neutral leptons in the low scale seesaw model*, *JHEP* **07** (2018) 105 [[arXiv:1801.04207](#)] [INSPIRE].
- [53] SHiP collaboration, *A facility to Search for Hidden Particles (SHiP) at the CERN SPS: addendum to technical proposal*, CERN-SPSC-2015-040, CERN, Geneva, Switzerland (2015) [SPSC-P-350-ADD-2] [INSPIRE].
- [54] SHiP collaboration, *Optimising the active muon shield for the SHiP experiment at CERN*, *J. Phys. Conf. Ser.* **934** (2017) 012050 [INSPIRE].
- [55] SHiP collaboration, *The active muon shield in the SHiP experiment*, *2017 JINST* **12** P05011 [[arXiv:1703.03612](#)] [INSPIRE].
- [56] I. Boiarska et al., *Scalars from Higgs bosons at FASER2*, to appear, (2019).
- [57] LHCb collaboration, *Observation of $B_c^+ \rightarrow D^0 K^+$ decays*, *Phys. Rev. Lett.* **118** (2017) 111803 [[arXiv:1701.01856](#)] [INSPIRE].
- [58] CDF collaboration, *Observation of the B_c meson in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV*, *Phys. Rev. Lett.* **81** (1998) 2432 [[hep-ex/9805034](#)] [INSPIRE].
- [59] CDF collaboration, *Observation of B_c mesons in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV*, *Phys. Rev. D* **58** (1998) 112004 [[hep-ex/9804014](#)] [INSPIRE].
- [60] K.-M. Cheung, *B_c meson production at the Tevatron revisited*, *Phys. Lett. B* **472** (2000) 408 [[hep-ph/9908405](#)] [INSPIRE].
- [61] A.V. Berezhnoy, *Color flows for the process $gg \rightarrow B_c + c + \bar{b}$* , *Phys. Atom. Nucl.* **68** (2005) 1866 [*Yad. Fiz.* **68** (2005) 1928] [[hep-ph/0407315](#)] [INSPIRE].
- [62] K. Kolodziej and R. Ruckl, *On the energy dependence of hadronic B_c production*, *Nucl. Instrum. Meth. A* **408** (1998) 33 [[hep-ph/9803327](#)] [INSPIRE].
- [63] K. Kolodziej, A. Leike and R. Ruckl, *Production of B_c mesons in hadronic collisions*, *Phys. Lett. B* **355** (1995) 337 [[hep-ph/9505298](#)] [INSPIRE].

- [64] SHiP collaboration, *Heavy flavour cascade production in a beam dump*, [CERN-SHiP-NOTE-2015-009](#), CERN, Geneva, Switzerland (2015).
- [65] CMS collaboration, *Measurement of the total and differential inclusive B^+ hadron cross sections in pp collisions at $\sqrt{s} = 13$ TeV*, *Phys. Lett. B* **771** (2017) 435 [[arXiv:1609.00873](#)] [[INSPIRE](#)].
- [66] ATLAS collaboration, *Measurement of $D^{*\pm}$, $D^\pm$ and $D_s^\pm$ meson production cross sections in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector*, *Nucl. Phys. B* **907** (2016) 717 [[arXiv:1512.02913](#)] [[INSPIRE](#)].
- [67] LHCb collaboration, *Measurement of B meson production cross-sections in proton-proton collisions at $\sqrt{s} = 7$ TeV*, *JHEP* **08** (2013) 117 [[arXiv:1306.3663](#)] [[INSPIRE](#)].
- [68] CMS collaboration, *Transverse-momentum and pseudorapidity distributions of charged hadrons in pp collisions at $\sqrt{s} = 7$ TeV*, *Phys. Rev. Lett.* **105** (2010) 022002 [[arXiv:1005.3299](#)] [[INSPIRE](#)].
- [69] M. Cacciari, S. Frixione, N. Houdeau, M.L. Mangano, P. Nason and G. Ridolfi, *Theoretical predictions for charm and bottom production at the LHC*, *JHEP* **10** (2012) 137 [[arXiv:1205.6344](#)] [[INSPIRE](#)].
- [70] M. Cacciari, M. Greco and P. Nason, *The p_T spectrum in heavy flavor hadroproduction*, *JHEP* **05** (1998) 007 [[hep-ph/9803400](#)] [[INSPIRE](#)].
- [71] M. Cacciari, S. Frixione and P. Nason, *The p_T spectrum in heavy flavor photoproduction*, *JHEP* **03** (2001) 006 [[hep-ph/0102134](#)] [[INSPIRE](#)].
- [72] M. Cacciari, M.L. Mangano and P. Nason, *Gluon PDF constraints from the ratio of forward heavy-quark production at the LHC at $\sqrt{s} = 7$ and 13 TeV*, *Eur. Phys. J. C* **75** (2015) 610 [[arXiv:1507.06197](#)] [[INSPIRE](#)].
- [73] ATLAS collaboration, *Search for scalar charm quark pair production in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector*, *Phys. Rev. Lett.* **114** (2015) 161801 [[arXiv:1501.01325](#)] [[INSPIRE](#)].
- [74] LHCb collaboration, *Measurements of prompt charm production cross-sections in pp collisions at $\sqrt{s} = 13$ TeV*, *JHEP* **03** (2016) 159 [Erratum *ibid.* **09** (2016) 013] [[arXiv:1510.01707](#)] [[INSPIRE](#)].
- [75] LHCb collaboration, *Search for rare $B_{(s)}^0 \rightarrow \mu^+ \mu^- \mu^+ \mu^-$ decays*, *Phys. Rev. Lett.* **110** (2013) 211801 [[arXiv:1303.1092](#)] [[INSPIRE](#)].
- [76] C.-H. Chang, C. Driouichi, P. Eerola and X.G. Wu, *BCVEGPY: an event generator for hadronic production of the B_c meson*, *Comput. Phys. Commun.* **159** (2004) 192 [[hep-ph/0309120](#)] [[INSPIRE](#)].
- [77] ATLAS collaboration, *Measurement of $W^\pm$ and Z -boson production cross sections in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Phys. Lett. B* **759** (2016) 601 [[arXiv:1603.09222](#)] [[INSPIRE](#)].
- [78] ATLAS collaboration, *Measurement of the transverse momentum distribution of W bosons in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector*, *Phys. Rev. D* **85** (2012) 012005 [[arXiv:1108.6308](#)] [[INSPIRE](#)].
- [79] CMS collaboration, *Measurement of the transverse momentum spectra of weak vector bosons produced in proton-proton collisions at $\sqrt{s} = 8$ TeV*, *JHEP* **02** (2017) 096 [[arXiv:1606.05864](#)] [[INSPIRE](#)].

- [80] J. Alwall et al., *The automated computation of tree-level and next-to-leading order differential cross sections and their matching to parton shower simulations*, *JHEP* **07** (2014) 079 [[arXiv:1405.0301](#)] [[INSPIRE](#)].
- [81] SHiP collaboration, *HNL sensitivity of SHiP experiment*, *Zenodo*, (2018).
- [82] SHiP collaboration, *Sensitivity of the SHiP experiment to dark scalars of GeV mass*, to appear, (2019).
- [83] B. Batell, M. Pospelov and A. Ritz, *Exploring portals to a hidden sector through fixed targets*, *Phys. Rev. D* **80** (2009) 095024 [[arXiv:0906.5614](#)] [[INSPIRE](#)].
- [84] J.D. Clarke, R. Foot and R.R. Volkas, *Phenomenology of a very light scalar ($100\text{ MeV} \leq m_h \leq 10\text{ GeV}$) mixing with the SM Higgs*, *JHEP* **02** (2014) 123 [[arXiv:1310.8042](#)] [[INSPIRE](#)].
- [85] K. Schmidt-Hoberg, F. Staub and M.W. Winkler, *Constraints on light mediators: confronting dark matter searches with B physics*, *Phys. Lett. B* **727** (2013) 506 [[arXiv:1310.6752](#)] [[INSPIRE](#)].
- [86] LHCb collaboration, *Measurement of b-hadron production fractions in 7 TeV pp collisions*, *Phys. Rev. D* **85** (2012) 032008 [[arXiv:1111.2357](#)] [[INSPIRE](#)].
- [87] F. Kling and S. Trojanowski, *Heavy neutral leptons at FASER*, *Phys. Rev. D* **97** (2018) 095016 [[arXiv:1801.08947](#)] [[INSPIRE](#)].
- [88] SHiP collaboration, *Mass dependence of branching ratios into HNL for FairShip*, *CERN-SHiP-NOTE-2016-001*, CERN, Geneva, Switzerland (2016).
- [89] D. Gorbunov and M. Shaposhnikov, *How to find neutral leptons of the νMSM ?*, *JHEP* **10** (2007) 015 [Erratum *ibid.* **11** (2013) 101] [[arXiv:0705.1729](#)] [[INSPIRE](#)].
- [90] A. Atre, T. Han, S. Pascoli and B. Zhang, *The search for heavy Majorana neutrinos*, *JHEP* **05** (2009) 030 [[arXiv:0901.3589](#)] [[INSPIRE](#)].
- [91] ATLAS collaboration, *Measurement of Υ production in 7 TeV pp collisions at ATLAS*, *Phys. Rev. D* **87** (2013) 052004 [[arXiv:1211.7255](#)] [[INSPIRE](#)].
- [92] CMS collaboration, *Measurement of the $\Upsilon(1S)$, $\Upsilon(2S)$ and $\Upsilon(3S)$ cross sections in pp collisions at $\sqrt{s} = 7\text{ TeV}$* , *Phys. Lett. B* **727** (2013) 101 [[arXiv:1303.5900](#)] [[INSPIRE](#)].
- [93] Z. Hu, N.T. Leonardo, T. Liu and M. Haytmyradov, *Review of bottomonium measurements from CMS*, *Int. J. Mod. Phys. A* **32** (2017) 1730015 [[arXiv:1708.02913](#)] [[INSPIRE](#)].
- [94] PARTICLE DATA GROUP collaboration, *Review of particle physics*, *Phys. Rev. D* **98** (2018) 030001 [[INSPIRE](#)].
- [95] H.-Y. Cheng, C.-K. Chua, K.-C. Yang and Z.-Q. Zhang, *Revisiting charmless hadronic B decays to scalar mesons*, *Phys. Rev. D* **87** (2013) 114001 [[arXiv:1303.4403](#)] [[INSPIRE](#)].
- [96] M. Spira, *QCD effects in Higgs physics*, *Fortsch. Phys.* **46** (1998) 203 [[hep-ph/9705337](#)] [[INSPIRE](#)].
- [97] J.C. Helo, S. Kovalenko and I. Schmidt, *Sterile neutrinos in lepton number and lepton flavor violating decays*, *Nucl. Phys. B* **853** (2011) 80 [[arXiv:1005.1607](#)] [[INSPIRE](#)].