RECEIVED: April 4, 2015

REVISED: May 28, 2015

ACCEPTED: July 23, 2015

PUBLISHED: September 1, 2015

Enhancing $t\bar{t}hh$ production through CP-violating top-Higgs interaction at the LHC and future colliders

Ning Liu,^{a,b} Yanming Zhang,^a Jinzhong Han^c and Bingfang Yang^a

^a*Institute of Theoretical Physics, Henan Normal University,
Xinxiang 453007, China*

^b*ARC Centre of Excellence for Particle Physics at the Terascale,
School of Physics, The University of Sydney,
Sydney, NSW 2006, Australia*

^c*School of Physics and Electromechanical Engineering, Zhoukou Normal University,
Zhoukou, 466001, China*

E-mail: wlln@mail.ustc.edu.cn, zhangyanming@henannu.edu.cn,
hanjinzhong@zknpu.edu.cn, yangbingfang@htu.edu.cn

ABSTRACT: The measurement of Higgs self-coupling is one of the most crucial physics goals at the future colliders. At the LHC, the di-Higgs production is a main way to measure the Higgs trilinear coupling. As a complementary to the di-Higgs production, $t\bar{t}hh$ process may open a new avenue to measure di-Higgs physics at the LHC and a future 100 TeV pp collider or a high energy e^+e^- collider since the extra $t\bar{t}$ in the final states may efficiently suppress the backgrounds. However, such a kind of process is also controlled by the top-Higgs coupling. In this work, we investigate the impact of CP-violating top-Higgs coupling on $t\bar{t}hh$ production at the LHC, e^+e^- and a 100 TeV hadron collider under the current Higgs data. Within 2σ Higgs data allowed parameter region, we find that the cross section of $t\bar{t}hh$ at the LHC-14 TeV, e^+e^- -1 TeV and VHE-LHC/SPPC-100 TeV can be enhanced up to 2.1 times the SM predictions. The future precise measurement of Higgs coupling will reveal the nature of top-Higgs interaction and improve the sensitivity of the determination of Higgs self-coupling through $t\bar{t}hh$ production.

KEYWORDS: Beyond Standard Model, Heavy Quark Physics

ARXIV EPRINT: [1503.08537](https://arxiv.org/abs/1503.08537)

Contents

1	Introduction	1
2	CP-violating Top-Higgs couplings	2
3	Numerical results and discussions	3
3.1	LHC and VHE-LHC/SPPC	4
3.2	e^+e^- and $\gamma\gamma$ collisions	7
4	Conclusion	11

1 Introduction

In 2012, a bosonic resonance with a mass around 125 GeV was independently observed by the ATLAS and CMS collaborations at the LHC [1, 2]. Up to now, most measurements of its properties are consistent with the predictions of the Standard Model (SM) Higgs boson [3–6]. However, the Higgs couplings with top quarks and with itself are still vacant and need to be verified at the future colliders.

The tree-level Higgs potential of the SM is given by,

$$V = -\mu^2(\phi^\dagger\phi) + \lambda(\phi^\dagger\phi)^2, \quad \phi = \frac{1}{\sqrt{2}}(0, v + h)^T, \quad (1.1)$$

which yields the trilinear and quartic Higgs self-couplings,

$$\lambda_{3h} = \frac{3m_h^2}{v}, \quad \lambda_{4h} = \frac{3m_h^2}{v^2}, \quad (1.2)$$

where the Higgs boson mass $m_h \simeq 125$ GeV have been well measured in the experiments and the vacuum expectation value of the Higgs field, $v = (\sqrt{2}G_F)^{-1/2} \simeq 246$ GeV. Hence, the determination of the Higgs self-coupling can directly test the relation eq. (1.2) obtained from the minimization of SM Higgs potential eq. (1.1). At the LHC, the main way to measure the Higgs self-coupling is the di-Higgs production, which is dominated by the gluon-gluon fusion mechanism and has small cross section (see examples in SM after 125 GeV Higgs boson [7–12]). Among various decay channels, the process $hh \rightarrow b\bar{b}\gamma\gamma$ is expected to have the most promising sensitivity due to the low backgrounds at the LHC [13–15] while the $4b$ final state has the largest fraction [16, 17]. Some applications of advanced analysis techniques to di-Higgs production have been proposed to improve the sensitivity of $b\bar{b}\tau^+\tau^-$ and $b\bar{b}W^+W^-$ channel [18–28]. However, one should keep in mind that, the main production process of di-Higgs $gg \rightarrow hh$ can also be triggered by the top-Higgs coupling itself through the box diagrams. So, any new physics in the top-Higgs coupling may significantly affect the measurement of Higgs self-coupling at the LHC [29–37].

In the SM, top quark has the strongest coupling to the Higgs boson and is widely speculated as a sensitive probe to the new physics beyond the SM. The most sensitive direct

way of measuring the top-Higgs coupling at the LHC is the associated production of the top pair with Higgs boson that has been extensively studied in literature [38–59]. However, since the cross section of $pp \rightarrow t\bar{t}h$ is about 130 fb at 8 TeV LHC [60]), current LHC luminosity and analysis are not yet sensitivity enough to observe such a signal. An upper limit on the signal strength $\mu_{t\bar{t}h} = \sigma_{t\bar{t}h}/\sigma_{t\bar{t}h}^{\text{SM}}$ to be $\mu_{t\bar{t}h} \lesssim 3.9$ at 95% C.L. limit has been set up by ATLAS collaboration through combining $h \rightarrow b\bar{b}$ and $h \rightarrow \gamma\gamma$ channels [61]. While the CMS collaboration gives a limit of $0.9 \lesssim \mu_{t\bar{t}h} \lesssim 3.5$ by using all search channels [62]. The other direct way is to observe the single top associated production with the Higgs boson. Such a process has a small cross section (18.28 fb at 8 TeV LHC) but is advocated to determine the sign of the top-Higgs coupling at the LHC [63–82]. Recently, the CMS collaboration has presented the result on thj searches in the $h \rightarrow \gamma\gamma$ channel, and obtained a weak bound on the cross section of events with inverted top-Higgs coupling [83].

In this work, we will investigate the effect of non-standard top-Higgs coupling in the $t\bar{t}hh$ production at the LHC, a e^+e^- and a 100 TeV hadron collider under the current Higgs data constraints. At the LHC, although the cross section of $pp \rightarrow t\bar{t}hh$ is about an order smaller than that of $pp \rightarrow hh$, the additional $t\bar{t}$ in the final states may suppress one order or orders more backgrounds [84, 85]. Besides, such a process has a cross section monotonically increasing with respect to Higgs self-coupling [11], which may complement the process $pp \rightarrow hh$ in measuring Higgs self-coupling, particularly for $\lambda \gg \lambda_{\text{SM}}$. So, the process $pp \rightarrow t\bar{t}hh$ opens a new avenue to measure di-Higgs physics at HL-LHC and a future 100 TeV pp -collider. On the other hand, given the limited precision of the LHC, an e^+e^- collider is crucial to scrutinize the detailed properties of the Higgs boson that might uncover the new physics beyond the SM [86–96]. In e^+e^- collision, the double Higgs strahlung $e^+e^- \rightarrow Zh h$ is considered as the main production process to measure the Higgs self-coupling. Besides, it should be noted that the di-Higgs bosons can also be radiated off from top quarks through the process $e^+e^- \rightarrow t\bar{t}hh$, or $\gamma\gamma \rightarrow t\bar{t}hh$ at photon collider, where the energetic electron beam to a photon beam is converted through the backward Compton scattering [97]. Similar to $pp \rightarrow t\bar{t}hh$, the processes $e^+e^-/\gamma\gamma \rightarrow t\bar{t}hh$ not only involve the Higgs self-coupling but also are governed by the top-Higgs coupling. So the investigation of $t\bar{t}hh$ production may lead us to obtain the nontrivial information on the Higgs potential and test the top-Higgs coupling [98, 99].

The structure of this paper is organized as follows. In section 2, we will briefly introduce the non-standard top-Higgs interaction and set up the calculations. In section 3, we present the numerical results and discuss the effects of non-standard top-Higgs coupling in the di-Higgs production at the LHC, e^+e^- and a 100 TeV hadron collider. Finally, we draw our conclusions in section 4.

2 CP-violating Top-Higgs couplings

In this study, we parameterize the top-Higgs couplings using the phenomenological Lagrangian:

$$\mathcal{L} = -\frac{y_t}{\sqrt{2}}\bar{t}(\cos\theta + i\gamma^5\sin\theta)th. \quad (2.1)$$

In the SM, y_t takes the value $y_t^{\text{SM}} = \sqrt{2}m_t/v$ and $\sin\theta = 0$, with $v \simeq 246$ GeV being the vacuum expectation value of the Higgs field. A pure pseudo-scalar interaction can be obtained by setting $\cos\theta = 0$. A CP violating interaction is realized if both $\cos\theta \neq 0$ and $\sin\theta \neq 0$. The exact values of these coefficients depend on the specific model. Here we are interested in a model-independent approach to determine the impact of general top-Higgs coupling on $t\bar{t}hh$ production. In the following calculations, we define two reduced couplings: $c_t = y_t \cos\theta/y_{t\text{SM}}$ and $\tilde{c}_t = y_t \sin\theta/y_{t\text{SM}}$ to discuss our results.

Since the CP-violating top-Higgs interaction can sizably alter the production rate of $gg \rightarrow h$ and decay width of $h \rightarrow \gamma\gamma$ through the loop effect, the most relevant indirect constraint on the couplings c_t and $\tilde{c}_t$ should be from the Higgs data. The signal strength of one specific analysis from a single Higgs boson is given by

$$\mu = \sum_i \mu_i \omega_i, \quad (2.2)$$

where the sum runs over all channels used in the experimental analysis. Each channel is characterized by one specific production and decay mode. The individual channel signal strength can be calculated by

$$\mu_i = \frac{[\sigma \times BR]_i}{[\sigma_{\text{SM}} \times BR_{\text{SM}}]_i}, \quad (2.3)$$

and the SM channel weight is

$$\omega_i = \frac{\epsilon_i [\sigma_{\text{SM}} \times BR_{\text{SM}}]_i}{\sum_j \epsilon_j [\sigma_{\text{SM}} \times BR_{\text{SM}}]_j}, \quad (2.4)$$

where ϵ_i is the relative experimental efficiencies for each channel. But these are rarely quoted in experimental publications. In this case, all channels considered in the analysis are treated equally, i.e. $\epsilon_i = 1$. The reduced effective coupling c_{hgg} and $c_{h\gamma\gamma}$ can be parameterized through the c_t and $\tilde{c}_t$ as following [100],

$$\begin{aligned} c_{hgg}^2 &\simeq c_t^2 + 2.6\tilde{c}_t^2 + 0.11c_t(c_t - 1), \\ c_{h\gamma\gamma}^2 &\simeq (1.28 - 0.28c_t)^2 + (0.43\tilde{c}_t)^2. \end{aligned} \quad (2.5)$$

We confront the effective coupling c_{hgg} and $c_{h\gamma\gamma}$ with the Higgs data by calculating the χ^2 of the Higgs sector with the public package **HiggsSignals-1.3.2** [101, 102]. We choose the mass-centered χ^2 method in the package **HiggsSignals**. Although the CP-violating top-Higgs interaction can contribute to the electric dipole moment (EDM), the bounds on the coupling $\tilde{c}_t$ strongly rely on the assumption of Higgs couplings to other light fermions [103, 104]. But such light quark Yukawa couplings are generally unobservable at the LHC, we do not impose EDM constraints in the study. Other low-energy physics constraints, such as $B_s - \bar{B}_s$ and $B \rightarrow X_s \gamma$, are still too weak due to the large uncertainty [100].

3 Numerical results and discussions

In our numerical calculations, we take the input parameters of the SM as [105, and 2013 partial update for the 2014 edition]

$$\begin{aligned} m_t &= 173.07 \text{ GeV}, & m_W &= 80.385, & m_Z &= 91.19 \text{ GeV}, \\ m_h &= 125.9 \text{ GeV}, & \sin^2 \theta_W &= 0.2228, & \alpha(m_Z)^{-1} &= 127.918. \end{aligned} \quad (3.1)$$

Figure 1. Feynman diagrams for the partonic process $gg \rightarrow t\bar{t}hh$ at hadron collider.

For the strong coupling constant $\alpha_s(\mu)$, we use its 2-loop evolution with QCD parameter $\Lambda^{n_f=5} = 226 \text{ MeV}$ and get $\alpha_s(m_Z) = 0.118$. We use CT10 parton distribution functions (PDF) for the calculation of $pp \rightarrow t\bar{t}hh$ [106]. The renormalization scale μ_R and factorization scale μ_F are chosen to be $\mu_R = \mu_F = (m_h + m_t)$. It should be noted that the higher order corrections are usually needed to improve the reliability of the leading order results. However, such calculations for four-body production involve complicated techniques and are beyond our study in this work. In the following we will use the ratios of the leading order cross sections $\sigma_{t\bar{t}hh}/\sigma_{t\bar{t}hh}^{\text{SM}}$ to present our results, which have the weak dependence on the variation of the scale. Besides, it is expected that the high order effect on the cross sections can be largely canceled in the ratios. We perform the numerical calculations by the package `calcchep-3.4` [107].

3.1 LHC and VHE-LHC/SPPC

At hadron collider, the main contributions to $t\bar{t}hh$ production are from the gluon fusion processes. The corresponding Feynman diagrams for the partonic process $gg \rightarrow t\bar{t}hh$ are shown in figure 1. The $q\bar{q}$ annihilation processes can be obtained by replacing the initial gluons with $q\bar{q}$ in the s -channel in figure 1. According to the production of di-Higgs bosons, we can classify amplitudes of $pp \rightarrow t\bar{t}hh$ into two categories: one is proportional to $\alpha_s y_t^2$; the other one is proportional to $\alpha_s y_t \lambda_{3h}$. In the SM, both amplitudes have the same sign and are constructive, which lead to the cross section of $pp \rightarrow t\bar{t}hh$ monotonically increases with respect to Higgs self-coupling λ_{3h} . However, it should be noted that each amplitude is

Figure 2. Ratios of $\sigma^{pp \rightarrow t\bar{t}hh} / \sigma_{SM}^{pp \rightarrow t\bar{t}hh}$ in the plane of $\tilde{c}_t - c_t$ at 14 TeV LHC and VHE-LHC/SPPC, where the dashed contours correspond to the 68% C.L. and 95% C.L. limits given by the current Higgs data fitting.

modulated by the top-Higgs coupling. So the measurement of Higgs self-coupling through $pp \rightarrow t\bar{t}hh$ strongly depends on the determination of top-Higgs coupling.

In figure 2, we respectively show the ratios of $\sigma^{pp \rightarrow t\bar{t}hh} / \sigma_{SM}^{pp \rightarrow t\bar{t}hh}$ in the plane of $\tilde{c}_t - c_t$ at 14 TeV LHC and VHE-LHC/SPPC, where the dashed contours correspond to the 68% C.L. and 95% C.L. limits given by the current Higgs data fitting. From figure 2, we can see that the negative scalar component c_t in eq. (2.1) is strongly disfavoured because the experimental measurement of the Higgs diphoton rate are consistent with the SM prediction. The pseudoscalar component in the range $|\tilde{c}_t| \gtrsim 0.6$ have been excluded at 95% C.L. by the Higgs data fitting. Besides, when $c_t = 1$ (or $y_t = y_t^{SM}$), a narrow region of pseudoscalar component $|\tilde{c}_t| \lesssim 0.3(0.44)$ is still allowed at 68% (95%) C.L..

As mentioned before, the cross section of $pp \rightarrow t\bar{t}hh$ not only depends on the magnitude of the top-Higgs coupling but also on the relative phase angle between c_t and $\tilde{c}_t$. (i) When $c_t = -1$ and $\tilde{c}_t = 0$ (or $y_t = -y_t^{SM}$), the ration of $\sigma^{pp \rightarrow t\bar{t}hh} / \sigma_{SM}^{pp \rightarrow t\bar{t}hh}$ is smaller than 1 due to the deconstructive interference between the processes with coupling $\alpha_s y_t^2$ and those with coupling $\alpha_s y_t \lambda_{3h}$; (ii) when $c_t = 0$ and $\tilde{c}_t = 1$ (or $y_t = i y_t^{SM}$), there is no interference between the terms with $\alpha_s y_t^2$ and those with $\alpha_s y_t \lambda_{3h}$, which will reduce the cross section of $pp \rightarrow t\bar{t}hh$. Finally, we find that the value of the ratio $\sigma^{pp \rightarrow t\bar{t}hh} / \sigma_{SM}^{pp \rightarrow t\bar{t}hh}$ can maximally reach about 1.5 (2.1) in the 68% (95%) C.L. allowed region at 14 TeV LHC and VHE-LHC/SPPC.

We implement the CP-violating interaction of $t\bar{t}h$ in eq. (2.1) with the package FeynRules [108]. In figure 3, we show the parton-level transverse momentum distributions of top quark and Higgs boson in the presence of the CP-violating interactions by using the package MadGraph5 [109] at 14 TeV LHC. From figure 3, we can see that the shapes of p_T

Figure 3. The parton-level transverse momentum distributions of top quark and Higgs boson at 14 TeV LHC.

Figure 4. The parton-level distributions of polar angle $\Delta\theta_{th}$ and the azimuthal angle $\Delta\phi_{th}$ between the top quark and the Higgs boson at 14 TeV LHC.

distributions of top quark and Higgs boson in the SM are almost the same as those in the CP-violating interactions.

Since the CP-violating $t\bar{t}h$ interaction in eq. (2.1) may affect the polarization states of the top quark, we present the parton-level distributions of polar angle $\Delta\theta_{th}$ and the azimuthal angle $\Delta\phi_{th}$ between the top quark and the Higgs boson. From figure 4, we can find that the CP-violating interactions slightly change the distributions of $\Delta\theta_{th}$ and $\Delta\phi_{th}$, as the comparison with the SM. To be specific, when $\Delta\theta_{th} \leq \pi/2$ ($\Delta\theta_{th} > \pi/2$), the SM has more (less) events than the CP-violating interactions. Similar results can be observed in the distribution of $\Delta\phi_{th}$. Here it should be mentioned that such differences will be more obvious if the pseudo-scalar component $\tilde{c}_t$ becomes large. However, as shown in figure 2, the current Higgs data has already tightly constrained the size of $\tilde{c}_t$. Besides, these differences will be further diluted when the detector effects are taken into account.

The LHC observability of $t\bar{t}hh$ in the SM including the parton shower and detector effects has been investigated through both Higgs bosons decaying $h \rightarrow b\bar{b}$ and semi-leptonic and hadronic top decays in ref. [84]. The main backgrounds are $t\bar{t}hb\bar{b}$, $t\bar{t}b\bar{b}b\bar{b}$ and $t\bar{t}ZZb\bar{b}$ productions. Due to the four b -jets in the signal $t\bar{t}hh$ coming from two Higgs bosons, the most efficient cut is the reconstruction of the Higgs mass through selecting the Higgs decay jets by minimizing, $\chi_{HH}^2 = \frac{(m_{b_i, b_j} - m_h)^2}{\Delta_h^2} + \frac{(m_{b_k, b_l} - m_h)^2}{\Delta_h^2}$, where $i \neq j \neq k \neq l$ run over all b -tagged jets and $\Delta_h = 20$ GeV. To remove the W +jets background, at least one top quark is required to be reconstructed by using $\chi_{t_l}^2 = \frac{(m_{b_i, l, \cancel{E}_T} - m_t)^2}{\Delta_t^2}$ for semi-leptonic and $\chi_{t_h}^2 = \frac{(m_{j_i, j_j, j_k} - m_t)^2}{\Delta_t^2}$ for hadronic top decays, respectively. Since the Higgs data has tightly constrained the CP-violating $t\bar{t}h$ interactions (see figure 2), the allowed values of c_t and $\tilde{c}_t$ can not sizably affects the polarization states of the top quark and will not change the distributions of the final states. Therefore, we can expect that the cuts efficiency obtained in the ref. [84] is applicable to our case, in particular for pure scalar $t\bar{t}h$ interaction. In this case, we can estimate the LHC sensitivity of our signals by normalizing the results in ref. [84]. For example, when $c_t = 1.12$ and $\tilde{c}_t = \pm 0.23$ (at 95% C.L. Higgs data allowed region), $\sigma^{pp \rightarrow t\bar{t}hh} / \sigma_{\text{SM}}^{pp \rightarrow t\bar{t}hh} = 2$ and $S/B = 30\%$. The corresponding statistical significance can reach 3σ at 14 TeV LHC with the luminosity $\mathcal{L} = 3000 \text{ fb}^{-1}$. So we may expect that the enhancement of $t\bar{t}hh$ arising from the CP-violating $t\bar{t}h$ interactions may be observed at the HL-LHC.

3.2 e^+e^- and $\gamma\gamma$ collisions

At a high energy e^+e^- collider, $e^+e^- \rightarrow t\bar{t}hh$ process can occur only through the s -channel. The corresponding Feynman diagrams for $e^+e^- \rightarrow t\bar{t}hh$ are shown in figure 5. Different from the process $pp \rightarrow t\bar{t}hh$, the di-Higgs bosons can be produced not only through the top-Higgs or Higgs self interaction but also by the Higgs gauge couplings hZZ or $hhZZ$. Since we assume that the Higgs gauge couplings and Higgs self-couplings be the SM values, any modifications in $t\bar{t}h$ coupling can change the interference behavior between the amplitudes with $\alpha_s y_t^2$ and with $\alpha_s y_t \lambda_{3h}$.

In figure 6, we show the ratios of $\sigma^{e^+e^- \rightarrow t\bar{t}hh} / \sigma_{\text{SM}}^{e^+e^- \rightarrow t\bar{t}hh}$ in the plane of $\tilde{c}_t - c_t$ at a e^+e^- collider with $\sqrt{s} = 1$ TeV. From figure 6, we can see that the cross section of $e^+e^- \rightarrow t\bar{t}hh$ monotonically increases with c_t when $\tilde{c}_t = 0$ and can maximally reach 4 times as large as the SM prediction. For a given c_t , the pseudo-scalar component in $t\bar{t}h$ can reduce or enhance the cross section of $e^+e^- \rightarrow t\bar{t}hh$. Although some of the subprocesses of $e^+e^- \rightarrow t\bar{t}hh$ that can be triggered by the Higgs gauge interactions, the cross section of $e^+e^- \rightarrow t\bar{t}hh$ is still dominated by top-Higgs coupling. Within the 68% (95%) C.L. Higgs data allowed region, we note that the value of the ratio $\sigma^{e^+e^- \rightarrow t\bar{t}hh} / \sigma_{\text{SM}}^{e^+e^- \rightarrow t\bar{t}hh}$ can maximally reach about 1.5 (2.1) at 1 TeV e^+e^- collider.

At a photon collider, the $\gamma\gamma$ collisions can be also achieved by the inverse Compton scattering of the incident electron- and the laser-beam, the number of events can be calculated by convoluting the cross section of $\gamma\gamma$ collision with the photon beam luminosity

Figure 5. Feynman diagrams for the process $e^+e^- \rightarrow t\bar{t}hh$.

Figure 6. Same as figure 2, but ratios of $\sigma^{e^+e^- \rightarrow t\bar{t}hh} / \sigma_{\text{SM}}^{e^+e^- \rightarrow t\bar{t}hh}$ at a e^+e^- collider with $\sqrt{s} = 1 \text{ TeV}$.

Figure 7. Feynman diagrams for the process $\gamma\gamma \rightarrow t\bar{t}hh$.

distribution:

$$N_{\gamma\gamma \rightarrow hh} = \int d\sqrt{s_{\gamma\gamma}} \frac{d\mathcal{L}_{\gamma\gamma}}{d\sqrt{s_{\gamma\gamma}}} \hat{\sigma}_{\gamma\gamma \rightarrow hh}(s_{\gamma\gamma}) \equiv \mathcal{L}_{e^+e^-} \sigma_{\gamma\gamma \rightarrow hh}(s) \quad (3.2)$$

where $d\mathcal{L}_{\gamma\gamma}/d\sqrt{s_{\gamma\gamma}}$ is the photon-beam luminosity distribution. $\sigma_{\gamma\gamma \rightarrow hh}(s)$ (s is the squared center-of-mass energy of e^+e^- collision) is defined as the effective cross section of $\gamma\gamma \rightarrow hh$, which can be written as [110, 111]

$$\sigma_{\gamma\gamma \rightarrow hh}(s) = \int_{\sqrt{a}}^{x_{\max}} 2z dz \hat{\sigma}_{\gamma\gamma \rightarrow hh}(s_{\gamma\gamma} = z^2 s) \int_{z^2/x_{\max}}^{x_{\max}} \frac{dx}{x} F_{\gamma/e}(x) F_{\gamma/e}\left(\frac{z^2}{x}\right). \quad (3.3)$$

Here $F_{\gamma/e}$ denotes the energy spectrum of the back-scattered photon for the unpolarized initial electron and laser photon beams, which is given by

$$F_{\gamma/e}(x) = \frac{1}{D(\xi)} \left[1 - x + \frac{1}{1-x} - \frac{4x}{\xi(1-x)} + \frac{4x^2}{\xi^2(1-x)^2} \right] \quad (3.4)$$

with

$$D(\xi) = \left(1 - \frac{4}{\xi} - \frac{8}{\xi^2} \right) \ln(1+\xi) + \frac{1}{2} + \frac{8}{\xi} - \frac{1}{2(1+\xi)^2}. \quad (3.5)$$

Figure 8. Same as figure 2, but ratios of $\sigma^{\gamma\gamma\rightarrow t\bar{t}hh}/\sigma_{\text{SM}}^{\gamma\gamma\rightarrow t\bar{t}hh}$ at a photon collider with $\sqrt{s} = 1$ TeV.

e^+e^-	$t\bar{t}hh$	$t\bar{t}h\bar{b}\bar{b}$	$t\bar{t}b\bar{b}b\bar{b}$	$t\bar{t}Zb\bar{b}$
σ (10^{-3} fb)	13.55	6.69	5.52	16.24

Table 1. The SM cross sections of signal $t\bar{t}hh$ and the backgrounds $t\bar{t}h\bar{b}\bar{b}$, $t\bar{t}b\bar{b}b\bar{b}$, $t\bar{t}Zb\bar{b}$ at a e^+e^- collider with $\sqrt{s} = 1$ TeV.

Here $\xi = 4E_e E_0/m_e^2$ (E_e is the incident electron energy, where E_0 is the initial laser photon energy) and $x = E/E_0$ with E being the energy of the scattered photon moving along the initial electron direction. In our calculations, we take the parameters as $\xi = 4.8$, $D(\xi) = 1.83$ and $x_{\text{max}} = 0.83$ [110, 111]. At photon collider, $\gamma\gamma \rightarrow t\bar{t}hh$ process can occur only through the t -channel. The corresponding Feynman diagrams for $\gamma\gamma \rightarrow t\bar{t}hh$ are shown in figure 7.

In figure 8, we present the ratios of $\sigma^{\gamma\gamma\rightarrow t\bar{t}hh}/\sigma_{\text{SM}}^{\gamma\gamma\rightarrow t\bar{t}hh}$ in the plane of $\tilde{c}_t - c_t$ at photon collider with $\sqrt{s} = 1$ TeV. Similar to the process $pp \rightarrow t\bar{t}hh$, the cross section of $\gamma\gamma \rightarrow t\bar{t}hh$ is determined by top-Higgs coupling. From figure 8, we can see that the allowed maximal cross section of $\gamma\gamma \rightarrow t\bar{t}hh$ can be 2 times as large as the SM prediction, which occurs at $c_t = 1.2$ and $\tilde{c}_t = 0$. The pseudo-scalar component in $t\bar{t}h$ always reduce the production rate of $\gamma\gamma \rightarrow t\bar{t}hh$.

In table 1, we present the cross sections of $e^+e^- \rightarrow t\bar{t}hh$ and the backgrounds $e^+e^- \rightarrow t\bar{t}h\bar{b}\bar{b}$, $t\bar{t}b\bar{b}b\bar{b}$, $t\bar{t}Zb\bar{b}$ for $\sqrt{s} = 1$ TeV. It can be seen that the cross section of $t\bar{t}hh$ can reach 13.55×10^{-3} fb in the SM due to the suppression of phase space. Given the enhancement effect in $t\bar{t}hh$ from the CP-violating top-Higgs couplings, if a e^+e^- collider could deliver an integrated luminosity up to 10 ab^{-1} , there will be about hundreds of $t\bar{t}hh$ events, which may be potentially used to test the anomalous top-Higgs couplings. However, the exact detector configurations are not finalized. Therefore, the background estimation and

signal extraction strategies would be largely dependent on the detector designs and trigger conditions [112], which is beyond the scope of our study. Besides, we find that the cross section of $\gamma\gamma \rightarrow t\bar{t}hh$ can only reach 3.74×10^{-4} fb in the SM for $\sqrt{s} = 1$ TeV and is hard to be detected at a photon collider.

4 Conclusion

As a complementary to the di-Higgs production, $t\bar{t}hh$ process may open a new avenue to measure di-Higgs physics at the LHC and a future 100 TeV pp collider or a high energy e^+e^- collider. In this paper, we studied the impact of CP-violating top-Higgs coupling on $t\bar{t}hh$ production at the LHC, a e^+e^- and a 100 TeV hadron collider under the current Higgs data. Within the allowed parameter region, we find that the cross section of di-Higgs production at the LHC-14 TeV, e^+e^- -1 TeV and VHE-LHC/SPPC-100 TeV can be enhanced up to about 2 times the SM predictions since a large deviation of top-Higgs coupling is still allowed. It is expected that the future precise measurement of Higgs coupling will reveal the nature of top-Higgs interaction and improve the sensitivity of the determination of Higgs self-coupling through $t\bar{t}hh$ production.

Acknowledgments

Ning Liu acknowledges Dr Archil Kobakhidze for his warm hospitality in the University of Sydney. This work is supported by the National Natural Science Foundation of China (NNSFC) under grants Nos. 11275057, 11305049 and 11405047, by Specialised Research Fund for the Doctoral Program of Higher Education under Grant No. 20134104120002 and by the Startup Foundation for Doctors of Henan Normal University under contract No.11112, by the China Postdoctoral Science Foundation under Grant No. 2014M561987 and the Joint Funds of the National Natural Science Foundation of China (U1404113).

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] ATLAS collaboration, *Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC*, *Phys. Lett. B* **716** (2012) 1 [[arXiv:1207.7214](https://arxiv.org/abs/1207.7214)] [[INSPIRE](#)].
- [2] CMS collaboration, *Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC*, *Phys. Lett. B* **716** (2012) 30 [[arXiv:1207.7235](https://arxiv.org/abs/1207.7235)] [[INSPIRE](#)].
- [3] ATLAS collaboration, *Measurements of the properties of the Higgs-like boson in the two photon decay channel with the ATLAS detector using 25 fb⁻¹ of proton-proton collision data*, *ATLAS-CONF-2013-012* (2013).
- [4] ATLAS collaboration, *Combined coupling measurements of the Higgs-like boson with the ATLAS detector using up to 25 fb⁻¹ of proton-proton collision data*, *ATLAS-CONF-2013-034* (2013).

- [5] CMS collaboration, *Updated measurements of the Higgs boson at 125 GeV in the two photon decay channel*, [CMS-PAS-HIG-13-001](#).
- [6] CMS collaboration, *Combination of standard model Higgs boson searches and measurements of the properties of the new boson with a mass near 125 GeV*, [CMS-PAS-HIG-13-005](#).
- [7] D.Y. Shao, C.S. Li, H.T. Li and J. Wang, *Threshold resummation effects in Higgs boson pair production at the LHC*, *JHEP* **07** (2013) 169 [[arXiv:1301.1245](#)] [[INSPIRE](#)].
- [8] D. de Florian and J. Mazzitelli, *Higgs Boson Pair Production at Next-to-Next-to-Leading Order in QCD*, *Phys. Rev. Lett.* **111** (2013) 201801 [[arXiv:1309.6594](#)] [[INSPIRE](#)].
- [9] X. Li and M.B. Voloshin, *Remarks on double Higgs boson production by gluon fusion at threshold*, *Phys. Rev. D* **89** (2014) 013012 [[arXiv:1311.5156](#)] [[INSPIRE](#)].
- [10] J. Grigo, K. Melnikov and M. Steinhauser, *Virtual corrections to Higgs boson pair production in the large top quark mass limit*, *Nucl. Phys. B* **888** (2014) 17 [[arXiv:1408.2422](#)] [[INSPIRE](#)].
- [11] R. Frederix et al., *Higgs pair production at the LHC with NLO and parton-shower effects*, *Phys. Lett. B* **732** (2014) 142 [[arXiv:1401.7340](#)] [[INSPIRE](#)].
- [12] L.-S. Ling, R.-Y. Zhang, W.-G. Ma, L. Guo, W.-H. Li and X.-Z. Li, *NNLO QCD corrections to Higgs pair production via vector boson fusion at hadron colliders*, *Phys. Rev. D* **89** (2014) 073001 [[arXiv:1401.7754](#)] [[INSPIRE](#)].
- [13] U. Baur, T. Plehn and D.L. Rainwater, *Determining the Higgs boson selfcoupling at hadron colliders*, *Phys. Rev. D* **67** (2003) 033003 [[hep-ph/0211224](#)] [[INSPIRE](#)].
- [14] U. Baur, T. Plehn and D.L. Rainwater, *Probing the Higgs selfcoupling at hadron colliders using rare decays*, *Phys. Rev. D* **69** (2004) 053004 [[hep-ph/0310056](#)] [[INSPIRE](#)].
- [15] J. Baglio, A. Djouadi, R. Gröber, M.M. Mühlleitner, J. Quevillon and M. Spira, *The measurement of the Higgs self-coupling at the LHC: theoretical status*, *JHEP* **04** (2013) 151 [[arXiv:1212.5581](#)] [[INSPIRE](#)].
- [16] D.E. Ferreira de Lima, A. Papaefstathiou and M. Spannowsky, *Standard model Higgs boson pair production in the $(b\bar{b})(b\bar{b})$ final state*, *JHEP* **08** (2014) 030 [[arXiv:1404.7139](#)] [[INSPIRE](#)].
- [17] D. Wardrope, E. Jansen, N. Konstantinidis, B. Cooper, R. Falla and N. Norjoharuddeen, *Non-resonant Higgs-pair production in the $b\bar{b}b\bar{b}$ final state at the LHC*, *Eur. Phys. J. C* **75** (2015) 219 [[arXiv:1410.2794](#)] [[INSPIRE](#)].
- [18] M.J. Dolan, C. Englert and M. Spannowsky, *Higgs self-coupling measurements at the LHC*, *JHEP* **10** (2012) 112 [[arXiv:1206.5001](#)] [[INSPIRE](#)].
- [19] F. Goertz, A. Papaefstathiou, L.L. Yang and J. Zurita, *Higgs Boson self-coupling measurements using ratios of cross sections*, *JHEP* **06** (2013) 016 [[arXiv:1301.3492](#)] [[INSPIRE](#)].
- [20] F. Goertz, A. Papaefstathiou, L.L. Yang and J. Zurita, *Higgs boson pair production in the $D = 6$ extension of the SM*, *JHEP* **04** (2015) 167 [[arXiv:1410.3471](#)] [[INSPIRE](#)].
- [21] A. Papaefstathiou, L.L. Yang and J. Zurita, *Higgs boson pair production at the LHC in the $b\bar{b}W^+W^-$ channel*, *Phys. Rev. D* **87** (2013) 011301 [[arXiv:1209.1489](#)] [[INSPIRE](#)].
- [22] F. Goertz, A. Papaefstathiou, L.L. Yang and J. Zurita, *Higgs Boson self-coupling measurements using ratios of cross sections*, *JHEP* **06** (2013) 016 [[arXiv:1301.3492](#)] [[INSPIRE](#)].

- [23] A.J. Barr, M.J. Dolan, C. Englert and M. Spannowsky, *Di-Higgs final states augMT2ed — selecting hh events at the high luminosity LHC*, *Phys. Lett. B* **728** (2014) 308 [[arXiv:1309.6318](#)] [[INSPIRE](#)].
- [24] M.J. Dolan, C. Englert, N. Greiner and M. Spannowsky, *Further on up the road: $hhjj$ production at the LHC*, *Phys. Rev. Lett.* **112** (2014) 101802 [[arXiv:1310.1084](#)] [[INSPIRE](#)].
- [25] P. Maierhöfer and A. Papaefstathiou, *Higgs Boson pair production merged to one jet*, *JHEP* **03** (2014) 126 [[arXiv:1401.0007](#)] [[INSPIRE](#)].
- [26] V. Barger, L.L. Everett, C.B. Jackson, A.D. Peterson and G. Shaughnessy, *New physics in resonant production of Higgs boson pairs*, *Phys. Rev. Lett.* **114** (2015) 011801 [[arXiv:1408.0003](#)] [[INSPIRE](#)].
- [27] V. Martín Lozano, J.M. Moreno and C.B. Park, *Resonant Higgs boson pair production in the $hh \rightarrow b\bar{b} WW \rightarrow b\bar{b}\ell^+\nu\ell^-\bar{\nu}$ decay channel*, *JHEP* **08** (2015) 004 [[arXiv:1501.03799](#)] [[INSPIRE](#)].
- [28] Q. Li, Z. Li, Q.-S. Yan and X. Zhao, *Probe Higgs boson pair production via the $3\ell 2j + \cancel{E}$ mode*, *Phys. Rev. D* **92** (2015) 014015 [[arXiv:1503.07611](#)] [[INSPIRE](#)].
- [29] N. Liu, S. Hu, B. Yang and J. Han, *Impact of top-Higgs couplings on Di-Higgs production at future colliders*, *JHEP* **01** (2015) 008 [[arXiv:1408.4191](#)] [[INSPIRE](#)].
- [30] F. Goertz, A. Papaefstathiou, L.L. Yang and J. Zurita, *Higgs boson pair production in the $D = 6$ extension of the SM*, *JHEP* **04** (2015) 167 [[arXiv:1410.3471](#)] [[INSPIRE](#)].
- [31] A. Pierce, J. Thaler and L.-T. Wang, *Disentangling Dimension Six Operators through Di-Higgs Boson Production*, *JHEP* **05** (2007) 070 [[hep-ph/0609049](#)] [[INSPIRE](#)].
- [32] R. Contino, M. Ghezzi, M. Moretti, G. Panico, F. Piccinini and A. Wulzer, *Anomalous Couplings in Double Higgs Production*, *JHEP* **08** (2012) 154 [[arXiv:1205.5444](#)] [[INSPIRE](#)].
- [33] L. Wu, J.M. Yang, C.-P. Yuan and M. Zhang, *Higgs self-coupling in the MSSM and NMSSM after the LHC Run 1*, *Phys. Lett. B* **747** (2015) 378 [[arXiv:1504.06932](#)] [[INSPIRE](#)].
- [34] C. Han, X. Ji, L. Wu, P. Wu and J.M. Yang, *Higgs pair production with SUSY QCD correction: revisited under current experimental constraints*, *JHEP* **04** (2014) 003 [[arXiv:1307.3790](#)] [[INSPIRE](#)].
- [35] K. Nishiwaki, S. Niyogi and A. Shivaji, *$t\bar{t}H$ Anomalous Coupling in Double Higgs Production*, *JHEP* **04** (2014) 011 [[arXiv:1309.6907](#)] [[INSPIRE](#)].
- [36] C.-R. Chen and I. Low, *Double take on new physics in double Higgs boson production*, *Phys. Rev. D* **90** (2014) 013018 [[arXiv:1405.7040](#)] [[INSPIRE](#)].
- [37] G. Bhattacharyya, D. Das and P.B. Pal, *Modified Higgs couplings and unitarity violation*, *Phys. Rev. D* **87** (2013) 011702 [[arXiv:1212.4651](#)] [[INSPIRE](#)].
- [38] ATLAS collaboration, *Search for the Standard Model Higgs boson produced in association with top quarks and decaying to $b\bar{b}$ in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector at the LHC*, *ATLAS-CONF-2014-011* (2014).
- [39] CMS collaboration, *Search for $t\bar{t}H$ production using the Matrix Element Method*, *CMS-PAS-HIG-14-010*.
- [40] W.J. Marciano and F.E. Paige, *Associated production of Higgs bosons with $t\bar{t}$ pairs*, *Phys. Rev. Lett.* **66** (1991) 2433 [[INSPIRE](#)].

- [41] J. Dai, J.F. Gunion and R. Vega, *Using b tagging to detect $t\bar{t}$ Higgs production with Higgs $\rightarrow b\bar{b}$* , *Phys. Rev. Lett.* **71** (1993) 2699 [[hep-ph/9306271](#)] [[INSPIRE](#)].
- [42] J. Goldstein, C.S. Hill, J. Incandela, S.J. Parke, D.L. Rainwater and D. Stuart, *$p\bar{p} \rightarrow t\bar{t}H$: A discovery mode for the Higgs boson at the Tevatron*, *Phys. Rev. Lett.* **86** (2001) 1694 [[hep-ph/0006311](#)] [[INSPIRE](#)].
- [43] A. Belyaev and L. Reina, *$pp \rightarrow t\bar{t}H$, $H \rightarrow \tau^+\tau^-$: Toward a model independent determination of the Higgs boson couplings at the LHC*, *JHEP* **08** (2002) 041 [[hep-ph/0205270](#)] [[INSPIRE](#)].
- [44] F. Maltoni, D.L. Rainwater and S. Willenbrock, *Measuring the top quark Yukawa coupling at hadron colliders via $t\bar{t}H$, $H \rightarrow W^+W^-$* , *Phys. Rev. D* **66** (2002) 034022 [[hep-ph/0202205](#)] [[INSPIRE](#)].
- [45] V. Drollinger, T. Müller and D. Denegri, *Searching for Higgs bosons in association with top quark pairs in the $H0 \rightarrow b\bar{b}$ decay mode*, [hep-ph/0111312](#) [[INSPIRE](#)].
- [46] W. Beenakker, S. Dittmaier, M. Krämer, B. Plumper, M. Spira and P.M. Zerwas, *NLO QCD corrections to $t\bar{t}H$ production in hadron collisions*, *Nucl. Phys. B* **653** (2003) 151 [[hep-ph/0211352](#)] [[INSPIRE](#)].
- [47] P. Agrawal, S. Bandyopadhyay and S.P. Das, *Multilepton Signatures of the Higgs Boson through its Production in Association with a Top-quark Pair*, *Phys. Rev. D* **88** (2013) 093008 [[arXiv:1308.3043](#)] [[INSPIRE](#)].
- [48] S. Biswas, R. Frederix, E. Gabrielli and B. Mele, *Enhancing the $t\bar{t}H$ signal through top-quark spin polarization effects at the LHC*, *JHEP* **07** (2014) 020 [[arXiv:1403.1790](#)] [[INSPIRE](#)].
- [49] M.V. Garzelli, A. Kardos, C.G. Papadopoulos and Z. Trócsányi, *Standard Model Higgs boson production in association with a top anti-top pair at NLO with parton showering*, *Europhys. Lett.* **96** (2011) 11001 [[arXiv:1108.0387](#)] [[INSPIRE](#)].
- [50] J. Cao, Z. Heng, L. Wu and J.M. Yang, *Top quark forward-backward asymmetry at the Tevatron: A comparative study in different new physics models*, *Phys. Rev. D* **81** (2010) 014016 [[arXiv:0912.1447](#)] [[INSPIRE](#)].
- [51] R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, R. Pittau and P. Torrielli, *Scalar and pseudoscalar Higgs production in association with a top-antitop pair*, *Phys. Lett. B* **701** (2011) 427 [[arXiv:1104.5613](#)] [[INSPIRE](#)].
- [52] J. Cao, K. Hikasa, L. Wang, L. Wu and J.M. Yang, *Testing new physics models by top charge asymmetry and polarization at the LHC*, *Phys. Rev. D* **85** (2012) 014025 [[arXiv:1109.6543](#)] [[INSPIRE](#)].
- [53] C. Degrande, J.M. Gerard, C. Grojean, F. Maltoni and G. Servant, *Probing Top-Higgs Non-Standard Interactions at the LHC*, *JHEP* **07** (2012) 036 [Erratum *ibid.* **1303** (2013) 032] [[arXiv:1205.1065](#)] [[INSPIRE](#)].
- [54] J. Cao, L. Wang, L. Wu and J.M. Yang, *Top quark forward-backward asymmetry, FCNC decays and like-sign pair production as a joint probe of new physics*, *Phys. Rev. D* **84** (2011) 074001 [[arXiv:1101.4456](#)] [[INSPIRE](#)].
- [55] N. Liu, L. Wu, P.W. Wu and J.M. Yang, *Complete one-loop effects of SUSY QCD in $b\bar{b}h$ production at the LHC under current experimental constraints*, *JHEP* **01** (2013) 161 [[arXiv:1208.3413](#)] [[INSPIRE](#)].

- [56] D. Curtin, J. Galloway and J.G. Wacker, *Measuring the $t\bar{t}h$ coupling from same-sign dilepton $+2b$ measurements*, *Phys. Rev. D* **88** (2013) 093006 [[arXiv:1306.5695](#)] [[INSPIRE](#)].
- [57] J. Adelman, A. Loginov, P. Tipton and J. Vasquez, *Study of $t\bar{t}H$ ($H \rightarrow \mu^+\mu^-$) in the three lepton channel at $\sqrt{s} = 14$ TeV; A Snowmass white paper*, [arXiv:1310.1132](#).
- [58] F. Boudjema, R.M. Godbole, D. Guadagnoli and K.A. Mohan, *Lab-frame observables for probing the top-Higgs interaction*, *Phys. Rev. D* **92** (2015) 015019 [[arXiv:1501.03157](#)] [[INSPIRE](#)].
- [59] X.-G. He, G.-N. Li and Y.-J. Zheng, *Probing Higgs Boson CP Properties with $t\bar{t}H$ at the LHC and the 100 TeV pp Collider*, [arXiv:1501.00012](#) [[INSPIRE](#)].
- [60] LHC HIGGS CROSS SECTION WORKING GROUP collaboration, J.R. Andersen et al., *Handbook of LHC Higgs Cross sections: 3. Higgs Properties*, [arXiv:1307.1347](#) [[INSPIRE](#)].
- [61] ATLAS collaboration, *Search for $H \rightarrow \gamma\gamma$ produced in association with top quarks and constraints on the top quark-Higgs boson Yukawa coupling using data taken at 7 TeV and 8 TeV with the ATLAS detector*, *ATLAS-CONF-2014-043* (2014).
- [62] CMS collaboration, *Search for the associated production of the Higgs boson with a top-quark pair*, *JHEP* **09** (2014) 087 [Erratum *ibid.* **1410** (2014) 106] [[arXiv:1408.1682](#)] [[INSPIRE](#)].
- [63] G. Bordes and B. van Eijk, *On the associate production of a neutral intermediate mass Higgs boson with a single top quark at the LHC and SSC*, *Phys. Lett. B* **299** (1993) 315 [[INSPIRE](#)].
- [64] A. Ballestrero and E. Maina, *$t\bar{b}H$ production for an intermediate mass Higgs*, *Phys. Lett. B* **299** (1993) 312 [[INSPIRE](#)].
- [65] W.J. Stirling and D.J. Summers, *Production of an intermediate mass Higgs boson in association with a single top quark at LHC and SSC*, *Phys. Lett. B* **283** (1992) 411 [[INSPIRE](#)].
- [66] J.L. Diaz-Cruz and O.A. Sampayo, *Associated production of the Higgs boson with $t\bar{b}$ at hadron colliders*, *Phys. Lett. B* **276** (1992) 211 [[INSPIRE](#)].
- [67] T.M.P. Tait and C.P. Yuan, *Single top quark production as a window to physics beyond the standard model*, *Phys. Rev. D* **63** (2000) 014018 [[hep-ph/0007298](#)] [[INSPIRE](#)].
- [68] F. Maltoni, K. Paul, T. Stelzer and S. Willenbrock, *Associated production of Higgs and single top at hadron colliders*, *Phys. Rev. D* **64** (2001) 094023 [[hep-ph/0106293](#)] [[INSPIRE](#)].
- [69] J. Cao, L. Wu and J.M. Yang, *New physics effects on top quark spin correlation and polarization at the LHC: a comparative study in different models*, *Phys. Rev. D* **83** (2011) 034024 [[arXiv:1011.5564](#)] [[INSPIRE](#)].
- [70] G. R. Lu and L. Wu, *Flavor-changing neutral-current production of t anti- c in association with a neutral top-Higgs at LHC in the topcolor-assisted technicolor model*, *Chin. Phys. Lett.* **27** (2010) 031401 [[INSPIRE](#)].
- [71] P. Agrawal, S. Mitra and A. Shivaji, *Effect of Anomalous Couplings on the Associated Production of a Single Top Quark and a Higgs Boson at the LHC*, *JHEP* **12** (2013) 077 [[arXiv:1211.4362](#)] [[INSPIRE](#)].
- [72] C. Han, N. Liu, L. Wu and J.M. Yang, *Probing topcolor-assisted technicolor from top charge asymmetry and triple-top production at the LHC*, *Phys. Lett. B* **714** (2012) 295 [[arXiv:1203.2321](#)] [[INSPIRE](#)].

- [73] Y. Wang, F.P. Huang, C.S. Li, B.H. Li, D.Y. Shao and J. Wang, *Constraints on flavor-changing neutral-current Htq couplings from the signal of tH associated production with QCD next-to-leading order accuracy at the LHC*, *Phys. Rev. D* **86** (2012) 094014 [[arXiv:1208.2902](#)] [[INSPIRE](#)].
- [74] M. Farina, C. Grojean, F. Maltoni, E. Salvioni and A. Thamm, *Lifting degeneracies in Higgs couplings using single top production in association with a Higgs boson*, *JHEP* **05** (2013) 022 [[arXiv:1211.3736](#)] [[INSPIRE](#)].
- [75] S. Biswas, E. Gabrielli and B. Mele, *Single top and Higgs associated production as a probe of the Htt coupling sign at the LHC*, *JHEP* **01** (2013) 088 [[arXiv:1211.0499](#)] [[INSPIRE](#)].
- [76] J. Ellis, D.S. Hwang, K. Sakurai and M. Takeuchi, *Disentangling Higgs-Top Couplings in Associated Production*, *JHEP* **04** (2014) 004 [[arXiv:1312.5736](#)] [[INSPIRE](#)].
- [77] C. Englert and E. Re, *Bounding the top Yukawa coupling with Higgs-associated single-top production*, *Phys. Rev. D* **89** (2014) 073020 [[arXiv:1402.0445](#)] [[INSPIRE](#)].
- [78] J. Chang, K. Cheung, J.S. Lee and C.-T. Lu, *Probing the Top-Yukawa Coupling in Associated Higgs production with a Single Top Quark*, *JHEP* **05** (2014) 062 [[arXiv:1403.2053](#)] [[INSPIRE](#)].
- [79] A. Kobakhidze, L. Wu and J. Yue, *Anomalous Top-Higgs Couplings and Top Polarisation in Single Top and Higgs Associated Production at the LHC*, *JHEP* **10** (2014) 100 [[arXiv:1406.1961](#)] [[INSPIRE](#)].
- [80] L. Wu, *Enhancing thj Production from Top-Higgs FCNC Couplings*, *JHEP* **02** (2015) 061 [[arXiv:1407.6113](#)] [[INSPIRE](#)].
- [81] J. Yue, *Enhanced thj signal at the LHC with $h \rightarrow \gamma\gamma$ decay and $\mathcal{CP}$ -violating top-Higgs coupling*, *Phys. Lett. B* **744** (2015) 131 [[arXiv:1410.2701](#)] [[INSPIRE](#)].
- [82] B. Yang, J. Han and N. Liu, *Associated production of single top and Higgs at the LHC in the lightest Higgs model with T -parity*, *JHEP* **04** (2015) 148 [[arXiv:1412.2927](#)] [[INSPIRE](#)].
- [83] CMS collaboration, *Search for associated production of a single top quark and a Higgs boson in events where the Higgs boson decays to two photons at $\sqrt{s} = 8$ TeV*, [CMS-PAS-HIG-14-001](#).
- [84] C. Englert, F. Krauss, M. Spannowsky and J. Thompson, *Di-Higgs phenomenology in $t\bar{t}hh$: The forgotten channel*, *Phys. Lett. B* **743** (2015) 93 [[arXiv:1409.8074](#)] [[INSPIRE](#)].
- [85] T. Liu and H. Zhang, *Measuring Di-Higgs Physics via the $t\bar{t}hh \rightarrow t\bar{t}b\bar{b}b\bar{b}$ Channel*, [arXiv:1410.1855](#) [[INSPIRE](#)].
- [86] D.M. Asner et al., *ILC Higgs White Paper*, [arXiv:1310.0763](#).
- [87] N. Craig, M. Farina, M. McCullough and M. Perelstein, *Precision Higgsstrahlung as a Probe of New Physics*, *JHEP* **03** (2015) 146 [[arXiv:1411.0676](#)] [[INSPIRE](#)].
- [88] J. Cao, C. Han, J. Ren, L. Wu, J.M. Yang and Y. Zhang, *SUSY effects in Higgs productions at high energy e^+e^- colliders*, [arXiv:1410.1018](#) [[INSPIRE](#)].
- [89] V. Barger, L.L. Everett, H.E. Logan and G. Shaughnessy, *Scrutinizing the 125 GeV Higgs boson in two Higgs doublet models at the LHC, ILC and Muon Collider*, *Phys. Rev. D* **88** (2013) 115003 [[arXiv:1308.0052](#)] [[INSPIRE](#)].
- [90] N. Liu, J. Ren, L. Wu, P. Wu and J.M. Yang, *Full one-loop electroweak corrections to $e^+e^- \rightarrow ZH\gamma$ at a Higgs factory*, *JHEP* **04** (2014) 189 [[arXiv:1311.6971](#)] [[INSPIRE](#)].

- [91] Z. Heng, L. Shang and P. Wan, *Pair production of a 125 GeV Higgs boson in MSSM and NMSSM at the ILC*, *JHEP* **10** (2013) 047 [[arXiv:1306.0279](#)] [[INSPIRE](#)].
- [92] S.L. Hu, N. Liu, J. Ren and L. Wu, *Revisiting Associated Production of 125 GeV Higgs Boson with a Photon at a Higgs Factory*, *J. Phys. G* **41** (2014) 125004 [[arXiv:1402.3050](#)] [[INSPIRE](#)].
- [93] L. Wang and X.-F. Han, *Study of the heavy CP-even Higgs with mass 125 GeV in two-Higgs-doublet models at the LHC and ILC*, *JHEP* **11** (2014) 085 [[arXiv:1404.7437](#)] [[INSPIRE](#)].
- [94] C. Han, A. Kobakhidze, N. Liu, L. Wu and B. Yang, *Constraining Top partner and Naturalness at the LHC and TLEP*, *Nucl. Phys. B* **890** (2014) 388 [[arXiv:1405.1498](#)] [[INSPIRE](#)].
- [95] J. Cao, Z. Heng, D. Li, L. Shang and P. Wu, *Higgs-strahlung production process $e^+e^- \rightarrow Zh$ at the future Higgs factory in the Minimal Dilaton Model*, *JHEP* **08** (2014) 138 [[arXiv:1405.4489](#)] [[INSPIRE](#)].
- [96] B. Yang, Z. Liu, N. Liu and J. Han, *Double Higgs production in the littlest Higgs Model with T-parity at high energy e^+e^- Colliders*, *Eur. Phys. J. C* **74** (2014) 3203 [[arXiv:1408.4295](#)] [[INSPIRE](#)].
- [97] I.F. Ginzburg, G.L. Kotkin, V.G. Serbo and V.I. Telnov, *Production of High-Energy Colliding gamma gamma and gamma e Beams with a High Luminosity at Vlepp Accelerators*, *JETP Lett.* **34** (1981) 491 [[INSPIRE](#)].
- [98] A. Gutierrez-Rodriguez, M.A. Hernandez-Ruiz, O.A. Sampayo, A. Chubykalo and A. Espinoza-Garrido, *The Triple Higgs Boson Self-Coupling at Future Linear e^+e^- Colliders Energies: ILC and CLIC*, *J. Phys. Soc. Jap.* **77** (2008) 094101 [[arXiv:0807.0663](#)] [[INSPIRE](#)].
- [99] A. Gutierrez-Rodriguez, J. Peressutti and O.A. Sampayo, *Higgs Boson Self-Coupling at High Energy $\gamma\gamma$ Collider*, *J. Phys. G* **38** (2011) 095002 [[arXiv:1107.0245](#)] [[INSPIRE](#)].
- [100] J. Brod, U. Haisch and J. Zupan, *Constraints on CP-violating Higgs couplings to the third generation*, *JHEP* **11** (2013) 180 [[arXiv:1310.1385](#)] [[INSPIRE](#)].
- [101] P. Bechtle, S. Heinemeyer, O. Stål, T. Stefaniak and G. Weiglein, *HiggsSignals: Confronting arbitrary Higgs sectors with measurements at the Tevatron and the LHC*, *Eur. Phys. J. C* **74** (2014) 2711 [[arXiv:1305.1933](#)] [[INSPIRE](#)].
- [102] P. Bechtle, O. Brein, S. Heinemeyer, G. Weiglein and K.E. Williams, *HiggsBounds: Confronting Arbitrary Higgs Sectors with Exclusion Bounds from LEP and the Tevatron*, *Comput. Phys. Commun.* **181** (2010) 138 [[arXiv:0811.4169](#)] [[INSPIRE](#)].
- [103] D. Atwood, S. Bar-Shalom, G. Eilam and A. Soni, *CP violation in top physics*, *Phys. Rept.* **347** (2001) 1 [[hep-ph/0006032](#)] [[INSPIRE](#)].
- [104] A. Kobakhidze, *Standard Model with a distorted Higgs sector and the enhanced Higgs diphoton decay rate*, [arXiv:1208.5180](#) [[INSPIRE](#)].
- [105] PARTICLE DATA GROUP collaboration, J. Beringer et al., *Review of Particle Physics (RPP)*, *Phys. Rev. D* **86** (2012) 010001 [[INSPIRE](#)].
- [106] H.-L. Lai et al., *New parton distributions for collider physics*, *Phys. Rev. D* **82** (2010) 074024 [[arXiv:1007.2241](#)] [[INSPIRE](#)].

- [107] A. Belyaev, N.D. Christensen and A. Pukhov, *CalcHEP 3.4 for collider physics within and beyond the Standard Model*, *Comput. Phys. Commun.* **184** (2013) 1729 [[arXiv:1207.6082](#)] [[INSPIRE](#)].
- [108] A. Alloul, N.D. Christensen, C. Degrande, C. Duhr and B. Fuks, *FeynRules 2.0 — A complete toolbox for tree-level phenomenology*, *Comput. Phys. Commun.* **185** (2014) 2250 [[arXiv:1310.1921](#)] [[INSPIRE](#)].
- [109] J. Alwall, M. Herquet, F. Maltoni, O. Mattelaer and T. Stelzer, *MadGraph 5: Going Beyond*, *JHEP* **06** (2011) 128 [[arXiv:1106.0522](#)] [[INSPIRE](#)].
- [110] I.F. Ginzburg et al., *Colliding γe and $\gamma\gamma$ beams based on single-pass e^+e^- accelerators II. Polarization effects, monochromatization improvement*, *Nucl. Instrum. Meth.* **219** (1984) 5.
- [111] V.I. Telnov, *Problems in obtaining $\gamma\gamma$ and γe colliding beams at linear colliders*, *Nucl. Instrum. Meth.* **294** (1990) 72.
- [112] S.T. Boogert and D.J. Miller, *Questions about the measurement of the e^+e^- luminosity spectrum at a future linear collider*, [hep-ex/0211021](#).