

Generalised holonomies and $K(E_9)$

Axel Kleinschmidt^{a,b} and Hermann Nicolai^a

^aMax-Planck-Institut für Gravitationsphysik (Albert-Einstein-Institut),
Am Mühlenberg 1, DE-14476 Potsdam, Germany

^bInternational Solvay Institutes,
ULB-Campus Plaine CP231, BE-1050 Brussels, Belgium

E-mail: axel.kleinschmidt@aei.mpg.de, hermann.nicolai@aei.mpg.de

ABSTRACT: The involutory subalgebra $K(\mathfrak{e}_9)$ of the affine Kac-Moody algebra \mathfrak{e}_9 was recently shown to admit an infinite sequence of unfaithful representations of ever increasing dimensions [1]. We revisit these representations and describe their associated ideals in more detail, with particular emphasis on two chiral versions that can be constructed for each such representation. For every such unfaithful representation we show that the action of $K(\mathfrak{e}_9)$ decomposes into a direct sum of two mutually commuting (‘chiral’ and ‘anti-chiral’) parabolic algebras with Levi subalgebra $\mathfrak{so}(16)_+ \oplus \mathfrak{so}(16)_-$. We also spell out the consistency conditions for uplifting such representations to unfaithful representations of $K(\mathfrak{e}_{10})$. From these results it is evident that the holonomy groups so far discussed in the literature are mere shadows (in a Platonic sense) of a much larger structure.

KEYWORDS: Extended Supersymmetry, M-Theory, Supergravity Models, Field Theories in Higher Dimensions

ARXIV EPRINT: [2107.02445](https://arxiv.org/abs/2107.02445)

Contents

1	Introduction	1
2	Parabolic and filtered algebras	5
2.1	$K(E_9)$ from E_9	5
2.2	Filtered and parabolic algebras	6
2.3	Representations from truncations	8
3	Ideals	9
3.1	Chiral ideals	10
3.2	Commuting chiral quotients	11
4	Examples and relation to root basis form	13
4.1	Spin- $\frac{1}{2}$	14
4.2	Spin- $\frac{3}{2}$	14
4.3	Spin- $\frac{5}{2}$	16
4.4	Spin- $\frac{7}{2}$	17
5	Extension to $K(\mathfrak{e}_{10})$	17

1 Introduction

In studies of supersymmetric solutions of supergravity, a central role is played by the Killing spinor equation that expresses the vanishing of the supersymmetry variation of the gravitino

$$\delta_\epsilon \Psi_M = \widehat{D}_M(\omega, F)\epsilon \equiv \left[\partial_M + \frac{1}{4} \omega_{MAB} \Gamma^{AB} + F \cdot \Gamma \right] \epsilon = 0. \quad (1.1)$$

Here, ϵ represents the spin- $\frac{1}{2}$ local supersymmetry parameter, ω_{MAB} the components of the spin connection and $F \cdot \Gamma$ is shorthand for flux contributions, where the components of the field strengths are contracted with elements of the gamma matrix algebra; the explicit form of this term depends on the theory in question. For maximal supergravity, the idea that the combination of the spin connection and the flux terms should be interpreted as a generalised connection goes back to [2, 3]. More recently, and in a more general context, it was pointed out that these flux terms extend the notion of holonomy of the spin bundle to that of a generalised holonomy [4–7]. The latter has been a subject of intense study in connection with classifying supersymmetric solutions, also in the context of (exceptional) generalised geometry [8–12]. In particular, the groups

$$\text{SO}(16)_+ \times \text{SO}(16)_- \subset \text{SO}(32) \subset \text{SL}(32, \mathbb{R}) \quad (1.2)$$

have been put forward as candidate generalised holonomy groups of maximal supergravity in [5, 6, 13], and we shall focus on this theory in the following. The basis for this conjecture and the chain of embeddings in (1.2) is that the $D = 11$ gamma matrices appearing in (1.1) generate the whole Lie algebra of $\mathfrak{sl}(32)$ upon commutation (together with its subalgebras $\mathfrak{so}(32)$ and $\mathfrak{so}(16) \oplus \mathfrak{so}(16)$), such that the 32-component parameter ϵ can be viewed as belonging to the fundamental representation of this group.

However, it was already pointed out in [14] that these conjectures are problematic at the group level in the sense that they do not contain the correct R-symmetry groups appearing upon dimensional reduction, notably the $\text{Spin}(16)$ symmetry in $D = 3$ maximal supergravity. This incompatibility of the group structures negates the idea that the proposed generalised holonomy groups could be true symmetry groups of $D = 11$ supergravity, although their properties continue to be useful for the study of supersymmetric solutions. A further argument against $\text{SO}(32)$ or $\text{SL}(32)$ as symmetries is that, while these groups can act on the $\text{spin-}\frac{1}{2}$ supersymmetry parameter ϵ , they cannot act on the propagating fermion of the theory, namely the gravitino: for maximal supergravity, the latter is a vector spinor (' $\text{spin-}\frac{3}{2}$ ') representation of the Lorentz group $\text{Spin}(1, 10)$, but there is no way to turn it into a representation of $\text{SL}(32)$.

A resolution of these issues was proposed in the context of studies of Kac-Moody symmetries of supergravity. It was found in [15–18] that the involutory subalgebra $K(\mathfrak{e}_{10})$ of the Kac-Moody Lie algebra \mathfrak{e}_{10} admits a 32-dimensional representation, dubbed $\text{spin-}\frac{1}{2}$ for obvious reasons, as well as a 320-component representation that corresponds to the $\text{spin-}\frac{3}{2}$ gravitino. These representations have the property that they lift correctly to the spin cover $\widetilde{K}(E_{10})$ of the corresponding group (see [19] for a discussion of the spin cover). Moreover, the representations have the property that they possess the correct branching to the other maximal supergravity theories, including the chiral fermions of type IIB, which cannot be obtained by usual dimensional reduction [20], and thus provide a common origin for both IIA and IIB fermions. Since the fermionic representations are finite-dimensional representations of an infinite-dimensional group, they are unfaithful and there is thus a huge kernel of the representation map. The corresponding quotient groups turn out to be $\text{SO}(32)$ for the $\text{spin-}\frac{1}{2}$ representation and the non-compact $\text{SO}(288, 32)$ for the $\text{spin-}\frac{3}{2}$ representation [21].¹ The significance of these results is that they represent incontrovertible evidence for the fact that (1.2) are not the actual generalised holonomy groups, but that the chain of embeddings (1.2) must be replaced by [23]

$$\widetilde{K}(E_9) \subset \widetilde{K}(E_{10}) \subset \widetilde{K}(E_{11}). \tag{1.3}$$

Accordingly, the finite-dimensional symmetry groups which have appeared so far must be interpreted as quotient groups which are obtained by dividing $\widetilde{K}(E_n)$ (for $n = 9, 10, 11$) by the annihilator ideals of the corresponding unfaithful spinor representations. Hence our claim that the holonomy groups identified so far are merely shadows of the full sequence

¹For the $\text{spin-}\frac{1}{2}$ and $\text{spin-}\frac{3}{2}$ representations of $K(\mathfrak{e}_{11})$ defined in [20] the corresponding quotient algebras are $\mathfrak{sl}(32)$ and $\mathfrak{sl}(352)$, respectively. See also [13] and [22] for work related to the $\text{spin-}\frac{1}{2}$ and $\text{spin-}\frac{3}{2}$ representations of $K(\mathfrak{e}_{11})$.

expressed by (1.3) (and their analogs for non-maximal supergravities). Since the actual group is $\widetilde{K}(E_n)$, the topological obstructions found in [14] disappear, as it is no longer necessary to embed the R-symmetry Spin(16) into SL(32) but it embeds correctly (as a quotient) into $\widetilde{K}(E_n)$. In this sense, the proposed Kac-Moody symmetries are much more promising candidates, see also [24] for a related discussion.

The purpose of this paper is to amplify this point, with special emphasis on the affine case, and in this way actually *prove* part of the claim (1.3). Since the finite-dimensional spin- $\frac{1}{2}$ and spin- $\frac{3}{2}$ representations furnish only infinitesimal glimpses of the huge Kac-Moody-type symmetry, it is important to understand the structure of its representations better. First steps were taken for $K(\mathfrak{e}_{10})$ by the construction of two new ‘higher spin’ representations in [25, 26] (see also [21] for a review), where the ‘spin’ actually refers to DeWitt superspace. In contrast to the spin- $\frac{1}{2}$ and spin- $\frac{3}{2}$ representations these can no longer be deduced from supergravity. A more systematic study in the case of affine \mathfrak{e}_9 , relevant in the context of $D = 2$ maximal supergravity [23, 27–29], was initiated in [1] where a general construction of infinite families of unfaithful representations was given. This construction, reviewed in section 2 below, hinges on some properties that are specific to the affine case, but it is hoped that this serves as a stepping stone to the more interesting hyperbolic algebra \mathfrak{e}_{10} . This hope stems from the fact that all known representations of $K(\mathfrak{e}_{10})$ split into two chiral halves when decomposed under $K(\mathfrak{e}_9) \subset K(\mathfrak{e}_{10})$ and the resulting $K(\mathfrak{e}_9)$ representations belong to the family of representations constructed in [1]. We stress that the converse is not true: not every $K(\mathfrak{e}_9)$ -representation can be ‘doubled’ and uplifted to a $K(\mathfrak{e}_{10})$ representation. It is one of the aims of this paper to find conditions when this is possible.

As one of our main results we will exhibit for the affine case the general structure of the quotient group Q of $\widetilde{K}(E_9)$ as a product of two mutually commuting (‘chiral’ and ‘anti-chiral’) parabolic groups (see (3.20) for the Lie algebra version of this statement)

$$Q = (\text{Spin}(16)_+ \ltimes U_+) \times (\text{Spin}(16)_- \ltimes U_-), \tag{1.4}$$

where the Levi subgroups of the two parabolic groups are $\text{Spin}(16)_\pm$ and the unipotent subgroups U_\pm are N -step unipotent, with N referring to a truncation condition in the construction of the unfaithful representations. This is the infinite series of groups that generalises the left-most entry in (1.2). The notion of chirality \pm in the construction is related mathematically to choosing one of two fixed points of an involution but in terms of $D = 2$ supergravity it corresponds to space-time chirality as we explain in more detail in section 5. We also stress that depending on details of the construction discussed in section 3, some of the finite \mathbb{Z}_2 factors in the center of Spin(16) can act trivially. From (1.4) we see that the proper generalisation of the holonomy group to non-trivial spinor representations therefore differs substantially from (1.2). In particular there is no limit on the size of the group Q , since the ‘cutoff parameter’ N can be taken arbitrarily large. For maximal supergravity the structure of (1.4) is in complete accord with the known action of $\widetilde{K}(E_9)$ on the supergravity fermions that was first exhibited in [23].

Our motivation for this detailed study of representations of $K(\mathfrak{e}_9)$ is twofold. Firstly, we can use the representations to obtain an ever more faithful description of the non-

Kac-Moody structure $K(\mathfrak{e}_9)$ by increasing the value of N . Secondly, the proposal of E_{10} symmetry in M-theory [30] is lacking a proper supersymmetric extension [31], a fact that can be possibly traced back to the imbalance between the faithful bosonic representation of \mathfrak{e}_{10} on the symmetric space $E_{10}/K(E_{10})$ and the unfaithful action of $K(\mathfrak{e}_{10})$ on the supergravity fermions. If the present construction of increasingly faithful representations of $K(\mathfrak{e}_9)$ leads to increasingly faithful representations of $K(\mathfrak{e}_{10})$, this can shed new light on the de-emergence of space and its replacement by algebraic concepts as discussed in [16].

Since $K(\mathfrak{e}_{10})$ differs from $K(\mathfrak{e}_9)$ by exactly one simple Berman generator (designated by x_1 with our labeling of the E_{10} Dynkin diagram, see figure 1 in section 5), representations of $K(\mathfrak{e}_9)$ that can be doubled to $K(\mathfrak{e}_{10})$ representations then must have a realisation of this Berman generator that mixes and interchanges the two parabolic subgroups in (1.4). The relations that this Berman generator must satisfy are known and spelt out in section 5. The problem of finding $K(\mathfrak{e}_{10})$ representation is thus greatly reduced, but it is worth stressing that the action of this extra Berman generator is not simply an interchange of the two chiral halves. As we shall see with the concrete examples of the known $K(E_{10})$ representations, the very existence of these representations relies on subtle consistency conditions that go beyond the affine construction in an essential way, and whose general form remains to be fully explored. We also note that the structure of (1.4) implies that $K(\mathfrak{e}_{10})$ representations obtained in this way must necessarily have *non-compact* quotients whenever the unipotent parts U_{\pm} are non-trivial. This explains one of the strangest features of our construction, namely the fact that these quotient groups are generically non-compact even though they descend from a ‘maximally compact’ subgroup [21].

However, the main challenge at this point concerns the extension of our results to $K(\mathfrak{e}_{10})$, with the idea that its corresponding unfaithful representations can be constructed by appropriately ‘gluing’ two chiral representations of $K(\mathfrak{e}_9)$, and by combining the two nilpotent subalgebras in (3.20), as we sketch in section 5. The latter strategy is suggested by the fact that all known unfaithful spinorial representations of $K(\mathfrak{e}_{10})$ are of this type. In section 5 we also state the consistency conditions that must be satisfied in order for such an ‘uplift’ to $K(\mathfrak{e}_{10})$ to work. This is important because we know that not every representation of $K(\mathfrak{e}_9)$ admits such an uplift to $K(\mathfrak{e}_{10})$ (an example is obtained by truncating a $\mathbf{16} \oplus \mathbf{16}$ from the spin- $\frac{3}{2}$ representation). Spinor representations for involutory subalgebras of other indefinite Kac-Moody algebras have also been discussed in [21, 32–34].

Let us comment on the possible relevance of our results in view of some of the Kac-Moody conjectures in the literature. Recently, E_n symmetries have been central in the context of exceptional field theory, see [35, 36] for reviews, and representations of $\widetilde{K(E_n)}$ enter when discussing fermions [37–39], where in particular the last reference discusses the Kac-Moody case $n = 11$. Exceptional field theories are duality *covariant* formulations of families of maximally supersymmetric theories. Focussing on a single maximally supersymmetric theory breaks the E_n symmetry but we expect the $\widetilde{K(E_n)}$ symmetry to remain intact, much in the same way as the reformulations of $D = 11$ supergravity in [3, 40] maintain a larger local symmetry.

By contrast, the conjectures of E_{10} [30] or E_{11} [41] symmetries of M-theory assert that the bosonic Kac-Moody symmetries are fully intact in (classical) M-theory, and at

this point it is not clear what the fermionic extension of these conjectures will be. In particular, it is not clear whether these theories are merely supposed to be (albeit very complicated) reformulations of maximal supergravities with essentially the same local dynamics, or whether the dynamics involve local degrees of freedom beyond the ones of maximal supergravity. The former point of view has been adopted in much of the literature on exceptional field theory. The second point of view is supported by the fact that there do exist theories ‘beyond’ string theory, and with different (non-perturbative) dynamics, namely the $D = 11$ supermembrane [42] and the associated maximally supersymmetric $SU(\infty)$ matrix theory [43]. We hope that the results presented in this paper can contribute to the clarification of these issues, and we expect that in either case further advances will make recourse to some of the structures derived in this paper.

The results presented in this paper are given for $K(\mathfrak{e}_9)$, but most of the general structure is applicable to any $K(\mathfrak{g})$ where \mathfrak{g} is a non-twisted affine algebra. One possible other arena for the considerations of this paper would therefore be pure $\mathcal{N} = 1$ supergravity reduced from $D = 4$ to $D = 2$ dimensions where the (Geroch) symmetry algebra \mathfrak{g} is affinised $\mathfrak{sl}(2)$ [44].

The structure of this article is as follows. We first review the general construction of $K(\mathfrak{e}_9)$ representations of [1] in section 2. In section 3, we discuss the structure of the associated ideals in more detail and in particular introduce the notion of chirality, showing that the chiral quotients commute. Section 4 connects the results of this paper to the explicit higher-spin representations of $K(\mathfrak{e}_{10})$ found in [25] and how they reduce to $K(\mathfrak{e}_9)$. In section 5 we then discuss the converse problem and discuss conditions for lifting $K(\mathfrak{e}_9)$ representations to $K(\mathfrak{e}_{10})$.

Acknowledgments

We are grateful to Ralf Köhl and Robin Lautenbacher for discussions. The work of H.N. has received funding from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (grant agreement No. 740209).

2 Parabolic and filtered algebras

In this section, we review the basic set-up from [1].

2.1 $K(E_9)$ from E_9

The Kac-Moody algebra \mathfrak{e}_9 can be defined in terms of a Chevalley-Serre basis and associated relations described by its Dynkin diagram. As for any affine Lie algebra it can be equivalently given in terms of a centrally extended loop algebra as

$$\mathfrak{e}_9 = \mathfrak{e}_8[t, t^{-1}] \oplus \mathbb{R}k \oplus \mathbb{R}d \tag{2.1}$$

where $\mathfrak{e}_8[t, t^{-1}] \cong \mathbb{R}[t, t^{-1}] \otimes \mathfrak{e}_8$ denotes \mathfrak{e}_8 -valued Laurent polynomials in a formal variable t and k and d are semi-simple elements of the algebra, with k being central. The commutation

relations are given by

$$\begin{aligned} [t^m \otimes x, t^n \otimes y] &= t^{m+n} \otimes [x, y] + m\delta_{m,n}(x, y)k, \\ [t^m \otimes x, k] &= [d, k] = 0, \quad [t^m \otimes x, d] = mt^m \otimes x \end{aligned} \quad (2.2)$$

for $m, n \in \mathbb{Z}$ and $x, y \in \mathfrak{e}_8$ and with (x, y) the canonically normalised Killing pairing on \mathfrak{e}_8 . We only consider the split real form of the algebra.

The 248-dimensional algebra \mathfrak{e}_8 has a Cartan-Chevalley involution $\hat{\omega}$ whose fixed point set is the 120-dimensional compact subalgebra $\mathfrak{so}(16)$ of \mathfrak{e}_8 . The remaining 128 generators transform in a spinor representation of $\mathfrak{so}(16)$. We shall denote the generators in the associated Cartan decomposition as (X^{IJ}, Y^A) with $X^{IJ} = X^{[IJ]}$ the 120 $\mathfrak{so}(16)$ generators while Y^A are the 128 coset generators. The commutation relations are

$$\begin{aligned} [X^{IJ}, X^{KL}] &= 2\delta^{K[J} X^{I]L} - 2\delta^{L[J} X^{I]K}, \\ [X^{IJ}, Y^A] &= -\frac{1}{2}\Gamma_{AB}^{IJ} Y^B, \\ [Y^A, Y^B] &= \frac{1}{4}\Gamma_{AB}^{IJ} X^{IJ}. \end{aligned} \quad (2.3)$$

Indices can be raised and lowered with the $\mathfrak{so}(16)$ -invariant metrics δ_{IJ} for fundamental indices and δ_{AB} for spinor indices. The matrices Γ_{AB}^{IJ} are taken from the $\mathfrak{so}(16)$ Clifford algebra and they are of size 128×128 for fixed I, J . Conjugate spinor representations of $\mathfrak{so}(16)$ will be denoted with indices \dot{A} . The involution $\hat{\omega}$ acts on these basis generators by

$$\hat{\omega}(X^{IJ}) = X^{IJ}, \quad \hat{\omega}(Y^A) = -Y^A. \quad (2.4)$$

The Cartan-Chevalley involution $\hat{\omega}$ on \mathfrak{e}_8 can be extended to an involution ω on \mathfrak{e}_9 by letting

$$\omega(t^n \otimes x) = t^{-n} \otimes \hat{\omega}(x), \quad \omega(d) = -d, \quad \omega(k) = -k. \quad (2.5)$$

We denote the invariant subalgebra under this involution by $K(\mathfrak{e}_9) \equiv \mathfrak{k}$.

2.2 Filtered and parabolic algebras

A basis for $K(\mathfrak{e}_9)$ can be given explicitly by

$$\mathcal{X}_n^{IJ} := \frac{1}{2}(t^n + t^{-n}) \otimes X^{IJ} \quad \text{for } n \geq 0, \quad (2.6a)$$

$$\mathcal{Y}_n^A := \frac{1}{2}(t^n - t^{-n}) \otimes Y^A \quad \text{for } n > 0. \quad (2.6b)$$

The elements d and k are projected out when descending from \mathfrak{e}_9 to $K(\mathfrak{e}_9)$.

We shall refer to the elements (2.6) as the *filtered basis* of $K(\mathfrak{e}_9)$. The reason for this terminology is that the commutation relations in this basis are no longer graded (as in (2.2)) but read

$$[\mathcal{X}_m^{IJ}, \mathcal{X}_n^{KL}] = 2\delta^{[I[K} (\mathcal{X}_{m+n}^{L]J]} + \mathcal{X}_{|m-n|}^{L]J]}, \quad (2.7a)$$

$$[\mathcal{X}_m^{IJ}, \mathcal{Y}_n^A] = -\frac{1}{4}\Gamma_{AB}^{IJ} (\mathcal{Y}_{m+n}^B - \text{sgn}(m-n)\mathcal{Y}_{|m-n|}^B), \quad (2.7b)$$

$$[\mathcal{Y}_m^A, \mathcal{Y}_n^B] = \frac{1}{8}\Gamma_{AB}^{IJ} (\mathcal{X}_{m+n}^{IJ} - \mathcal{X}_{|m-n|}^{IJ}). \quad (2.7c)$$

As the involution (2.5) sends $t \rightarrow t^{-1}$ one can also consider a different algebra that is more adapted to the fixed points of this involution on Laurent polynomials $\mathbb{R}[t, t^{-1}]$. The fixed points are given by $t_* = \pm 1$ and we consider the explicit change of variables

$$u = \frac{1 \mp t}{1 \pm t} \iff t = \pm \frac{1-u}{1+u} \tag{2.8}$$

for the two choices of fixed points. Around these fixed points we define the Lie algebra

$$\mathfrak{N} = \mathbb{R}[[u^2]] \otimes \langle X^{IJ} \rangle \oplus u\mathbb{R}[[u^2]] \otimes \langle Y^A \rangle \tag{2.9}$$

with the bracket defined in the obvious way and allowing for arbitrary formal power series. Written explicitly, a basis of this Lie algebra is given by

$$\begin{aligned} \mathcal{A}_{2n}^{IJ} &:= u^{2n} \otimes X^{IJ} && \text{for } n \geq 0, \\ \mathcal{S}_{2n+1}^A &:= u^{2n+1} \otimes Y^A && \text{for } n \geq 0. \end{aligned} \tag{2.10}$$

We stress that the Lie algebra (2.9) is allowed to contain arbitrary formal power series in u . Since only positive powers of u arises, the commutation relations of the basis elements are graded according to

$$[\mathcal{A}_{2m}^{IJ}, \mathcal{A}_{2n}^{KL}] = \frac{1}{2} \delta^{JK} \mathcal{A}_{2(m+n)}^{IL} - \frac{1}{2} \delta^{IK} \mathcal{A}_{2(m+n)}^{JL} - \frac{1}{2} \delta^{JL} \mathcal{A}_{2(m+n)}^{IK} + \frac{1}{2} \delta^{IL} \mathcal{A}_{2(m+n)}^{JK}, \tag{2.11a}$$

$$[\mathcal{A}_{2m}^{IJ}, \mathcal{S}_{2n+1}^{KL}] = -\frac{1}{4} \Gamma_{AB}^{IJ} \mathcal{S}_{2(m+n)+1}^B, \tag{2.11b}$$

$$[\mathcal{S}_{2m+1}^A, \mathcal{S}_{2n+1}^B] = \frac{1}{8} \Gamma_{AB}^{IJ} \mathcal{A}_{2(m+n+1)}^{IJ}. \tag{2.11c}$$

We refer to this algebra as a *parabolic* Lie algebra as the Levi part $\mathfrak{so}(16)$, generated by \mathcal{A}_0^{IJ} , acts on the (generalised) nilpotent part with positive subscripts.

One of the main results of [1] was that there are injective Lie algebra homomorphisms $\rho_{\pm} : K(\mathfrak{e}_9) \rightarrow \mathfrak{N}$. These can be obtained by expanding out the explicit change of variables (2.8) according to

$$t^n + t^{-n} = (\pm 1)^n \sum_{k \geq 0} a_{2k}^{(n)} u^{2k}, \quad t^n - t^{-n} = (\pm 1)^n \sum_{k \geq 0} a_{2k+1}^{(n)} u^{2k+1} \tag{2.12}$$

for $n \geq 0$ and $n > 0$, respectively. The two choices of sign correspond to the two choices in (2.8). The two Lie algebra homomorphisms are then defined explicitly by

$$\rho_{\pm}(\mathcal{X}_n^{IJ}) = (\pm 1)^n \frac{1}{2} \sum_{k \geq 0} a_{2k}^{(n)} \mathcal{A}_{2k}^{IJ}, \quad \rho_{\pm}(\mathcal{Y}_n^A) = (\pm 1)^n \frac{1}{2} \sum_{k \geq 0} a_{2k+1}^{(n)} \mathcal{S}_{2k+1}^A. \tag{2.13}$$

As explained in [1], the homomorphisms are not surjective as one would require power series in $\mathbb{R}[[t, t^{-1}]]$ that, however, do not behave well under multiplication.

Explicit formulas for the coefficients are

$$a_{2k}^{(n)} = 2 \sum_{\ell=0}^n \binom{2n}{2\ell} \binom{k-\ell+n-1}{k-\ell} \quad \text{for } n \geq 1 \text{ and } k \geq 1, \tag{2.14a}$$

$$a_{2k+1}^{(n)} = -2 \sum_{\ell=0}^{n-1} \binom{2n}{2\ell+1} \binom{k-\ell+n-1}{k-\ell} \quad \text{for } n \geq 0 \text{ and } k \geq 0. \tag{2.14b}$$

Also $a_{2k}^{(0)} = 0$ for all $k \geq 1$ and $a_0^{(n)} = 2$ for all $n \geq 0$.

2.3 Representations from truncations

It is easy to construct finite-dimensional representations for the parabolic algebra \mathfrak{N} defined in (2.9) by considering quotients of the algebra. Examples can be obtained by quotienting the ring of power series $\mathbb{R}[[u]]$ by the ideal $u^{N+1}\mathbb{R}[[u]]$ of power series whose lowest order term is u^{N+1} . As a vector space, the quotient corresponds to polynomials of degree at most N , but the quotient construction also endows the vector space with a product structure. This is given by working modulo terms of order $\mathcal{O}(u^{N+1})$. In terms of the parabolic generators this amounts to setting

$$\mathcal{A}_{2k}^{IJ} = 0 \text{ for } k > \lfloor N/2 \rfloor, \quad \mathcal{S}_{2k+1}^A = 0 \text{ for } k > \lfloor (N-1)/2 \rfloor. \quad (2.15)$$

There are two cases to be distinguished here, according to whether the highest ‘active’ generator is of type \mathcal{A}_{2k} or of type \mathcal{S}_{2k+1} . Since these two cases are largely analogous we will for definiteness assume $N = 2k_0$ even and

$$\mathcal{A}_{2k}^{IJ} = \mathcal{S}_{2k-1}^A = 0 \quad \text{for } k > k_0 = \frac{N}{2} \quad (2.16)$$

in the remainder, such that the highest ‘active’ generators are $\mathcal{A}_{2k_0}^{IJ}$ and $\mathcal{S}_{2k_0-1}^A$. This quotient on the ring of power series induces a quotient Lie algebra of \mathfrak{N} that we denote by \mathfrak{N}_N . Representations of this quotient Lie algebra can be constructed by considering all elements of degree at most N in the universal enveloping algebra of \mathfrak{N}_N acting on a given $\mathfrak{so}(16)$ module \mathfrak{W}_0 . In practice, this means that we are considering the graded representation

$$\mathfrak{W}_N = \bigoplus_{i=0}^N \mathfrak{W}_i \quad (2.17)$$

with the constituent $\mathfrak{so}(16)$ modules (with $N = 2k_0$)

$$\begin{aligned} \mathfrak{W}_0 & \\ \mathfrak{W}_1 &= \langle \mathcal{S}_1^A \mathfrak{W}_0 \rangle \\ \mathfrak{W}_2 &= \langle \mathcal{S}_1^A \mathcal{S}_1^B \mathfrak{W}_0 \rangle \oplus \langle \mathcal{A}_2^{IJ} \mathfrak{W}_0 \rangle \end{aligned} \quad (2.18)$$

$$\begin{aligned} & \vdots \\ \mathfrak{W}_N &= \langle \mathcal{S}_1^{A_1} \dots \mathcal{S}_1^{A_N} \mathfrak{W}_0 \rangle \oplus \dots \oplus \langle \mathcal{S}_1^A \mathcal{S}_{N-1}^B \mathfrak{W}_0 \rangle \oplus \langle \mathcal{A}_N^{IJ} \mathfrak{W}_0 \rangle \end{aligned} \quad (2.19)$$

where the generators must be ordered in accordance with the Poincaré-Birkhoff-Witt theorem. As these are representations of a quotient \mathfrak{N}_N of the parabolic Lie algebra \mathfrak{N} into which $K(\mathfrak{e}_9)$ embeds injectively, the representations can be pulled back to finite-dimensional representations of $K(\mathfrak{e}_9)$ using the homomorphisms ρ_{\pm} [1]. They provide a plethora of new representations.

Yet more representations can be obtained from (2.18) by further truncating away sub-representations at any level together with their associated submodules of \mathfrak{W}_N , as was already explained in [1]. Unlike the quotient to \mathfrak{W}_N , removing further $\mathfrak{so}(16)$ representations within one of the constituent \mathfrak{W}_i may require a careful check that the resulting space

is a representation of $K(\mathfrak{e}_9)$. Nevertheless, the known examples of $K(\mathfrak{e}_{10})$ representations suggest that such extra truncations may be necessary for the existence of the over-extended Berman generator x_1 and for a consistent uplift to $K(E_{10})$ [26].

3 Ideals

The representations \mathfrak{W}_N defined in the previous section are characterised by the fact that $\mathcal{A}_{2k}^{IJ} = 0$ and $\mathcal{S}_{2k-1}^A = 0$ for $k > k_0$, where we fix k_0 throughout this section. The case when the highest active generator is of type \mathcal{S} can be dealt with similarly.

We therefore have the finite sum relations valid in these representations

$$\rho_{\pm}(\mathcal{X}_n^{IJ}) = (\pm 1)^n \frac{1}{2} \sum_{k=0}^{k_0} a_{2k}^{(n)} \mathcal{A}_{2k}^{IJ}, \quad (3.1a)$$

$$\rho_{\pm}(\mathcal{Y}_n^A) = (\pm 1)^n \frac{1}{2} \sum_{k=1}^{k_0} a_{2k-1}^{(n)} \mathcal{S}_{2k-1}^A. \quad (3.1b)$$

We can consider the first relation for $0 \leq n \leq k_0$ and the second one for $1 \leq n \leq k_0$. From the explicit form of the $a_{2k}^{(n)}$ one can check that these represent linear systems of relations with unique solutions for \mathcal{A}_{2k} and \mathcal{S}_{2k-1} in terms of \mathcal{X}_n and \mathcal{Y}_n for $n \leq k_0$. Substituting this solution with the \mathcal{A}_{2k}^{IJ} and \mathcal{S}_{2k+1}^A in terms of the corresponding \mathcal{X}_n^{IJ} and \mathcal{Y}_n^A back into (3.1) we can express any of the higher \mathcal{X}_n^{IJ} and \mathcal{Y}_n^A for $n > k_0$ in terms of these. Explicitly, we have

$$(\pm 1)^n \mathcal{X}_n^{IJ} = \sum_{m=0}^{k_0} (\pm 1)^m c_{n,m} \mathcal{X}_m^{IJ}, \quad (3.2a)$$

$$(\pm 1)^n \mathcal{Y}_n^A = \sum_{m=1}^{k_0} (\pm 1)^m d_{n,m} \mathcal{Y}_m^A \quad (3.2b)$$

where the coefficients are given by

$$c_{n,m} \equiv c_{n,m}^{(k_0)} := \prod_{\substack{0 \leq j \leq k_0 \\ j \neq m}} \frac{n^2 - j^2}{m^2 - j^2}, \quad d_{n,m} \equiv d_{n,m}^{(k_0)} := \frac{n}{m} \prod_{\substack{1 \leq j \leq k_0 \\ j \neq m}} \frac{n^2 - j^2}{m^2 - j^2}, \quad (3.3)$$

where always $m \leq k_0$ (below we will often suppress the superscript k_0). For $0 \leq n \leq k_0$, the relations (3.2) become tautologies, as in this range $c_{n,m}^{(k_0)} = d_{n,m}^{(k_0)} = \delta_{n,m}$ (remember that the second index m is always restricted to a finite range). Consequently, only the \mathcal{X}_n^{IJ} for $0 \leq n \leq k_0$ and the \mathcal{Y}_n^A for $1 \leq n \leq k_0$ are independent, whereas the higher index objects depend linearly on them via the above ideal relations. Put differently, for \mathcal{X}_n^{IJ} with $n > k_0$ and \mathcal{Y}_n^A with $n > k_0$ the relations (3.2) are non-trivial relations among the representation matrices of $K(\mathfrak{e}_9)$. Importantly, these are valid *only* in the representations obtained by truncating the parabolic algebra. For ease of notation we omit the homomorphisms ρ_{\pm} and representation maps from the formulæ that are always to be understood in the representation \mathfrak{W}_N .

The above coefficients are the unique solutions to the problem of finding coefficients that satisfy the conditions that $c_{n,m}^{(k_0)} = d_{n,m}^{(k_0)} = \delta_{n,m}$ for small $0 \leq n \leq k_0$, and are polynomial of the right degree and parity in n .

3.1 Chiral ideals

We denote representations where the relations (3.2) are satisfied with a definite choice of sign as *chiral* representations, and write them as S_{\pm} . In the following discussion we shall not write out the $\mathfrak{so}(16)$ indices IJ and A on \mathcal{X}_m^{IJ} and \mathcal{Y}_m^A as they are spectators in the whole discussion. Corresponding to the two choices of sign in (3.2) we define the following elements

$$\begin{aligned}\mathcal{I}_n^{\pm} &:= \mathcal{X}_n - \sum_{m=0}^{k_0} (\pm 1)^{n+m} c_{n,m} \mathcal{X}_m, \\ \mathcal{J}_n^{\pm} &:= \mathcal{Y}_n - \sum_{m=1}^{k_0} (\pm 1)^{n+m} d_{n,m} \mathcal{Y}_m.\end{aligned}\tag{3.4}$$

The \mathcal{I}_n^{\pm} and the \mathcal{J}_n^{\pm} are only non-zero for $n > k_0$. These are represented trivially in the truncated representations for any n , and therefore we have the two ideals

$$\mathfrak{i}_{\pm} = \bigoplus_{n > k_0} \mathbb{R} \mathcal{I}_n^{\pm} \oplus \bigoplus_{n > k_0} \mathbb{R} \mathcal{J}_n^{\pm} \subset \mathfrak{k}.\tag{3.5}$$

The non-direct sum $\mathfrak{i}_+ + \mathfrak{i}_- \subset \mathfrak{k}$ is consequently spanned by the even and odd combinations

$$\begin{aligned}\mathcal{X}_{2n} - \sum_{m=0}^{\lfloor \frac{k_0}{2} \rfloor} c_{2n,2m}^{(k_0)} \mathcal{X}_{2m} \quad \text{and} \quad \mathcal{Y}_{2n} - \sum_{m=1}^{\lfloor \frac{k_0}{2} \rfloor} d_{2n,2m}^{(k_0)} \mathcal{Y}_{2m}, \\ \mathcal{X}_{2n+1} - \sum_{m=0}^{\lfloor \frac{k_0-1}{2} \rfloor} c_{2n+1,2m+1}^{(k_0)} \mathcal{X}_{2m+1} \quad \text{and} \quad \mathcal{Y}_{2n+1} - \sum_{m=0}^{\lfloor \frac{k_0-1}{2} \rfloor} d_{2n+1,2m+1}^{(k_0)} \mathcal{Y}_{2m+1},\end{aligned}\tag{3.6}$$

and the combinations

$$\begin{aligned}\sum_{m=0}^{\lfloor \frac{k_0}{2} \rfloor} c_{2n+1,2m}^{(k_0)} \mathcal{X}_{2m} \quad \text{and} \quad \sum_{m=1}^{\lfloor \frac{k_0}{2} \rfloor} d_{2n+1,2m}^{(k_0)} \mathcal{Y}_{2m}, \\ \sum_{m=0}^{\lfloor \frac{k_0-1}{2} \rfloor} c_{2n,2m+1}^{(k_0)} \mathcal{X}_{2m+1} \quad \text{and} \quad \sum_{m=0}^{\lfloor \frac{k_0-1}{2} \rfloor} d_{2n,2m+1}^{(k_0)} \mathcal{Y}_{2m+1}.\end{aligned}\tag{3.7}$$

These combinations are, respectively, obtained by adding and subtracting the above ideal components for even and for odd n .

We can now show that these combinations together span all of $\mathfrak{k} = K(\mathfrak{e}_9)$.

Lemma. We have

$$\mathfrak{i}_+ + \mathfrak{i}_- = \mathfrak{k}.\tag{3.8}$$

Proof. We first note that the relations (3.7) only involve the generators \mathcal{X}_m and \mathcal{Y}_m for $m \leq k_0$. Consequently we have an *infinite set of relations* for finitely many objects, and these relations are not of finite corank due to the structure of the coefficients (3.3). From this we conclude that all \mathcal{X}_m for $0 \leq m \leq k_0$ and all \mathcal{Y}_m for $1 \leq m \leq k_0$ can be obtained by suitable linear combinations, and are therefore contained in $\mathfrak{i}_+ + \mathfrak{i}_-$. Taking linear combinations with (3.6) we then see that all \mathcal{X}_m and \mathcal{Y}_m for any index are contained in $\mathfrak{i}_+ + \mathfrak{i}_-$ which therefore equals \mathfrak{k} . \square

3.2 Commuting chiral quotients

We now consider the representation of \mathfrak{k} that is given by the direct sum $S_+ \oplus S_-$ of two chiral representations S_\pm with ideals \mathfrak{i}_\pm and associated quotient algebras

$$\mathfrak{q}_\pm = \mathfrak{k}/\mathfrak{i}_\pm. \tag{3.9}$$

Note that the ideals do not fix the representations and the following statements should be correct for any choice of representations with the same ideal. The representations S_+ and S_- are related to one another by exchanging $\mathcal{X}_m \leftrightarrow (\pm 1)^m \mathcal{X}_m$ and $\mathcal{Y}_m \leftrightarrow (\pm 1)^m \mathcal{Y}_m$, and in this sense equivalent. From the explicit form of the ideals we see that representatives of the quotient algebras can be given solely in terms of the generators $\mathcal{X}_0, \mathcal{X}_1, \dots, \mathcal{X}_{k_0}$ and $\mathcal{Y}_1, \mathcal{Y}_2, \dots, \mathcal{Y}_{k_0}$.

The ideal of the representation $S_+ \oplus S_-$ is given by the intersection of ideals $\mathfrak{i}_+ \cap \mathfrak{i}_-$ and the quotient algebra therefore by

$$\mathfrak{q} := \mathfrak{k}/\mathfrak{i}, \quad \mathfrak{i} := \mathfrak{i}_+ \cap \mathfrak{i}_-. \tag{3.10}$$

We first describe the intersection ideal \mathfrak{i} in more detail. This intersection consists of linear combinations of elements in \mathfrak{i}_\pm given in (3.5) that can be formed such that the choice of sign disappears. We discuss this for the case when k_0 is even for concreteness; other cases can be analysed similarly.

The spanning elements (3.4) have different signs depending on the chirality and the \mathcal{I}_n^\pm (resp. \mathcal{J}_n^\pm) therefore do not lie in the intersection ideal \mathfrak{i} . We need to form linear combinations that are common to \mathfrak{i}_+ and \mathfrak{i}_- where these alternating signs do not appear and we describe this in detail for the \mathcal{X}_m . This can be done by choosing $k_0 + 1$ appropriate elements \mathcal{I}_n^\pm that we take to be $\mathcal{I}_{k_0+1}^\pm, \mathcal{I}_{k_0+2}^\pm, \dots, \mathcal{I}_{2k_0+1}^\pm$. The $(\pm 1)^m \mathcal{X}_m$ with $m \leq k_0$ can be expressed in terms of these elements so that we have found expressions that do not involve any alternating sign and these elements belong to the intersection ideal. Because they involve combinations of evenly indexed \mathcal{X}_m with $0 \leq m \leq 2k_0 + 1$, the range of indices is extended over twice the previous range. We therefore conclude that the intersection ideal $\mathfrak{i} = \mathfrak{i}_+ \cap \mathfrak{i}_-$ is given by

$$\begin{aligned} \mathfrak{i} = & \bigoplus_{n > k_0} \mathbb{R} \left(\mathcal{X}_{2n} - \sum_{m=0}^{k_0} c_{2n,2m}^{(k_0)} \mathcal{X}_{2m} \right) \oplus \bigoplus_{n > k_0+1} \mathbb{R} \left(\mathcal{Y}_{2n} - \sum_{m=1}^{k_0} d_{2n,2m}^{(k_0)} \mathcal{Y}_{2m} \right) \\ & \oplus \bigoplus_{n > k_0} \mathbb{R} \left(\mathcal{X}_{2n+1} - \sum_{m=0}^{k_0} c_{2n+1,2m+1}^{(k_0)} \mathcal{X}_{2m+1} \right) \oplus \bigoplus_{n > k_0-1} \mathbb{R} \left(\mathcal{Y}_{2n+1} - \sum_{m=0}^{k_0-1} d_{2n+1,2m+1}^{(k_0)} \mathcal{Y}_{2m+1} \right) \end{aligned} \tag{3.11}$$

with the *same* coefficients as in (3.3). However, the indices are doubled and, compared to (3.6), so is the summation range. Therefore the quotient algebra \mathfrak{q} defined in (3.10) has representatives given in terms of $\mathcal{X}_0, \mathcal{X}_1, \dots, \mathcal{X}_{2k_0+1}$ and $\mathcal{Y}_1, \mathcal{Y}_2, \dots, \mathcal{Y}_{2k_0}$, exactly twice as many as the chiral quotients \mathfrak{q}_\pm .

As an example we consider $k_0 = 2$. The \mathcal{I}_n^\pm elements occurring in \mathfrak{i}_\pm are

$$\begin{aligned} \mathcal{I}_{2n}^\pm &= \mathcal{X}_{2n} - \frac{1}{4}(4n^2 - 1)(4n^2 - 4)\mathcal{X}_0 \pm \frac{4}{3}n^2(4n^2 - 4)\mathcal{X}_1 - \frac{1}{3}n^2(4n^2 - 1)\mathcal{X}_2, \\ \mathcal{I}_{2n+1}^\pm &= \mathcal{X}_{2n+1} \mp (n^2 + n)(4n^2 + 4n - 3)\mathcal{X}_0 + \frac{1}{3}(2n + 1)^2(4n^2 + 4n - 3)\mathcal{X}_1 \\ &\quad \mp \frac{1}{3}(n^2 + n)(2n + 1)^2\mathcal{X}_2. \end{aligned} \tag{3.12}$$

For \mathcal{I}_n^\pm with $n \in \{0, 1, 2\}$ they vanish identically. For other values of n we can use this to solve for $\pm\mathcal{X}_1$, $\mp\mathcal{X}_0$ and $\mp\mathcal{X}_2$. For example, as \mathcal{I}_4^\pm is given by

$$\mathcal{I}_4^\pm = \mathcal{X}_4 - 45\mathcal{X}_0 \pm 64\mathcal{X}_1 - 20\mathcal{X}_2, \tag{3.13}$$

we have

$$\pm\mathcal{X}_1 = \frac{1}{64} \left(\mathcal{I}_4^\pm + 45\mathcal{X}_0 + 20\mathcal{X}_2 - \mathcal{X}_4 \right). \tag{3.14}$$

Substituting this back into (3.12) leads to expressions that are identical in \mathfrak{i}_+ and \mathfrak{i}_- . For the even case we get

$$\mathcal{I}_{2n}^\pm - \frac{1}{48}n^2(4n^2 - 4)\mathcal{I}_4^\pm = \mathcal{X}_{2n} - \frac{1}{4}(n^2 - 1)(n^2 - 4)\mathcal{X}_0 + \frac{1}{3}n^2(n^2 - 4)\mathcal{X}_2 - \frac{1}{12}n^2(n^2 - 1)\mathcal{X}_4 \tag{3.15}$$

as elements of \mathfrak{i}_\pm . As the right-hand side does not depend on the choice of chirality, these elements belong to the intersection ideal. Among the even \mathcal{X}_n we therefore remain with \mathcal{X}_0 , \mathcal{X}_2 and \mathcal{X}_4 in the quotient algebra (3.10). Performing the same analysis for the elements with odd indices, we arrive at the conclusion that \mathfrak{q} has representatives given by $\mathcal{X}_0, \mathcal{X}_1, \dots, \mathcal{X}_5$ and $\mathcal{Y}_1, \mathcal{Y}_2, \dots, \mathcal{Y}_4$. (The case when the highest active generator is of \mathcal{S} -type can be treated analogously.)

We next discuss the relation between the quotients \mathfrak{q}_\pm and \mathfrak{q} of \mathfrak{k} . Since we have trivially $\mathfrak{i} \subset \mathfrak{i}_\pm$ as an ideal, we can view \mathfrak{q}_\pm also canonically as quotients of \mathfrak{q} via

$$\mathfrak{q}_\pm \cong \mathfrak{q}/(\mathfrak{i}_\pm/\mathfrak{i}). \tag{3.16}$$

From the truncation construction it also follows that \mathfrak{q}_\pm are actually subalgebras of \mathfrak{q} . Using the relation (3.8) above we also obtain

$$\mathfrak{q}_\pm = (\mathfrak{i}_+ + \mathfrak{i}_-)/\mathfrak{i}_\pm = \mathfrak{i}_\mp/\mathfrak{i}. \tag{3.17}$$

Proposition. For any finite-dimensional representation of \mathfrak{k} defined by the cutoff conditions (2.15) the quotient algebra \mathfrak{q} decomposes into a direct sum of two mutually commuting subalgebras

$$\mathfrak{q} = \mathfrak{q}_+ \oplus \mathfrak{q}_-, \quad [\mathfrak{q}_+, \mathfrak{q}_-] = 0 \quad \text{as subalgebras of } \mathfrak{q}. \tag{3.18}$$

Both algebras are parabolic, with $\mathfrak{so}(16)_\pm$ as their Levi subalgebras. In the supergravity realisation \mathfrak{q}_\pm correspond to chiral and anti-chiral subalgebras [23] (as we will make explicit below).

Proof. We only need to show the subalgebras $\mathfrak{q}_\pm \subset \mathfrak{q}$ commute since this will imply their direct sum structures. In order to see that they commute we take recourse to (3.17) which tells us that we can find representatives of \mathfrak{q}_\pm of the form $x_\mp + \mathfrak{i}$ with $x_\mp \in \mathfrak{i}_\mp$. Then the commutator is

$$[x_- + \mathfrak{i}, x_+ + \mathfrak{i}] \in \mathfrak{i}_+ \cap \mathfrak{i}_-, \tag{3.19}$$

which is equal to zero in \mathfrak{q} . □

These observations imply that *for all finite-dimensional* $K(\mathfrak{e}_9)$ representations the action of the algebra splits into a direct sum of mutually commuting chiral and anti-chiral parabolic algebras. Hence we conclude that the quotient algebra (*alias* the generalised holonomy algebra) has the form

$$\mathfrak{q} = (\mathfrak{so}(16)_+ \oplus \mathfrak{n}_+) \oplus (\mathfrak{so}(16)_- \oplus \mathfrak{n}_-) \tag{3.20}$$

where the nilpotent algebras \mathfrak{n}_\pm are N -step nilpotent. We repeat that this structure holds *only* for the finite-dimensional representations considered here, but is not true generally because neither $\mathfrak{so}(16)_+ \oplus \mathfrak{so}(16)_-$ nor its parabolic extensions are subalgebras of $K(\mathfrak{e}_9)$. They only arise as quotients via the above construction and depend on the choice of ideal through a representation. The group-theoretic version of (3.20) is encapsulated in (1.4) of the introduction. We note that under the exponential map one obtains Spin(16) or SO(16) realised faithfully, depending on whether the starting representation \mathfrak{Q}_0 is a representation of SO(16) or its spin cover.

4 Examples and relation to root basis form

In this section we exemplify the abstract considerations above in terms of concretely known representations, referred to as spin- $\frac{1}{2}$, spin- $\frac{3}{2}$, spin- $\frac{5}{2}$ and spin- $\frac{7}{2}$. These are actually representations of $K(E_{10})$ [21, 25, 26], but we will here focus on the affine subalgebra $K(E_9)$ by restricting the E_{10} roots to the $K(E_9)$ root system (see [45] for a more detailed analysis of this embedding). With the E_8 roots $\hat{\alpha}, \hat{\beta}, \dots$ and the affine null root δ , the roots of affine E_9 are $m\delta$ (null roots) and $m\delta + \hat{\alpha}$ (real roots) for $m \in \mathbb{Z}$. The E_9 generators are thus

$$E(m\delta + \hat{\alpha}) \quad \text{and} \quad H^i(m\delta) \quad \text{for } m \in \mathbb{Z} \tag{4.1}$$

plus the central charge, which will however drop out. The corresponding $K(E_9)$ generators are

$$\begin{aligned} J(m\delta + \hat{\alpha}) &:= E(m\delta + \hat{\alpha}) - E(-m\delta - \hat{\alpha}) && \text{for } m \geq 0 \\ J^i(m\delta) &:= H^i(m\delta) - H^i(-m\delta) && \text{for } m \geq 1 \end{aligned} \tag{4.2}$$

so that

$$J(\gamma) = -J(-\gamma) \tag{4.3}$$

for all E_9 roots γ .

4.1 Spin- $\frac{1}{2}$

Recall that the generators of $\widetilde{K(E_8)} \equiv \text{Spin}(16)$ can be expressed by the gamma matrices $\Gamma(\dot{\alpha})$, where $\dot{\alpha}$ runs over the *positive* roots of E_8 (see [21, 25, 26] for details). These descend from the real 32-by-32 gamma matrices $\Gamma(\alpha)$ for $K(E_{10})$ (where α is a root of E_{10}), but decompose into two blocks of 16-by-16 matrices for $K(E_8)$. Furthermore we have the mod 2 property (also valid for $K(E_{10})$)

$$\Gamma(\dot{\alpha}) = \Gamma(-\dot{\alpha}) = \Gamma(\dot{\alpha} + 2n\dot{\beta}) \quad \forall n \in \mathbb{Z} \tag{4.4}$$

for any elements $\dot{\alpha}$ and $\dot{\beta}$ of the E_8 root lattice. To ensure (4.3) one would have to include an extra co-boundary factor

$$\tilde{\Gamma}(\dot{\alpha}) \equiv \sigma_{\dot{\alpha}} \Gamma(\dot{\alpha}), \quad \sigma_{\dot{\alpha}} \sigma_{-\dot{\alpha}} = -1 \tag{4.5}$$

but this subtlety can be ignored as long as we are dealing only with positive roots (as will be the case in the remainder).

The general form of real $K(E_{10})$ generators in this language for the representations found in [21, 25, 26] is then

$$J(\alpha) = -2X(\alpha) \otimes \Gamma(\alpha), \tag{4.6}$$

where the ‘polarisation tensor’ $X(\alpha)$ is constructed only out of data associated with the real root α . For spin- $\frac{1}{2}$, the tensor $X(\alpha) = -\frac{1}{4}$ is a pure number.

For $K(E_9)$ we need the null root δ as an extra root in addition to the E_8 roots. From the gamma matrix relations derived before we have (with the wall basis of [25, 26])

$$\Gamma(\delta) \equiv \Gamma^2 \dots \Gamma^{10} = \Gamma^0 \Gamma^1 \tag{4.7}$$

which is just the two-dimensional chirality matrix (note that $\Gamma(2m\delta) \equiv \mathbf{1}$). Hence, in this representation, $K(E_9)$ decomposes into mutually commuting chiral and anti-chiral Spin(16) algebras, with generators (recall that $\alpha \cdot \delta = 0$ for all E_9 roots)

$$\frac{1}{2}(\mathbf{1} \pm \Gamma(\delta))\Gamma(\dot{\alpha}) \equiv \frac{1}{2}(\Gamma(\dot{\alpha}) \pm \Gamma(\dot{\alpha} + \delta)) \quad (\dot{\alpha} > 0) \tag{4.8}$$

In terms of the previous abstract construction the chiral doubling is already apparent from the sign ambiguity in (2.13) where the positive (negative) chirality gets tied to the two fixed points of the involution in the spectral parameter plane [23].

4.2 Spin- $\frac{3}{2}$

From [25] we recall the correspondence with the generators written in spin- $\frac{3}{2}$ form

$$\begin{aligned} J(m\delta + \dot{\alpha}) &\cong X_{\mathbf{ab}}(m\delta + \dot{\alpha})\Gamma(m\delta + \dot{\alpha}) \\ J^i(m\delta) &\cong m \xi_{[\mathbf{a}}^i \delta_{\mathbf{b}]} \Gamma(m\delta) \end{aligned} \tag{4.9}$$

where the symmetric 10×10 matrices $X_{\mathbf{ab}}$ are defined by

$$X_{\mathbf{ab}}(\alpha) := -\frac{1}{2}\alpha_{\mathbf{a}}\alpha_{\mathbf{b}} + \frac{1}{4}G_{\mathbf{ab}} = X_{\mathbf{ab}}(-\alpha) \tag{4.10}$$

with the Lorentzian metric $G_{ab} \equiv 1 - \delta_{ab}$, and for any real E_9 root α (as we said this formula is actually valid also for E_{10}).

For the further correspondence with the generators \mathcal{X}_m^{IJ} and \mathcal{Y}_m^A we now take $\hat{\alpha} > 0$ to be a positive E_8 root, and identify²

$$\begin{aligned} \mathcal{X}_m(\hat{\alpha}) &= \frac{1}{2} \left[X(m\delta + \hat{\alpha}) + X(m\delta - \hat{\alpha}) \right] \Gamma(m\delta + \hat{\alpha}) & (\hat{\alpha} > 0, m \geq 0) \\ \mathcal{Y}_m(\hat{\alpha}) &= \frac{1}{2} \left[X(m\delta + \hat{\alpha}) - X(m\delta - \hat{\alpha}) \right] \Gamma(m\delta + \hat{\alpha}) & (\hat{\alpha} > 0, m \geq 1) \\ &\text{and } m(\xi^i \wedge \delta) \Gamma(m\delta) & (\text{for } m \geq 1) \end{aligned} \quad (4.11)$$

To determine the ideal relations directly from these formulas we note the elementary identities which follow directly from (4.10)

$$\begin{aligned} \frac{1}{2} \left[X(n\delta + \hat{\alpha}) + X(n\delta - \hat{\alpha}) \right] &= n^2 \cdot \frac{1}{2} \left[X(\delta + \hat{\alpha}) + X(\delta - \hat{\alpha}) \right] + (1 - n^2) X(\hat{\alpha}), \\ \frac{1}{2} \left[X(n\delta + \hat{\alpha}) - X(n\delta - \hat{\alpha}) \right] &= n \cdot \frac{1}{2} \left[X(\delta + \hat{\alpha}) - X(\delta - \hat{\alpha}) \right]. \end{aligned} \quad (4.12)$$

Here, we already recognise some of the previously derived relations, but we still need to keep track of the extra Γ -matrix factors in (4.11) which differ according to whether δ is multiplied by an even or an odd integer. Not mixing even and odd multiples of δ , we thus derive the relations

$$\begin{aligned} \mathcal{X}_{2n} &= n^2 \mathcal{X}_2 + (1 - n^2) \mathcal{X}_0, \\ \mathcal{X}_{2n+1} &= \frac{1}{8} [(2n+1)^2 - 1] \mathcal{X}_3 - \frac{1}{8} [(2n+1)^2 - 9] \mathcal{X}_1, \\ \mathcal{Y}_{2n} &= n \mathcal{Y}_2, \quad \mathcal{Y}_{2n+1} = (2n+1) \mathcal{Y}_1. \end{aligned} \quad (4.13)$$

For $n \geq 2$ these relations correspond precisely to the combinations in (3.6) which span the intersection ideal $\mathfrak{i}_+ \cap \mathfrak{i}_-$. As we saw above, in the parabolic basis this corresponds to the truncation

$$\mathcal{A}_4 = \mathcal{A}_6 = \dots = 0, \quad \mathcal{S}_3 = \mathcal{S}_5 = \dots = 0. \quad (4.14)$$

To get the chiral combinations we must now combine the X factors with the ‘wrong’ Γ -matrices by means of these relations, and re-express them in terms of \mathcal{X} generators; idem for \mathcal{Y} . For instance,

$$\frac{1}{2} \left[X(\delta + \hat{\alpha}) + X(\delta - \hat{\alpha}) \right] \Gamma(\hat{\alpha}) = \frac{1}{4} \left(\frac{1}{2} \left[X(2\delta + \hat{\alpha}) + X(2\delta - \hat{\alpha}) \right] + 3X(\hat{\alpha}) \right) \Gamma(\hat{\alpha}) \quad (4.15)$$

from which obtain the chiral combinations

$$\frac{1}{2} \left[X(\delta + \hat{\alpha}) + X(\delta - \hat{\alpha}) \right] \Gamma(\delta + \hat{\alpha}) (\mathbf{1} \pm \Gamma(\delta)) = \mathcal{X}_1 \pm \frac{1}{4} (\mathcal{X}_2 + 3\mathcal{X}_0) \quad (4.16)$$

which are, respectively, elements of \mathfrak{i}_+ and \mathfrak{i}_- . Similarly,

$$X(\hat{\alpha}) \Gamma(\hat{\alpha}) (\mathbf{1} \pm \Gamma(\delta)) = \mathcal{X}_0 \mp \frac{1}{8} (\mathcal{X}_3 - 9\mathcal{X}_1). \quad (4.17)$$

²There is no need to spell out the explicit relation between the generators \mathcal{X}_m^{IJ} and \mathcal{Y}_m^A and the root basis here, as we hope the notation is self-explanatory.

For the \mathcal{Y} generators the relevant relations are even simpler: we have

$$\frac{1}{2} \left[X(\delta + \dot{\alpha}) - X(\delta - \dot{\alpha}) \right] \Gamma(\delta + \dot{\alpha}) (\mathbf{1} \pm \Gamma(\delta)) = \mathcal{Y}_1 \pm \frac{1}{2} \mathcal{Y}_2. \quad (4.18)$$

Hence, all chiral and ant-chiral generators can be expressed as linear combinations of \mathcal{X}_0 , \mathcal{X}_1 , \mathcal{X}_2 , \mathcal{X}_3 and \mathcal{Y}_1 , \mathcal{Y}_2 , giving a total of $2 \times (120 + 128 + 120)$ generators, which is the correct count corresponding to $\mathfrak{k}/\mathfrak{i}_+ \cap \mathfrak{i}_-$, and in complete agreement with the supergravity analysis [23].

To check that chiral and anti-chiral transformations commute, we compute (for instance)

$$\begin{aligned} \left[\mathcal{Y}_1 + \frac{1}{2} \mathcal{Y}_2, \mathcal{Y}_1 - \frac{1}{2} \mathcal{Y}_2 \right] &= \frac{1}{4} (-\mathcal{X}_4 + 4\mathcal{X}_2 - 3\mathcal{X}_0), \\ \left[\mathcal{X}_1 + \frac{1}{4} (\mathcal{X}_2 + 3\mathcal{X}_0), \mathcal{X}_1 - \frac{1}{4} (\mathcal{X}_2 + 3\mathcal{X}_0) \right] &= \frac{1}{16} (-\mathcal{X}_4 + 4\mathcal{X}_2 - 3\mathcal{X}_0). \end{aligned} \quad (4.19)$$

Both commutators vanish by the ideal relation (4.13) (actually identically in this explicit representation!), so chiral and anti-chiral transformations indeed commute by virtue of the ideal relations. In the explicit representation with chiral projectors this commutation property is of course obvious, and merely confirms the abstract argument given before.

As shown in the preceding section one can therefore abstract from such concrete realisations, and define the commuting chiral algebras entirely in terms of the \mathcal{X}_n and \mathcal{Y}_n and ideal relations, such that closure of the given subalgebra depends on the respective ideal relations.

4.3 Spin- $\frac{5}{2}$

For spin- $\frac{5}{2}$ we have a similar representation as in (4.9), except that the matrices $X_{\mathbf{ab|cd}}$ now act in a 55-dimensional space [25], where

$$X_{\mathbf{ab,cd}}(\alpha) := \frac{1}{2} \alpha_{\mathbf{a}} \alpha_{\mathbf{b}} \alpha_{\mathbf{c}} \alpha_{\mathbf{d}} - \alpha_{(\mathbf{a}} G_{\mathbf{b})(\mathbf{c}} \alpha_{\mathbf{d})} + \frac{1}{4} G_{\mathbf{a}(\mathbf{c}} G_{\mathbf{d})\mathbf{b}}. \quad (4.20)$$

The formula (4.20) now yields the relations

$$\begin{aligned} \frac{1}{2} \left[X(n\delta + \dot{\alpha}) + X(n\delta - \dot{\alpha}) \right] &= \frac{1}{12} n^2 (n^2 - 1) \cdot \frac{1}{2} \left[X(2\delta + \dot{\alpha}) + X(2\delta - \dot{\alpha}) \right] \\ &\quad - \frac{1}{3} n^2 (n^2 - 4) \cdot \frac{1}{2} \left[X(\delta + \dot{\alpha}) + X(\delta - \dot{\alpha}) \right] + \frac{1}{4} (n^4 - 5n^2 + 4) X(\dot{\alpha}) \end{aligned} \quad (4.21)$$

and

$$\begin{aligned} \frac{1}{2} \left[X(n\delta + \dot{\alpha}) - X(n\delta - \dot{\alpha}) \right] &= \frac{1}{24} (n^3 - n) \cdot \frac{1}{2} \left[X(3\delta + \dot{\alpha}) - X(3\delta - \dot{\alpha}) \right] \\ &\quad + \frac{1}{8} (-n^3 + 9n) \cdot \frac{1}{2} \left[X(\delta + \dot{\alpha}) - X(\delta - \dot{\alpha}) \right]. \end{aligned} \quad (4.22)$$

These are exactly the relations we have found above in (3.2) with $k_0 = 2$, i.e., for

$$\mathcal{A}_6 = \mathcal{A}_8 = \dots = 0, \quad \mathcal{S}_5 = \mathcal{S}_7 = \dots = 0. \quad (4.23)$$

4.4 Spin- $\frac{7}{2}$

For spin- $\frac{7}{2}$ the polarisation matrices $X_{\mathbf{a}_1\mathbf{a}_2\mathbf{a}_3|\mathbf{b}_1\mathbf{b}_2\mathbf{b}_3}(\alpha) = X_{(\mathbf{a}_1\mathbf{a}_2\mathbf{a}_3)|(\mathbf{b}_1\mathbf{b}_2\mathbf{b}_3)}(\alpha)$ are given by [25]

$$\begin{aligned}
 X(\alpha)^{\mathbf{a}_1\mathbf{a}_2\mathbf{a}_3}_{\mathbf{b}_1\mathbf{b}_2\mathbf{b}_3} &= -\frac{1}{3}\alpha^{\mathbf{a}_1}\alpha^{\mathbf{a}_2}\alpha^{\mathbf{a}_3}\alpha_{\mathbf{b}_1}\alpha_{\mathbf{b}_2}\alpha_{\mathbf{b}_3} + \frac{3}{2}\alpha^{(\mathbf{a}_1}\alpha^{\mathbf{a}_2}\delta_{(\mathbf{b}_1}^{\mathbf{a}_3)}\alpha_{\mathbf{b}_2}\alpha_{\mathbf{b}_3)} - \frac{3}{2}\alpha^{(\mathbf{a}_1}\delta_{(\mathbf{b}_1}^{\mathbf{a}_2}\delta_{\mathbf{b}_2}^{\mathbf{a}_3)}\alpha_{\mathbf{b}_3)} \\
 &+ \frac{1}{4}\delta_{(\mathbf{b}_1}^{(\mathbf{a}_1}\delta_{\mathbf{b}_2}^{\mathbf{a}_2}\delta_{\mathbf{b}_3}^{\mathbf{a}_3)} + \frac{1}{12}(2 - \sqrt{3})\alpha^{(\mathbf{a}_1}G^{\mathbf{a}_2\mathbf{a}_3)}G_{(\mathbf{b}_1\mathbf{b}_2}\alpha_{\mathbf{b}_3)} \\
 &+ \frac{1}{12}(-1 + \sqrt{3})\left(\alpha^{(\mathbf{a}_1}\alpha^{\mathbf{a}_2}\alpha^{\mathbf{a}_3)}G_{(\mathbf{b}_1\mathbf{b}_2}\alpha_{\mathbf{b}_3)} + \alpha^{(\mathbf{a}_1}G^{\mathbf{a}_2\mathbf{a}_3)}\alpha_{(\mathbf{b}_1}\alpha_{\mathbf{b}_2}\alpha_{\mathbf{b}_3)}\right), \tag{4.24}
 \end{aligned}$$

where as before we raise (and lower) indices with the DeWitt metric $G_{\mathbf{ab}}$. Restricting from E_{10} roots α to affine roots $m\delta \pm \hat{\alpha}$ one can deduce the following relations from (4.24)

$$\begin{aligned}
 &\frac{1}{2}\left[X_{\mathbf{a}_1\mathbf{a}_2\mathbf{a}_3|\mathbf{b}_1\mathbf{b}_2\mathbf{b}_3}(n\delta + \hat{\alpha}) + X_{\mathbf{a}_1\mathbf{a}_2\mathbf{a}_3|\mathbf{b}_1\mathbf{b}_2\mathbf{b}_3}(n\delta - \hat{\alpha})\right] \\
 &= -\frac{1}{36}(n^2 - 9)(n^2 - 4)(n^2 - 1)X_{\mathbf{a}_1\mathbf{a}_2\mathbf{a}_3|\mathbf{b}_1\mathbf{b}_2\mathbf{b}_3}(\hat{\alpha}) \\
 &+ \frac{1}{48}(n^2 - 9)(n^2 - 4)n^2\left[X_{\mathbf{a}_1\mathbf{a}_2\mathbf{a}_3|\mathbf{b}_1\mathbf{b}_2\mathbf{b}_3}(\delta + \hat{\alpha}) + X_{\mathbf{a}_1\mathbf{a}_2\mathbf{a}_3|\mathbf{b}_1\mathbf{b}_2\mathbf{b}_3}(\delta - \hat{\alpha})\right] \\
 &- \frac{1}{120}(n^2 - 9)(n^2 - 1)n^2\left[X_{\mathbf{a}_1\mathbf{a}_2\mathbf{a}_3|\mathbf{b}_1\mathbf{b}_2\mathbf{b}_3}(2\delta + \hat{\alpha}) + X_{\mathbf{a}_1\mathbf{a}_2\mathbf{a}_3|\mathbf{b}_1\mathbf{b}_2\mathbf{b}_3}(2\delta - \hat{\alpha})\right] \\
 &+ \frac{1}{720}(n^2 - 4)(n^2 - 1)n^2\left[X_{\mathbf{a}_1\mathbf{a}_2\mathbf{a}_3|\mathbf{b}_1\mathbf{b}_2\mathbf{b}_3}(3\delta + \hat{\alpha}) + X_{\mathbf{a}_1\mathbf{a}_2\mathbf{a}_3|\mathbf{b}_1\mathbf{b}_2\mathbf{b}_3}(3\delta - \hat{\alpha})\right] \tag{4.25}
 \end{aligned}$$

and

$$\begin{aligned}
 &\frac{1}{2}\left[X_{\mathbf{a}_1\mathbf{a}_2\mathbf{a}_3|\mathbf{b}_1\mathbf{b}_2\mathbf{b}_3}(n\delta + \hat{\alpha}) - X_{\mathbf{a}_1\mathbf{a}_2\mathbf{a}_3|\mathbf{b}_1\mathbf{b}_2\mathbf{b}_3}(n\delta - \hat{\alpha})\right] \\
 &= \frac{1}{48}(n^2 - 9)(n^2 - 4)n\left[X_{\mathbf{a}_1\mathbf{a}_2\mathbf{a}_3|\mathbf{b}_1\mathbf{b}_2\mathbf{b}_3}(\delta + \hat{\alpha}) - X_{\mathbf{a}_1\mathbf{a}_2\mathbf{a}_3|\mathbf{b}_1\mathbf{b}_2\mathbf{b}_3}(\delta - \hat{\alpha})\right] \\
 &- \frac{1}{60}(n^2 - 9)(n^2 - 1)n\left[X_{\mathbf{a}_1\mathbf{a}_2\mathbf{a}_3|\mathbf{b}_1\mathbf{b}_2\mathbf{b}_3}(2\delta + \hat{\alpha}) - X_{\mathbf{a}_1\mathbf{a}_2\mathbf{a}_3|\mathbf{b}_1\mathbf{b}_2\mathbf{b}_3}(2\delta - \hat{\alpha})\right] \\
 &+ \frac{1}{240}(n^2 - 4)(n^2 - 1)n\left[X_{\mathbf{a}_1\mathbf{a}_2\mathbf{a}_3|\mathbf{b}_1\mathbf{b}_2\mathbf{b}_3}(3\delta + \hat{\alpha}) - X_{\mathbf{a}_1\mathbf{a}_2\mathbf{a}_3|\mathbf{b}_1\mathbf{b}_2\mathbf{b}_3}(3\delta - \hat{\alpha})\right]. \tag{4.26}
 \end{aligned}$$

These correspond precisely to the relations obtained by truncation with $k_0 = 3$ in (3.2). Remarkably, the square root appearing in (4.24) that is needed for satisfying the Berman relations does not affect the ideal relations. The appearance of ‘strange’ prefactors in (4.24) (which are absolutely necessary for the $K(E_{10})$ Berman relations to work, see following section) is another clear indication of subtleties that go beyond the affine construction.

5 Extension to $K(\mathfrak{e}_{10})$

As we already mentioned the ultimate goal of the present investigation is a better understanding of the representation theory for the maximally hyperbolic algebra \mathfrak{e}_{10} and its maximal compact subalgebra $K(\mathfrak{e}_{10})$. The $E_{10}/K(E_{10})$ model of [30] is conjectured to provide a framework for (the bosonic sector of) M-theory in a small tension limit, with (de-)emergent space at the singularity, such that a precise correspondence with the field content and equations of motion of $D = 11$ supergravity holds if the field theory is restricted to first order spatial gradients, and the Lie algebra expansion to levels $\ell \leq 3$ (or




Figure 1. Dynkin diagram of E_{10} with nodes labelled.

roots of height $\text{ht}(\alpha) \leq 29$) in a decomposition of \mathfrak{e}_{10} under its \mathfrak{gl}_{10} subalgebra. Consequently the main challenge that remains is to extend this correspondence to higher order spatial gradients and the exponentially growing tower of $\ell > 3$ representations, and this is a problem that has defied solution until now. It is here that the study of fermions could make a crucial difference, because the existence of unfaithful $\widetilde{K}(\mathfrak{e}_{10})$ representations of ever increasing dimensions may afford a more ‘global’ view of $\widetilde{K}(E_{10})$, and thus also E_{10} , which is unachievable with the tools usually employed in the theory of Kac-Moody algebras and Kac-Moody groups (partition functions, level decompositions, local constructions, etc.). This final section is meant to provide a sketchy account of how progress on the representation theory of $K(\mathfrak{e}_9)$ could be possibly exploited towards a better understanding of $K(\mathfrak{e}_{10})$. Namely, the available evidence indicates that this can be accomplished by ‘gluing’ chiral and anti-chiral representations of $K(\mathfrak{e}_9)$ by means of an explicit action of the one remaining Berman generator x_1 on the given representations. However, the consistency conditions that must be satisfied (cf. (5.6), (5.7) and (5.8) below) remain to be explored.

In order to discuss the extension from $K(\mathfrak{e}_9)$ to $K(\mathfrak{e}_{10})$ we make use of the labelling of the Dynkin diagram shown in figure 1. The algebra $K(\mathfrak{e}_{10})$ is generated by ten Berman generators x_1, \dots, x_{10} with Berman relations [46]

$$\begin{aligned} [x_i, x_j] &= 0 && \text{if } i \text{ and } j \text{ disconnected in the diagram,} \\ [x_i [x_i, x_j]] &= -x_j && \text{if } i \text{ and } j \text{ connected in the diagram.} \end{aligned} \quad (5.1)$$

The $K(\mathfrak{e}_9)$ subalgebra is obtained by restricting to the abstract generators x_2, \dots, x_{10} .

For any finite-dimensional representation of $K(\mathfrak{e}_9)$, the Berman generators x_2, \dots, x_{10} are represented by specific matrices X_2, \dots, X_{10} that can be written as

$$X_i := \frac{1}{4} \left(\gamma_{\alpha\beta}^{i,i+1} \mathcal{X}_0^{\alpha\beta} + \gamma_{\dot{\alpha}\dot{\beta}}^{i,i+1} \mathcal{X}_0^{\dot{\alpha}\dot{\beta}} \right) \quad (5.2)$$

for $i = 3, \dots, 9$, and

$$X_{10} := -\frac{1}{2} \gamma_{\alpha\beta}^{8,9,10} \mathcal{X}_0^{\alpha\beta} \quad \text{and} \quad X_2 := -\frac{1}{2} \gamma_{\alpha\beta}^2 \left(\mathcal{X}_1^{\alpha\beta} + \mathcal{Y}_1^{\alpha\beta} \right), \quad (5.3)$$

where all \mathcal{X}_n and \mathcal{Y}_n are realised as *finite-dimensional* matrices in the concrete unfaithful representation under consideration. In the above equations we use the notation of [45] as well as the decomposition of the $\text{SO}(16)$ indices $[IJ]$ and A under the $\text{SO}(8) \times \text{SO}(8)$ subgroup of $\text{SO}(16)$ explained there.

We then seek a realisation of the $K(\mathfrak{e}_{10})$ Berman generators for the ‘doubled’ representation $S_+ \oplus S_-$ by means of the ansatz

$$x_1 = \begin{pmatrix} 0 & B \\ C & 0 \end{pmatrix}, \quad x_2 = \begin{pmatrix} X_2 & 0 \\ 0 & X_2 \end{pmatrix}, \quad x_i = \begin{pmatrix} X_i & 0 \\ 0 & X_i \end{pmatrix}, \quad x_{10} = \begin{pmatrix} X_{10} & 0 \\ 0 & -X_{10} \end{pmatrix} \quad (5.4)$$

with $i = 3, \dots, 9$ as before. The relation $[x_1, x_j] = 0$ for $j = 3, \dots, 10$ requires

$$[B, X_j] = [C, X_j] = 0 \tag{5.5}$$

or $j = 3, \dots, 9$, and $\{B, X_{10}\} = \{C, X_{10}\} = 0$. Therefore, B and C must be $\text{SO}(16)$ singlets. Since the Berman relations for $j = 2, \dots, 10$ are satisfied by construction, the critical Berman relations are the two remaining ones which read

$$[x_1, [x_1, x_2]] = -x_2, \quad [x_2, [x_2, x_1]] = -x_1 \tag{5.6}$$

These follow if

$$\{BC, X_2\} - 2BX_2C \stackrel{!}{=} -X_2, \quad \{CB, X_2\} - 2CX_2B \stackrel{!}{=} -X_2 \tag{5.7}$$

and

$$[X_2, [X_2, B]] \stackrel{!}{=} -B, \quad [X_2, [X_2, C]] \stackrel{!}{=} -C. \tag{5.8}$$

If x_1 simply rotated the two chiralities, we would obtain that $B = -C = \frac{1}{2}$. While this solves (5.7), it clearly does not solve (5.8) and therefore this is not a viable solution. From the physics perspective this is clear, because the Berman generator x_1 is associated to a spatial $\mathfrak{so}(2)$ rotation, whereas the chirality refers to the behaviour under a non-compact space-time $\mathfrak{so}(1, 1)$ (the group $\text{SO}(1, 1)$ scales the chiral and antichiral parts with inverse factors).

This is borne out in the example of spin- $\frac{1}{2}$ where one check that the equations (5.7) and (5.8) are indeed satisfied with $B = -C = -\frac{1}{2}\gamma^2$ and $X_2 = \frac{1}{2}\gamma^{23}$. For the higher spin examples $s = \frac{3}{2}, \frac{5}{2}, \frac{7}{2}$ we can similarly read off the solution from the explicit formulas of the foregoing section, with the result

$$B = -C = -2X(\alpha_1) \otimes \gamma^2 \tag{5.9}$$

where α_1 is the over-extended (left-most) simple root in the E_{10} Dynkin diagram. This formula is valid for all representations of $K(\mathfrak{e}_{10})$ constructed as in (4.6). The relations (5.7) and (5.8) are then implied by the properties of the tensor $X(\alpha_1)$ together with $X_2 = -2X(\alpha_2) \otimes \gamma^{23}$.

For any given unfaithful representation of $K(E_9)$ the problem of uplifting it to an unfaithful representation of $K(E_{10})$ is thus reduced to finding finite-dimensional matrices B and C solving the relations (5.7) and (5.8). Consider for example the (chiral) $K(\mathfrak{e}_9)$ representation that consists of $\mathfrak{V}_0 \cong \mathbf{16}$ and $\mathfrak{V}_1 \cong \mathbf{128}_c$. Doubling this chiral representation leads to a 288-dimensional space that can *not* be turned into a $K(\mathfrak{e}_{10})$ representation, i.e., no x_1 satisfying the relations (5.6) exists, because the spin- $\frac{3}{2}$ representation of $K(\mathfrak{e}_{10})$ requires a second $\mathfrak{V}_2 \cong \mathbf{16}$ in the chiral $K(\mathfrak{e}_9)$ representation [45]. We therefore see from the explicit examples that it is by no means trivial that matrices B and C exist, and it is furthermore quite possible that a more general ansatz than (5.9) may be needed.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] A. Kleinschmidt, R. Köhl, R. Lautenbacher and H. Nicolai, *Representations of involutory subalgebras of affine Kac-Moody algebras*, [arXiv:2102.00870](#) [[INSPIRE](#)].
- [2] E. Cremmer and B. Julia, *The SO(8) Supergravity*, *Nucl. Phys. B* **159** (1979) 141.
- [3] B. de Wit and H. Nicolai, *$d = 11$ Supergravity With Local SU(8) Invariance*, *Nucl. Phys. B* **274** (1986) 363 [[INSPIRE](#)].
- [4] M.J. Duff and K.S. Stelle, *Multimembrane solutions of $D = 11$ supergravity*, *Phys. Lett. B* **253** (1991) 113 [[INSPIRE](#)].
- [5] M.J. Duff and J.T. Liu, *Hidden space-time symmetries and generalized holonomy in M-theory*, *Nucl. Phys. B* **674** (2003) 217 [[hep-th/0303140](#)] [[INSPIRE](#)].
- [6] C. Hull, *Holonomy and symmetry in M-theory*, [hep-th/0305039](#) [[INSPIRE](#)].
- [7] H. Lü, C.N. Pope and K.S. Stelle, *Generalised holonomy for higher-order corrections to supersymmetric backgrounds in string and M-theory*, *Nucl. Phys. B* **741** (2006) 17 [[hep-th/0509057](#)] [[INSPIRE](#)].
- [8] M. Graña, *Flux compactifications in string theory: A Comprehensive review*, *Phys. Rept.* **423** (2006) 91 [[hep-th/0509003](#)] [[INSPIRE](#)].
- [9] M. Graña, R. Minasian, M. Petrini and A. Tomasiello, *Generalized structures of $N = 1$ vacua*, *JHEP* **11** (2005) 020 [[hep-th/0505212](#)] [[INSPIRE](#)].
- [10] M. Gabella, J.P. Gauntlett, E. Palti, J. Sparks and D. Waldram, *AdS_5 Solutions of Type IIB Supergravity and Generalized Complex Geometry*, *Commun. Math. Phys.* **299** (2010) 365 [[arXiv:0906.4109](#)] [[INSPIRE](#)].
- [11] A. Coimbra, C. Strickland-Constable and D. Waldram, *Supersymmetric Backgrounds and Generalised Special Holonomy*, *Class. Quant. Grav.* **33** (2016) 125026 [[arXiv:1411.5721](#)] [[INSPIRE](#)].
- [12] A. Coimbra and C. Strickland-Constable, *Supersymmetric Backgrounds, the Killing Superalgebra, and Generalised Special Holonomy*, *JHEP* **11** (2016) 063 [[arXiv:1606.09304](#)] [[INSPIRE](#)].
- [13] P.C. West, *E_{11} , SL(32) and central charges*, *Phys. Lett. B* **575** (2003) 333 [[hep-th/0307098](#)] [[INSPIRE](#)].
- [14] A. Keurentjes, *The Topology of U duality (sub)groups*, *Class. Quant. Grav.* **21** (2004) 1695 [[hep-th/0309106](#)] [[INSPIRE](#)].
- [15] S. de Buyl, M. Henneaux and L. Paulot, *Hidden symmetries and Dirac fermions*, *Class. Quant. Grav.* **22** (2005) 3595 [[hep-th/0506009](#)] [[INSPIRE](#)].
- [16] T. Damour, A. Kleinschmidt and H. Nicolai, *Hidden symmetries and the fermionic sector of eleven-dimensional supergravity*, *Phys. Lett. B* **634** (2006) 319 [[hep-th/0512163](#)] [[INSPIRE](#)].
- [17] S. de Buyl, M. Henneaux and L. Paulot, *Extended E_8 invariance of 11-dimensional supergravity*, *JHEP* **02** (2006) 056 [[hep-th/0512292](#)] [[INSPIRE](#)].
- [18] T. Damour, A. Kleinschmidt and H. Nicolai, *$K(E_{10})$, Supergravity and Fermions*, *JHEP* **08** (2006) 046 [[hep-th/0606105](#)] [[INSPIRE](#)].
- [19] P. Harring and R. Köhl, *Fundamental groups of split real Kac-Moody groups and generalized flag manifolds*, [arXiv:1905.13444](#).

- [20] A. Kleinschmidt and H. Nicolai, *IIA and IIB spinors from $K(E_{10})$* , *Phys. Lett. B* **637** (2006) 107 [[hep-th/0603205](#)] [[INSPIRE](#)].
- [21] A. Kleinschmidt, H. Nicolai and A. Viganò, *On spinorial representations of involutory subalgebras of Kac-Moody algebras*, [arXiv:1811.11659](#) [[INSPIRE](#)].
- [22] D. Steele and P. West, *E_{11} and Supersymmetry*, *JHEP* **02** (2011) 101 [[arXiv:1011.5820](#)] [[INSPIRE](#)].
- [23] H. Nicolai and H. Samtleben, *On $K(E_9)$* , *Q. J. Pure Appl. Math.* **1** (2005) 180 [[hep-th/0407055](#)] [[INSPIRE](#)].
- [24] A. Kleinschmidt, *Unifying R-symmetry in M-theory*, in *15th International Congress on Mathematical Physics*, V. Sidoravičius ed., Springer (2009) [[DOI](#)] [[hep-th/0703262](#)] [[INSPIRE](#)].
- [25] A. Kleinschmidt and H. Nicolai, *On higher spin realizations of $K(E_{10})$* , *JHEP* **08** (2013) 041 [[arXiv:1307.0413](#)] [[INSPIRE](#)].
- [26] A. Kleinschmidt and H. Nicolai, *Higher spin representations of $K(E_{10})$* , in *International Workshop on Higher Spin Gauge Theories*, pp. 25–38, L. Brink, M. Henneaux and M. Vasiliev eds., World Scientific (2017) [[DOI](#)] [[arXiv:1602.04116](#)] [[INSPIRE](#)].
- [27] B. Julia, *Group disintegrations*, *Conf. Proc. C* **8006162** (1980) 331 [[INSPIRE](#)].
- [28] B. Julia, *Kac-Moody Symmetry of Gravitation and Supergravity Theories*, in *Applications of Group Theory in Physics and Mathematical Physics*, M. Flato, P. Sally and G. Zuckerman eds., Lectures in Applied Mathematics, vol. 21, Am. Math. Soc., Providence (1985), pp. 355–374 [[LPTENS-82-22](#)] [[INSPIRE](#)].
- [29] H. Nicolai, *The Integrability of $N = 16$ Supergravity*, *Phys. Lett. B* **194** (1987) 402 [[INSPIRE](#)].
- [30] T. Damour, M. Henneaux and H. Nicolai, *E_{10} and a ‘small tension expansion’ of M-theory*, *Phys. Rev. Lett.* **89** (2002) 221601 [[hep-th/0207267](#)] [[INSPIRE](#)].
- [31] A. Kleinschmidt, H. Nicolai and N.K. Chidambaram, *Canonical structure of the E_{10} model and supersymmetry*, *Phys. Rev. D* **91** (2015) 085039 [[arXiv:1411.5893](#)] [[INSPIRE](#)].
- [32] T. Damour and C. Hillmann, *Fermionic Kac-Moody Billiards and Supergravity*, *JHEP* **08** (2009) 100 [[arXiv:0906.3116](#)] [[INSPIRE](#)].
- [33] G. Hainke, R. Köhl and P. Levy, *Generalized spin representations*, with an appendix by M. Horn and R. Köhl, *Münster J. Math.* **8** (2015) 181 [[arXiv:1110.5576](#)].
- [34] R. Lautenbacher and R. Köhl *Extending generalized spin representations*, *J. Lie Theory* **28** (2018) 915 [[arXiv:1705.00118](#)].
- [35] O. Hohm and H. Samtleben, *The many facets of exceptional field theory*, *PoS CORFU2018* (2019) 098 [[arXiv:1905.08312](#)] [[INSPIRE](#)].
- [36] D.S. Berman and C.D.A. Blair, *The Geometry, Branes and Applications of Exceptional Field Theory*, *Int. J. Mod. Phys. A* **35** (2020) 2030014 [[arXiv:2006.09777](#)] [[INSPIRE](#)].
- [37] H. Godazgar, M. Godazgar, O. Hohm, H. Nicolai and H. Samtleben, *Supersymmetric $E_{7(7)}$ Exceptional Field Theory*, *JHEP* **09** (2014) 044 [[arXiv:1406.3235](#)] [[INSPIRE](#)].
- [38] A. Baguet and H. Samtleben, *$E_{8(8)}$ Exceptional Field Theory: Geometry, Fermions and Supersymmetry*, *JHEP* **09** (2016) 168 [[arXiv:1607.03119](#)] [[INSPIRE](#)].

- [39] G. Bossard, A. Kleinschmidt and E. Sezgin, *On supersymmetric E_{11} exceptional field theory*, *JHEP* **10** (2019) 165 [[arXiv:1907.02080](#)] [[INSPIRE](#)].
- [40] H. Nicolai, *$D = 11$ Supergravity With Local $SO(16)$ Invariance*, *Phys. Lett. B* **187** (1987) 316 [[INSPIRE](#)].
- [41] P.C. West, *E_{11} and M-theory*, *Class. Quant. Grav.* **18** (2001) 4443 [[hep-th/0104081](#)] [[INSPIRE](#)].
- [42] E. Bergshoeff, E. Sezgin and P.K. Townsend, *Supermembranes and Eleven-Dimensional Supergravity*, *Phys. Lett. B* **189** (1987) 75 [[INSPIRE](#)].
- [43] B. de Wit, J. Hoppe and H. Nicolai, *On the Quantum Mechanics of Supermembranes*, *Nucl. Phys. B* **305** (1988) 545 [[INSPIRE](#)].
- [44] P. Breitenlohner and D. Maison, *On the Geroch Group*, *Ann. Inst. H. Poincaré Phys. Theor.* **46** (1987) 215 [[INSPIRE](#)].
- [45] A. Kleinschmidt, H. Nicolai and J. Palmkvist, *$K(E_9)$ from $K(E_{10})$* , *JHEP* **06** (2007) 051 [[hep-th/0611314](#)] [[INSPIRE](#)].
- [46] S. Berman, *On generators and relations for certain involutory subalgebras of Kac-Moody Lie algebras*, *Comm. Algebra* **17** (1989) 3165.