

Evaluating four-loop conformal Feynman integrals by D -dimensional differential equations

Burkhard Eden^a and Vladimir A. Smirnov^b

^a*Institut für Mathematik und Physik, Humboldt-Universität zu Berlin,
Zum großen Windkanal 6, 12489 Berlin, Germany*

^b*Skobeltsyn Institute of Nuclear Physics, Moscow State University,
119992 Moscow, Russia*

E-mail: eden@math.hu-berlin.de, smirnov@theory.sinp.msu.ru

ABSTRACT: We evaluate a four-loop conformal integral, i.e. an integral over four four-dimensional coordinates, by turning to its dimensionally regularized version and applying differential equations for the set of the corresponding 213 master integrals. To solve these linear differential equations we follow the strategy suggested by Henn and switch to a uniformly transcendental basis of master integrals. We find a solution to these equations up to weight eight in terms of multiple polylogarithms. Further, we present an analytical result for the given four-loop conformal integral considered in four-dimensional space-time in terms of single-valued harmonic polylogarithms. As a by-product, we obtain analytical results for all the other 212 master integrals within dimensional regularization, i.e. considered in D dimensions.

KEYWORDS: Conformal Field Theory, Scattering Amplitudes, Supersymmetric gauge theory

ARXIV EPRINT: [1607.06427](https://arxiv.org/abs/1607.06427)

Contents

1	Introduction	1
2	Master integrals and differential equations	3
3	Solving canonical differential equations	5
4	Conclusions	7

1 Introduction

Correlation functions of half-BPS operators in $\mathcal{N} = 4$ super Yang-Mills (SYM) theory — in particular of the stress-tensor multiplet — have been extensively studied because their strong coupling regime is accessible via the AdS/CFT correspondence. Subsequently, the integrability of the model’s spectrum problem was discovered, and in an initially unrelated effort, many results were obtained for scattering amplitudes and a dual set of Wilson loops. These two developments were brought together by the construction of an integrable system [1, 2] for the so-called remainder function in amplitudes.

The stress-tensor correlators came back to center stage when it became clear that they can act as generating objects for both scattering amplitudes and the dual polygonal Wilson loops [3, 4]. It is then a natural question whether these correlators can also be analyzed from an integrable systems perspective. In [5] such ideas have been put forward for general three-point functions in weak coupling perturbation theory. Unfortunately, comparison to perturbative “data” is possible only indirectly because it is a fairly hard task to obtain exact field theory results for non-trivial three-point functions at higher loops. However, three-point couplings for two half-BPS and one twist operator are available from OPE limits of half-BPS four-point functions. The explicit result for the two-loop four-point stress-tensor correlator from a decade ago [6, 7] has been a guideline for the construction of the “hexagon proposal” of [5], and more recent work [8–11] for the three-loop part of the four-point function has been successfully compared to the hexagon prediction [12, 13]; a vital test of the proposal.

At the next order it is not yet clear how to handle the hexagon due to problems with a double pole. On the field theory side, the integrand of the four-point function of stress-tensor multiplets has been elaborated in [8, 14–16] up to eight loops. It takes the form of a kinematic factor [17] times a sum of scalar conformal integrals in a propagator representation. At four loops (and beyond), the largest part of the integrals has not yet been evaluated: there are 26 genuine four-loop integrals in the planar part of the correlator (so the part relevant to integrability), of which five can be related to the ladder with four rungs by flip identities on subintegrals. One further integral could be solved in [10] as it




Figure 1. A four-loop conformal integral.

obeys a Laplace equation, see below. The recent paper [18] considers the leading terms in an asymptotic expansion for the entire set of integrals.

A possible way to exactly evaluate conformal integrals is to follow the idea advocated in [19] and to try to perform integration over Feynman parameters in an appropriate order. If it turns out that there is an order in which the dependence of the denominator of the integrand on the Feynman parameters is linear then the whole integral can be solved in terms of multiple polylogarithms. This strategy was successfully applied for example in [20–24] and implemented as the computer code `HyperInt` in [25].

We initiate here the study of the remaining 20 integrals by the method of differential equations, choosing at will the simplest looking diagram. The goal of this paper is thus to evaluate the coordinate-space Feynman integral associated with the graph of figure 1. Although this integral is linearly reducible, in the sense of [19], and an analytical result can be obtained with `HyperInt` in [25], we are going to evaluate it with differential equations, keeping in mind that many cases in our set of 20 remaining four-loop conformal integrals will be linearly irreducible, although knowledge on polynomial reduction is constantly increasing. Indeed, it is helpful to be able to test our result against a different method.

The previous paper [10] chiefly aimed at three loops, but it also contains the aforementioned application of the Laplace equation to one of the 26 four-loop integrals. Upon flipping a subintegral, one external vertex is connected to the rest of this diagram only by a single line. Acting by the operator $\square = \frac{\partial}{\partial x_\mu} \frac{\partial}{\partial x^\mu}$, where x is the four-coordinate of this vertex, is described in graph-theoretical language as the contraction of the corresponding line, so that we easily obtain a differential equation in four dimensions.

In figure 1, conformal invariance allows us to send the point x_4 to infinity to reduce to the Feynman integral whose graph is shown in figure 2. Here two lines are incident to each of the three external vertices so that we cannot derive a differential equation of the same type as in the previous case.

The approach based on differential equations and integration by parts substantially enlarges any problem, because instead of a single integral one considers a family of master integrals closed under IBP relations. A four-dimensional version of the method [26] is




Figure 2. A vertex integral obtained from our four-loop conformal integral.

more economical with respect to the size of the basis of integrals. However, an obvious complication for our set of conformal integrals is that there are potentially ultraviolet and infrared divergences, and it is not straightforward to choose a subspace of Feynman integrals which includes a given conformal integral and other finite integrals (both in the ultraviolet and infrared sense) closed under four-dimensional integration by parts relations.

We believe that the strategy outlined in [26] can be adjusted to the case of conformal integrals as well. In this paper, however, we decided to employ the well-known straightforward technique within dimensional regularization based on D -dimensional integration by parts [27] and differential equations [28–34].

On the one hand, we certainly make the situation more complicated, because dimensional regularization means involving a fairly large number of integrals. Indeed, as we will see shortly, there are 213 master integrals in the corresponding extended family of Feynman integrals; the integral of figure 2 is only one of them. On the other hand, we obtain the possibility to apply the very well-known powerful machinery of differential equations.

To solve these linear differential equations we follow the strategy suggested by Henn [34] (see also [35]) and first applied in [36–38] and then in many other papers. To this end we switch to a uniformly transcendental basis of master integrals. We obtain a solution to these equations up to weight eight in terms of multiple polylogarithms [39]. An analytical result for the given four-loop conformal integral considered in four-dimensional space-time is given in terms of harmonic polylogarithms [40] of weight eight. This result can naturally be represented in terms of single-valued harmonic polylogarithms [41]. As a by-product we obtain analytical results for all the other 212 master integrals considered in D dimensions.

In the next section, we present definitions, describe master integrals and differential equations and explain how we arrived at a canonical basis, in the sense of [34]. In section 3, we solve the differential equations in our canonical basis and describe our results. We discuss perspectives in the conclusions.

2 Master integrals and differential equations

Since after going to $D = 4 - 2\epsilon$ dimensions we have to deal with a complete set of integrals closed under integration by parts relations, we introduce eight more numerators (in the form of propagators) in addition to the ten existing propagators. Thus, we arrive at the

following family of integrals

$$\begin{aligned}
 F_{a_1, \dots, a_{18}} = & \int \cdots \int \frac{d^D x_5 d^D x_6 d^D x_7 d^D x_8}{[-x_5^2]^{a_1} [-x_6^2]^{a_2} [-(x_1 - x_5)^2]^{a_3} [-(x_1 - x_7)^2]^{a_4} [-(x_2 - x_6)^2]^{a_5}} \\
 & \times \frac{[-(x_2 - x_5)^2]^{-a_{11}} [-(x_1 - x_6)^2]^{-a_{12}} [-(x_2 - x_7)^2]^{-a_{13}} [-(x_6 - x_7)^2]^{-a_{14}} [-x_7^2]^{-a_{15}}}{[-(x_2 - x_8)^2]^{a_6} [-(x_5 - x_6)^2]^{a_7} [-(x_5 - x_7)^2]^{a_8}} \\
 & \times \frac{[-(x_1 - x_8)^2]^{-a_{16}} [-(x_5 - x_8)^2]^{-a_{17}} [-x_8^2]^{-a_{18}}}{[-(x_6 - x_8)^2]^{a_9} [-(x_7 - x_8)^2]^{a_{10}}}.
 \end{aligned} \tag{2.1}$$

The powers of the propagators (indices) a_i are integer and $a_i \leq 0$ for $i \geq 11$. Although the situation is of Euclidean type we prefer to deal with propagators in Minkowski space, with $x^2 = x_0^2 - \vec{x}^2$ because some computer codes are oriented at Minkowski propagators.

This is a family of vertex Feynman integrals depending on x_1, x_2 and x_3 . In the last formula translation invariance was used to set x_3 to zero. We put $(x_1 - x_2)^2 = -1$ and introduce standard conformal variables by

$$x_1^2 = -z\bar{z}, x_2^2 = -(1-z)(1-\bar{z}). \tag{2.2}$$

Using FIRE [42–44] combined with LiteRed [45] we reveal 213 master integrals. In particular, there are four master integrals in the top sector, i.e. with all the first ten indices positive. The integral number 210 is the integral $F_{1, \dots, 1, 0, \dots}$ equal to the conformal integral in figure 2 which was our starting point. The full list of primary master integrals is present in a file attached to this paper.

The derivation of differential equations for a family of master integrals is a straightforward procedure. We take derivatives of the master integrals in z and \bar{z} with the help of the package LiteRed [45] and then apply FIRE to reduce the resulting integrals to master integrals. As a result we obtain two systems of linear differential equations.

$$\frac{\partial}{\partial z} f = A_1(z, \bar{z}, \epsilon) f, \tag{2.3}$$

$$\frac{\partial}{\partial \bar{z}} f = A_2(z, \bar{z}, \epsilon) f, \tag{2.4}$$

where f is the vector of primary master integrals and A_1, A_2 are 213×213 -matrices.

We follow the strategy suggested in [34] and turn to a new basis where the differential equations take the form

$$\frac{\partial}{\partial z} f = \epsilon \bar{A}_1(z, \bar{z}) f, \tag{2.5}$$

$$\frac{\partial}{\partial \bar{z}} f = \epsilon \bar{A}_2(z, \bar{z}) f, \tag{2.6}$$

where the matrices \bar{A}_1, \bar{A}_2 are independent of ϵ .

In differential form, we have

$$d f(z, \bar{z}) = \epsilon (d \tilde{A}(z, \bar{z})) f(z, \bar{z}),$$

where

$$\tilde{A} = \sum_k \tilde{A}_{\alpha_k} \log(\alpha_k). \tag{2.7}$$

The matrices \tilde{A}_{α_k} are *constant* matrices and the arguments of the logarithms α_i (*letters*) are functions of z, \bar{z} . One calls this form of differential equations canonical.

In our case, the list of letters is

$$\{z, 1 - z, \bar{z}, 1 - \bar{z}, -z + \bar{z}, 1 - z - \bar{z}, 1 - z\bar{z}, z + \bar{z} - z\bar{z}\}. \tag{2.8}$$

In the case of one variable there exists an algorithm [46] which provides the possibility of arriving at a canonical basis.¹ In our case there are two variables. We followed recipes formulated in [34, 35] and successfully applied in [36–38, 47] and many other papers. In particular, one tries to choose basis integrals that have constant leading singularities² [48]. Sometimes we also used small additional rotations of the basis to ‘integrate out’ terms with an ϵ^0 dependence, as it was done in many cases, see e.g. [36–38, 47].

We have implemented these recipes in a code which we successfully applied in the case at hand. A description of this code will be given in a future publication together with results on more complicated four-loop conformal integrals. Our canonical basis is presented in the supplementary material attached to this paper.

3 Solving canonical differential equations

We solve the linear system (2.5), (2.6), in a power expansion in ϵ according to the strategy described in detail in [49, 50]. We solve the first linear system, eq. (2.5), which results into multiple polylogarithms of the argument z , up to unknown functions of \bar{z} .

The multiple polylogarithms are defined recursively by

$$G(a_1, \dots, a_n; z) = \int_0^z \frac{dt}{t - a_1} G(a_2, \dots, a_n; t) \tag{3.1}$$

with $a_i, z \in \mathbb{C}$ and $G(z) = 1$. In the special case where $a_i = 0$ for all i one has by definition

$$G(0, \dots, 0; x) = \frac{1}{n!} \ln^n x. \tag{3.2}$$

Then we substitute this solution into the second system, eq. (2.6), check that the dependence on z drops out and solve the resulting linear system depending only on \bar{z} , up to constants of weights $w \leq 8$. To fix these 213×9 unknown constants, we match our results for the canonical basis in terms of multiple polylogarithms to the leading order asymptotic behaviour of the solution of eqs. (2.5), (2.6) in the Euclidean limit $x_1 \rightarrow 0$, or equivalently $z, \bar{z} \rightarrow 0$, found by solving the differential equations in the limit.

¹In fact, some parts of it can be applied in the case of several variables as well.

²Leading singularities are multidimensional residues of the integrand. They determine rational factors in front of the otherwise logarithmic functions in explicit expressions for Feynman integrals.

The corresponding terms of the expansion can be written in the well-known graph-theoretical language — see, e.g., [51]. Alternatively, they can be described in the language of expansion by regions [51, 52]. The crucial point in the matching is that the leading-order contributions are classified according to the power dependence on the small parameter of the limit, i.e. $\sim z\bar{z}$. This parameter enters with powers of the form $-k\epsilon$ where $k = 0, 1, 2, 3$, and this is seen both from the point of view of differential equations and expansion by regions.

Upon fixing all the constants in our solution we obtain analytic results for all the 213 elements of the canonical basis. These are presented in two files (for contributions up to weight 7 and of weight 8, respectively) which can be downloaded from: <http://theory.sinp.msu.ru/~smirnov/ci4>.

Element number 210 of the basis is

$$f_{210} = \epsilon^8 (z - \bar{z})^2 F_{1,\dots,1,0,\dots} \tag{3.3}$$

Its ϵ -expansion starts from order ϵ^8 . We obtain the following result for the original conformal integral

$$\begin{aligned}
 F_{1,\dots,1,0,\dots} &= \frac{4}{(z - \bar{z})^2} \times \tag{3.4} \\
 &[-\mathcal{L}_{\{3,5\}} + \mathcal{L}_{\{5,3\}} - \mathcal{L}_{\{1,2,5\}} + \mathcal{L}_{\{1,3,4\}} + \mathcal{L}_{\{1,4,3\}} - \mathcal{L}_{\{1,5,2\}} \\
 &- \mathcal{L}_{\{2,1,5\}} + \mathcal{L}_{\{2,5,0\}} + \mathcal{L}_{\{2,5,1\}} - \mathcal{L}_{\{3,1,4\}} - \mathcal{L}_{\{3,4,1\}} + \mathcal{L}_{\{4,1,3\}} \\
 &- \mathcal{L}_{\{4,3,0\}} - \mathcal{L}_{\{4,3,1\}} + \mathcal{L}_{\{5,1,2\}} + \mathcal{L}_{\{5,2,1\}} - \mathcal{L}_{\{1,1,1,5\}} + \mathcal{L}_{\{1,1,2,4\}} \\
 &- \mathcal{L}_{\{1,1,4,2\}} + \mathcal{L}_{\{1,1,5,0\}} + \mathcal{L}_{\{1,1,5,1\}} - \mathcal{L}_{\{1,2,3,2\}} - \mathcal{L}_{\{1,2,4,1\}} + 2\mathcal{L}_{\{1,3,1,3\}} \\
 &+ \mathcal{L}_{\{1,4,2,1\}} - \mathcal{L}_{\{1,5,0,0\}} - \mathcal{L}_{\{1,5,1,0\}} - \mathcal{L}_{\{1,5,1,1\}} - \mathcal{L}_{\{2,1,1,4\}} - \mathcal{L}_{\{2,1,2,3\}} \\
 &+ \mathcal{L}_{\{2,1,4,0\}} + \mathcal{L}_{\{2,3,2,1\}} + \mathcal{L}_{\{2,4,1,0\}} + \mathcal{L}_{\{2,4,1,1\}} - \mathcal{L}_{\{3,1,3,0\}} - 2\mathcal{L}_{\{3,1,3,1\}} \\
 &+ \mathcal{L}_{\{3,2,1,2\}} + \mathcal{L}_{\{3,3,0,0\}} + \mathcal{L}_{\{3,3,1,0\}} + \mathcal{L}_{\{4,1,1,2\}} - \mathcal{L}_{\{4,1,2,0\}} - \mathcal{L}_{\{4,2,1,0\}} \\
 &- \mathcal{L}_{\{4,2,1,1\}} + \mathcal{L}_{\{5,1,1,1\}} - \mathcal{L}_{\{1,1,1,2,3\}} - \mathcal{L}_{\{1,1,1,3,2\}} + \mathcal{L}_{\{1,1,1,4,0\}} + \mathcal{L}_{\{1,1,2,1,3\}} \\
 &+ \mathcal{L}_{\{1,1,2,3,0\}} + \mathcal{L}_{\{1,1,2,3,1\}} - \mathcal{L}_{\{1,1,3,1,2\}} + \mathcal{L}_{\{1,1,3,2,1\}} - \mathcal{L}_{\{1,1,4,0,0\}} + \mathcal{L}_{\{1,2,1,1,3\}} \\
 &- \mathcal{L}_{\{1,2,1,3,1\}} - \mathcal{L}_{\{1,2,3,1,1\}} + \mathcal{L}_{\{1,3,1,1,2\}} + \mathcal{L}_{\{1,3,1,2,1\}} - \mathcal{L}_{\{1,3,2,1,0\}} - \mathcal{L}_{\{1,3,2,1,1\}} \\
 &- \mathcal{L}_{\{1,4,1,0,0\}} - \mathcal{L}_{\{1,4,1,1,0\}} - \mathcal{L}_{\{2,1,1,1,3\}} - \mathcal{L}_{\{2,1,1,3,1\}} + \mathcal{L}_{\{2,1,3,0,0\}} + 2\mathcal{L}_{\{2,1,3,1,0\}} \\
 &+ \mathcal{L}_{\{2,1,3,1,1\}} - \mathcal{L}_{\{2,2,1,2,0\}} - \mathcal{L}_{\{2,3,0,0,0\}} - \mathcal{L}_{\{2,3,1,0,0\}} + \mathcal{L}_{\{2,3,1,1,1\}} + \mathcal{L}_{\{3,1,1,1,2\}} \\
 &- \mathcal{L}_{\{3,1,1,2,0\}} - \mathcal{L}_{\{3,1,1,2,1\}} + \mathcal{L}_{\{3,1,2,0,0\}} - \mathcal{L}_{\{3,1,2,1,1\}} + \mathcal{L}_{\{3,2,1,0,0\}} + \mathcal{L}_{\{3,2,1,1,0\}} \\
 &+ \mathcal{L}_{\{3,2,1,1,1\}} - \mathcal{L}_{\{4,1,1,1,0\}} + \mathcal{L}_{\{1,1,1,1,3,0\}} - \mathcal{L}_{\{1,1,1,2,1,2\}} + \mathcal{L}_{\{1,1,1,3,1,0\}} + \mathcal{L}_{\{1,1,2,1,2,1\}} \\
 &- \mathcal{L}_{\{1,1,3,0,0,0\}} - 2\mathcal{L}_{\{1,1,3,1,0,0\}} - \mathcal{L}_{\{1,1,3,1,1,0\}} + \mathcal{L}_{\{1,2,1,1,1,2\}} + \mathcal{L}_{\{1,2,1,2,0,0\}} - \mathcal{L}_{\{1,2,1,2,1,1\}} \\
 &+ \mathcal{L}_{\{1,3,0,0,0,0\}} + \mathcal{L}_{\{1,3,1,0,0,0\}} - \mathcal{L}_{\{1,3,1,1,1,0\}} - \mathcal{L}_{\{2,1,1,1,2,0\}} - \mathcal{L}_{\{2,1,1,1,2,1\}} + \mathcal{L}_{\{2,1,1,2,0,0\}} \\
 &+ \mathcal{L}_{\{2,1,1,2,1,0\}} - \mathcal{L}_{\{2,1,2,0,0,0\}} + \mathcal{L}_{\{2,1,2,1,1,0\}} + \mathcal{L}_{\{2,1,2,1,1,1\}} - \mathcal{L}_{\{2,2,1,0,0,0\}} - \mathcal{L}_{\{2,2,1,1,0,0\}} \\
 &- \mathcal{L}_{\{2,2,1,1,1,0\}} + \mathcal{L}_{\{3,1,1,1,0,0\}} + \mathcal{L}_{\{1,1,1,1,2,0,0\}} + \mathcal{L}_{\{1,1,1,1,2,1,0\}} - \mathcal{L}_{\{1,1,1,2,0,0,0\}} \\
 &- \mathcal{L}_{\{1,1,1,2,1,0,0\}} + \mathcal{L}_{\{1,1,2,0,0,0,0\}} - \mathcal{L}_{\{1,1,2,1,1,0,0\}} - \mathcal{L}_{\{1,1,2,1,1,1,0\}} + \mathcal{L}_{\{1,2,1,0,0,0,0\}} \\
 &+ \mathcal{L}_{\{1,2,1,1,0,0,0\}} + \mathcal{L}_{\{1,2,1,1,1,0,0\}} - \mathcal{L}_{\{2,1,1,1,0,0,0\}} + \mathcal{L}_{\{1,1,1,1,0,0,0,0\}}]
 \end{aligned}$$

which is, of course, of weight 8. In the last formula \mathcal{L} denotes a single-valued harmonic polylogarithm

$$\mathcal{L}_{\{a_1, \dots, a_8\}} = (-1)^{\sum a_i} G(a_1, \dots, a_8; z) + \sum c_{ij} G(\underline{a}_i; z) G(\underline{a}_j; \bar{z}) \quad (3.5)$$

where $\underline{a}_i \cup \underline{a}_j$ has length 8 and \underline{a}_j is never the empty word. The coefficients c_{ij} are polynomials of multiple zeta values such that all branch cuts cancel. The entries in the weight vectors are in the set $\{0, 1\}$ and we are using the “condensed notation” $\dots 0, 0, 0, 1 \dots = \dots 4 \dots$ etc. These are the SVHPL’s introduced by Brown in [41].

Note that (3.4) takes a much simpler form after flipping points $x_2 \leftrightarrow x_3$ (i.e. $z \rightarrow 1/z$, $\bar{z} \rightarrow 1/\bar{z}$) followed by $x_1 \leftrightarrow x_2$ (which implies $z \rightarrow 1-z$, $\bar{z} \rightarrow 1-\bar{z}$): the pure function in the square bracket transforms as

$$[\dots] \rightarrow -\mathcal{L}_{\{3,5\}} + \mathcal{L}_{\{5,3\}} + \mathcal{L}_{\{2,5,0\}} - \mathcal{L}_{\{4,3,0\}} - \mathcal{L}_{\{1,5,0,0\}} + \mathcal{L}_{\{3,3,0,0\}} - \mathcal{L}_{\{2,3,0,0,0\}} + \mathcal{L}_{\{1,3,0,0,0,0\}} \cdot \quad (3.6)$$

We checked our result (3.4) by a numerical calculation with FIESTA [53], as well as those for some other elements in the basis. Our result (3.4) is in agreement with a calculation by different means [54] about which we knew in advance and with a calculation³ based on HyperInt [25].

4 Conclusions

To evaluate a four-loop conformal integral we applied powerful techniques designed for dimensionally regularized Feynman integrals, although we still believe that one can develop an efficient purely four-dimensional technique to evaluate conformal integrals. We have also evaluated a family of four-loop vertex master integrals with all the differences of the external coordinates off the light-cone. The corresponding dual momentum space Feynman integrals are vertex integrals with all the end-points off the light-cone. We believe that this is a first example of a complete calculation of such a family of vertex integrals at the level of four loops. We hope to report on further results on four-loop conformal integrals obtained with the help of D -dimensional differential equations.

Acknowledgments

We thank Johannes Henn for various pieces of advice, Oliver Schnetz for comparison of our results with his which were obtained by an independent technique, and Johannes Broedel and Claude Duhr for help in performing computer manipulations with multiple polylogarithms. We thank Paul Heslop, Vladimir Mitev, Erik Panzer and Oliver Schnetz for careful reading of the draft of the paper. VS is grateful to Matthias Staudacher for kind hospitality at the Humboldt University of Berlin where this project started. The work of VS was partially supported by the Alexander von Humboldt Foundation (Humboldt Forschungspreis). BE acknowledges support by SFB 647 of the DFG, and the Cluster of Excellence “Image, Knowledge, Gestaltung” at Humboldt-University Berlin, funded by the Excellence Initiative and DFG.

³Thanks to Erik Panzer for the comparison.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] L.F. Alday, D. Gaiotto and J. Maldacena, *Thermodynamic Bubble Ansatz*, *JHEP* **09** (2011) 032 [[arXiv:0911.4708](https://arxiv.org/abs/0911.4708)] [[INSPIRE](#)].
- [2] B. Basso, A. Sever and P. Vieira, *Spacetime and Flux Tube S-Matrices at Finite Coupling for $N = 4$ Supersymmetric Yang-Mills Theory*, *Phys. Rev. Lett.* **111** (2013) 091602 [[arXiv:1303.1396](https://arxiv.org/abs/1303.1396)] [[INSPIRE](#)].
- [3] L.F. Alday, B. Eden, G.P. Korchemsky, J. Maldacena and E. Sokatchev, *From correlation functions to Wilson loops*, *JHEP* **09** (2011) 123 [[arXiv:1007.3243](https://arxiv.org/abs/1007.3243)] [[INSPIRE](#)].
- [4] B. Eden, G.P. Korchemsky and E. Sokatchev, *From correlation functions to scattering amplitudes*, *JHEP* **12** (2011) 002 [[arXiv:1007.3246](https://arxiv.org/abs/1007.3246)] [[INSPIRE](#)].
- [5] B. Basso, S. Komatsu and P. Vieira, *Structure Constants and Integrable Bootstrap in Planar $N = 4$ SYM Theory*, [arXiv:1505.06745](https://arxiv.org/abs/1505.06745) [[INSPIRE](#)].
- [6] B. Eden, C. Schubert and E. Sokatchev, *Three loop four point correlator in $N = 4$ SYM*, *Phys. Lett. B* **482** (2000) 309 [[hep-th/0003096](https://arxiv.org/abs/hep-th/0003096)] [[INSPIRE](#)].
- [7] M. Bianchi, S. Kovacs, G. Rossi and Y.S. Stanev, *Anomalous dimensions in $N = 4$ SYM theory at order g^4* , *Nucl. Phys. B* **584** (2000) 216 [[hep-th/0003203](https://arxiv.org/abs/hep-th/0003203)] [[INSPIRE](#)].
- [8] B. Eden, P. Heslop, G.P. Korchemsky and E. Sokatchev, *Hidden symmetry of four-point correlation functions and amplitudes in $N = 4$ SYM*, *Nucl. Phys. B* **862** (2012) 193 [[arXiv:1108.3557](https://arxiv.org/abs/1108.3557)] [[INSPIRE](#)].
- [9] B. Eden, *Three-loop universal structure constants in $N = 4$ SUSY Yang-Mills theory*, [arXiv:1207.3112](https://arxiv.org/abs/1207.3112) [[INSPIRE](#)].
- [10] J. Drummond, C. Duhr, B. Eden, P. Heslop, J. Pennington and V.A. Smirnov, *Leading singularities and off-shell conformal integrals*, *JHEP* **08** (2013) 133 [[arXiv:1303.6909](https://arxiv.org/abs/1303.6909)] [[INSPIRE](#)].
- [11] D. Chicherin, J. Drummond, P. Heslop and E. Sokatchev, *All three-loop four-point correlators of half-BPS operators in planar $\mathcal{N} = 4$ SYM*, *JHEP* **08** (2016) 053 [[arXiv:1512.02926](https://arxiv.org/abs/1512.02926)] [[INSPIRE](#)].
- [12] B. Eden and A. Sfondrini, *Three-point functions in $\mathcal{N} = 4$ SYM: the hexagon proposal at three loops*, *JHEP* **02** (2016) 165 [[arXiv:1510.01242](https://arxiv.org/abs/1510.01242)] [[INSPIRE](#)].
- [13] B. Basso, V. Gonçalves, S. Komatsu and P. Vieira, *Gluing Hexagons at Three Loops*, *Nucl. Phys. B* **907** (2016) 695 [[arXiv:1510.01683](https://arxiv.org/abs/1510.01683)] [[INSPIRE](#)].
- [14] B. Eden, P. Heslop, G.P. Korchemsky and E. Sokatchev, *Constructing the correlation function of four stress-tensor multiplets and the four-particle amplitude in $N = 4$ SYM*, *Nucl. Phys. B* **862** (2012) 450 [[arXiv:1201.5329](https://arxiv.org/abs/1201.5329)] [[INSPIRE](#)].
- [15] R.G. Ambrosio, B. Eden, T. Goddard, P. Heslop and C. Taylor, *Local integrands for the five-point amplitude in planar $N = 4$ SYM up to five loops*, *JHEP* **01** (2015) 116 [[arXiv:1312.1163](https://arxiv.org/abs/1312.1163)] [[INSPIRE](#)].

- [16] J.L. Bourjaily, P. Heslop and V.-V. Tran, *Perturbation Theory at Eight Loops: Novel Structures and the Breakdown of Manifest Conformality in $N = 4$ Supersymmetric Yang-Mills Theory*, *Phys. Rev. Lett.* **116** (2016) 191602 [[arXiv:1512.07912](#)] [[INSPIRE](#)].
- [17] B. Eden, A.C. Petkou, C. Schubert and E. Sokatchev, *Partial nonrenormalization of the stress tensor four point function in $N = 4$ SYM and AdS/CFT*, *Nucl. Phys. B* **607** (2001) 191 [[hep-th/0009106](#)] [[INSPIRE](#)].
- [18] V. Gonçalves, *Extracting OPE coefficient of Konishi at four loops*, [arXiv:1607.02195](#) [[INSPIRE](#)].
- [19] F. Brown, *The massless higher-loop two-point function*, *Commun. Math. Phys.* **287** (2009) 925 [[arXiv:0804.1660](#)] [[INSPIRE](#)].
- [20] E. Panzer, *On the analytic computation of massless propagators in dimensional regularization*, *Nucl. Phys. B* **874** (2013) 567 [[arXiv:1305.2161](#)] [[INSPIRE](#)].
- [21] E. Panzer, *On hyperlogarithms and Feynman integrals with divergences and many scales*, *JHEP* **03** (2014) 071 [[arXiv:1401.4361](#)] [[INSPIRE](#)].
- [22] A. von Manteuffel, E. Panzer and R.M. Schabinger, *A quasi-finite basis for multi-loop Feynman integrals*, *JHEP* **02** (2015) 120 [[arXiv:1411.7392](#)] [[INSPIRE](#)].
- [23] F. Brown and O. Schnetz, *Proof of the zig-zag conjecture*, [arXiv:1208.1890](#) [[INSPIRE](#)].
- [24] O. Schnetz, *Graphical functions and single-valued multiple polylogarithms*, *Commun. Num. Theor. Phys.* **08** (2014) 589 [[arXiv:1302.6445](#)] [[INSPIRE](#)].
- [25] E. Panzer, *Algorithms for the symbolic integration of hyperlogarithms with applications to Feynman integrals*, *Comput. Phys. Commun.* **188** (2015) 148 [[arXiv:1403.3385](#)] [[INSPIRE](#)].
- [26] S. Caron-Huot and J.M. Henn, *Iterative structure of finite loop integrals*, *JHEP* **06** (2014) 114 [[arXiv:1404.2922](#)] [[INSPIRE](#)].
- [27] K.G. Chetyrkin and F.V. Tkachov, *Integration by Parts: The Algorithm to Calculate β -functions in 4 Loops*, *Nucl. Phys. B* **192** (1981) 159 [[INSPIRE](#)].
- [28] A.V. Kotikov, *Differential equations method: New technique for massive Feynman diagrams calculation*, *Phys. Lett. B* **254** (1991) 158 [[INSPIRE](#)].
- [29] A.V. Kotikov, *Differential equation method: The calculation of N point Feynman diagrams*, *Phys. Lett. B* **267** (1991) 123 [*Erratum ibid.* **B 295** (1992) 409] [[INSPIRE](#)].
- [30] E. Remiddi, *Differential equations for Feynman graph amplitudes*, *Nuovo Cim. A* **110** (1997) 1435 [[hep-th/9711188](#)] [[INSPIRE](#)].
- [31] T. Gehrmann and E. Remiddi, *Differential equations for two loop four point functions*, *Nucl. Phys. B* **580** (2000) 485 [[hep-ph/9912329](#)] [[INSPIRE](#)].
- [32] T. Gehrmann and E. Remiddi, *Two loop master integrals for $\gamma^* \rightarrow 3$ jets: The planar topologies*, *Nucl. Phys. B* **601** (2001) 248 [[hep-ph/0008287](#)] [[INSPIRE](#)].
- [33] T. Gehrmann and E. Remiddi, *Two loop master integrals for $\gamma^* \rightarrow 3$ jets: The nonplanar topologies*, *Nucl. Phys. B* **601** (2001) 287 [[hep-ph/0101124](#)] [[INSPIRE](#)].
- [34] J.M. Henn, *Multiloop integrals in dimensional regularization made simple*, *Phys. Rev. Lett.* **110** (2013) 251601 [[arXiv:1304.1806](#)] [[INSPIRE](#)].
- [35] J.M. Henn, *Lectures on differential equations for Feynman integrals*, *J. Phys. A* **48** (2015) 153001 [[arXiv:1412.2296](#)] [[INSPIRE](#)].

- [36] J.M. Henn, A.V. Smirnov and V.A. Smirnov, *Analytic results for planar three-loop four-point integrals from a Knizhnik-Zamolodchikov equation*, *JHEP* **07** (2013) 128 [[arXiv:1306.2799](#)] [[INSPIRE](#)].
- [37] J.M. Henn and V.A. Smirnov, *Analytic results for two-loop master integrals for Bhabha scattering I*, *JHEP* **11** (2013) 041 [[arXiv:1307.4083](#)] [[INSPIRE](#)].
- [38] J.M. Henn, A.V. Smirnov and V.A. Smirnov, *Evaluating single-scale and/or non-planar diagrams by differential equations*, *JHEP* **03** (2014) 088 [[arXiv:1312.2588](#)] [[INSPIRE](#)].
- [39] A.B. Goncharov, *Multiple polylogarithms, cyclotomy and modular complexes*, *Math. Res. Lett.* **5** (1998) 497 [[arXiv:1105.2076](#)] [[INSPIRE](#)].
- [40] E. Remiddi and J.A.M. Vermaseren, *Harmonic polylogarithms*, *Int. J. Mod. Phys. A* **15** (2000) 725 [[hep-ph/9905237](#)] [[INSPIRE](#)].
- [41] F.C.S. Brown, *Single-valued multiple polylogarithms in one variable*, *Compt. Rendus Math.* **338** (2004) 527.
- [42] A.V. Smirnov, *Algorithm FIRE — Feynman Integral REduction*, *JHEP* **10** (2008) 107 [[arXiv:0807.3243](#)] [[INSPIRE](#)].
- [43] A.V. Smirnov and V.A. Smirnov, *FIRE4, LiteRed and accompanying tools to solve integration by parts relations*, *Comput. Phys. Commun.* **184** (2013) 2820 [[arXiv:1302.5885](#)] [[INSPIRE](#)].
- [44] A.V. Smirnov, *FIRE5: a C++ implementation of Feynman Integral REduction*, *Comput. Phys. Commun.* **189** (2015) 182 [[arXiv:1408.2372](#)] [[INSPIRE](#)].
- [45] R.N. Lee, *Presenting LiteRed: a tool for the Loop InTEgrals REDuction*, [arXiv:1212.2685](#) [[INSPIRE](#)].
- [46] R.N. Lee, *Reducing differential equations for multiloop master integrals*, *JHEP* **04** (2015) 108 [[arXiv:1411.0911](#)] [[INSPIRE](#)].
- [47] J.M. Henn, A.V. Smirnov, V.A. Smirnov and M. Steinhauser, *A planar four-loop form factor and cusp anomalous dimension in QCD*, *JHEP* **05** (2016) 066 [[arXiv:1604.03126](#)] [[INSPIRE](#)].
- [48] F. Cachazo, *Sharpening The Leading Singularity*, [arXiv:0803.1988](#) [[INSPIRE](#)].
- [49] J.M. Henn, K. Melnikov and V.A. Smirnov, *Two-loop planar master integrals for the production of off-shell vector bosons in hadron collisions*, *JHEP* **05** (2014) 090 [[arXiv:1402.7078](#)] [[INSPIRE](#)].
- [50] F. Caola, J.M. Henn, K. Melnikov and V.A. Smirnov, *Non-planar master integrals for the production of two off-shell vector bosons in collisions of massless partons*, *JHEP* **09** (2014) 043 [[arXiv:1404.5590](#)] [[INSPIRE](#)].
- [51] V.A. Smirnov, *Applied asymptotic expansions in momenta and masses*, *Springer Tracts Mod. Phys.* **177** (2002) 1.
- [52] M. Beneke and V.A. Smirnov, *Asymptotic expansion of Feynman integrals near threshold*, *Nucl. Phys. B* **522** (1998) 321 [[hep-ph/9711391](#)] [[INSPIRE](#)].
- [53] A.V. Smirnov, *FIESTA4: Optimized Feynman integral calculations with GPU support*, *Comput. Phys. Commun.* **204** (2016) 189 [[arXiv:1511.03614](#)] [[INSPIRE](#)].
- [54] O. Schnetz, to be published.