RECEIVED: July 27, 2020

REVISED: September 10, 2020

ACCEPTED: September 24, 2020

PUBLISHED: October 28, 2020

Reggeon field theory and self duality: making ends meet

Alex Kovner,^a Eugene Levin,^{b,c} Ming Li^a and Michael Lublinsky^d

^a*Physics Department, University of Connecticut,
2152 Hillside Road, Storrs, CT 06269, U.S.A.*

^b*Departamento de Física, Universidad Técnica Federico Santa María,
and Centro Científico-Tecnológico de Valparaíso,
Avda. España 1680, Casilla 110-V, Valparaíso, Chile*

^c*Department of Particle Physics,
Tel Aviv University, Tel Aviv 69978, Israel*

^d*Physics Department, Ben-Gurion University of the Negev,
Beer Sheva 84105, Israel*

E-mail: kovner@phys.uconn.edu, leving@tauex.tau.ac.il, mli48@ncsu.edu,
lublinm@bgu.ac.il

ABSTRACT: Motivated by the question of unitarity of Reggeon Field Theory, we use the effective field theory philosophy to find possible Reggeon Field Theory Hamiltonians H_{RFT} . We require that H_{RFT} is self dual, reproduce all known limits (dilute-dense and dilute-dilute) and exhibits all the symmetries of the JIMWLK Hamiltonian. We find a family of Hamiltonians which satisfy all the above requirements. One of these is identical in form to the so called “diamond action” discussed in [67, 68]. However we show by explicit calculation that the so called “diamond condition” is not satisfied beyond leading perturbative order.

KEYWORDS: Perturbative QCD, Resummation, Effective Field Theories

ARXIV EPRINT: [2007.12132](https://arxiv.org/abs/2007.12132)

Contents

1	Introduction	1
2	The Reggeon Field Theory: scattering amplitudes and field algebra	4
3	The RFT Hamiltonian	6
3.1	JIMWLK/KLWMIJ Hamiltonians	6
3.2	The self dual extension	9
3.2.1	The symmetries	9
3.2.2	The “left” and “right” Wilson lines	10
3.2.3	Constructing H_{RFT}	12
4	The dense-dilute limit	14
5	Continuous symmetries	19
5.1	The vector $SU_V(N)$ symmetry	19
5.2	Is $SU_L(N)$ there?	20
6	Is this the “Diamond action”?	21
7	Discussion	25
A	Checking $SU_L(N)$	28
A.1	$[Q_L^a, \bar{V}_L(\mathbf{x})]$	28
A.2	Checking $[Q_L^a, H_{RFT}] = 0$	35

1 Introduction

Reggeon Field Theory (RFT) of Quantum Chromodynamics (QCD) is a putative effective theory that is meant to describe scattering at asymptotically high energies. Development of this theory during the last three decades lead to understanding of many features of high energy scattering as well as phenomenological applications to HERA, RHIC and LHC data. Nevertheless, much work notwithstanding the theoretical framework of RFT is incomplete.

The basic pre-QCD ideas of RFT go back to Gribov [1], who considered a very general picture and properties of high energy exchanges in a local field theory. These ideas have been adopted to QCD and further developed over the years in many works [2–39]. Direct derivation of some elements of RFT from QCD has been given. In particular the Hamiltonian of RFT that governs the evolution of physical scattering amplitudes with energy has been derived in two limits - the dilute-dilute limit, where both scattering objects (the projectile and the target) are considered to be small and perturbative (which we refer to

as “dilute”), and the dilute-dense limit, where one of the objects is dilute and the other one is “dense”, i.e. contains a nonperturbatively large gluonic density. The appropriate evolution in the first limit is given by the BFKL equation [2, 3], while in the second by the so-called JIMWLK equation [40–45, 47–49] (and its dual KLWMIJ [46]). The direct relation between the JIMWLK and BK evolution equations [34, 35, 40–45], or Color Glass Condensate (CGC) [47–49] and the RFT has been recognized in [50].

The JIMWLK evolution equation is derived directly from QCD in the leading order perturbative expansion in the dense-dilute regime. As such it does not contain some important effects, like higher order perturbative corrections and the so called Pomeron loops. The NLO corrections to JIMWLK have been derived [51–55] with the conformal part of the kernel known today at the three loop level [56, 57].

The hunt after Pomeron loops on the other hand has not concluded yet. The Pomeron loops are important when both, the density effects in the wave function and multiple effects in scattering are equally important. Some 15 years ago the activity aimed at incorporating the effects of the Pomeron loops into the CGC framework has been very lively [38, 58–65]. Some interesting progress has been made to include both the “splitting” and the “merging” Pomeron processes into the high energy evolution. This activity unfortunately has not converged to a universally accepted form of high energy evolution and RFT.

JIMWLK evolution is valid only in a limited domain of rapidities, i.e. only as long as one of the colliding objects is dilute. The limitation of the JIMWLK evolution to a dense-dilute scattering is a genuine physical restriction. Even though nominally the JIMWLK equation applies to the evolution of a dense system, the fact that the scattering of this large system is allowed to be perturbative (target is dilute) leads to some paradoxical features. For example, as was anticipated in [21] and explicitly demonstrated in [20], when interpreted as the evolution of QCD wave function of a dense object, JIMWLK evolution leads to appearance of negative probabilities. The negative probabilities accompany states arising in the evolution with smaller number of gluons than the number of gluons at the outset of the evolution. Physically one expects of course that the number of gluons in the QCD wave function **increases** with energy, while within the JIMWLK framework the number decreases but the low gluon number states appear with negative probability. This strange behavior nevertheless produces correct energy dependence of the S-matrix but only as long as one of the colliding objects is dilute. The violation of unitarity is a precursor of the eventual breakdown of the JIMWLK evolution at high enough energy. At high energy the Pomeron loops must become important and their effect on the evolution must be significant.

This issue of the unitarity violation in the JIMWLK limit motivates us to reconsider the problem of including Pomeron loops. More precisely we take up a limited goal to try and extend H_{JIMWLK} in a way that it becomes consistent with a very important property of RFT - the self duality. It has been established in [66] that the Hamiltonian that generates the high energy evolution must be invariant under the dense-dilute duality transformation. Physically the self duality has a very simple meaning. It expresses the fact that a scattering amplitude for a scattering of any two hadrons does not depend on which one of them is right moving and which one is left moving, i.e. which one of them we

call the target and which one the projectile. As discussed many times in the literature, the JIMWLK evolution explicitly violates the self duality property which one expects to hold in RFT, since within the domain of validity of JIMWLK the target and the projectile are very different and thus are explicitly treated differently in H_{JIMWLK} .

Although self duality alone may not be sufficient to restore unitarity of the evolution, in a zero dimensional toy model addressed in [21] it was shown that the unitary Hamiltonian is indeed self-dual. Motivated by this, in the present paper we explore possible generalization of the JIMWLK Hamiltonian which restores self duality. Our approach here does not rely on direct derivation from QCD, but instead is akin to typical effective field theory (EFT) attitude: identify relevant degrees of freedom and impose appropriate symmetries. We also require that in the dense-dilute limit the Hamiltonian reproduces both H_{JIMWLK} and H_{KLWMIJ} . We find a family of such Hamiltonians which all reduce to H_{JIMWLK} in the dense-dilute limit and are self dual. We note that one of these Hamiltonians is similar in structure to the so called “diamond action” introduced some years ago in [67] and discussed in [68]. However a more detailed analysis presented below shows that our construction does not support the condition imposed on the product of Wilson loops in [67], which was crucial in the approach of [67] to maintain self duality. Thus our current suggestion is not equivalent to the diamond action of [67]. Additionally we note that our approach relies on the development of RFT formalism in [20], and thus provides directly an algorithm for calculation of scattering amplitudes once the Hamiltonian H_{RFT} is specified.

We thus find a family of self-dual RFT Hamiltonians that reproduces all the known limits. Unfortunately it turns out to be technically involved to check whether the evolution generated by these Hamiltonians is unitary and we are currently unable to answer this question. We are nevertheless encouraged by many similarities with the zero dimensional toy model where the very analogous construction provided a solution to the unitarity problem. The quantitative analysis of this question is left for further research.

The plan of this paper is as follows. In section 2 we recap the formulation of RFT, its algebra of operators and Hilbert space structure discussed in [20]. In section 3 we present the construction of H_{RFT} imposing the discrete symmetries of H_{JIMWLK} in addition to self duality. In section 4 we show that in the dense-dilute limit our H_{RFT} reproduces the JIMWLK and KLWMIJ evolutions. In section 5 we discuss the continuous symmetries of H_{RFT} . This discussion is perturbative, and we conclude that the continuous symmetry group of our H_{RFT} is somewhat surprisingly $SU(N) \times SU(N) \times SU(N)$.¹ In section 6 we consider the relation with the diamond action [67], and show that the so called “diamond condition” on the Wilson lines is violated at second order in g . We conclude with discussion in section 7.

¹We have abused the notation here somewhat. The symmetry group is not in fact a direct product of three factors of $SU(N)$. The more appropriate way to characterize it is to say that the generators contain three linearly independent sets of generators of $SU(N)$. The commutation relations between some of these generators are quite complicated to calculate and thus the full group structure is not known. We will expand on this in the body of the paper.

2 The Reggeon Field Theory: scattering amplitudes and field algebra

In this section we briefly recap the general formulation of the Hamiltonian Reggeon Field Theory given in [20].

Consider an S matrix element S_{fi} for scattering from the initial QCD state $|\Psi_i\rangle = |\mathbf{x}_1, a_1; \dots; \mathbf{x}_N, a_N\rangle_T |\mathbf{y}_1, c_1; \dots; \mathbf{y}_M, c_M\rangle_P$ to the final state $|\Psi_f\rangle = |\mathbf{x}_1, b_1; \dots; \mathbf{x}_N, b_N\rangle_T |\mathbf{y}_1, d_1; \dots; \mathbf{y}_M, d_M\rangle_P$. Here the target state (subscript T) contains N gluons, and the projectile state (subscript P) contains M gluons. The states are labeled by the transverse coordinates and color indexes of the gluons. At high energy in the eikonal approximation this is given by

$$S_{if} \equiv \langle \Psi_i | \hat{S} | \Psi_f \rangle = \langle L | U^{a_1 b_1}(\mathbf{x}_1) \dots U^{a_N b_N}(\mathbf{x}_N) \bar{U}^{c_1 d_1}(\mathbf{y}_1) \dots \bar{U}^{c_M d_M}(\mathbf{y}_M) | R \rangle \quad (2.1)$$

where the left and right RFT Fock vacuum states satisfy

$$\langle L | \bar{U}_{ab} = \delta_{ab} \langle L |; \quad U_{ab} | R \rangle = \delta_{ab} | R \rangle. \quad (2.2)$$

The projectile and target adjoint Wilson line operators are defined in terms of the projectile color charge density $\rho^a(\mathbf{x})$ as

$$\bar{U}(\mathbf{x}) = e^{T^a \frac{\delta}{\delta \rho^a(\mathbf{x})}}; \quad U(\mathbf{x}) = e^{ig T^a \int_y \phi(\mathbf{x}-\mathbf{y}) \rho^a(\mathbf{y})} \quad (2.3)$$

with

$$\alpha^a(\mathbf{x}) = \int_{\mathbf{y}} \phi(\mathbf{x}-\mathbf{y}) \rho^a(\mathbf{y}); \quad \phi(\mathbf{x}-\mathbf{y}) = \frac{g}{2\pi} \ln \frac{|\mathbf{x}-\mathbf{y}|}{L}. \quad (2.4)$$

Here α^a is the potential at point x produced by the charge distribution of the target. The scale L is arbitrary and does not enter calculations of any physical quantities. The $SU(N)$ generators in the adjoint representation are defined in terms of the $SU(N)$ structure constants as

$$T_{bc}^a = -if_{abc}. \quad (2.5)$$

These equations imply non-trivial commutation relations, between U and $\bar{U}$, which constitute the algebra of the RFT in analogy with Heisenberg algebra of fields in the ordinary QFT. In order to calculate the scattering amplitude eq. (2.1) one uses the algebra of U and $\bar{U}$ to commute the factors of U to the right of $\bar{U}$, at which point they disappear by virtue of eq. (2.2).

This algebra encodes the diagrammatic calculation of scattering amplitudes in the operator language. Consider for example the scattering of one gluon on one gluon. The scattering amplitude up to second order in α_s is given by

$$\begin{aligned} \langle L | U^{ab}(\mathbf{x}) \bar{U}^{cd}(\mathbf{y}) | R \rangle = \\ \delta^{ab} \delta^{cd} - ig \phi(\mathbf{x}-\mathbf{y}) T_{ab}^i T_{cd}^i + \left[\frac{1}{2!} ig \phi(\mathbf{x}-\mathbf{y}) \right]^2 (T^i T^j)_{ab} [(T^i T^j)_{cd} + (T^j T^i)_{cd}] + \dots \end{aligned} \quad (2.6)$$

This corresponds to the sum of one and two gluon exchange diagrams in figure 1-a. In fact as was shown in [20], higher order terms organize themselves into all possible diagrams

Figure 1. The one, two (a) and three (b) gluon exchange contributions to the algebra.

where the relative order of the vertices on the target gluon line is permuted in all possible ways. These are the relevant diagrams for eikonal scattering in the Lorentz gauge. The $O(\alpha_s^3)$ contributions correspond to the three gluon exchange diagrams (figure 1-b).

With the algebra encoded in eq. (2.3) and the rule for calculating scattering amplitudes eq. (2.1), the framework of the QCD RFT is defined. To complete the RFT framework one needs to specify the Hamiltonian H_{RFT} that generates the evolution of the scattering amplitude in energy. We will spend some time discussing this Hamiltonian below. But before setting along this route let us recap unitarity constraints on any RFT state as derived in [20]. These constraints must be preserved by energy evolution of the scattering amplitudes. This implies a non-trivial constraint on H_{RFT} [20].

Eq. (2.1) is easily extended for scattering of a state which is a superposition of states with fixed number of gluons. For example, starting with the initial QCD projectile state

$$|\Psi_i\rangle_P = \sum_{n; \mathbf{x}_i; a_i} C_{a_1, a_2 \dots a_n} |\mathbf{x}_1, a_1; \dots; \mathbf{x}_n, a_n\rangle \quad (2.7)$$

the eikonal scattering can only produce a state of the form

$$|\Psi_f\rangle_P = \sum_{n; \mathbf{x}_i; b_i} C_{b_1, b_2 \dots b_n} |\mathbf{x}_1, b_1; \dots; \mathbf{x}_n, b_n\rangle. \quad (2.8)$$

The same holds for the target

$$|\Psi_i\rangle_T = \sum_{m; \mathbf{y}_j; c_j} \bar{C}_{c_1, c_2 \dots c_m} |\mathbf{y}_1, c_1; \dots; \mathbf{y}_m, c_m\rangle \quad (2.9)$$

the eikonal scattering can only produce a state of the form

$$|\Psi_f\rangle_T = \sum_{m; \mathbf{y}_j; d_i} \bar{C}_{d_1, d_2 \dots d_m} |\mathbf{y}_1, d_1; \dots; \mathbf{y}_m, d_m\rangle \quad (2.10)$$

The S-matrix element is given by

$$S_{if} = \langle L | W_T[U] W_P[\bar{U}] | R \rangle \quad (2.11)$$

with

$$W_P = \sum_{n, \{a, b; \mathbf{x}\}} F^n(\{a, b; \mathbf{x}\}) \prod_{i=1}^n [\bar{U}^{a_i b_i}(\mathbf{x}_i)] \quad (2.12)$$

$$F^n(\{a, b; \mathbf{x}\}) = C_{a_1, a_2 \dots a_n}(\mathbf{x}_1 \dots \mathbf{x}_n) C_{b_1, b_2 \dots b_n}^*(\mathbf{x}_1 \dots \mathbf{x}_n) \quad (2.13)$$

and

$$W_T = \sum_{m, \{c, d; \mathbf{y}\}} \bar{F}^n(\{c, d; \mathbf{y}\}) \prod_{i=1}^m [U^{c_i, d_i}(\mathbf{y}_i)] \quad (2.14)$$

$$\bar{F}^m(\{c, d; \mathbf{y}\}) = \bar{C}_{c_1, c_2 \dots c_m}(\mathbf{y}_1 \dots \mathbf{y}_m) \bar{C}_{d_1, d_2 \dots d_m}^*(\mathbf{y}_1 \dots \mathbf{y}_m). \quad (2.15)$$

As is obvious from eqs. (2.13), (2.15), the functions F and $\bar{F}$ must satisfy the properties of s-channel unitarity [20]

$$F^n(\{a, a; \mathbf{x}\}) \geq 0; \quad \sum_{n, \{a\}} \int_{\{\mathbf{x}\}} F^n(\{a, a; \mathbf{x}\}) = 1; \quad (2.16)$$

and the same for $\bar{F}$.

As shown in [20] some these conditions are violated in JIMWLK evolution, which leads to negative probabilities $\bar{F}$ when evolving the state of a dense target.

3 The RFT Hamiltonian

The subject of RFT is the evolution of scattering amplitudes with energy. In general the energy evolution is generated by the action of the RFT Hamiltonian $H_{RFT}[U, \bar{U}]$. The S-matrix element of eq. (2.1) evolved to rapidity Y is given by

$$S_{if}(Y) = \langle L | U^{a_1 b_1}(\mathbf{x}_1) \dots U^{a_N b_N}(\mathbf{x}_N) e^{Y H_{RFT}[U, \bar{U}]} \bar{U}^{c_1 d_1}(\mathbf{y}_1) \dots \bar{U}^{c_M d_M}(\mathbf{y}_M) | R \rangle. \quad (3.1)$$

3.1 JIMWLK/KLWMIJ Hamiltonians

Exploring the functional form of H_{RFT} is the subject of this paper. Ideally we would like to derive it directly from a QCD calculation. This has been achieved in the dense-dilute limit, where one of the scattering objects is dense and the other one is dilute. The two versions of the Hamiltonian related by the duality transformation have been derived in [40–49].

When the target is dense and the projectile dilute, the relevant limit is the JIMWLK Hamiltonian:

$$H_{JIMWLK} = \frac{\alpha_s}{2\pi^2} \int_{\mathbf{x}, \mathbf{y}, \mathbf{z}} \frac{(\mathbf{x} - \mathbf{z}) \cdot (\mathbf{y} - \mathbf{z})}{(\mathbf{x} - \mathbf{z})^2 (\mathbf{y} - \mathbf{z})^2} \left[2 \mathcal{J}_L^a(\mathbf{x}) \mathcal{J}_R^b(\mathbf{y}) \bar{U}^{ab}(\mathbf{z}) - \mathcal{J}_L^a(\mathbf{x}) \mathcal{J}_L^a(\mathbf{y}) - \mathcal{J}_R^a(\mathbf{x}) \mathcal{J}_R^a(\mathbf{y}) \right]. \quad (3.2)$$

Here the right and left rotation operators are defined as [69]

$$\begin{aligned} \mathcal{J}_L^a(\mathbf{x}) &= \left[\frac{1}{2} T^e \frac{\delta}{\delta \rho^e(\mathbf{x})} \left(\coth \left[\frac{1}{2} T^e \frac{\delta}{\delta \rho^e(\mathbf{x})} \right] - 1 \right) \right]^{ba} \rho^b(\mathbf{x}), \\ \mathcal{J}_R^a(\mathbf{x}) &= \left[\frac{1}{2} T^e \frac{\delta}{\delta \rho^e(\mathbf{x})} \left(\coth \left[\frac{1}{2} T^e \frac{\delta}{\delta \rho^e(\mathbf{x})} \right] + 1 \right) \right]^{ba} \rho^b(\mathbf{x}), \end{aligned} \quad (3.3)$$

The function on the right hand side as usual should be understood as a power series expansion. For a single variable t we have

$$\begin{aligned} M_L(t) &\equiv \frac{t}{2} \left(\coth \frac{t}{2} - 1 \right) = \frac{t}{e^t - 1} = \sum_{m=0}^{\infty} \frac{B_m^- t^m}{m!} = \sum_{m=0}^{\infty} C_m^- t^m \\ M_R(t) &\equiv \frac{t}{2} \left(\coth \frac{t}{2} + 1 \right) = \frac{t}{1 - e^{-t}} = \sum_{m=0}^{\infty} \frac{B_m^+ t^m}{m!} = \sum_{m=0}^{\infty} C_m^+ t^m. \end{aligned} \quad (3.4)$$

Here B_m^- and B_m^+ are Bernoulli numbers. They have the properties that $B_{2n}^- = B_{2n}^+$ for all even integers $2n$ while $B_{2n+1}^- = B_{2n+1}^+ = 0$ for all odd integers $2n+1$ except $B_1^- = -\frac{1}{2} = -B_1^+$. Also the relations $M_L(t) = M_R(t)e^{-t}$ and $M_R(t) = M_L(t)e^t$ can be readily verified.

The operators $\mathcal{J}_L^a(\mathbf{x}), \mathcal{J}_R^a(\mathbf{x})$ act as left rotation and right rotation on the Wilson line $\bar{U}^{mn}(\mathbf{x})$,

$$\begin{aligned} [\mathcal{J}_L^a(\mathbf{x}), \bar{U}^{mn}(\mathbf{y})] &= -(T^a \bar{U}(\mathbf{y}))^{mn} \delta(\mathbf{x} - \mathbf{y}), \\ [\mathcal{J}_R^a(\mathbf{x}), \bar{U}^{mn}(\mathbf{y})] &= -(\bar{U}(\mathbf{y}) T^a)^{mn} \delta(\mathbf{x} - \mathbf{y}). \end{aligned} \quad (3.5)$$

One seemingly peculiar feature of these definitions is that when considered as operators on the standard Hilbert space of functions of ρ , the operators $\mathcal{J}_{L(R)}$ are not Hermitian

$$\mathcal{J}_L^\dagger \neq \mathcal{J}_L; \quad \mathcal{J}_R^\dagger \neq \mathcal{J}_R. \quad (3.6)$$

However one has to keep in mind that the operation of Hermitian conjugation of the operators in QCD Hilbert space does not correspond to naive Hermitian conjugation in the RFT space. Without going into detailed discussion here, we refer the reader to [70] where it was shown that the RFT transformation that corresponds to Hermitian conjugation in the QCD Hilbert space is

$$[\text{QCD operator}]^\dagger \rightarrow (L \leftrightarrow R)^* \quad (3.7)$$

Under this transformation indeed we have

$$\mathcal{J}_L \rightarrow \mathcal{J}_R^* = \mathcal{J}_L; \quad \mathcal{J}_R \rightarrow \mathcal{J}_L^* = \mathcal{J}_R \quad (3.8)$$

as is required for Hermitian operators in the QCD Hilbert space.

The evolution in the reverse situation (dilute target and dense projectile) is governed by the so called KLWMIJ Hamiltonian,

$$H_{KLWMIJ} = \frac{\alpha_s}{2\pi^2} \int_{\mathbf{x}, \mathbf{y}, \mathbf{z}} \frac{(\mathbf{x} - \mathbf{z}) \cdot (\mathbf{y} - \mathbf{z})}{(\mathbf{x} - \mathbf{z})^2 (\mathbf{y} - \mathbf{z})^2} \left[2\mathcal{I}_L^a(\mathbf{x}) \mathcal{I}_R^b(\mathbf{y}) U^{ab}(\mathbf{z}) - \mathcal{I}_L^a(\mathbf{x}) \mathcal{I}_L^a(\mathbf{y}) - \mathcal{I}_R^a(\mathbf{x}) \mathcal{I}_R^a(\mathbf{y}) \right] \quad (3.9)$$

where $\mathcal{I}_{L(R)}$ are defined as

$$\begin{aligned} \mathcal{I}_L^a(\mathbf{x}) &= \frac{-i}{g} \frac{\delta}{\delta \alpha^b(\mathbf{x})} \left[\frac{1}{2} T^e i g \alpha^e(\mathbf{x}) \left(\coth \left[\frac{1}{2} T^e i g \alpha^e(\mathbf{x}) \right] - 1 \right) \right]^{ba}, \\ \mathcal{I}_R^a(\mathbf{x}) &= \frac{-i}{g} \frac{\delta}{\delta \alpha^b(\mathbf{x})} \left[\frac{1}{2} T^e i g \alpha^e(\mathbf{x}) \left(\coth \left[\frac{1}{2} T^e i g \alpha^e(\mathbf{x}) \right] + 1 \right) \right]^{ba}, \end{aligned} \quad (3.10)$$

with $\alpha^a(\mathbf{x})$ defined in eq. (2.4). These satisfy

$$\begin{aligned} [U^{mn}(\mathbf{y}), \mathcal{I}_L^a(\mathbf{x})] &= -(T^a \mathbf{U}(\mathbf{y}))^{mn} \delta(\mathbf{x} - \mathbf{y}), \\ [U^{mn}(\mathbf{y}), \mathcal{I}_R^a(\mathbf{x})] &= -(\mathbf{U}(\mathbf{y}) T^a)^{mn} \delta(\mathbf{x} - \mathbf{y}). \end{aligned} \quad (3.11)$$

The two sets of operators satisfy two copies of $SU(N) \times SU(N)$ commutation relations:

$$\begin{aligned} [\mathcal{J}_L^a(\mathbf{x}), \mathcal{J}_L^b(\mathbf{y})] &= i f^{abc} \mathcal{J}_L^c(\mathbf{x}) \delta(\mathbf{x} - \mathbf{y}), \\ [\mathcal{J}_R^a(\mathbf{x}), \mathcal{J}_R^b(\mathbf{y})] &= -i f^{abc} \mathcal{J}_R^c(\mathbf{x}) \delta(\mathbf{x} - \mathbf{y}) \\ [\mathcal{J}_L^a(\mathbf{x}), \mathcal{J}_R^a(\mathbf{y})] &= 0. \end{aligned} \quad (3.12)$$

and

$$\begin{aligned} [\mathcal{I}_L^a(\mathbf{x}), \mathcal{I}_L^b(\mathbf{y})] &= -i f^{abc} \mathcal{I}_L^c(\mathbf{x}) \delta(\mathbf{x} - \mathbf{y}), \\ [\mathcal{I}_R^a(\mathbf{x}), \mathcal{I}_R^b(\mathbf{y})] &= i f^{abc} \mathcal{I}_R^c(\mathbf{x}) \delta(\mathbf{x} - \mathbf{y}) \\ [\mathcal{I}_L^a(\mathbf{x}), \mathcal{I}_R^a(\mathbf{y})] &= 0. \end{aligned} \quad (3.13)$$

The commutation relations between $\mathcal{J}$ and $\mathcal{I}$ are rather complicated and we will not attempt to derive them here.

The Hamiltonian of RFT must possess a property of self duality, i.e. it has to be invariant under the transformation that interchanges the projectile and the target. This is obvious from the point of view of QCD, since it is immaterial which one of the colliding objects we call the target, and which one the projectile. Thus scattering of an N gluon projectile on an M gluon target is the same as scattering of an M gluon projectile on an N gluon target. The JIMWLK (and likewise KLWMIJ) Hamiltonian is not self dual, since it is only meant to be valid in the very asymmetric regime where one of the colliding objects is dense and one is dilute. This lack of self duality means among other things, that JIMWLK cannot be used at asymptotically high energies, where the projectile becomes dense as well. It is thus clearly desirable to find a self dual extension of H_{JIMWLK} .

Some years ago a considerable effort has been dedicated to a search for a self dual extension of the Hamiltonian. One such extension in the context of large N_c Pomeron theory was suggested by Braun [31–33]. The solutions to the Braun theory however exhibit a nonphysical bifurcating behavior [71] which was an original motivation for the study of [21]. It was shown in [21] that Braun’s theory suffers from unitarity violation. Other attempts based on the QCD path integral approach were reported in [67, 68]. Those works have proposed the so called “diamond action” as a self dual effective action of RFT. Although the question has not been settled, in recent years this effort has only been simmering on a back burner.

Here we return to this problem motivated by considerations of unitarity. As we showed in [20], the JIMWLK Hamiltonian violates QCD unitarity constraints when acting on the dense target wave function. In view of the discussion in [21] of the zero dimensional toy model, it seems likely that the self duality of H_{RFT} is necessary in order to restore unitarity. In this section we present a self dual H_{RFT} and show that it reduces to H_{JIMWLK} and H_{KLWMIJ} in the appropriate dense-dilute limit.

3.2 The self dual extension

3.2.1 The symmetries

Our strategy in this paper is similar to that of EFT: we are not going to attempt to derive H_{RFT} from first principles, but will rather construct a family of Hamiltonians which on the one hand reduce to H_{JIMWLK} and H_{KLWMIJ} in the appropriate limits, and on the other hand are symmetric under the known symmetries of H_{JIMWLK} in addition to being self dual.

The symmetries of H_{JIMWLK} have been analyzed for example in [50] and [72]. H_{JIMWLK} possesses the continuous symmetry group $SU_L(N) \times SU_R(N)$ generated by $\mathcal{J}_{L(R)}$. In addition it has the discrete $Z_2^S \times Z_2^C$ symmetry group with the two discrete transformations acting in the following way:

1. The signature Z_2^S

$$SUS^\dagger = U^\dagger; \quad S\bar{U}S^\dagger = \bar{U}^\dagger; \quad S\mathcal{J}_L S^\dagger = -\mathcal{J}_R; \quad S\mathcal{I}_L S^\dagger = -\mathcal{I}_R. \quad (3.14)$$

2. The charge conjugation Z_2^C .

For simplicity we choose to work in the basis where the generators in the fundamental representation t^a are either real and symmetric or imaginary and antisymmetric. In this basis the charge conjugation symmetry corresponds to changing the sign of the real generators since this has the effect $t^a \rightarrow -t^{a*}$ which interchanges the generators in fundamental and anti fundamental representations. Defining the matrix

$$c_{ab} = -2\text{tr}[t^a t^{*b}] \quad (3.15)$$

the “second quantized” form of the transformation is

$$C\mathcal{J}_{L(R)}^a C^\dagger = c^{ab}\mathcal{J}_{L(R)}^b; \quad C\mathcal{I}_{L(R)}^a C^\dagger = c^{ab}\mathcal{I}_{L(R)}^b. \quad (3.16)$$

The eikonal factors in fundamental (U_F) and adjoint (U) representations transform as

$$CU_F C^\dagger = U_F^*; \quad C\bar{U}_F C^\dagger = \bar{U}_F^*. \quad (3.17)$$

$$CU_{ab} C^\dagger = c_{ac}U_{cd}c_{db}; \quad C\bar{U}_{ab} C^\dagger = c_{ac}\bar{U}_{cd}c_{db}. \quad (3.18)$$

We expect both the discrete symmetries of H_{JIMWLK} to remain the symmetries of the general H_{RFT} since they directly reflect the symmetries of QCD. The situation with $SU_L(N) \times SU_R(N)$ is less clear. It is certainly true that we expect the diagonal vector subgroup $SU_V(N)$ to be a symmetry of H_{RFT} , since it descends directly from the global color group of QCD as it rotates simultaneously the initial and final scattering states. The left rotation acts only on the initial states and may be an accidental symmetry of the dense-dilute limit. Thus we will not insist on $SU_L(N)$ and $SU_R(N)$ to be separate symmetries but will return to this question later.

In addition to these symmetries which are symmetries of $JIMWLK$ limit, we will also require H_{RFT} to be invariant under the dense dilute duality Z_2^D . To understand how the

duality transformation acts on the field variables in the current RFT setup, we recall that physically it simply interchanges the projectile and the target. In other words for basic scattering amplitude we should have

$$\begin{aligned} \langle L|U^{a_1 b_1}(\mathbf{x}_1) \dots U^{a_N b_N}(\mathbf{x}_N) \bar{U}^{c_1 d_1}(\mathbf{y}_1) \dots \bar{U}^{c_M d_M}(\mathbf{y}_M)|R\rangle &\rightarrow \\ \langle L|U^{d_1 c_1}(\mathbf{y}_1) \dots U^{d_M c_M}(\mathbf{y}_M) \bar{U}^{b_1 a_1}(\mathbf{x}_1) \dots \bar{U}^{b_N a_N}(\mathbf{x}_N)|R\rangle. \end{aligned} \quad (3.19)$$

Self duality, or invariance, under Z_2^D is a realization of the fact that the two amplitudes must be equal at any collision energy

$$\begin{aligned} \langle L|U^{a_1 b_1}(\mathbf{x}_1) \dots U^{a_N b_N}(\mathbf{x}_N) \bar{U}^{c_1 d_1}(\mathbf{y}_1) \dots \bar{U}^{c_M d_M}(\mathbf{y}_M)|R\rangle \\ = \langle L|U^{d_1 c_1}(\mathbf{y}_1) \dots U^{d_M c_M}(\mathbf{y}_M) \bar{U}^{b_1 a_1}(\mathbf{x}_1) \dots \bar{U}^{b_N a_N}(\mathbf{x}_N)|R\rangle. \end{aligned} \quad (3.20)$$

When considered as a transformation acting on a function of the basic fields ρ and $\frac{\delta}{\delta\rho}$, the Z_2^D transformation can be written as

$$F[\rho, \frac{\delta}{\delta\rho}] \rightarrow F^\dagger[-\frac{i}{g} \frac{\delta}{\delta\alpha^a}, ig\alpha^a]. \quad (3.21)$$

In terms of individual operators this is [66]

$$\begin{aligned} \rho^a &\rightarrow -\frac{i}{g} \frac{\delta}{\delta\alpha^a}, & \frac{\delta}{\delta\rho^a} &\rightarrow -ig\alpha^a \\ U &\rightarrow \bar{U}; & \mathcal{J}_{L(R)} &\rightarrow \mathcal{I}_{R(L)}^\dagger. \end{aligned} \quad (3.22)$$

However, in addition to this action one has to take an overall Hermitian conjugation of the whole expression which is being transformed. Note that due to this additional action of Hermitian conjugation the duality transformation Z_2^D cannot be represented by an action of a unitary operator on the RFT Hilbert space. This is similar to time reversal in quantum mechanics, which is not a unitary but an anti unitary transformation. Recall that anti unitary transformation involves complex conjugation of an operator function in addition to the transformation of basic variables. The duality is not an anti unitary transformation either, since it involves hermitian conjugation rather than a simple complex conjugation of a function F . Nevertheless, just like the time reversal in quantum mechanics, it is a bona fide linear transformation in the Hilbert space and thus should be considered on par with other symmetries of the theory.

3.2.2 The “left” and “right” Wilson lines

To construct H_{RFT} let us introduce the following Wilson line like operators in the *fundamental representation*

$$\begin{aligned} V_L(\mathbf{x}) &= \text{Exp} \left\{ i \int_{\mathbf{y}} g\phi(\mathbf{x} - \mathbf{y}) t^e \mathcal{J}_L^e(\mathbf{y}) \right\} \\ V_R(\mathbf{x}) &= \text{Exp} \left\{ -i \int_{\mathbf{y}} g\phi(\mathbf{x} - \mathbf{y}) t^e \mathcal{J}_R^e(\mathbf{y}) \right\} \\ \bar{V}_L(\mathbf{x}) &= \text{Exp} \left\{ i \int_{\mathbf{y}} g\phi(\mathbf{x} - \mathbf{y}) t^e \mathcal{I}_L^e(\mathbf{y}) \right\} \\ \bar{V}_R(\mathbf{x}) &= \text{Exp} \left\{ -i \int_{\mathbf{y}} g\phi(\mathbf{x} - \mathbf{y}) t^e \mathcal{I}_R^e(\mathbf{y}) \right\}. \end{aligned} \quad (3.23)$$

These expressions resemble our reggeized gluon operators U and $\bar{U}$. However they are defined in terms of $SU(N)$ generators $\mathcal{J}_{L(R)}$ and $\mathcal{I}_{L(R)}$ rather than commuting variables ρ .

The reason to introduce these operators is that they look like appropriate building blocks for H_{RFT} . Recall that we need H_{RFT} to reduce to H_{JIMWLK} in the dense dilute limit, i.e. in the leading order of expansion in powers of ρ . Now H_{JIMWLK} is a simple function when written in terms of $\mathcal{J}_{L(R)}$ rather than the regular Wilson line operators U . It therefore seems likely that in order to extend it beyond the dense-dilute limit the basic building blocks also should be simple function of $\mathcal{J}$'s. On the other hand H_{JIMWLK} is also a simple function of $\bar{U}$. Given that we want to impose self duality on H_{RFT} it is reasonable to choose our building blocks to be in some way similar to Wilson lines. Hence the motivation to introduce the operators in eq. (3.23). We chose to discuss these operators in fundamental representation for simplicity. As we will show later, the construction we propose works with an arbitrary representation of $SU(N)$, thus providing an infinite set of Hamiltonians that satisfy our requirements.

When calculating the RFT “correlators” of these operators with U and $\bar{U}$, the ordering of the vertices is important, unlike in the calculation of correlators of U 's and $\bar{U}$'s among themselves. For example consider the simplest correlator

$$\langle L|V_L^{\alpha\beta}(\mathbf{x})\bar{U}^{cd}(\mathbf{y})|R\rangle = \delta^{\alpha\beta}\delta^{cd} - ig\phi(\mathbf{x}-\mathbf{y})t_{\alpha\beta}^iT_{cd}^i + \frac{1}{2!}[ig\phi(\mathbf{x}-\mathbf{y})]^2(t^it^j)_{\alpha\beta}(T^jT^i)_{cd} + \dots \quad (3.24)$$

where the ellipsis denotes contributions of order g^6 and higher, i.e. three and higher gluon exchange diagrams. For comparison, a similar correlator for the fundamental Wilson line defined as

$$V(\mathbf{x}) = U_F(\mathbf{x}) = \text{Exp} \left\{ i \int_{\mathbf{y}} \phi(\mathbf{x}-\mathbf{y}) t^e \rho^e(\mathbf{y}) \right\}$$

is

$$\begin{aligned} \langle L|V^{\alpha\beta}(\mathbf{x})\bar{U}^{cd}(\mathbf{y})|R\rangle = \\ \delta^{\alpha\beta}\delta^{cd} - ig\phi(\mathbf{x}-\mathbf{y})t_{\alpha\beta}^iT_{cd}^i + \left[\frac{1}{2!}ig\phi(\mathbf{x}-\mathbf{y}) \right]^2 (t^it^j)_{\alpha\beta}[(T^iT^j)_{cd} + (T^jT^i)_{cd}] + \dots \end{aligned} \quad (3.25)$$

At the two gluon exchange level the difference between the two is

$$\langle L|V_L^{\alpha\beta}(\mathbf{x})\bar{U}^{cd}(\mathbf{y})|R\rangle - \langle L|V^{\alpha\beta}(\mathbf{x})\bar{U}^{cd}(\mathbf{y})|R\rangle = -\frac{1}{4}[ig\phi(\mathbf{x}-\mathbf{y})]^2(t^it^j)_{\alpha\beta}[T^i, T^j]_{cd} \quad (3.26)$$

which corresponds to the diagram in figure 2. Note that this difference is a two gluon exchange in the octet channel, and may be viewed simply as the reggeization correction to a single gluon exchange.

In general if one thinks about V_L as representing a fundamentally charged parton in the target wave function, the parton in question would be something of a black sheep. It would always scatter on the projectile only after all the other partons have had their day. As an example, a sample diagram corresponding to the calculation of the correlator $\langle L|U(\mathbf{x}_1)U(\mathbf{x}_2)V_L(\mathbf{z})\bar{U}(\mathbf{y})|R\rangle$ is depicted on figure 3. Note that all the gluons exchanged

Figure 2. Reggeization corrections to a single gluon exchange.

Figure 3. A sample diagram for a correlator that includes V_L .

between $\bar{U}$ and V_L attach to the $\bar{U}$ line to the left of any gluon exchanged between $\bar{U}$ and any of the U 's. This follows since V_L contains only left rotation generators of $\bar{U}$. Similarly, V_R only contains right rotation operators, and therefore in a scattering diagram always exchanges gluons with the projectile **before** any other exchanges with target gluons.

Also note that operatorially V_L and V_R **do not commute** with U , although they commute with each other. Similar comments apply to $\bar{V}_{L(R)}$.

3.2.3 Constructing H_{RFT}

Let us now consider the following expression

$$\begin{aligned}
 H_{RFT}^{(1)} &= \frac{1}{\pi g^2} \int d^2 \mathbf{x} \text{Tr} [\partial^2 V_L(\mathbf{x}) \bar{V}_L(\mathbf{x}) V_R(\mathbf{x}) \bar{V}_R(\mathbf{x}) + V_L(\mathbf{x}) \bar{V}_L(\mathbf{x}) \partial^2 V_R(\mathbf{x}) \bar{V}_R(\mathbf{x}) \\
 &\quad + 2 \partial_i V_L(\mathbf{x}) \bar{V}_L(\mathbf{x}) \partial_i V_R(\mathbf{x}) \bar{V}_R(\mathbf{x})] \\
 &= \frac{1}{\pi g^2} \int d^2 \mathbf{x} \bar{V}_L^{\beta\gamma}(\mathbf{x}) \bar{V}_R^{\delta\alpha}(\mathbf{x}) \partial^2 [V_L^{\alpha\beta}(\mathbf{x}) V_R^{\gamma\delta}(\mathbf{x})] \\
 &= \frac{1}{\pi g^2} \int d^2 \mathbf{x} \partial^2 [\bar{V}_L^{\beta\gamma}(\mathbf{x}) \bar{V}_R^{\delta\alpha}(\mathbf{x})] V_L^{\alpha\beta}(\mathbf{x}) V_R^{\gamma\delta}(\mathbf{x})
 \end{aligned} \tag{3.27}$$

where in the last line we have integrated by parts assuming that the boundary terms vanish. Note that the order of factors is important, since the operators V and $\bar{V}$ do not commute with each other. In (3.27) all factors V_L, V_R are understood as positioned to the right of any factor $\bar{V}_L, \bar{V}_R$. The diagram that schematically represents the color flow between the four Wilson lines is shown in figure 4.

We start with this expression since as we will see shortly it reproduces both, the JIMWLK and the KLWMIJ Hamiltonians in the appropriate dense-dilute limit. Following our EFT like strategy we would like to impose on H_{RFT} the discrete symmetries discussed above. It turns out that it is quite easy to do.

We start with the duality transformation Z_2^D . We perform the transformation in two steps. First we perform the canonical transformation

$$\rho^a \leftrightarrow -\frac{i}{g} \frac{\delta}{\delta \alpha^a}, \quad \frac{\delta}{\delta \rho^a} \leftrightarrow -ig \alpha^a \tag{3.28}$$

Figure 4. The Reggeon field theory Hamiltonian $H_{RFT}^{(1)}$. The arrows indicate the directions of color charge flow.

under which

$$\mathcal{J}_L^a \leftrightarrow \mathcal{I}_R^{a\dagger}, \quad \mathcal{J}_R^a \leftrightarrow \mathcal{I}_L^{a\dagger} \quad (3.29)$$

or equivalently,

$$\begin{aligned} V_L &\rightarrow \bar{V}_R^\dagger, & V_R &\rightarrow \bar{V}_L^\dagger, \\ \bar{V}_L &\rightarrow V_R^\dagger, & \bar{V}_R &\rightarrow V_L^\dagger. \end{aligned} \quad (3.30)$$

Second, in accordance with eq. (3.21) we take the Hermitian conjugation of the transformed Hamiltonian to obtain

$$\begin{aligned} H_{RFT}^{(1)dual} &= \left(\frac{1}{\pi g^2} \int d\mathbf{x} V_R^{\dagger\beta\gamma}(\mathbf{x}) V_L^{\dagger\delta\alpha}(\mathbf{x}) \partial^2 [\bar{V}_R^{\dagger\alpha\beta}(\mathbf{x}) \bar{V}_L^{\dagger\gamma\delta}(\mathbf{x})] \right)^\dagger \\ &= \frac{1}{\pi g^2} \int d\mathbf{x} \partial^2 [\bar{V}_R^{\beta\alpha}(\mathbf{x}) \bar{V}_L^{\delta\gamma}(\mathbf{x})] V_R^{\gamma\beta}(\mathbf{x}) V_L^{\alpha\delta}(\mathbf{x}) = H_{RFT}^{(1)}. \end{aligned} \quad (3.31)$$

Thus we find that $H_{RFT}^{(1)}$ is self dual already.

The next in line is the signature transformation eq. (3.14)

$$V_L \leftrightarrow V_R, \quad \bar{V}_L \leftrightarrow \bar{V}_R, \quad (3.32)$$

It is easily seen that $H_{RFT}^{(1)}$ is invariant under this transformation.

The only remaining discrete symmetry is charge conjugation. Although $H_{RFT}^{(1)}$ itself is not invariant it is easy to rectify this.

According to eq. (3.16) the charge conjugation transformation acts on the left and right Wilson lines. From the definition of $\mathcal{J}_{L(R)}^a$ and $\mathcal{I}_{L(R)}^a$, taking complex conjugate, one obtains

$$\begin{aligned} CV_L C^\dagger &\equiv V_L^c = \exp \left\{ -ig \int_{\mathbf{y}} \phi(\mathbf{x} - \mathbf{y}) t^{e*} \mathcal{J}_L^e(\mathbf{y}) \right\}, \\ CV_R C^\dagger &\equiv V_R^c = \exp \left\{ ig \int_{\mathbf{y}} \phi(\mathbf{x} - \mathbf{y}) t^{e*} \mathcal{J}_R^e(\mathbf{y}) \right\}, \\ C\bar{V}_L C^\dagger &\equiv \bar{V}_L^c = \exp \left\{ -ig \int_{\mathbf{y}} \phi(\mathbf{x} - \mathbf{y}) t^{e*} \mathcal{I}_L^e(\mathbf{y}) \right\}, \\ C\bar{V}_R C^\dagger &\equiv \bar{V}_R^c = \exp \left\{ ig \int_{\mathbf{y}} \phi(\mathbf{x} - \mathbf{y}) t^{e*} \mathcal{I}_R^e(\mathbf{y}) \right\}. \end{aligned} \quad (3.33)$$

Applying the charge conjugation on $H_{RFT}^{(1)}$ we obtain

$$CH_{RFT}^{(1)}C^\dagger \equiv H_{RFT}^{(1)c} = \frac{1}{\pi g^2} \int d\mathbf{x} \bar{V}_L^{c,\beta\gamma}(\mathbf{x}) \bar{V}_R^{c,\delta\alpha}(\mathbf{x}) \partial^2 \left[V_L^{c,\alpha\beta}(\mathbf{x}) V_R^{c,\gamma\delta}(\mathbf{x}) \right]. \quad (3.34)$$

It is easy to see that $H_{RFT}^{(1)c}$ is by itself invariant under the signature and duality transformations. Therefore, the following Hamiltonian is invariant under all relevant discrete symmetries:

$$\begin{aligned} H_{RFT} &= \frac{1}{2} \left(H_{RFT}^{(1)} + H_{RFT}^{(1)c} \right) \\ &= \frac{1}{2\pi g^2} \int d^2\mathbf{x} \left(\bar{V}_L^{\beta\gamma}(\mathbf{x}) \bar{V}_R^{\delta\alpha}(\mathbf{x}) \partial^2 \left[V_L^{\alpha\beta}(\mathbf{x}) V_R^{\gamma\delta}(\mathbf{x}) \right] \right. \\ &\quad \left. + \bar{V}_L^{c,\beta\gamma}(\mathbf{x}) \bar{V}_R^{c,\delta\alpha}(\mathbf{x}) \partial^2 \left[V_L^{c,\alpha\beta}(\mathbf{x}) V_R^{c,\gamma\delta}(\mathbf{x}) \right] \right). \end{aligned} \quad (3.35)$$

So far we have not discussed the continuous symmetries of H_{RFT} . We will postpone this discussion to section 5 after we consider the dense-dilute limit.

We have found a candidate RFT Hamiltonian which is self dual. In fact the construction above defines a family of self dual Hamiltonians. In particular rather than using the fundamental representation for defining $V_{L(R)}$ and $\bar{V}_{L(R)}$ we could have used any representation of the color group. Any one of these variations is self dual and, as we will see later reduces to the JIMWLK Hamiltonian in the dense-dilute limit. We do not have any a priori reason to prefer one of these versions to another, although it may seem unnatural to involve very high representations of the color group. One should also note that for representations that have vanishing N -ality, like the adjoint representation one has $H_{RFT}^{(1)} = H_{RFT}^{(1)c}$ which is a simplifying feature.

In this paper we will be working with the fundamental representation defined in eq. (3.23) when deriving the JIMWLK and KLWMIJ limits so that not to lose generality. We will show that $H_{RFT}^{(1)}$ and $H_{RFT}^{(1)c}$ separately reduce to H_{JIMWLK} and H_{KLWMIJ} in appropriate limits and that this feature extends to any representation of $SU(N)$.

4 The dense-dilute limit

The most important test for H_{RFT} is that it must reproduce H_{JIMWLK} in the dense-dilute limit. In this section we demonstrate explicitly that this is indeed the case.

The dense-dilute limit arises when the number of gluons in the projectile is of order one, while the number of gluons in the target is large, parametrically $n \sim O(1/\alpha_s^2)$. Thus we are considering the amplitude in eq. (2.1) and eq. (3.1) where the number of factors $\bar{U}$ is of order one, and the number of factors U is of order $1/\alpha_s^2$. In this limit several simplifications occur.

We will first give a simplified argument, and then complete the mathematical details of the demonstration.

First of all, note that at weak coupling any given projectile gluon can exchange at most two gluons with any given target gluon. However, since the number of gluons in the target is large, a projectile gluon can multiply scatter on many gluons of the target. A

Figure 5. A representative diagram for scattering of a single projectile gluon in the dense-dilute limit. Here we have drawn the projectile gluon on the top of the figure and the target gluons at the bottom.

Figure 6. A representative diagram for scattering of a single target gluon in the dense-dilute limit. Here we have drawn the target gluon on the bottom of the figure and the projectile gluons at the top.

representative diagram for scattering of a single projectile gluon is depicted on figure 5. The diagram in figure 5 contains single and double gluon exchanges between individual pairs of gluons. If a single gluon exchange is present such a diagram contributes to an inelastic amplitude as the final state of the scattering process is necessarily different from the initial state. The elastic amplitude has contribution only from two gluon exchanges where the two gluons are in the color singlet. Since every two gluon exchange carries a factor α_s^2 , and there are in total $O(1/\alpha_s^2)$ target partons that can participate in the scattering, the total elastic scattering amplitude in the dense-dilute limit is of order unity.²

On the other hand since the projectile is dilute, every target gluon can only scatter either on one or two projectile gluons. The appropriate diagrams are represented on figure 6. Technically this means that in the dense-dilute limit all factors of U have to be expanded to second order in ρ . This insures that once two gluons are exchanged between a target gluon and the projectile, the target gluon does not participate in any further scattering.

Now consider the diagrams as in figure 6 but which, instead of one of the factors $\bar{U}$ contain a factor $\bar{V}_L$ that appears in the RFT Hamiltonian.

As we have discussed above, the only difference between these two sets of diagrams is that all the gluons exchanged between $\bar{V}_L$ and any given factor U connect to the left of any other gluons that might be exchanged by this U and a different factor of $\bar{U}$ present in the amplitude. However any given U can exchange at most two gluons. If these two gluons

²Single gluon exchanges behave a little differently. One does not add single gluon exchange amplitudes between a given projectile gluon and different target gluons since those lead to different final states of the target and do not contribute to the same S matrix element. Instead the single gluon exchanges with distinct target gluons lead to appearance of many nonvanishing off diagonal matrix elements of the S matrix albeit each such matrix element is of order α_s . The number of such nonvanishing matrix elements is $O(1/\alpha_s^2)$.

Figure 7. An α_s suppressed correction to a correlator containing $\bar{V}_L$ which is negligible in the weak coupling limit.

are exchanged between U and $\bar{V}_L$, no further gluons are exchanged and the action of $\bar{V}_L$ is identical to the action of $\bar{V}$. If U exchanges only one gluon with $\bar{V}_L$ and another gluon with some other factor of $\bar{U}$, it is still true that as far as elastic amplitude is concerned the action of $\bar{V}_L$ and $\bar{V}$ is identical. The difference only appears in the inelastic amplitude, but here again it appears as α_s suppressed correction through a diagram analogous to that of figure 2, see figure 7. This correction is not enhanced by the number of target gluons, and thus is indeed negligible in the dense-dilute limit. We therefore conclude that in the dense dilute limit we can safely replace $\bar{V}_L$ by $\bar{V}$. The same is obviously true for $\bar{V}_R$. Thus in the dense-dilute limit in H_{RFT} we can replace

$$\bar{V}_L \rightarrow \bar{V}; \quad \bar{V}_R \rightarrow \bar{V}^\dagger \quad (4.1)$$

Another simplification follows since any factor of U , V_L or V_R can be expanded to second order as only two gluons can be exchanged by any of the target gluons. Thus in the dense-dilute limit we have

$$\begin{aligned} V_L(\mathbf{x}) &= 1 + \int_{\mathbf{y}} ig\phi(\mathbf{x}-\mathbf{y})t^e\mathcal{J}_L^e(\mathbf{y}) - \frac{g^2}{2} \int_{\mathbf{y},\mathbf{z}} \phi(\mathbf{x}-\mathbf{y})\phi(\mathbf{x}-\mathbf{z})t^et^d\mathcal{J}_L^e(\mathbf{y})\mathcal{J}_L^d(\mathbf{z}), \\ V_R(\mathbf{x}) &= 1 - \int_{\mathbf{y}} ig\phi(\mathbf{x}-\mathbf{y})t^e\mathcal{J}_R^e(\mathbf{y}) - \frac{g^2}{2} \int_{\mathbf{y},\mathbf{z}} \phi(\mathbf{x}-\mathbf{y})\phi(\mathbf{x}-\mathbf{z})t^et^d\mathcal{J}_R^e(\mathbf{y})\mathcal{J}_R^d(\mathbf{z}). \end{aligned} \quad (4.2)$$

With these simplification we now consider the RFT Hamiltonian. Let us concentrate on $H_{RFT}^{(1)}$

$$H_{RFT}^{(1)} \approx \frac{1}{\pi g^2} \int_{\mathbf{x}} \bar{V}^{\beta\lambda}(\mathbf{x})\bar{V}^{\dagger\kappa\alpha}(\mathbf{x})\partial^2 \left[V_L^{\alpha\beta}(\mathbf{x})V_R^{\lambda\kappa}(\mathbf{x}) \right]. \quad (4.3)$$

with the understanding that V_L, V_R are expanded to second order. The zeroth order in expansion, the product $V_L V_R$ is a constant and does not contribute to the Hamiltonian due to derivative acting on it. The first order also vanishes because it involves a factor $\text{Tr}(t^a) = 0$. At second order there are three terms

$$\begin{aligned} V_L^{\alpha\beta}(\mathbf{x})V_R^{\lambda\kappa}(\mathbf{x}) &= \frac{g^2}{2} \int_{\mathbf{y},\mathbf{z}} i\phi(\mathbf{x}-\mathbf{y})i\phi(\mathbf{x}-\mathbf{z}) \left[-2t_{\alpha\beta}^et_{\lambda\kappa}^d\mathcal{J}_L^e(\mathbf{y})\mathcal{J}_R^d(\mathbf{z}) + (t^et^d)_{\alpha\beta}\delta^{\lambda\kappa}\mathcal{J}_L^e(\mathbf{y})\mathcal{J}_L^d(\mathbf{z}) \right. \\ &\quad \left. + (t^et^d)_{\lambda\kappa}\delta^{\alpha\beta}\mathcal{J}_R^e(\mathbf{y})\mathcal{J}_R^d(\mathbf{z}) \right]. \end{aligned} \quad (4.4)$$

Substituting the above expression into H_{RFT} , one obtains

$$\begin{aligned}
 H_{RFT}^{(1)} &= \frac{1}{2\pi} \int_{\mathbf{x}, \mathbf{y}, \mathbf{z}} \partial_{\mathbf{x}}^2 [i\phi(\mathbf{x}-\mathbf{y})i\phi(\mathbf{x}-\mathbf{z})] \left(-2\bar{V}^{\beta\lambda}(\mathbf{x})\bar{V}^{\dagger\kappa\alpha}(\mathbf{x})[t^e \mathcal{J}_L^e(\mathbf{y})]_{\alpha\beta}[t^d \mathcal{J}_R^d(\mathbf{z})]_{\lambda\kappa} \right. \\
 &\quad \left. + [t^e \mathcal{J}_L^e(\mathbf{y})]_{\alpha\beta}[t^d \mathcal{J}_L^d(\mathbf{z})]_{\beta\alpha} + [t^e \mathcal{J}_R^e(\mathbf{y})]_{\alpha\beta}[t^d \mathcal{J}_R^d(\mathbf{z})]_{\beta\alpha} \right) \\
 &= \frac{1}{4\pi} \int_{\mathbf{x}, \mathbf{y}, \mathbf{z}} \partial_{\mathbf{x}}^2 [i\phi(\mathbf{x}-\mathbf{y})i\phi(\mathbf{x}-\mathbf{z})] \left[-2\bar{U}^{ed}(\mathbf{x})\mathcal{J}_L^e(\mathbf{y})\mathcal{J}_R^d(\mathbf{z}) + \mathcal{J}_L^e(\mathbf{y})\mathcal{J}_L^e(\mathbf{z}) + \mathcal{J}_R^e(\mathbf{y})\mathcal{J}_R^e(\mathbf{z}) \right] \\
 &= \frac{1}{2\pi} \int_{\mathbf{x}, \mathbf{y}, \mathbf{z}} [i\partial_{\mathbf{x}}\phi(\mathbf{x}-\mathbf{y})i\partial_{\mathbf{x}}\phi(\mathbf{x}-\mathbf{z})] \left[-2\bar{U}^{ed}(\mathbf{x})\mathcal{J}_L^e(\mathbf{y})\mathcal{J}_R^d(\mathbf{z}) + \mathcal{J}_L^e(\mathbf{y})\mathcal{J}_L^e(\mathbf{z}) + \mathcal{J}_R^e(\mathbf{y})\mathcal{J}_R^e(\mathbf{z}) \right].
 \end{aligned} \tag{4.5}$$

Note that the spatial derivatives generate other terms

$$\partial_{\mathbf{x}}^2(\phi(\mathbf{x}-\mathbf{y})\phi(\mathbf{x}-\mathbf{z})) = [\partial_{\mathbf{x}}^2\phi(\mathbf{x}-\mathbf{y})]\phi(\mathbf{x}-\mathbf{z}) + \phi(\mathbf{x}-\mathbf{y})[\partial_{\mathbf{x}}^2\phi(\mathbf{x}-\mathbf{z})] + 2\partial_{\mathbf{x}}\phi(\mathbf{x}-\mathbf{y})\partial_{\mathbf{x}}\phi(\mathbf{x}-\mathbf{z}). \tag{4.6}$$

However, with $\partial_{\mathbf{x}}^2\phi(\mathbf{x}-\mathbf{y}) = g\delta(\mathbf{x}-\mathbf{y})$ and $\partial_{\mathbf{x}}^2\phi(\mathbf{x}-\mathbf{z}) = g\delta(\mathbf{x}-\mathbf{z})$, performing the integration over $\mathbf{x}$ and using the relations $\bar{U}^{ed}(\mathbf{y})\mathcal{J}_L^e(\mathbf{y}) = \mathcal{J}_R^d(\mathbf{y})$ and $\bar{U}^{ed}(\mathbf{z})\mathcal{J}_R^d(\mathbf{z}) = \mathcal{J}_L^e(\mathbf{z})$, these additional terms cancel each other. Thus only the term where the two derivatives separately act on $\phi(\mathbf{x}-\mathbf{y})$ and $\phi(\mathbf{x}-\mathbf{z})$ survives. Performing the same calculation for $H_{RFT}^{(1)c}$ we find to this order an identical result. Thus in the dense-dilute approximation we get

$$\begin{aligned}
 H_{RFT} &\rightarrow H_{JIMWLK} \\
 &= \frac{1}{2\pi} \int_{\mathbf{x}, \mathbf{y}, \mathbf{z}} [i\partial_{\mathbf{x}}\phi(\mathbf{x}-\mathbf{y})i\partial_{\mathbf{x}}\phi(\mathbf{x}-\mathbf{z})] \left[-2\bar{U}^{ed}(\mathbf{x})\mathcal{J}_L^e(\mathbf{y})\mathcal{J}_R^d(\mathbf{z}) + \mathcal{J}_L^e(\mathbf{y})\mathcal{J}_L^e(\mathbf{z}) + \mathcal{J}_R^e(\mathbf{y})\mathcal{J}_R^e(\mathbf{z}) \right].
 \end{aligned} \tag{4.7}$$

There is one subtlety in this derivation which we need to address, i.e. at what order does the correction to eq. (4.1) affect the calculation. To answer this we need to develop a controlled expansion of H_{RFT} in the dense-dilute limit. To do this we note that although we have justified eqs. (4.1) and (4.2) by analyzing the contributions to the S-matrix generated by exchanges of at most two gluon, the same result can be obtained formally by taking the limit of small ρ . It is obvious that at small ρ , the operators V_L and V_R should be simply expanded in power series in $\mathcal{J}_{L(R)}$ to the leading order to which the Hamiltonian does not vanish, leading to eq. (4.2). On the other hand at small ρ we should also expand $\mathcal{I}_{L(R)}$ to leading order in ρ , which gives

$$\mathcal{I}_L^a = \mathcal{I}_R^a = \frac{-i}{g} \frac{\delta}{\delta\alpha^a(\mathbf{x})}; \quad \bar{V}_L = V; \quad \bar{V}_R = \bar{V}^\dagger. \tag{4.8}$$

In fact expansion in powers of ρ is the proper formal way to derive the form of the Hamiltonian in the dense-dilute limit.

Formally expanding $H_{RFT}^{(1)}$ in powers of ρ we see that H_{JIMWLK} arises at order ρ^2 by multiplying the $O(1)$ term in $\bar{V}_L\bar{V}_R$ and $O(\rho^2)$ term in V_LV_R . However we also have to consider a possible contribution arising from $O(\rho)$ term in $\bar{V}_L\bar{V}_R$ (the first order correction

to eq. (4.8)) multiplied by $O(\rho)$ term in $V_L V_R$. We write this additional term as

$$\begin{aligned}\delta H_{JIMWLK} &= \frac{1}{\pi g^2} \int_{\mathbf{x}} \left(\bar{V}_L^{\beta\lambda}(\mathbf{x}) \bar{V}^{\dagger\kappa\alpha} + \bar{V}^{\beta\lambda} \bar{V}_R^{\kappa\alpha}(\mathbf{x}) \right) \partial^2 \left[V_L^{\alpha\beta}(\mathbf{x}) \delta^{\lambda\kappa} + \delta^{\alpha\beta} V_R^{\lambda\kappa}(\mathbf{x}) \right] \\ &= \frac{1}{\pi g^2} \int_{\mathbf{x}} \left((\bar{V}_L \bar{V}^\dagger)^{\beta\alpha} + (\bar{V} \bar{V}_R)^{\beta\alpha} \right) \left[i g^2 t_{\alpha\beta}^e \mathcal{J}_L^e(\mathbf{x}) \right] \\ &\quad + \left((\bar{V}^\dagger \bar{V}_L)^{\kappa\lambda} + (\bar{V}_R \bar{V})^{\kappa\lambda} \right) \left[-i g^2 t_{\lambda\kappa}^d \mathcal{J}_R^d(\mathbf{x}) \right]\end{aligned}\tag{4.9}$$

Here $\bar{V}_L$ and $\bar{V}_R$ are understood as expanded to $O(g)$, however we will not need the explicit form of this expansion, since we will show that this expression vanishes.

We use the two identities

$$\bar{V}_{\lambda\alpha}^\dagger t_{\alpha\beta}^e \bar{V}_{\beta\gamma} = \bar{U}^{ed} t_{\lambda\gamma}^d\tag{4.10}$$

and

$$\bar{U}^{\dagger de}(\mathbf{x}) \mathcal{J}_L^e(\mathbf{x}) = \mathcal{J}_R^d(\mathbf{x}).\tag{4.11}$$

Here, as before $\bar{V}^\dagger = \exp\{-t^a \frac{\delta}{\delta \rho^a}\}$ and $\bar{V} = \exp\{t^a \frac{\delta}{\delta \rho^a}\}$ are defined in the fundamental representation while $\bar{U} = \exp\{T^a \frac{\delta}{\delta \rho^a}\}$ is defined in the adjoint representation. We then calculate

$$(\bar{V}_L \bar{V}^\dagger)^{\beta\alpha} t_{\alpha\beta}^e \mathcal{J}_L^e = \bar{V}_L^{\beta\lambda} \left[\bar{U}^{ed} t_{\lambda\gamma}^d \bar{V}^{\dagger\gamma\beta} \right] \mathcal{J}_L^e = \mathcal{J}_R^d t_{\lambda\gamma}^d (\bar{V}^\dagger \bar{V}_L)^{\gamma\lambda}\tag{4.12}$$

and

$$(\bar{V} \bar{V}_R)^{\beta\alpha} t_{\alpha\beta}^e \mathcal{J}_L^e = \left[\bar{U}^{ed} \bar{V}_{\alpha\lambda} t_{\lambda\gamma}^d \right] \bar{V}_R^{\gamma\alpha} \mathcal{J}_L^e = \mathcal{J}_R^d t_{\lambda\gamma}^d (\bar{V}_R \bar{V})^{\gamma\lambda}.\tag{4.13}$$

Thus the four terms in eq. (4.9) pairwise cancel.

We have thus proved that when expanded to second order in ρ , the Hamiltonian $H_{RFT}^{(1)}$ reproduces H_{JIMWLK} . It is obvious that the same is true for $H_{RFT}^{(1)c}$, since H_{JIMWLK} is charge conjugation invariant.

If instead of expanding in powers of ρ , we expand in powers of $\delta/\delta\rho$, the leading order expansion gives H_{KLWMIJ} , eq. (3.9). This is easily done explicitly, but the final result is obvious by duality.

Finally we note that the exact same result is obtained if we were to use the left and right Wilson lines not in the fundamental but in any other representation of $SU(N)$. The only property of the $SU(N)$ matrices that is needed to derive H_{JIMWLK} in eq. (4.5) is

$$\text{Tr}[\bar{V}^\dagger t^e \bar{V} t^d] = \frac{1}{2} \bar{U}^{ed}\tag{4.14}$$

for a fundamental matrix $\bar{V}$ and an adjoint matrix $\bar{U}$. However a similar relation holds for $SU(N)$ matrices in any representation D

$$\text{Tr}[\bar{U}_D^\dagger T_D^a \bar{U}_D T_D^b] = \frac{C_2(D) R_D}{N^2 - 1} \bar{U}^{ab}.\tag{4.15}$$

Here $\bar{U}_D$ is a matrix and T_D is a generator in an arbitrary representation D of $SU(N)$, and $C_2(D)$ and R_D are the second Casimir and the dimensionality of D respectively. Thus using U_D and $\bar{U}_D$ in any representation in the definition of H_{RFT} will reproduce H_{JIMWLK} and H_{KLWMIJ} in expansion once the overall normalization is adjusted.

5 Continuous symmetries

Let us now discuss the continuous symmetries of H_{RFT} . As we have mentioned above, both the JIMWLK and the KLWMIJ Hamiltonians have a continuous $SU(N) \times SU(N)$ symmetry, albeit those are distinct symmetry transformations. The $SU(N) \times SU(N)$ symmetry of H_{JIMWLK} is generated by the charges

$$Q_L^a = \int d^2\mathbf{z} \mathcal{J}_L^a(\mathbf{z}); \quad Q_R^a = \int d^2\mathbf{z} \mathcal{J}_R^a(\mathbf{z}) \quad (5.1)$$

while the $SU(N) \times SU(N)$ symmetry of H_{KLWMIJ} by

$$\bar{Q}_L^a = \int d^2\mathbf{z} \mathcal{I}_L^a(\mathbf{z}); \quad \bar{Q}_R^a = \int d^2\mathbf{z} \mathcal{I}_R^a(\mathbf{z}). \quad (5.2)$$

It is an interesting question which of these symmetries are also the symmetries of the self dual H_{RFT} eq. (3.35). The question is not completely straightforward to answer even though we do have an explicit representation of the charge operators on the RFT Hilbert space. The reason is that the commutation relations between $\mathcal{J}_{L(R)}$ and $\bar{V}_{L(R)}$ as well as between $\mathcal{I}_{L(R)}$ and $V_{L(R)}$ are quite complicated. We will nevertheless try to answer this question, using a perturbative expansion. Our answer is somewhat surprising: the symmetry of H_{RFT} appears to be $SU(N) \times SU(N) \times SU(N)$.³

We start with discussing the vector part of the group, which is the easiest and can be analyzed without recourse to perturbation theory.

To better organize the calculation, we rescale the charge density $\tilde{\rho}^a(\mathbf{x}) = g\rho^a(\mathbf{x})$ and also introduce $\tilde{\phi}(\mathbf{x} - \mathbf{y}) = \frac{1}{g}\phi(\mathbf{x} - \mathbf{y})$. Then $\mathcal{J}_L^a, \mathcal{J}_R^a, \mathcal{I}_L^a, \mathcal{I}_R^a$ can be Taylor expanded by counting the powers of the coupling constant g . We will use this expansion in this and the next sections. We will refer to this counting in powers of the coupling constant as the “BFKL counting”, since it is equivalent to simultaneous expansion in powers of ρ and $\delta/\delta\rho$.

5.1 The vector $SU_V(N)$ symmetry

The analysis of the vector symmetry is facilitated by the following simple observation

$$Q_L - Q_R = \bar{Q}_R - \bar{Q}_L. \quad (5.3)$$

To prove this we note that

$$\mathcal{J}_L^a(\mathbf{z}) - \mathcal{J}_R^a(\mathbf{z}) = \tilde{\rho}^b(\mathbf{z}) T_{bc}^a \frac{\delta}{\delta \tilde{\rho}^c(\mathbf{z})} \quad (5.4)$$

$$\mathcal{I}_R^a(\mathbf{z}) - \mathcal{I}_L^a(\mathbf{z}) = \tilde{\alpha}^b(\mathbf{z}) T_{bc}^a \frac{\delta}{\delta \tilde{\alpha}^c(\mathbf{z})} = \int d^2\mathbf{y} \tilde{\phi}(\mathbf{z} - \mathbf{y}) \tilde{\rho}^b(\mathbf{y}) T_{bc}^a \partial_{\mathbf{z}}^2 \frac{\delta}{\delta \tilde{\rho}^c(\mathbf{z})}. \quad (5.5)$$

Integrating by parts we find

$$\bar{Q}_R - \bar{Q}_L = \int d^2\mathbf{z} d^2\mathbf{y} \tilde{\phi}(\mathbf{z} - \mathbf{y}) \tilde{\rho}^b(\mathbf{y}) T_{bc}^a \partial_{\mathbf{z}}^2 \frac{\delta}{\delta \tilde{\rho}^c(\mathbf{z})} = \int d^2\mathbf{z} \tilde{\rho}^b(\mathbf{z}) T_{bc}^a \frac{\delta}{\delta \tilde{\rho}^c(\mathbf{z})} = Q_L - Q_R. \quad (5.6)$$

³To be precise, while $SU(N) \times SU(N)$ is there, the third $SU(N)$ does not necessary form a direct product with the first two. We have not attempted to write down the full algebra of the currents, which appears to be quite complicated.

It is now straightforward to check that the vector $SU_V(N)$ transformation generated by $Q_L - Q_R$ is the symmetry of H_{RFT} . By virtue of eq. (5.3) the charge $Q_V^a \equiv Q_L^a - Q_R^a$ acts as a rotation generator on all the currents, i.e.

$$\begin{aligned} [Q_V^a, \mathcal{J}_L^b] &= i f^{abc} \mathcal{J}_L^c, \\ [Q_V^a, \mathcal{J}_R^b] &= i f^{abc} \mathcal{J}_R^c, \\ [Q_V^a, \mathcal{I}_L^b] &= i f^{abc} \mathcal{I}_L^c, \\ [Q_V^a, \mathcal{I}_R^b] &= i f^{abc} \mathcal{I}_R^c. \end{aligned} \tag{5.7}$$

It then follows that for a finite group transformation

$$\hat{W} = \exp \{ i \lambda^a Q_V^a \} \tag{5.8}$$

we have

$$\hat{W}^\dagger \mathcal{J}_{L(R)}^a(\mathbf{x}) \hat{W} = \mathcal{W}_A^{ab} \mathcal{J}_{L(R)}^b(\mathbf{x}); \quad \hat{W}^\dagger \mathcal{I}_{L(R)}^a(\mathbf{x}) \hat{W} = \mathcal{W}_A^{ab} \mathcal{I}_{L(R)}^b(\mathbf{x}); \tag{5.9}$$

with

$$\mathcal{W}_A^{ab} = \left[e^{i \lambda^d T^d} \right]^{ab}. \tag{5.10}$$

As a consequence

$$\hat{W}^\dagger V_L^{\alpha\beta}(\mathbf{x}) \hat{W} = \left(\mathcal{W}_F V_L(\mathbf{x}) \mathcal{W}_F^\dagger \right)^{\alpha\beta} \tag{5.11}$$

with the fundamental representation matrix

$$\mathcal{W}_F^{\kappa\beta} = \left[e^{i \lambda^e t^e} \right]^{\kappa\beta}. \tag{5.12}$$

The same transformation as in eq. (5.11) applies to $V_R(\mathbf{x})$, as well as to $\bar{V}_{L(R)}$. It is now obvious that H_{RFT} is invariant under $SU_V(N)$.

5.2 Is $SU_L(N)$ there?

Let us now consider other transformations generated by the left and right charges. The analysis for all of them is similar, and we will concentrate on Q_L . The question we are asking, does Q_L commute with H_{RFT} ?

What is the action of Q_L^a on the building blocks of H_{RFT} ? The answer for V_L and V_R is obvious. Under the $SU_L(N)$ transformation

$$\hat{S} = \exp \{ i \lambda^a Q_L^a \} \tag{5.13}$$

we have

$$\hat{S}^\dagger \mathcal{J}_L^e(\mathbf{z}) \hat{S} = \mathcal{S}_A^{ed} \mathcal{J}_L^d(\mathbf{z}); \quad \hat{S}^\dagger \mathcal{J}_R^e(\mathbf{z}) \hat{S} = \mathcal{J}_R^e(\mathbf{z}) \tag{5.14}$$

with

$$\mathcal{S}_A^{ed}(\mathbf{x}) = \left[e^{i \lambda^a T^a} \right]^{ed}. \tag{5.15}$$

As a consequence,

$$\hat{S}^\dagger V_L^{\alpha\beta}(\mathbf{x}) \hat{S} = \mathcal{S}_F^{\alpha\gamma} V_L^{\gamma\kappa}(\mathbf{x}) \mathcal{S}_F^{\dagger\kappa\beta}; \quad \hat{S}^\dagger V_R^{\alpha\beta}(\mathbf{x}) \hat{S} = V_R^{\alpha\beta}(\mathbf{x}) \quad (5.16)$$

with

$$S_F^{\kappa\beta} = \left[e^{i\lambda^e t^e} \right]^{\kappa\beta}. \quad (5.17)$$

What is the transformation of $\bar{V}_L$ and $\bar{V}_R$? Examining the expression for H_{RFT} we see that if the transformation was

$$\hat{S}^\dagger \bar{V}_L^{\beta\gamma}(\mathbf{x}) \hat{S} = \mathcal{S}_F^{\beta\kappa} \bar{V}_L^{\kappa\gamma}(\mathbf{x}); \quad \hat{S}^\dagger \bar{V}_R^{\beta\gamma}(\mathbf{x}) \hat{S} = \bar{V}_R^{\beta\kappa} \mathcal{S}_F^{\kappa\gamma}(\mathbf{x}); \quad (\text{true or false ?}) \quad (5.18)$$

the Hamiltonian would be invariant under $SU_L(N)$. Indeed if instead of $\bar{V}_L$ and $\bar{V}_R$ we had $\bar{V}$ and $\bar{V}^\dagger$, this would be the case. This is precisely what happens in the JIMWLK limit.

The transformation eq. (5.18) is equivalent to the commutation relation

$$\left[Q_L^a, \bar{V}_L^{\alpha\beta}(\mathbf{x}) \right] = - (t^a \bar{V}_L(\mathbf{x}))^{\alpha\beta} \quad (\text{true or false?}) \quad (5.19)$$

and similarly for $\bar{V}_R$.

We were unable to calculate the commutation relation in eq. (5.19) in a closed form. However we were able to calculate first several orders in perturbative expansion in g . We performed the calculation in the BFKL counting of orders of g . The details of the calculation are presented in the appendix A. Our results are the following.

We have calculated the commutator between Q_L^a and $\bar{V}_L$ up to order g^3 and found that relation eq. (5.19) holds up to order g^2 , but is violated at order g^3 .

We have also calculated the commutator of Q_L^a with the Hamiltonian $[Q_L^a, H_{RFT}]$ up to order g^6 . We have found that this commutator vanishes up to this order. This leads us to believe that even though eq. (5.19) is not satisfied, the $SU_L(N)$ is indeed a symmetry of H_{RFT} . We stress that we do not have a closed form proof of this, but only perturbative calculation to order g^6 .

The analysis of $\bar{Q}_L^a$ is identical, since Q and $\bar{Q}$ are related by duality transformation. Thus we believe that $\bar{Q}_L^a$ also commutes with the Hamiltonian.

If this is indeed the case, the continuous symmetry of H_{RFT} is at least $SU(N) \times SU(N) \times SU(N)$. In fact the symmetry could be even larger since we have not calculated the commutators $[Q_L, \bar{Q}_L]$. If this commutator does not close on any of the four charges (or their products) $Q_{L(R)}, \bar{Q}_{L(R)}$ the symmetry group is larger. We have not investigated this question any further.

6 Is this the “Diamond action”?

The family of Hamiltonians that we have identified carries uncanny resemblance to the so called “Diamond action” suggested in [67] and also discussed in [68]. There is of course a host of differences between our approach and that of [67] and [68]. On the technical level we are dealing with the Hamiltonian formulation of RFT together with the accompanying field algebra and the structure of the RFT Hilbert space, while these references strive

to derive the effective action in terms of certain Wilson line functions. On the other hand [67] and [68] derive the action directly from QCD (although in both cases certain not entirely straightforward approximations are utilized) whereas our expression is an ansatz constrained by the expected symmetries and the appropriate limiting forms.

Nevertheless, abstracting ourselves from these differences we can compare H_{RFT} with the effective action of [67]. We concentrate on the Hamiltonian eq. (3.35) defined with Wilson line in the *adjoint representation*.

$$\begin{aligned}
 U_L(\mathbf{x}) &= \text{Exp} \left\{ i \int_{\mathbf{y}} g \phi(\mathbf{x} - \mathbf{y}) T^e \mathcal{J}_L^e(\mathbf{y}) \right\} \\
 U_R(\mathbf{x}) &= \text{Exp} \left\{ -i \int_{\mathbf{y}} g \phi(\mathbf{x} - \mathbf{y}) T^e \mathcal{J}_R^e(\mathbf{y}) \right\} \\
 \bar{U}_L(\mathbf{x}) &= \text{Exp} \left\{ i \int_{\mathbf{y}} g \phi(\mathbf{x} - \mathbf{y}) T^e \mathcal{I}_L^e(\mathbf{y}) \right\} \\
 \bar{U}_R(\mathbf{x}) &= \text{Exp} \left\{ -i \int_{\mathbf{y}} g \phi(\mathbf{x} - \mathbf{y}) T^e \mathcal{I}_R^e(\mathbf{y}) \right\}.
 \end{aligned} \tag{6.1}$$

In this case the two terms in eq. (3.35) are equal and we have

$$H_{RFT}^A = \frac{1}{2\pi g^2 N} \int d^2 \mathbf{x} \partial^2 [\bar{U}_L^{bc}(\mathbf{x}) \bar{U}_R^{da}(\mathbf{x})] U_L^{ab}(\mathbf{x}) U_R^{cd}(\mathbf{x}). \tag{6.2}$$

It is easily checked that with the correspondence

$$U_L \rightarrow W_{-\infty}, \quad U_R \rightarrow W_{\infty}^{\dagger}, \quad \bar{U}_L \rightarrow V_{-\infty}, \quad \bar{U}_R \rightarrow V_{\infty}^{\dagger} \tag{6.3}$$

our eq. (6.2) looks identical to the effective action suggested in [67]. However beyond the looks there are significant differences between the two. In particular in [67] the four Wilson lines are not independent, but satisfy the so called diamond condition

$$V_{\infty}^{\dagger} W_{-\infty} V_{-\infty} W_{\infty}^{\dagger} = 1. \tag{6.4}$$

This relation was essential in the derivation of [67] and only using this relation the effective action obtained in [67] could be written in the form eq. (6.2). On the other hand in our framework, although all four Wilson line operators are expressible in terms of ρ and $\delta/\delta\rho$, there is no such condition that constrains the four.

We can check eq. (6.4) explicitly, expanding all the operators $U_{L,R}$ and $\bar{U}_{L,R}$ to first order in the respective left and right charge densities. In our notations eq. (6.4) corresponds to

$$\bar{U}_R(\mathbf{x}) U_L(\mathbf{x}) \bar{U}_L(\mathbf{x}) U_R(\mathbf{x}) = 1. \tag{6.5}$$

We will calculate the l.h.s. of eq. (6.5) to second order in g . To this order we need

$$\begin{aligned}\mathcal{J}_L^a(\mathbf{x}) &= \frac{1}{g} \left[\frac{1}{2} g T^e \frac{\delta}{\delta \tilde{\rho}^e(\mathbf{x})} \left(\coth \left[\frac{1}{2} g T^e \frac{\delta}{\delta \tilde{\rho}^e(\mathbf{x})} \right] - 1 \right) \right]^{ba} \tilde{\rho}^b(\mathbf{x}), \\ &= \frac{1}{g} \tilde{\rho}^a(\mathbf{x}) - \frac{1}{2} \tilde{\rho}^b(\mathbf{x}) T_{ba}^e \frac{\delta}{\delta \tilde{\rho}^e(\mathbf{x})} + \mathcal{O}(g)\end{aligned}\quad (6.6)$$

$$\begin{aligned}\mathcal{J}_R^a(\mathbf{x}) &= \frac{1}{g} \left[\frac{1}{2} g T^e \frac{\delta}{\delta \tilde{\rho}^e(\mathbf{x})} \left(\coth \left[\frac{1}{2} g T^e \frac{\delta}{\delta \tilde{\rho}^e(\mathbf{x})} \right] + 1 \right) \right]^{ba} \tilde{\rho}^b(\mathbf{x}), \\ &= \frac{1}{g} \tilde{\rho}^a(\mathbf{x}) + \frac{1}{2} \tilde{\rho}^b(\mathbf{x}) T_{ba}^e \frac{\delta}{\delta \tilde{\rho}^e(\mathbf{x})} + \mathcal{O}(g).\end{aligned}\quad (6.7)$$

$$\begin{aligned}\mathcal{I}_L^a(\mathbf{x}) &= \frac{-i}{g} \partial^2 \frac{\delta}{\delta \tilde{\rho}^b(\mathbf{x})} \left[\frac{1}{2} i g T^e \frac{1}{\partial^2} \tilde{\rho}^e(\mathbf{x}) \left(\coth \left[\frac{1}{2} i g T^e \frac{1}{\partial^2} \tilde{\rho}^e(\mathbf{x}) \right] - 1 \right) \right]^{ba}, \\ &= \frac{-i}{g} \partial^2 \frac{\delta}{\delta \tilde{\rho}^a(\mathbf{x})} - \frac{1}{2} T_{ba}^e \partial^2 \frac{\delta}{\delta \tilde{\rho}^b(\mathbf{x})} \int_{\mathbf{z}} \tilde{\phi}(\mathbf{x} - \mathbf{z}) \tilde{\rho}^e(\mathbf{z}) + \mathcal{O}(g).\end{aligned}\quad (6.8)$$

$$\begin{aligned}\mathcal{I}_R^a(\mathbf{x}) &= \frac{-i}{g^2} \partial^2 \frac{\delta}{\delta \rho^b(\mathbf{x})} \left[\frac{1}{2} T^e i g^2 \frac{1}{\partial^2} \rho^e(\mathbf{x}) \left(\coth \left[\frac{1}{2} T^e i g^2 \frac{1}{\partial^2} \rho^e(\mathbf{x}) \right] + 1 \right) \right]^{ba}, \\ &= \frac{-i}{g} \partial^2 \frac{\delta}{\delta \tilde{\rho}^a(\mathbf{x})} + \frac{1}{2} T_{ba}^e \partial^2 \frac{\delta}{\delta \tilde{\rho}^b(\mathbf{x})} \int_{\mathbf{z}} \tilde{\phi}(\mathbf{x} - \mathbf{z}) \tilde{\rho}^e(\mathbf{z}) + \mathcal{O}(g).\end{aligned}\quad (6.9)$$

From the definition of $\mathcal{I}_L^a$ and $\mathcal{I}_R^a$, one obtains

$$\begin{aligned}& i g^2 \int_{\mathbf{y}} \tilde{\phi}(\mathbf{x} - \mathbf{y}) T^e \mathcal{I}_L^e(\mathbf{y}) \\ &= i g^2 \int_{\mathbf{y}} \tilde{\phi}(\mathbf{x} - \mathbf{y}) T^e \left(\frac{-i}{g} \partial_{\mathbf{y}}^2 \frac{\delta}{\delta \tilde{\rho}^e(\mathbf{y})} - \frac{1}{2} \partial_{\mathbf{y}}^2 \frac{\delta}{\delta \tilde{\rho}^b(\mathbf{y})} T_{be}^a \int_{\mathbf{z}} \tilde{\phi}(\mathbf{y} - \mathbf{z}) \tilde{\rho}^a(\mathbf{z}) + \mathcal{O}(g) \right) \\ &= g T^e \frac{\delta}{\delta \tilde{\rho}^e(\mathbf{x})} - \frac{1}{2} i g^2 T^e T_{ab}^e \int_{\mathbf{y}, \mathbf{z}} \tilde{\phi}(\mathbf{x} - \mathbf{y}) \tilde{\phi}(\mathbf{y} - \mathbf{z}) \partial_{\mathbf{y}}^2 \frac{\delta}{\delta \tilde{\rho}^b(\mathbf{y})} \tilde{\rho}^a(\mathbf{z}) + \mathcal{O}(g^3) \\ &= g T^e \frac{\delta}{\delta \tilde{\rho}^e(\mathbf{x})} - \frac{1}{2} i g^2 T^e T_{ab}^e \int_{\mathbf{y}} \tilde{\phi}(\mathbf{x} - \mathbf{y}) \frac{\delta}{\delta \tilde{\rho}^b(\mathbf{y})} \tilde{\rho}^a(\mathbf{y}) - \frac{1}{2} i g^2 T^e T_{ab}^e \int_{\mathbf{z}} \tilde{\phi}(\mathbf{x} - \mathbf{z}) \frac{\delta}{\delta \tilde{\rho}^b(\mathbf{x})} \tilde{\rho}^a(\mathbf{z}) \\ &\quad - i g^2 T^e T_{ab}^e \int_{\mathbf{y}, \mathbf{z}} \partial_{\mathbf{y}} \tilde{\phi}(\mathbf{x} - \mathbf{y}) \partial_{\mathbf{y}} \tilde{\phi}(\mathbf{y} - \mathbf{z}) \frac{\delta}{\delta \tilde{\rho}^b(\mathbf{y})} \tilde{\rho}^a(\mathbf{z}) + \mathcal{O}(g^3).\end{aligned}\quad (6.10)$$

We have used integration by parts. As a consequence

$$\begin{aligned}\bar{U}_L(\mathbf{x}) &= \exp \left\{ i g^2 \int_{\mathbf{y}} \tilde{\phi}(\mathbf{x} - \mathbf{y}) T^e \mathcal{I}_L^e(\mathbf{y}) \right\} \\ &= 1 + g T^e \frac{\delta}{\delta \tilde{\rho}^e(\mathbf{x})} - \frac{1}{2} i g^2 T^e T_{ab}^e \int_{\mathbf{y}} \tilde{\phi}(\mathbf{x} - \mathbf{y}) \frac{\delta}{\delta \tilde{\rho}^b(\mathbf{y})} \tilde{\rho}^a(\mathbf{y}) - \frac{1}{2} i g^2 T^e T_{ab}^e \int_{\mathbf{z}} \tilde{\phi}(\mathbf{x} - \mathbf{z}) \frac{\delta}{\delta \tilde{\rho}^b(\mathbf{x})} \tilde{\rho}^a(\mathbf{z}) \\ &\quad - i g^2 T^e T_{ab}^e \int_{\mathbf{y}, \mathbf{z}} \partial_{\mathbf{y}} \tilde{\phi}(\mathbf{x} - \mathbf{y}) \partial_{\mathbf{y}} \tilde{\phi}(\mathbf{y} - \mathbf{z}) \frac{\delta}{\delta \tilde{\rho}^b(\mathbf{y})} \tilde{\rho}^a(\mathbf{z}) + \frac{1}{2!} g^2 T^e T^d \frac{\delta}{\delta \tilde{\rho}^e(\mathbf{x})} \frac{\delta}{\delta \tilde{\rho}^d(\mathbf{x})} + \mathcal{O}(g^3)\end{aligned}\quad (6.11)$$

and

$$\begin{aligned}
 \bar{U}_R(\mathbf{x}) &= \exp \left\{ -ig^2 \int_{\mathbf{y}} \tilde{\phi}(\mathbf{x}-\mathbf{y}) T^e \mathcal{I}_R^e(\mathbf{y}) \right\} \\
 &= 1 - gT^e \frac{\delta}{\delta \tilde{\rho}^e(\mathbf{x})} - \frac{1}{2} ig^2 T^e T_{ab}^e \int_{\mathbf{y}} \tilde{\phi}(\mathbf{x}-\mathbf{y}) \frac{\delta}{\delta \tilde{\rho}^b(\mathbf{y})} \tilde{\rho}^a(\mathbf{y}) - \frac{1}{2} ig^2 T^e T_{ab}^e \int_{\mathbf{z}} \tilde{\phi}(\mathbf{x}-\mathbf{z}) \frac{\delta}{\delta \tilde{\rho}^b(\mathbf{x})} \tilde{\rho}^a(\mathbf{z}) \\
 &\quad - ig^2 T^e T_{ab}^e \int_{\mathbf{y}, \mathbf{z}} \partial_{\mathbf{y}} \tilde{\phi}(\mathbf{x}-\mathbf{y}) \partial_{\mathbf{y}} \tilde{\phi}(\mathbf{y}-\mathbf{z}) \frac{\delta}{\delta \tilde{\rho}^b(\mathbf{y})} \tilde{\rho}^a(\mathbf{z}) + \frac{1}{2!} g^2 T^e T^d \frac{\delta}{\delta \tilde{\rho}^e(\mathbf{x})} \frac{\delta}{\delta \tilde{\rho}^d(\mathbf{x})} + \mathcal{O}(g^3).
 \end{aligned} \tag{6.12}$$

On the other hand, from

$$\begin{aligned}
 &ig^2 \int_{\mathbf{y}} \tilde{\phi}(\mathbf{x}-\mathbf{y}) T^e \mathcal{J}_L^e(\mathbf{y}) \\
 &= ig^2 \int_{\mathbf{y}} \tilde{\phi}(\mathbf{x}-\mathbf{y}) T^e \left(\frac{1}{g} \tilde{\rho}^e(\mathbf{y}) - \frac{1}{2} \tilde{\rho}^b(\mathbf{y}) T_{be}^a \frac{\delta}{\delta \tilde{\rho}^a(\mathbf{y})} + \mathcal{O}(g^3) \right) \\
 &= ig \int_{\mathbf{y}} \tilde{\phi}(\mathbf{x}-\mathbf{y}) T^e \tilde{\rho}^e(\mathbf{y}) - \frac{1}{2} ig^2 T^e T_{ab}^e \int_{\mathbf{y}} \tilde{\phi}(\mathbf{x}-\mathbf{y}) \tilde{\rho}^b(\mathbf{y}) \frac{\delta}{\delta \tilde{\rho}^a(\mathbf{y})} + \mathcal{O}(g^3).
 \end{aligned} \tag{6.13}$$

one obtains

$$\begin{aligned}
 U_L(\mathbf{x}) &= \exp \left\{ ig^2 \int_{\mathbf{y}} \tilde{\phi}(\mathbf{x}-\mathbf{y}) T^e \mathcal{J}_L^e(\mathbf{y}) \right\} \\
 &= 1 + ig \int_{\mathbf{y}} \tilde{\phi}(\mathbf{x}-\mathbf{y}) T^e \tilde{\rho}^e(\mathbf{y}) - \frac{1}{2} ig^2 T^e T_{ab}^e \int_{\mathbf{y}} \tilde{\phi}(\mathbf{x}-\mathbf{y}) \tilde{\rho}^b(\mathbf{y}) \frac{\delta}{\delta \tilde{\rho}^a(\mathbf{y})} \\
 &\quad + \frac{1}{2!} (ig)^2 \int_{\mathbf{y}, \mathbf{z}} \tilde{\phi}(\mathbf{x}-\mathbf{y}) \tilde{\phi}(\mathbf{x}-\mathbf{z}) T^e T^d \tilde{\rho}^e(\mathbf{y}) \tilde{\rho}^d(\mathbf{z}) + \mathcal{O}(g^3).
 \end{aligned} \tag{6.14}$$

$$\begin{aligned}
 U_R(\mathbf{x}) &= \exp \left\{ -ig^2 \int_{\mathbf{y}} \tilde{\phi}(\mathbf{x}-\mathbf{y}) T^e \mathcal{J}_R^e(\mathbf{y}) \right\} \\
 &= 1 - ig \int_{\mathbf{y}} \tilde{\phi}(\mathbf{x}-\mathbf{y}) T^e \tilde{\rho}^e(\mathbf{y}) - \frac{1}{2} ig^2 T^e T_{ab}^e \int_{\mathbf{y}} \tilde{\phi}(\mathbf{x}-\mathbf{y}) \tilde{\rho}^b(\mathbf{y}) \frac{\delta}{\delta \tilde{\rho}^a(\mathbf{y})} \\
 &\quad + \frac{1}{2!} (ig)^2 \int_{\mathbf{y}, \mathbf{z}} \tilde{\phi}(\mathbf{x}-\mathbf{y}) \tilde{\phi}(\mathbf{x}-\mathbf{z}) T^e T^d \tilde{\rho}^e(\mathbf{y}) \tilde{\rho}^d(\mathbf{z}) + \mathcal{O}(g^3).
 \end{aligned} \tag{6.15}$$

At order $\mathcal{O}(g)$ it is obvious that eq. (6.5) is satisfied, and the first nontrivial check of the relation is at $\mathcal{O}(g^2)$. At this order we obtain

$$\begin{aligned}
 &\bar{U}_R(\mathbf{x}) U_L(\mathbf{x}) \bar{U}_L(\mathbf{x}) U_R(\mathbf{x}) \\
 &= 1 - ig^2 T^e T_{ab}^e \int_{\mathbf{z}} \tilde{\phi}(\mathbf{x}-\mathbf{z}) \frac{\delta}{\delta \tilde{\rho}^b(\mathbf{x})} \tilde{\rho}^a(\mathbf{z}) - 2ig^2 T^e T_{ab}^e \int_{\mathbf{y}, \mathbf{z}} \partial_{\mathbf{y}} \tilde{\phi}(\mathbf{x}-\mathbf{y}) \partial_{\mathbf{y}} \tilde{\phi}(\mathbf{y}-\mathbf{z}) \frac{\delta}{\delta \tilde{\rho}^b(\mathbf{y})} \tilde{\rho}^a(\mathbf{z}) \\
 &\quad + ig \int_{\mathbf{y}} \tilde{\phi}(\mathbf{x}-\mathbf{y}) T^a \tilde{\rho}^a(\mathbf{y}) gT^b \frac{\delta}{\delta \tilde{\rho}^b(\mathbf{x})} + gT^b \frac{\delta}{\delta \tilde{\rho}^b(\mathbf{x})} (-ig) \int_{\mathbf{y}} \tilde{\phi}(\mathbf{x}-\mathbf{y}) T^a \tilde{\rho}^a(\mathbf{y}) \\
 &= 1 - 2ig^2 T^e T_{ab}^e \int_{\mathbf{y}, \mathbf{z}} \partial_{\mathbf{y}} \tilde{\phi}(\mathbf{x}-\mathbf{y}) \partial_{\mathbf{y}} \tilde{\phi}(\mathbf{y}-\mathbf{z}) \frac{\delta}{\delta \tilde{\rho}^b(\mathbf{y})} \tilde{\rho}^a(\mathbf{z}) - 2ig^2 T^e T_{ab}^e \int_{\mathbf{z}} \tilde{\phi}(\mathbf{x}-\mathbf{z}) \frac{\delta}{\delta \tilde{\rho}^b(\mathbf{x})} \tilde{\rho}^a(\mathbf{z}) \neq 1.
 \end{aligned} \tag{6.16}$$

Thus we have established that at order $\mathcal{O}(g^2)$, the diamond condition is not satisfied by our Wilson line like operators.

We thus conclude that in spite of certain similarities, the self dual RFT Hamiltonian eq. (6.2) is not the same as the effective action of [67]. The status of this comparison is further discussed in the next section.

7 Discussion

In this paper we have revisited the problem of constructing a self dual Reggeon Field Theory Hamiltonian H_{RFT} . We have followed the EFT strategy by imposing the relevant symmetries and also required that H_{RFT} reduces to H_{JIMWLK} (or H_{KLWMIJ}) in the dense-dilute limit.

As a result we have found a family of Hamiltonians that satisfy these requirements. These Hamiltonians are constructed from Wilson line - like operators in different representations of the $SU(N)$ group. We note that any of these Hamiltonians in addition to reproducing the dense dilute limit, also generates correct Pomeron loops. The simplest way to see this is to perform the coupling constant expansion using the BFKL counting introduced in section 5. This is equivalent to simultaneous expansion in powers of ρ and $\delta/\delta\rho$. At order α_s the Hamiltonian reduced to H_{BFKL} , while at order α_s^2 it contains both splitting and merging vertices ($\rho^2(\delta/\delta\rho)^4$ and $\rho^4(\delta/\delta\rho)^2$) with correct coefficients. As discussed in [38, 58–65, 69] these vertices are responsible both for a certain set of reggeization corrections, and for the QCD Pomeron loops.

We have analyzed the continuous symmetries of H_{RFT} . This is an interesting question since both H_{JIMWLK} and H_{KLWMIJ} possess an $SU_L(N) \times SU_R(N)$ symmetry group, but the generators of these transformations are not the same in the two dense-dilute cases. For H_{RFT} we are able to show nonperturbatively the existence of one $SU_V(N)$ symmetry, which is the diagonal subgroup of the symmetry group in both JIMWLK and KLWMIJ limits. We established the fact that the two diagonal subgroups are identical explicitly using the algebra of the generators in the RFT Hilbert space. We have also shown that H_{RFT} is invariant under the left and right rotations at least to $O(g^6)$ in perturbative expansion. This is a strong indication that the continuous symmetry group is at least $SU(N) \times SU(N) \times SU(N)$.

One member of the family of the Hamiltonians we found is very similar to the “diamond action” [67, 68]. Our Hamiltonian RFT framework is different from the effective action approach of [67, 68] which somewhat hampers direct comparison. Nevertheless if we juxtapose our H_{RFT} defined in terms of adjoint Wilson lines directly with the effective action of [67, 68], the two look identical. There is however one significant difference between our result and that of [67]. Namely the action in [67] is written in terms of four Wilson loops that satisfy the diamond condition, eq. (6.4). This condition played a very important role in [67]. In fact the effective action derived in [67] directly from QCD is equivalent to the “KLWMIJ+” Hamiltonian suggested in [73, 74], whereby KLWMIJ Hamiltonian is generalized by including nonlinear corrections in the solution for classical field. This Hamiltonian is not explicitly self dual, and only with the help of the diamond condition it

was recast in [67] in the form which looks self dual, at least superficially. However whether the “diamond action” is in fact self dual or not remained an open question. To check the self duality one has to verify that the duality transformation is canonical, or in the quantum sense a linear transformation on the RFT Hilbert space. This was not possible to do with the tools of [67], as no operator realization of the algebra of Wilson lines was explicitly presented. In the present paper we operate within the RFT Hilbert space with well defined operator algebra; and therefore we have explicit realization of the duality transformation in the Hilbert space. We find within this consistent framework that the diamond action (RFT Hamiltonian) is self dual, but the diamond condition between the Wilson lines is not satisfied. The condition is violated starting with order $O(g^2)$ in perturbative expansion. In this sense our paper is closer to [68], where the diamond action is derived as a self dual form of the action in the dense-dilute limit without assuming the diamond constraint between the Wilson lines. In [68] the constraint was shown to hold in the first order in perturbation theory, which is consistent with our conclusion here, but was not checked at higher orders.

Our “bottom up” approach does not allow us to decide which one of the candidate hamiltonians we have found is the right one, and in fact whether any one of them is the correct QCD RFT Hamiltonian. Even though we have used the EFT methodology to determine possible terms in H_{RFT} , we are at a disadvantage here compared to standard applications of EFT in quantum field theory. The generic situation is that one is searching for local operators that can be incorporated into the EFT Lagrangian (or Hamiltonian) in the situation where there is only a finite number of possible operators of a given dimension. The higher the dimension of the operator the stronger the suppression of its contribution to low energy observables. Thus EFT organizes the possible operators according to their importance in the interesting kinematics. In our case the situation appears to be different. Although RFT is the effective theory of QCD at high energy, all the operators we have found may contribute at leading order in E^{-1} . We do not see any obvious parameter which would order the possible contributions. The similarity with the diamond action may suggest that one should work with the Wilson lines in the adjoint representation. However as is clear from the derivation in [68] the diamond action is not the full story, but is only a leading term in an expansion away from the abelian limit. Thus it is possible that the other candidate terms we have found also play a role in the full RFT Hamiltonian.

It would be interesting to find a criterion which could discriminate between the possible terms. One possibility is to compare H_{RFT} with NLO JIMWLK. Although we have no reason to expect that H_{RFT} contains all, or even most NLO terms, it does contain some such terms. Comparing those to NLO JIMWLK could be instructive and possibly discriminatory.

Another interesting question is the unitarity of H_{RFT} . As we have mentioned in the introduction, our main motivation to search for the self dual H_{RFT} was the unitarity violation in H_{JIMWLK} . The question of unitarity really has two parts: the t-channel unitarity and the s-channel unitarity.

Although we have not studied this in detail here, it is broadly believed that the t-channel unitarity, which has been a cornerstone of Gribov’s RFT is ensured by the self-duality of H_{RFT} . This connection is rooted in boost invariance of the scattering amplitudes.

On one hand Lorentz invariance requires self-duality of RFT [66], and at the same time, boost invariance has been argued to be equivalent to the t -channel unitarity, see [75] for latest discussion. On the technical level we note that the coupling constant expansion of H_{RFT} (in the BFKL counting discussed above) in the large N_c limit generates the Gribov Pomeron calculus. Scattering amplitudes are then represented in terms of the exchanges of the BFKL Pomerons and their interactions via the “merging” and “splitting” three Pomeron vertexes. Such a theory is known to satisfy the t -channel unitarity, and we are therefore confident that our H_{RFT} indeed is t -channel unitary.

As for the s -channel unitarity, the situation here is more complex. We have formulated the conditions for s -channel unitarity in [20]. Given H_{RFT} one can in principle follow the procedure explained in [20] to determine whether its action corresponds to unitarity evolution of QCD states in energy.

This entails taking a generic QCD projectile state

$$|\Psi_i\rangle_P = C_{b_1, b_2 \dots b_m}^{\text{in}} |\mathbf{y}_1, b_1; \dots; \mathbf{y}_m, b_m\rangle$$

and evolving it to infinitesimally higher energy. The result of the evolution in general can be represented in the form:

$$|\Psi\rangle \rightarrow \sum_{n; \mathbf{x}_i; a_i} C_{a_1, a_2 \dots a_n} |\mathbf{x}_1, a_1; \dots; \mathbf{x}_n, a_n\rangle$$

The energy evolution of the scattering amplitude of this evolved state on a fixed target is given by the action of H_{RFT} as in eq. (3.1). Next, one has to construct a probability function F defined in (2.15) and verify the unitarity condition (2.16). The unitarity should hold for any initial state $|\Psi_i\rangle_P$.

In principle one should be able to pursue this calculation, since the algebra of RFT is explicitly known, and therefore the action of H_{RFT} on an unevolved amplitude is completely defined. Unfortunately analyzing the unitarity conditions beyond the JIMWLK limit is technically a complicated problem, due to complicated algebra of the Wilson lines, which at this point we are not able to solve. We believe it is a very important question and are planning to address it in future work.

Acknowledgments

We thank our colleagues at Tel Aviv university and UTFSM for encouraging discussions. AK and Ming Li were supported by the NSF Nuclear Theory grants 1614640 and 1913890. EL was supported by ANID PIA/APOYO AFB180002 (Chile) and Fondecyt (Chile) grant # 1180118. ML was supported by the Israeli Science Foundation (ISF) grant #1635/16. ML and AK were also supported by the Binational Science Foundation grants #2015626, #2018722, and the Horizon 2020 RISE “Heavy ion collisions: collectivity and precision in saturation physics” under grant agreement No. 824093. This work has been performed in the framework of COST Action CA15213 “Theory of hot matter and relativistic heavy-ion collisions” (THOR).

A Checking $SU_L(N)$

In this appendix we calculate perturbatively the commutator of Q_L^a with the Hamiltonian.

A.1 $[Q_L^a, \bar{V}_L(\mathbf{x})]$

We start by trying to verify the conjectured commutation relation:

$$\left[Q_L^a, \bar{V}_L^{\alpha\beta}(\mathbf{x})\right] = -\left(t^a \bar{V}_L(\mathbf{x})\right)^{\alpha\beta} (???) \quad (\text{A.1})$$

We calculate the commutator perturbatively using the BFKL counting. We express

$$\mathcal{J}_L^a(z) = \frac{1}{g} B_{(-1)}^a(\mathbf{z}) + B_{(0)}^a(\mathbf{z}) + g B_{(1)}^a(\mathbf{z}) + g^3 B_{(3)}^a(\mathbf{z}) + \dots \quad (\text{A.2})$$

with

$$\begin{aligned} B_{(-1)}^a &= \tilde{\rho}^a(\mathbf{z}), \\ B_{(0)}^a &= -\frac{1}{2} \tilde{\rho}^b(\mathbf{z}) T_{ba}^e \frac{\delta}{\delta \tilde{\rho}^e(\mathbf{z})}, \\ B_{(1)}^a &= \frac{1}{12} \tilde{\rho}_{\mathbf{z}}^b (T^{e_1} T^{e_2})_{ba} \frac{\delta}{\delta \tilde{\rho}_{\mathbf{z}}^{e_1}} \frac{\delta}{\delta \tilde{\rho}_{\mathbf{z}}^{e_2}}, \\ B_{(3)}^a &= -\frac{1}{720} \tilde{\rho}_{\mathbf{z}}^b (T^{e_1} T^{e_2} T^{e_3} T^{e_4})_{ba} \frac{\delta}{\delta \tilde{\rho}_{\mathbf{z}}^{e_1}} \frac{\delta}{\delta \tilde{\rho}_{\mathbf{z}}^{e_2}} \frac{\delta}{\delta \tilde{\rho}_{\mathbf{z}}^{e_3}} \frac{\delta}{\delta \tilde{\rho}_{\mathbf{z}}^{e_4}}, \end{aligned} \quad (\text{A.3})$$

To expand $\bar{V}_L$ we need

$$\mathcal{I}_L^e(\mathbf{y}) = \frac{1}{g} E_{(-1)}^e(\mathbf{y}) + E_{(0)}^e(\mathbf{y}) + g E_{(1)}^e(\mathbf{y}) + g^3 E_{(3)}^e(\mathbf{y}) + \dots \quad (\text{A.4})$$

with

$$\begin{aligned} E_{(-1)}^a &= -i \partial_{\mathbf{y}}^2 \frac{\delta}{\delta \tilde{\rho}_{\mathbf{y}}^a}, \\ E_{(0)}^a &= -\frac{1}{2} \partial_{\mathbf{y}}^2 \frac{\delta}{\delta \tilde{\rho}_{\mathbf{y}}^b} T_{ba}^e \frac{1}{\partial^2} \tilde{\rho}_{\mathbf{y}}^e, \\ E_{(1)}^a &= \frac{i}{12} \partial_{\mathbf{y}}^2 \frac{\delta}{\delta \tilde{\rho}_{\mathbf{y}}^b} (T^{e_1} T^{e_2})_{ba} \frac{1}{\partial^2} \tilde{\rho}_{\mathbf{y}}^{e_1} \frac{1}{\partial^2} \tilde{\rho}_{\mathbf{y}}^{e_2}, \\ E_{(3)}^a &= \frac{i}{720} \partial_{\mathbf{y}}^2 \frac{\delta}{\delta \tilde{\rho}_{\mathbf{y}}^b} (T^{e_1} T^{e_2} T^{e_3} T^{e_4})_{ba} \frac{1}{\partial^2} \tilde{\rho}_{\mathbf{y}}^{e_1} \frac{1}{\partial^2} \tilde{\rho}_{\mathbf{y}}^{e_2} \frac{1}{\partial^2} \tilde{\rho}_{\mathbf{y}}^{e_3} \frac{1}{\partial^2} \tilde{\rho}_{\mathbf{y}}^{e_4}. \end{aligned} \quad (\text{A.5})$$

Then $\bar{V}_L$ is expanded as

$$\begin{aligned} \bar{V}_L &= e^{g A_{(1)} + g^2 A_{(2)} + g^3 A_{(3)} + g^5 A_{(5)} + \dots} \\ &= 1 + g A_{(1)} + g^2 \left(A_{(2)} + \frac{1}{2} A_{(1)} A_{(1)} \right) + g^3 \left(A_{(3)} + \frac{1}{2} (A_{(1)} A_{(2)} + A_{(2)} A_{(1)}) + \frac{1}{3!} (A_{(1)})^3 \right) \\ &\quad + g^4 \left(\frac{1}{2} (A_{(2)})^2 + \frac{1}{2} (A_{(1)} A_{(3)} + A_{(3)} A_{(1)}) + \frac{1}{3!} (A_{(1)} A_{(1)} A_{(2)} + A_{(1)} A_{(2)} A_{(1)} \right. \\ &\quad \left. + A_{(2)} A_{(1)} A_{(1)}) + \frac{1}{4!} (A_{(1)})^4 \right) + \dots \end{aligned} \quad (\text{A.6})$$

with

$$A_{(1)} = i \int_{\mathbf{y}} \tilde{\phi}(\mathbf{x} - \mathbf{y}) t^e \left(-i \partial_{\mathbf{y}}^2 \frac{\delta}{\delta \tilde{\rho}_{\mathbf{y}}^e} \right) = t^e \frac{\delta}{\delta \tilde{\rho}_{\mathbf{x}}^e} \quad (\text{A.7})$$

$$\begin{aligned} A_{(2)} &= i \int_{\mathbf{y}} \tilde{\phi}(\mathbf{x} - \mathbf{y}) t^d \left(-\frac{1}{2} \partial_{\mathbf{y}}^2 \frac{\delta}{\delta \tilde{\rho}_{\mathbf{y}}^b} T_{bd}^e \frac{1}{\partial^2} \tilde{\rho}^e(\mathbf{y}) \right) \\ &= \frac{i}{2} (t^e t^b - t^b t^e) \left(\frac{\delta}{\delta \tilde{\rho}_{\mathbf{x}}^b} \int_{\mathbf{w}} \tilde{\phi}(\mathbf{x} - \mathbf{w}) \tilde{\rho}_{\mathbf{w}}^e + \int_{\mathbf{y}} \tilde{\phi}(\mathbf{x} - \mathbf{y}) \frac{\delta}{\delta \tilde{\rho}_{\mathbf{y}}^b} \tilde{\rho}_{\mathbf{y}}^e \right. \\ &\quad \left. + 2 \int_{\mathbf{y}} \partial_{\mathbf{y}} \tilde{\phi}(\mathbf{x} - \mathbf{y}) \frac{\delta}{\delta \tilde{\rho}_{\mathbf{y}}^b} \int_{\mathbf{w}} \partial_{\mathbf{y}} \tilde{\phi}(\mathbf{y} - \mathbf{w}) \tilde{\rho}_{\mathbf{w}}^e \right) \end{aligned} \quad (\text{A.8})$$

$$\begin{aligned} A_{(3)} &= i \int_{\mathbf{y}} \tilde{\phi}(\mathbf{x} - \mathbf{y}) t^d \left(\frac{i}{12} \partial_{\mathbf{y}}^2 \frac{\delta}{\delta \tilde{\rho}^b} (T^{e_1} T^{e_1})_{bd} \frac{1}{\partial^2} \tilde{\rho}^{e_1} \frac{1}{\partial^2} \tilde{\rho}^{e_2} \right) \\ &= -\frac{1}{12} (t^{e_1} t^{e_2} t^b - 2 t^{e_1} t^b t^{e_2} + t^b t^{e_1} t^{e_2}) \int_{\mathbf{y}} \tilde{\phi}(\mathbf{x} - \mathbf{y}) \partial_{\mathbf{y}}^2 \frac{\delta}{\delta \tilde{\rho}_{\mathbf{y}}^b} \int_{\mathbf{z}_1} \tilde{\phi}(\mathbf{y} - \mathbf{z}_1) \tilde{\rho}_{\mathbf{z}_1}^{e_1} \int_{\mathbf{z}_2} \tilde{\phi}(\mathbf{y} - \mathbf{z}_2) \tilde{\rho}_{\mathbf{z}_2}^{e_2}. \end{aligned} \quad (\text{A.9})$$

In terms of coupling constant g , we check the commutator eq. (A.1) order by order.

- $O(g^0)$ is satisfied.

$$\left[\int_{\mathbf{z}} B_{(-1)}^a(\mathbf{z}), A_{(1)}(\mathbf{x}) \right] = -t^a. \quad (\text{A.10})$$

- $O(g)$, the relation to be checked is

$$\left[\int_{\mathbf{z}} B_{(-1)}^a(\mathbf{z}), A_{(2)} + \frac{1}{2} (A_{(1)})^2 \right] + \left[\int_{\mathbf{z}} B_{(0)}^a(\mathbf{z}), A_{(1)}(\mathbf{x}) \right] = -t^a A_{(1)}. \quad (\text{A.11})$$

First note that each individual term is

$$\begin{aligned} \left[\int_{\mathbf{z}} B_{(0)}^a(\mathbf{z}), A_{(1)}(\mathbf{x}) \right] &= \frac{1}{2} t^b T_{ba}^e \frac{\delta}{\delta \tilde{\rho}_{\mathbf{x}}^e} = -\frac{1}{2} (t^a t^e - t^e t^a) \frac{\delta}{\delta \tilde{\rho}_{\mathbf{x}}^e} \\ &= -\frac{1}{2} (t^a A_{(1)} - A_{(1)} t^a) \\ &= -\frac{1}{2} [t^a, A_{(1)}]. \end{aligned} \quad (\text{A.12})$$

$$\begin{aligned} \left[\int_{\mathbf{z}} B_{(-1)}^a(\mathbf{z}), \frac{1}{2} (A_{(1)})^2 \right] &= -\frac{1}{2} (t^e t^a + t^a t^e) \frac{\delta}{\delta \tilde{\rho}_{\mathbf{x}}^e} \\ &= -\frac{1}{2} (t^a A_{(1)} + A_{(1)} t^a). \end{aligned} \quad (\text{A.13})$$

We have one additional term but it vanishes.

$$\left[\int_{\mathbf{z}} B_{(-1)}^a(\mathbf{z}), A_{(2)} \right] = - \int_{\mathbf{z}} \left(-\frac{i}{2} \right) t^d T_{bd}^e \int_{\mathbf{y}} \tilde{\phi}(\mathbf{x} - \mathbf{y}) [\partial_{\mathbf{y}}^2 \delta(\mathbf{y} - \mathbf{z}) \delta^{ba}] \int_{\mathbf{w}} \tilde{\phi}(\mathbf{y} - \mathbf{w}) \tilde{\rho}_{\mathbf{w}}^e = 0. \quad (\text{A.14})$$

This vanishes due to $\partial_{\mathbf{y}}^2 \int_{\mathbf{z}} \delta(\mathbf{y} - \mathbf{z}) = 0$.

- $O(g^2)$, the relation to be checked is

$$\begin{aligned}
 & \int_{\mathbf{z}} \left[B_{(-1)}^a(\mathbf{z}), A_{(3)} + \frac{1}{2}(A_{(1)}A_{(2)} + A_{(2)}A_{(1)}) + \frac{1}{3!}(A_{(1)})^3 \right] \\
 & + \int_{\mathbf{z}} \left[B_{(0)}(\mathbf{z}), A_{(2)} + \frac{1}{2}A_{(1)}A_{(1)} \right] + \int_{\mathbf{z}} [B_{(1)}(\mathbf{z}), A_{(1)}] \\
 & = -t^a \left(A_{(2)} + \frac{1}{2}A_{(1)}A_{(1)} \right). \tag{A.15}
 \end{aligned}$$

First note that

$$\begin{aligned}
 \int_{\mathbf{z}} [B_{(-1)}^a(\mathbf{z}), A_{(3)}] &= 0, \\
 \int_{\mathbf{z}} [B_{(-1)}^a(\mathbf{z}), A_{(2)}] &= 0. \tag{A.16}
 \end{aligned}$$

for the same reason as eq. (A.14). This is obviously a general property. Now we evaluate each term.

$$\int_{\mathbf{z}} \left[B_{(-1)}^a(\mathbf{z}), \frac{1}{2}(A_{(1)}A_{(2)} + A_{(2)}A_{(1)}) \right] = -\frac{1}{2}(t^a A_{(2)} + A_{(2)}t^a) \tag{A.17}$$

$$\begin{aligned}
 \int_{\mathbf{z}} \left[B_{(-1)}^a(\mathbf{z}), \frac{1}{3!}(A_{(1)})^3 \right] &= -\frac{1}{6}(t^a t^{e_1} t^{e_2} + t^{e_1} t^a t^{e_2} + t^{e_1} t^{e_2} t^a) \frac{\delta}{\delta \tilde{\rho}_{\mathbf{x}}^{e_1}} \frac{\delta}{\delta \tilde{\rho}_{\mathbf{x}}^{e_2}} \\
 &= -\frac{1}{6}(t^a A_{(1)} A_{(1)} + A_{(1)} t^a A_{(1)} + A_{(1)} A_{(1)} t^a) \tag{A.18}
 \end{aligned}$$

$$\begin{aligned}
 \int_{\mathbf{z}} [B_{(1)}(\mathbf{z}), A_{(1)}] &= -\frac{1}{12} t^e (T^{e_1} T^{e_2})_{ea} \frac{\delta}{\delta \tilde{\rho}_{\mathbf{x}}^{e_1}} \frac{\delta}{\delta \tilde{\rho}_{\mathbf{x}}^{e_2}} \\
 &= -\frac{1}{12} (t^a t^{e_1} t^{e_2} - 2t^{e_1} t^a t^{e_2} + t^{e_1} t^{e_2} t^a) \frac{\delta}{\delta \tilde{\rho}_{\mathbf{x}}^{e_1}} \frac{\delta}{\delta \tilde{\rho}_{\mathbf{x}}^{e_2}} \\
 &= -\frac{1}{12} (t^a A_{(1)} A_{(1)} - 2A_{(1)} t^a A_{(1)} + A_{(1)} A_{(1)} t^a) \\
 &= -\frac{1}{12} [[t^a, A_{(1)}], A_{(1)}] \tag{A.19}
 \end{aligned}$$

$$\begin{aligned}
 \int_{\mathbf{z}} \left[B_{(0)}(\mathbf{z}), \frac{1}{2}A_{(1)}A_{(1)} \right] &= -\frac{1}{4}(t^a t^{e_1} t^{e_2} - t^{e_1} t^{e_2} t^a) \frac{\delta}{\delta \tilde{\rho}_{\mathbf{x}}^{e_1}} \frac{\delta}{\delta \tilde{\rho}_{\mathbf{x}}^{e_2}} \\
 &= -\frac{1}{4}(t^a A_{(1)} A_{(1)} - A_{(1)} A_{(1)} t^a). \tag{A.20}
 \end{aligned}$$

Adding eqs. (A.18), (A.19), (A.20), one obtains

$$-\frac{1}{2}(t^a t^{e_1} t^{e_2}) \frac{\delta}{\delta \tilde{\rho}_{\mathbf{x}}^{e_1}} \frac{\delta}{\delta \tilde{\rho}_{\mathbf{x}}^{e_2}} = -t^a \left(\frac{1}{2}A_{(1)}A_{(1)} \right) \tag{A.21}$$

which is part of the right hand side of the relation eq. (A.15). To continue

$$\begin{aligned}
 \int_{\mathbf{z}} [B_{(0)}(\mathbf{z}), A_{(2)}] &= \int_{\mathbf{z}} \left[-\frac{1}{2} \tilde{\rho}_{\mathbf{z}}^p T_{pa}^q \frac{\delta}{\delta \tilde{\rho}_{\mathbf{z}}^q}, -\frac{i}{2} t^d T_{bd}^e \int_{\mathbf{y}} \tilde{\phi}(\mathbf{x} - \mathbf{y}) \partial_{\mathbf{y}}^2 \frac{\delta}{\delta \tilde{\rho}_{\mathbf{y}}^b} \int_{\mathbf{w}} \tilde{\phi}(\mathbf{y} - \mathbf{w}) \tilde{\rho}_{\mathbf{w}}^e \right] \\
 &= \left(-\frac{i}{4} T_{pa}^m T_{pd}^n + \frac{i}{4} T_{na}^e T_{md}^e \right) t^d \int_{\mathbf{y}} \tilde{\phi}(\mathbf{x} - \mathbf{y}) \partial_{\mathbf{y}}^2 \frac{\delta}{\delta \tilde{\rho}_{\mathbf{y}}^m} \int_{\mathbf{w}} \tilde{\phi}(\mathbf{y} - \mathbf{w}) \tilde{\rho}_{\mathbf{w}}^n \\
 &= \frac{1}{2} (A_{(2)} t^a - t^a A_{(2)}).
 \end{aligned} \tag{A.22}$$

So eqs. (A.17), (A.22) adds up gives $-t^a A_{(2)}$, which is exactly the last piece on the right hand side of eq. (A.15). To second order in g eq. (A.1) holds.

- $O(g^3)$. The relation to be proved is

$$\begin{aligned}
 &\int_{\mathbf{z}} \left[B_{(-1)}^a(\mathbf{z}), \frac{1}{2} ((A_{(2)})^2 + A_{(1)} A_{(3)} + A_{(3)} A_{(1)}) \right. \\
 &\quad \left. + \frac{1}{3!} (A_{(1)} A_{(1)} A_{(2)} + A_{(1)} A_{(2)} A_{(1)} + A_{(2)} A_{(1)} A_{(1)}) + \frac{1}{4!} (A_{(1)})^4 \right] \\
 &+ \int_{\mathbf{z}} \left[B_{(0)}^a(\mathbf{z}), A_{(3)} + \frac{1}{2} (A_{(1)} A_{(2)} + A_{(2)} A_{(1)}) + \frac{1}{3!} (A_{(1)})^3 \right] \\
 &+ \int_{\mathbf{z}} \left[B_{(1)}^a(\mathbf{z}), A_{(2)} + \frac{1}{2} A_{(1)} A_{(1)} \right] \\
 &= -t^a \left(A_{(3)} + \frac{1}{2} (A_{(1)} A_{(2)} + A_{(2)} A_{(1)}) + \frac{1}{3!} (A_{(1)})^3 \right).
 \end{aligned} \tag{A.23}$$

We calculate each commutator separately. The first one is easy to compute as we know that

$$\begin{aligned}
 \int_{\mathbf{z}} [B_{(-1)}^a(\mathbf{z}), A_{(1)}] &= -t^a, \\
 \int_{\mathbf{z}} [B_{(-1)}^a(\mathbf{z}), A_{(n)}] &= 0, \quad \text{for } n \geq 2
 \end{aligned} \tag{A.24}$$

Using this relation, one obtains

$$\begin{aligned}
 &\int_{\mathbf{z}} \left[B_{(-1)}^a(\mathbf{z}), \frac{1}{2} ((A_{(2)})^2 + A_{(1)} A_{(3)} + A_{(3)} A_{(1)}) \right. \\
 &\quad \left. + \frac{1}{3!} (A_{(1)} A_{(1)} A_{(2)} + A_{(1)} A_{(2)} A_{(1)} + A_{(2)} A_{(1)} A_{(1)}) + \frac{1}{4!} (A_{(1)})^4 \right] \\
 &= -\frac{1}{2} (t^a A_{(3)} + A_{(3)} t^a) - \frac{1}{6} (t^a A_{(1)} A_{(2)} + A_{(1)} t^a A_{(2)} + t^a A_{(2)} A_{(1)} + A_{(1)} A_{(2)} t^a \\
 &\quad + A_{(2)} t^a A_{(1)} + A_{(2)} A_{(1)} t^a) - \frac{1}{24} (t^a A_{(1)} A_{(1)} A_{(1)} + A_{(1)} t^a A_{(1)} A_{(1)} \\
 &\quad + A_{(1)} A_{(1)} t^a A_{(1)} + A_{(1)} A_{(1)} A_{(1)} t^a).
 \end{aligned} \tag{A.25}$$

We also notice that

$$\int_{\mathbf{z}} [B_{(0)}^a(\mathbf{z}), A_{(n)}] = \frac{1}{2} (A_{(n)} t^a - t^a A_{(n)}). \tag{A.26}$$

for $n = 1, 2, 3$. It is possible that this relation holds for all the relevant n . Using this relation, we calculate

$$\begin{aligned}
 & \int_{\mathbf{z}} \left[B_{(0)}^a(\mathbf{z}), A_{(3)} + \frac{1}{2}(A_{(1)}A_{(2)} + A_{(2)}A_{(1)}) + \frac{1}{3!}(A_{(1)})^3 \right] \\
 &= \frac{1}{2}(A_{(3)}t^a - t^a A_{(3)}) + \frac{1}{4}(A_{(1)}t^a - t^a A_{(1)})A_{(2)} + \frac{1}{4}A_{(1)}(A_{(2)}t^a - t^a A_{(2)}) \\
 & \quad + \frac{1}{4}(A_{(2)}t^a - t^a A_{(2)})A_{(1)} + \frac{1}{4}A_{(2)}(A_{(1)}t^a - t^a A_{(1)}) + \frac{1}{12}(A_{(1)}t^a - t^a A_{(1)})A_{(1)}A_{(1)} \\
 & \quad + \frac{1}{12}A_{(1)}(A_{(1)}t^a - t^a A_{(1)})A_{(1)} + \frac{1}{12}A_{(1)}A_{(1)}(A_{(1)}t^a - t^a A_{(1)}) \\
 &= \frac{1}{2}(A_{(3)}t^a - t^a A_{(3)}) + \frac{1}{4} \left(A_{(1)}A_{(2)}t^a + A_{(2)}A_{(1)}t^a - t^a A_{(1)}A_{(2)} - t^a A_{(2)}A_{(1)} \right) \\
 & \quad + \frac{1}{12} \left(A_{(1)}A_{(1)}A_{(1)}t^a - t^a A_{(1)}A_{(1)}A_{(1)} \right).
 \end{aligned} \tag{A.27}$$

From eq. (A.19), we know that

$$\int_{\mathbf{z}} [B_{(1)}(\mathbf{z}), A_{(1)}] = -\frac{1}{12}(t^a A_{(1)}A_{(1)} - 2A_{(1)}t^a A_{(1)} + A_{(1)}A_{(1)}t^a). \tag{A.28}$$

Using this relation, one can compute

$$\begin{aligned}
 & \int_{\mathbf{z}} \left[B_{(1)}^a(\mathbf{z}), \frac{1}{2}A_{(1)}A_{(1)} \right] \\
 &= -\frac{1}{24} \left(t^a A_{(1)}A_{(1)}A_{(1)} - 2A_{(1)}t^a A_{(1)}A_{(1)} + A_{(1)}A_{(1)}t^a A_{(1)} \right. \\
 & \quad \left. + A_{(1)}t^a A_{(1)}A_{(1)} - 2A_{(1)}A_{(1)}t^a A_{(1)} + A_{(1)}A_{(1)}A_{(1)}t^a \right). \\
 &= -\frac{1}{24} \left(t^a A_{(1)}A_{(1)}A_{(1)} - A_{(1)}t^a A_{(1)}A_{(1)} - A_{(1)}A_{(1)}t^a A_{(1)} + A_{(1)}A_{(1)}A_{(1)}t^a \right).
 \end{aligned} \tag{A.29}$$

The last piece we need to calculate is

$$\begin{aligned}
 & \int_{\mathbf{z}} \left[B_{(1)}^a(\mathbf{z}), A_{(2)}(\mathbf{x}) \right] \\
 &= \int_{\mathbf{z}} \left[\frac{1}{12} \tilde{\rho}_{\mathbf{z}}^c (T^{e_1} T^{e_2})_{ca} \frac{\delta}{\delta \tilde{\rho}_{\mathbf{z}}^{e_1}} \frac{\delta}{\delta \tilde{\rho}_{\mathbf{z}}^{e_2}}, \frac{i}{2} (t^e t^b - t^b t^e) \int_{\mathbf{y}} \tilde{\phi}(\mathbf{x} - \mathbf{y}) \partial_{\mathbf{y}}^2 \frac{\delta}{\delta \tilde{\rho}_{\mathbf{y}}^b} \int_{\mathbf{w}} \tilde{\phi}(\mathbf{y} - \mathbf{w}) \tilde{\rho}_{\mathbf{w}}^e \right] \\
 &= \frac{1}{12} (T^{e_1} T^{e_2})_{ba} \frac{i}{2} T_{be}^p t^p \left(- \int_{\mathbf{z}} \frac{\delta}{\delta \tilde{\rho}_{\mathbf{z}}^{e_1}} \frac{\delta}{\delta \tilde{\rho}_{\mathbf{z}}^{e_2}} \int_{\mathbf{y}} \tilde{\phi}(\mathbf{x} - \mathbf{y}) \partial_{\mathbf{y}}^2 \delta(\mathbf{y} - \mathbf{z}) \int_{\mathbf{w}} \tilde{\phi}(\mathbf{y} - \mathbf{w}) \tilde{\rho}_{\mathbf{w}}^e \right) \\
 &\quad + \frac{1}{12} (T^e T^d + T^d T^e)_{ha} \frac{i}{2} T_{be}^p t^p \left(\int_{\mathbf{z}} \tilde{\rho}_{\mathbf{z}}^h \frac{\delta}{\delta \tilde{\rho}_{\mathbf{z}}^d} \int_{\mathbf{y}} \tilde{\phi}(\mathbf{x} - \mathbf{y}) \partial_{\mathbf{y}}^2 \frac{\delta}{\delta \tilde{\rho}_{\mathbf{y}}^b} \tilde{\phi}(\mathbf{y} - \mathbf{z}) \right) \\
 &= \frac{1}{12} (T^b T^d)_{qa} \frac{i}{2} T_{qh}^p t^p \left(- \int_{\mathbf{z}} \frac{\delta}{\delta \tilde{\rho}_{\mathbf{z}}^b} \frac{\delta}{\delta \tilde{\rho}_{\mathbf{z}}^d} \int_{\mathbf{y}} \tilde{\phi}(\mathbf{x} - \mathbf{y}) \partial_{\mathbf{y}}^2 \delta(\mathbf{y} - \mathbf{z}) \int_{\mathbf{w}} \tilde{\phi}(\mathbf{y} - \mathbf{w}) \tilde{\rho}_{\mathbf{w}}^h \right) \\
 &\quad + \frac{1}{12} (T^e T^d + T^d T^e)_{ha} \frac{i}{2} T_{be}^p t^p \left(\int_{\mathbf{z}} \tilde{\rho}_{\mathbf{z}}^h \frac{\delta}{\delta \tilde{\rho}_{\mathbf{z}}^d} \int_{\mathbf{y}} \tilde{\phi}(\mathbf{x} - \mathbf{y}) \partial_{\mathbf{y}}^2 \frac{\delta}{\delta \tilde{\rho}_{\mathbf{y}}^b} \tilde{\phi}(\mathbf{y} - \mathbf{z}) \right) \\
 &= \frac{i}{24} \left(- (T^p T^b T^d)_{ha} t^p \right) \left(- \frac{\delta}{\delta \tilde{\rho}_{\mathbf{x}}^b} \frac{\delta}{\delta \tilde{\rho}_{\mathbf{x}}^d} \int_{\mathbf{w}} \tilde{\phi}(\mathbf{x} - \mathbf{w}) \tilde{\rho}_{\mathbf{w}}^h - \int_{\mathbf{z}} \tilde{\phi}(\mathbf{x} - \mathbf{z}) \frac{\delta}{\delta \tilde{\rho}_{\mathbf{z}}^b} \frac{\delta}{\delta \tilde{\rho}_{\mathbf{z}}^d} \tilde{\rho}_{\mathbf{z}}^h \right. \\
 &\quad \left. - 2 \int_{\mathbf{z}} \frac{\delta}{\delta \tilde{\rho}_{\mathbf{z}}^b} \frac{\delta}{\delta \tilde{\rho}_{\mathbf{z}}^d} \partial_{\mathbf{z}} \tilde{\phi}(\mathbf{x} - \mathbf{z}) \int_{\mathbf{w}} \partial_{\mathbf{z}} \tilde{\phi}(\mathbf{z} - \mathbf{w}) \tilde{\rho}_{\mathbf{w}}^h \right) \\
 &\quad + \frac{i}{24} \left((T^p T^h T^a)_{bd} t^p + (T^p T^a T^h)_{bd} t^p \right) \left(\frac{\delta}{\delta \tilde{\rho}_{\mathbf{x}}^b} \int_{\mathbf{z}} \tilde{\phi}(\mathbf{x} - \mathbf{z}) \tilde{\rho}_{\mathbf{z}}^h \frac{\delta}{\delta \tilde{\rho}_{\mathbf{z}}^d} \right. \\
 &\quad \left. + \int_{\mathbf{z}} \tilde{\phi}(\mathbf{x} - \mathbf{z}) \tilde{\rho}_{\mathbf{z}}^h \frac{\delta}{\delta \tilde{\rho}_{\mathbf{z}}^d} \frac{\delta}{\delta \tilde{\rho}_{\mathbf{z}}^b} + 2 \int_{\mathbf{z}} \tilde{\rho}_{\mathbf{z}}^h \frac{\delta}{\delta \tilde{\rho}_{\mathbf{z}}^d} \int_{\mathbf{y}} \partial_{\mathbf{y}} \tilde{\phi}(\mathbf{x} - \mathbf{y}) \frac{\delta}{\delta \tilde{\rho}_{\mathbf{y}}^b} \partial_{\mathbf{y}} \tilde{\phi}(\mathbf{y} - \mathbf{z}) \right). \tag{A.30}
 \end{aligned}$$

We have calculated the color structures for the two parts

$$(T^b T^d)_{qa} T_{qh}^p t^p = - (T^p T^b T^d)_{ha} t^p \tag{A.31}$$

$$\begin{aligned}
 (T^e T^d + T^d T^e)_{ha} T_{be}^p t^p &= - (T^p T^h T^d)_{ba} t^p + (T^d T^a T^p)_{hb} t^p \\
 &= (T^p T^h T^a)_{bd} t^p + (T^p T^a T^h)_{bd} t^p \tag{A.32}
 \end{aligned}$$

and performed the integration by parts

$$\begin{aligned}
 & - \int_{\mathbf{z}} \frac{\delta}{\delta \tilde{\rho}_{\mathbf{z}}^b} \frac{\delta}{\delta \tilde{\rho}_{\mathbf{z}}^d} \int_{\mathbf{y}} \tilde{\phi}(\mathbf{x} - \mathbf{y}) \partial_{\mathbf{y}}^2 \delta(\mathbf{y} - \mathbf{z}) \int_{\mathbf{w}} \tilde{\phi}(\mathbf{y} - \mathbf{w}) \tilde{\rho}_{\mathbf{w}}^h \\
 &= - \frac{\delta}{\delta \tilde{\rho}_{\mathbf{x}}^b} \frac{\delta}{\delta \tilde{\rho}_{\mathbf{x}}^d} \int_{\mathbf{w}} \tilde{\phi}(\mathbf{x} - \mathbf{w}) \tilde{\rho}_{\mathbf{w}}^h - \int_{\mathbf{z}} \tilde{\phi}(\mathbf{x} - \mathbf{z}) \frac{\delta}{\delta \tilde{\rho}_{\mathbf{z}}^b} \frac{\delta}{\delta \tilde{\rho}_{\mathbf{z}}^d} \tilde{\rho}_{\mathbf{z}}^h \\
 &\quad - 2 \int_{\mathbf{z}} \frac{\delta}{\delta \tilde{\rho}_{\mathbf{z}}^b} \frac{\delta}{\delta \tilde{\rho}_{\mathbf{z}}^d} \partial_{\mathbf{z}} \tilde{\phi}(\mathbf{x} - \mathbf{z}) \int_{\mathbf{w}} \partial_{\mathbf{z}} \tilde{\phi}(\mathbf{z} - \mathbf{w}) \tilde{\rho}_{\mathbf{w}}^h \tag{A.33}
 \end{aligned}$$

and

$$\begin{aligned}
 & \int_{\mathbf{z}} \tilde{\rho}_{\mathbf{z}}^h \frac{\delta}{\delta \tilde{\rho}_{\mathbf{z}}^d} \int_{\mathbf{y}} \tilde{\phi}(\mathbf{x} - \mathbf{y}) \partial_{\mathbf{y}}^2 \frac{\delta}{\delta \tilde{\rho}_{\mathbf{y}}^b} \tilde{\phi}(\mathbf{y} - \mathbf{z}) \\
 &= \frac{\delta}{\delta \tilde{\rho}_{\mathbf{x}}^b} \int_{\mathbf{z}} \tilde{\phi}(\mathbf{x} - \mathbf{z}) \tilde{\rho}_{\mathbf{z}}^h \frac{\delta}{\delta \tilde{\rho}_{\mathbf{z}}^d} + \int_{\mathbf{z}} \tilde{\phi}(\mathbf{x} - \mathbf{z}) \tilde{\rho}_{\mathbf{z}}^h \frac{\delta}{\delta \tilde{\rho}_{\mathbf{z}}^d} \frac{\delta}{\delta \tilde{\rho}_{\mathbf{z}}^b} \\
 &\quad + 2 \int_{\mathbf{z}} \tilde{\rho}_{\mathbf{z}}^h \frac{\delta}{\delta \tilde{\rho}_{\mathbf{z}}^d} \int_{\mathbf{y}} \partial_{\mathbf{y}} \tilde{\phi}(\mathbf{x} - \mathbf{y}) \frac{\delta}{\delta \tilde{\rho}_{\mathbf{y}}^b} \partial_{\mathbf{y}} \tilde{\phi}(\mathbf{y} - \mathbf{z}). \tag{A.34}
 \end{aligned}$$

The commutation $\int_{\mathbf{z}} [B_{(1)}^a(\mathbf{z}), A_{(2)}(\mathbf{x})]$, for the relation eq. (A.23) to be correct, should be equal to

$$\begin{aligned}
 & -\frac{1}{12} (t^a A_{(1)} A_{(2)} + t^a A_{(2)} A_{(1)} + A_{(1)} A_{(2)} t^a + A_{(2)} A_{(1)} t^a - 2A_{(1)} t^a A_{(2)} - 2A_{(2)} t^a A_{(1)}) \\
 & = -\frac{1}{12} ([t^a, A_{(1)}], A_{(2)}) + ([t^a, A_{(2)}], A_{(1)}) \\
 & = -\frac{1}{12} ([t^a, t^b], t^m) + ([t^a, t^m], t^b) T_{dh}^m \times \frac{i}{2} \left(\frac{\delta}{\delta \tilde{\rho}_{\mathbf{x}}^b} \frac{\delta}{\delta \tilde{\rho}_{\mathbf{x}}^d} \int_{\mathbf{w}} \tilde{\phi}(\mathbf{x} - \mathbf{w}) \tilde{\rho}_{\mathbf{w}}^h \right. \\
 & \quad \left. + \frac{\delta}{\delta \tilde{\rho}_{\mathbf{x}}^b} \int_{\mathbf{y}} \tilde{\phi}(\mathbf{x} - \mathbf{y}) \frac{\delta}{\delta \tilde{\rho}_{\mathbf{y}}^d} \tilde{\rho}_{\mathbf{y}}^h + 2 \frac{\delta}{\delta \tilde{\rho}_{\mathbf{x}}^b} \int_{\mathbf{y}} \partial_{\mathbf{y}} \tilde{\phi}(\mathbf{x} - \mathbf{y}) \frac{\delta}{\delta \tilde{\rho}_{\mathbf{y}}^d} \int_{\mathbf{w}} \partial_{\mathbf{y}} \tilde{\phi}(\mathbf{y} - \mathbf{w}) \tilde{\rho}_{\mathbf{w}}^h \right) \\
 & = -\frac{1}{12} ((T^a T^p T^h)_{bd} + (T^p T^a T^h)_{bd}) t^p \times \frac{i}{2} \left(\frac{\delta}{\delta \tilde{\rho}_{\mathbf{x}}^b} \frac{\delta}{\delta \tilde{\rho}_{\mathbf{x}}^d} \int_{\mathbf{w}} \tilde{\phi}(\mathbf{x} - \mathbf{w}) \tilde{\rho}_{\mathbf{w}}^h \right. \\
 & \quad \left. + \frac{\delta}{\delta \tilde{\rho}_{\mathbf{x}}^b} \int_{\mathbf{y}} \tilde{\phi}(\mathbf{x} - \mathbf{y}) \frac{\delta}{\delta \tilde{\rho}_{\mathbf{y}}^d} \tilde{\rho}_{\mathbf{y}}^h + 2 \frac{\delta}{\delta \tilde{\rho}_{\mathbf{x}}^b} \int_{\mathbf{y}} \partial_{\mathbf{y}} \tilde{\phi}(\mathbf{x} - \mathbf{y}) \frac{\delta}{\delta \tilde{\rho}_{\mathbf{y}}^d} \int_{\mathbf{w}} \partial_{\mathbf{y}} \tilde{\phi}(\mathbf{y} - \mathbf{w}) \tilde{\rho}_{\mathbf{w}}^h \right). \tag{A.35}
 \end{aligned}$$

We used

$$\begin{aligned}
 [[t^a, t^b], t^m] + [[t^a, t^m], t^b] T_{dh}^m &= (T_{ba}^s [t^s, t^m] + T_{ma}^s [t^s, t^b]) T_{dh}^m \\
 &= (T_{ba}^s T_{ms}^p t^p + T_{ma}^s T_{bs}^p) T_{dh}^m \\
 &= ((T^a T^p T^h)_{bd} + (T^p T^a T^h)_{bd}) t^p. \tag{A.36}
 \end{aligned}$$

The difference is

$$\begin{aligned}
 & \int_{\mathbf{z}} [B_{(1)}^a(\mathbf{z}), A_{(2)}(\mathbf{x})] - \left(-\frac{1}{12} ([t^a, A_{(1)}], A_{(2)}) + ([t^a, A_{(2)}], A_{(1)}) \right) \\
 &= \frac{i}{24} \frac{\delta}{\delta \tilde{\rho}_{\mathbf{x}}^b} \frac{\delta}{\delta \tilde{\rho}_{\mathbf{x}}^d} \int_{\mathbf{w}} \tilde{\phi}(\mathbf{x} - \mathbf{w}) \tilde{\rho}_{\mathbf{w}}^h ((T^a T^p T^h)_{bd} + (T^p T^a T^h)_{bd} + (T^p T^b T^d)_{ha}) t^p \\
 & \quad + \frac{i}{24} \frac{\delta}{\delta \tilde{\rho}_{\mathbf{x}}^b} \int_{\mathbf{y}} \tilde{\phi}(\mathbf{x} - \mathbf{y}) \frac{\delta}{\delta \tilde{\rho}_{\mathbf{y}}^d} \tilde{\rho}_{\mathbf{y}}^h ((T^a T^p T^h)_{bd} + 2(T^p T^a T^h)_{bd} + (T^p T^h T^a)_{bd}) t^p \\
 & \quad + \frac{i}{24} \int_{\mathbf{z}} \tilde{\phi}(\mathbf{x} - \mathbf{z}) \tilde{\rho}_{\mathbf{z}}^h \frac{\delta}{\delta \tilde{\rho}_{\mathbf{z}}^d} \frac{\delta}{\delta \tilde{\rho}_{\mathbf{z}}^b} ((T^p T^h T^a)_{bd} + (T^p T^a T^h)_{bd} + (T^p T^b T^d)_{ha}) t^p \\
 & \quad + \frac{i}{12} \frac{\delta}{\delta \tilde{\rho}_{\mathbf{x}}^b} \int_{\mathbf{y}} \partial_{\mathbf{y}} \tilde{\phi}(\mathbf{x} - \mathbf{y}) \frac{\delta}{\delta \tilde{\rho}_{\mathbf{y}}^d} \int_{\mathbf{w}} \partial_{\mathbf{y}} \tilde{\phi}(\mathbf{y} - \mathbf{w}) \tilde{\rho}_{\mathbf{w}}^h ((T^a T^p T^h)_{bd} + (T^p T^a T^h)_{bd}) t^p \\
 & \quad + \frac{i}{12} \int_{\mathbf{z}} \frac{\delta}{\delta \tilde{\rho}_{\mathbf{z}}^b} \frac{\delta}{\delta \tilde{\rho}_{\mathbf{z}}^d} \partial_{\mathbf{z}} \tilde{\phi}(\mathbf{x} - \mathbf{z}) \int_{\mathbf{w}} \partial_{\mathbf{z}} \tilde{\phi}(\mathbf{z} - \mathbf{w}) \tilde{\rho}_{\mathbf{w}}^h (T^p T^b T^d)_{ha} t^p \\
 & \quad + \frac{i}{12} \int_{\mathbf{z}} \tilde{\rho}_{\mathbf{z}}^h \frac{\delta}{\delta \tilde{\rho}_{\mathbf{z}}^d} \int_{\mathbf{y}} \partial_{\mathbf{y}} \tilde{\phi}(\mathbf{x} - \mathbf{y}) \frac{\delta}{\delta \tilde{\rho}_{\mathbf{y}}^b} \partial_{\mathbf{y}} \tilde{\phi}(\mathbf{y} - \mathbf{z}) ((T^p T^h T^a)_{bd} + (T^p T^a T^h)_{bd}) t^p. \tag{A.37}
 \end{aligned}$$

This clearly does not vanish, and so eq. (A.1) is violated at $O(g^3)$.

A.2 Checking $[Q_L^a, H_{RFT}] = 0$

We now directly calculate the commutator of Q_L^a with H_{RFT} .

The calculation is organized as expansion in powers of g

$$\begin{aligned} [Q_L^a, H_{RFT}] &= \sum_{n=0}^{\infty} g^n [Q_L^a, H_{RFT}]_{(n)} \\ &= [Q_L^a, H_{RFT}]_{(0)} + g [Q_L^a, H_{RFT}]_{(1)} + g^2 [Q_L^a, H_{RFT}]_{(2)} + \dots \end{aligned} \quad (\text{A.38})$$

where the subscript “ (n) ” indicates the n -th order in g .

Using $[Q_L^a, V_R] = 0$ and $[Q_L^a, V_L^{\alpha\beta}(\mathbf{x})] = -(t^a V_L)^{\alpha\beta} + (V_L t^a)^{\alpha\beta}$, we can write

$$\begin{aligned} &[Q_L^a, H_{RFT}] \\ &= \int_{\mathbf{x}} \left[Q_L^a, \bar{V}_L^{\beta\gamma}(\mathbf{x}) \right] \bar{V}_R^{\delta\alpha}(\mathbf{x}) \partial_{\mathbf{x}}^2 \left(V_L^{\alpha\beta}(\mathbf{x}) V_R^{\gamma\delta}(\mathbf{x}) \right) + \bar{V}_L^{\beta\gamma}(\mathbf{x}) \left[Q_L^a, \bar{V}_R^{\delta\alpha}(\mathbf{x}) \right] \partial_{\mathbf{x}}^2 \left(V_L^{\alpha\beta}(\mathbf{x}) V_R^{\gamma\delta}(\mathbf{x}) \right) \\ &\quad + \left((t^a \bar{V}_L)^{\beta\gamma} \bar{V}_R^{\delta\alpha}(\mathbf{x}) - \bar{V}_L^{\beta\gamma}(\mathbf{x}) (\bar{V}_R t^a)^{\delta\alpha} \right) \partial_{\mathbf{x}}^2 \left(V_L^{\alpha\beta}(\mathbf{x}) V_R^{\gamma\delta}(\mathbf{x}) \right) \\ &= \int_{\mathbf{x}} \left\{ \left([Q_L^a, \bar{V}_L^{\beta\gamma}] + (t^a \bar{V}_L)^{\beta\gamma} \right) \bar{V}_R^{\delta\alpha} + \bar{V}_L^{\beta\gamma} \left([Q_L^a, \bar{V}_R^{\delta\alpha}] - (\bar{V}_R t^a)^{\delta\alpha} \right) \right\} \partial_{\mathbf{x}}^2 \left(V_L^{\alpha\beta}(\mathbf{x}) V_R^{\gamma\delta}(\mathbf{x}) \right) \end{aligned} \quad (\text{A.39})$$

Here we have rescaled H_{RFT} by the overall factor πg^2 for simplicity.

Let us denote

$$\begin{aligned} d_L^{\beta\gamma} &= [Q_L^a, \bar{V}_L^{\beta\gamma}] + (t^a \bar{V}_L)^{\beta\gamma} \\ d_R^{\delta\alpha} &= [Q_L^a, \bar{V}_R^{\delta\alpha}] - (\bar{V}_R t^a)^{\delta\alpha} \end{aligned} \quad (\text{A.40})$$

then

$$[Q_L^a, H_{RFT}] = \int_{\mathbf{x}} \left(d_L^{\beta\gamma} \bar{V}_R^{\delta\alpha} + \bar{V}_L^{\beta\gamma} d_R^{\delta\alpha} \right) \partial_{\mathbf{x}}^2 \left(V_L^{\alpha\beta}(\mathbf{x}) V_R^{\gamma\delta}(\mathbf{x}) \right) \quad (\text{A.41})$$

Symbolically we write

$$\begin{aligned} \bar{V}_L &= \exp \left\{ g A_{(1)} + g^2 A_{(2)} + g^3 A_{(3)} + g^5 A_{(5)} + \dots \right\}, \\ \bar{V}_R &= \exp \left\{ -g A_{(1)} + g^2 A_{(2)} - g^3 A_{(3)} - g^5 A_{(5)} - \dots \right\}, \end{aligned} \quad (\text{A.42})$$

and

$$\begin{aligned} V_L &= \exp \left\{ g X_{(1)} + g^2 X_{(2)} + g^3 X_{(3)} + g^5 X_{(5)} + \dots \right\}, \\ V_R &= \exp \left\{ -g X_{(1)} + g^2 X_{(2)} - g^3 X_{(3)} - g^5 X_{(5)} - \dots \right\}, \end{aligned} \quad (\text{A.43})$$

Expansion of $\partial_{\mathbf{x}}^2 \left(V_L^{\alpha\beta}(\mathbf{x}) V_R^{\gamma\delta}(\mathbf{x}) \right)$ starts at order g

$$\begin{aligned}
 & \partial_{\mathbf{x}}^2 \left(V_L^{\alpha\beta}(\mathbf{x}) V_R^{\gamma\delta}(\mathbf{x}) \right) \\
 &= g \partial^2 \left(-\delta^{\alpha\beta} X_{(1)}^{\gamma\delta} + X_{(1)}^{\alpha\beta} \delta^{\gamma\delta} \right) \\
 &+ g^2 \partial^2 \left(\delta^{\alpha\beta} (X_{(2)}^{\gamma\delta} + \frac{1}{2} (X_{(1)}^2)^{\gamma\delta}) + (X_{(2)}^{\alpha\beta} + \frac{1}{2} (X_{(1)}^2)^{\alpha\beta}) \delta^{\gamma\delta} - X_{(1)}^{\alpha\beta} X_{(1)}^{\gamma\delta} \right) \\
 &+ g^3 \partial^2 \left(-\delta^{\alpha\beta} \left(X_{(3)} + \frac{1}{2} (X_{(1)} X_{(2)} + X_{(2)} X_{(1)}) + \frac{1}{3!} X_{(1)}^3 \right)^{\gamma\delta} \right. \\
 &\quad \left. + \left(X_{(3)} + \frac{1}{2} (X_{(1)} X_{(2)} + X_{(2)} X_{(1)}) + \frac{1}{3!} X_{(1)}^3 \right)^{\alpha\beta} \delta^{\gamma\delta} \right. \\
 &\quad \left. + X_{(1)}^{\alpha\beta} \left(X_{(2)} + \frac{1}{2} X_{(1)}^2 \right)^{\gamma\delta} - \left(X_{(2)} + \frac{1}{2} X_{(1)}^2 \right)^{\alpha\beta} X_{(1)}^{\gamma\delta} \right) \\
 &+ \dots
 \end{aligned} \tag{A.44}$$

Thus the expansion of the Hamiltonian H_{RFT} starts at order $O(g)$. On the other hand recall that

$$Q_L^a = \frac{1}{g} \bar{B}_{(-1)} + \bar{B}_{(0)} + g \bar{B}_{(1)} + g^3 \bar{B}_{(3)} + \dots \tag{A.45}$$

So the commutation relation $[Q_L^a, H_{RFT}]$ formally starts at order $O(1)$, but from eq. (A.39) it is obvious that at $O(1)$ the commutator vanishes.

The results of the previous subsection we have calculated explicitly $d_{L(n)}$ for $n \leq 3$. Although we have not explicitly calculated $d_{R(n)}$, this calculation up to $n = 3$ is identical to that of $d_{L(n)}$ and thus we have

$$\begin{aligned}
 d_{L(2)} &= d_{L(1)} = d_{L(0)} = 0, \\
 d_{R(2)} &= d_{R(1)} = d_{R(0)} = 0. \\
 d_{R(3)} &= d_{L(3)}
 \end{aligned} \tag{A.46}$$

As a consequence

$$\begin{aligned}
 [Q_L^a, H_{RFT}]_{(1)} &= 0, \\
 [Q_L^a, H_{RFT}]_{(2)} &= 0, \\
 [Q_L^a, H_{RFT}]_{(3)} &= 0.
 \end{aligned} \tag{A.47}$$

- $[Q_L^a, H_{RFT}]_{(4)}$.

At order g^4 , the possible contributions are

$$[Q_L^a, H_{RFT}]_{(4)} = \int_{\mathbf{x}} \left(d_{L(3)}^{\beta\gamma} \delta^{\delta\alpha} + \delta^{\beta\gamma} d_{R(3)}^{\delta\alpha} \right) \partial_{\mathbf{x}}^2 \left(V_L^{\alpha\beta}(\mathbf{x}) V_R^{\gamma\delta}(\mathbf{x}) \right)_{(1)} \tag{A.48}$$

However, this expression vanishes after substituting the first order result in eq. (A.44) and using the identity

$$(-\delta^{\alpha\beta} X_{(1)}^{\gamma\delta} + X_{(1)}^{\alpha\beta} \delta^{\gamma\delta}) \delta^{\delta\alpha} = 0. \tag{A.49}$$

Therefore

$$[Q_L^a, H_{RFT}]_{(4)} = 0 \quad (\text{A.50})$$

- $[Q_L^a, H_{RFT}]_{(5)}$.

The possible contributions at order g^5 are

$$\begin{aligned} & [Q_L^a, H_{RFT}]_{(5)} \\ &= \int_{\mathbf{x}} \left(d_{L(3)}^{\beta\gamma} \delta^{\delta\alpha} + \delta^{\beta\gamma} d_{R(3)}^{\delta\alpha} \right) \partial_{\mathbf{x}}^2 \left(V_L^{\alpha\beta} V_R^{\gamma\delta} \right)_{(2)} \\ &+ \int_{\mathbf{x}} \left(d_{L(3)}^{\beta\gamma} (-A_{(1)}^{\delta\alpha}) + A_{(1)}^{\beta\gamma} d_{R(3)}^{\delta\alpha} \right) \partial_{\mathbf{x}}^2 \left(V_L^{\alpha\beta} V_R^{\gamma\delta} \right)_{(1)} \\ &+ \int_{\mathbf{x}} \left(d_{L(4)}^{\beta\gamma} \delta^{\delta\alpha} + \delta^{\beta\gamma} d_{R(4)}^{\delta\alpha} \right) \partial_{\mathbf{x}}^2 \left(V_L^{\alpha\beta} V_R^{\gamma\delta} \right)_{(1)} \end{aligned} \quad (\text{A.51})$$

Note that the last term vanishes due to eq. (A.49). Let us focus on the other two terms. From eq. (A.44), one obtains

$$\begin{aligned} & \int_{\mathbf{x}} \left(d_{L(3)}^{\beta\gamma} \delta^{\delta\alpha} + \delta^{\beta\gamma} d_{R(3)}^{\delta\alpha} \right) \partial_{\mathbf{x}}^2 \left(V_L^{\alpha\beta} V_R^{\gamma\delta} \right)_{(2)} \\ &= \int_{\mathbf{x}} 2\text{Tr} \left[(d_{L(3)} + d_{R(3)}) \partial^2 X_{(2)} \right] \end{aligned} \quad (\text{A.52})$$

and

$$\begin{aligned} & \int_{\mathbf{x}} \left(d_{L(3)}^{\beta\gamma} (-A_{(1)}^{\delta\alpha}) + A_{(1)}^{\beta\gamma} d_{R(3)}^{\delta\alpha} \right) \partial_{\mathbf{x}}^2 \left(V_L^{\alpha\beta} V_R^{\gamma\delta} \right)_{(1)} \\ &= \int_{\mathbf{x}} \text{Tr} \left[(d_{L(3)} + d_{R(3)}) (\partial^2 X_{(1)} A_{(1)} - A_{(1)} \partial^2 X_{(1)}) \right] \end{aligned} \quad (\text{A.53})$$

Recall the expressions

$$\begin{aligned} A_{(1)}(\mathbf{x}) &= t^a \frac{\delta}{\delta \tilde{\rho}^a(\mathbf{x})}, \quad X_{(1)}(\mathbf{x}) = i \int_{\mathbf{y}} \phi(\mathbf{x} - \mathbf{y}) t^a \tilde{\rho}^a(\mathbf{y}), \\ X_{(2)} &= i \int_{\mathbf{y}} \phi(\mathbf{x} - \mathbf{y}) t^a \left(-\frac{1}{2} \frac{\delta}{\delta \tilde{\rho}^e} T_{ba}^e \rho^b \right). \end{aligned} \quad (\text{A.54})$$

Using this one obtains

$$\begin{aligned} \partial^2 X_{(1)} A_{(1)} - A_{(1)} \partial^2 X_{(1)} &= i(t^b t^e - t^e t^b) \tilde{\rho}^b(\mathbf{x}) \frac{\delta}{\delta \tilde{\rho}^e(\mathbf{x})} = iT_{eb}^a t^a \tilde{\rho}^b(\mathbf{x}) \frac{\delta}{\delta \tilde{\rho}^e(\mathbf{x})} \\ &= -2\partial^2 X_{(2)}. \end{aligned} \quad (\text{A.55})$$

As a consequence the contributions in eq. (A.52) and eq. (A.53) cancel each other.

We have proved that $[Q_L^a, H_{RFT}]_{(5)} = 0$.

- $[Q_L^a, H_{RFT}]_{(6)}$.

The possible contributions at order g^6 are

$$\begin{aligned}
 & [Q_L^a, H_{RFT}]_{(6)} \\
 &= \int_{\mathbf{x}} \left(d_{L(5)}^{\beta\gamma} \delta^{\delta\alpha} + \delta^{\beta\gamma} d_{R(5)}^{\delta\alpha} \right) \partial_{\mathbf{x}}^2 \left(V_L^{\alpha\beta} V_R^{\gamma\delta} \right)_{(1)} \\
 &+ \int_{\mathbf{x}} \left(d_{L(4)}^{\beta\gamma} \bar{V}_{R(1)}^{\delta\alpha} + \bar{V}_{L(1)}^{\beta\gamma} d_{R(4)}^{\delta\alpha} \right) \partial_{\mathbf{x}}^2 \left(V_L^{\alpha\beta} V_R^{\gamma\delta} \right)_{(1)} \\
 &+ \int_{\mathbf{x}} \left(d_{L(4)}^{\beta\gamma} \delta^{\delta\alpha} + \delta^{\beta\gamma} d_{R(4)}^{\delta\alpha} \right) \partial_{\mathbf{x}}^2 \left(V_L^{\alpha\beta} V_R^{\gamma\delta} \right)_{(2)} \\
 &+ \int_{\mathbf{x}} \left(d_{L(3)}^{\beta\gamma} \bar{V}_{R(2)}^{\delta\alpha} + \bar{V}_{L(2)}^{\beta\gamma} d_{R(3)}^{\delta\alpha} \right) \partial_{\mathbf{x}}^2 \left(V_L^{\alpha\beta} V_R^{\gamma\delta} \right)_{(1)} \\
 &+ \int_{\mathbf{x}} \left(d_{L(3)}^{\beta\gamma} \bar{V}_{R(1)}^{\delta\alpha} + \bar{V}_{L(1)}^{\beta\gamma} d_{R(3)}^{\delta\alpha} \right) \partial_{\mathbf{x}}^2 \left(V_L^{\alpha\beta} V_R^{\gamma\delta} \right)_{(2)} \\
 &+ \int_{\mathbf{x}} \left(d_{L(3)}^{\beta\gamma} \delta^{\delta\alpha} + \delta^{\beta\gamma} d_{R(3)}^{\delta\alpha} \right) \partial_{\mathbf{x}}^2 \left(V_L^{\alpha\beta} V_R^{\gamma\delta} \right)_{(3)}
 \end{aligned} \tag{A.56}$$

The first term vanishes due to eq. (A.49). The second and third terms add up to zero because of eq. (A.55). Now we focus on the fourth, fifth and sixth terms.

For the sixth term, note that

$$\begin{aligned}
 \delta^{\delta\alpha} \left(V_L^{\alpha\beta} V_R^{\gamma\delta} \right)_{(3)} &= (X_{(2)} X_{(1)} - X_{(1)} X_{(2)})^{\gamma\beta} \\
 \delta^{\beta\gamma} \left(V_L^{\alpha\beta} V_R^{\gamma\delta} \right)_{(3)} &= (X_{(1)} X_{(2)} - X_{(2)} X_{(1)})^{\alpha\delta}
 \end{aligned} \tag{A.57}$$

Using these relations, one calculates

$$\begin{aligned}
 & \int_{\mathbf{x}} \left(d_{L(3)}^{\beta\gamma} \delta^{\delta\alpha} + \delta^{\beta\gamma} d_{R(3)}^{\delta\alpha} \right) \partial_{\mathbf{x}}^2 \left(V_L^{\alpha\beta} V_R^{\gamma\delta} \right)_{(3)} \\
 &= \int_{\mathbf{x}} \text{Tr} \left[(d_{L(3)} - d_{R(3)}) \partial_{\mathbf{x}}^2 (X_{(2)} X_{(1)} - X_{(1)} X_{(2)}) \right] \\
 &= 0
 \end{aligned} \tag{A.58}$$

Here we have used the relation $d_{L(3)} = d_{R(3)}$.

For the fifth term,

$$\begin{aligned}
 d_{L(3)}^{\beta\gamma} \bar{V}_{R(1)}^{\delta\alpha} \partial^2 (V_L^{\alpha\beta} V_R^{\gamma\delta})_{(2)} &= -\text{Tr} \left[d_{L(3)} \left(X_{(2)} + \frac{1}{2} X_{(1)}^2 \right) A_{(1)} \right] \\
 &\quad - \text{Tr} \left[d_{L(3)} A_{(1)} \left(X_{(2)} + \frac{1}{2} X_{(1)}^2 \right) \right] + \text{Tr} [d_{L(3)} X_{(1)} A_{(1)} X_{(1)}]
 \end{aligned} \tag{A.59}$$

and

$$\begin{aligned}
 \bar{V}_{L(1)}^{\beta\gamma} d_{R(3)}^{\delta\alpha} (V_L^{\alpha\beta} V_R^{\gamma\delta})_{(2)} &= \text{Tr} \left[d_{R(3)} A_{(1)} \left(X_{(2)} + \frac{1}{2} X_{(1)}^2 \right) \right] \\
 &\quad + \text{Tr} \left[d_{R(3)} \left(X_{(2)} + \frac{1}{2} X_{(1)}^2 \right) A_{(1)} \right] - \text{Tr} [d_{R(3)} X_{(1)} A_{(1)} X_{(1)}]
 \end{aligned} \tag{A.60}$$

The sum of these two terms vanishes due to equality $d_{L(3)} = d_{R(3)}$.

For the fourth term

$$d_{L(3)}^{\beta\gamma} \bar{V}_{R(2)}^{\delta\alpha} (V_L^{\alpha\beta} V_R^{\gamma\delta})_{(1)} = -\text{Tr} \left[d_{L(3)} X_{(1)} (A_{(2)} + \frac{1}{2} A_{(1)}^2) \right] + \text{Tr} \left[d_{L(3)} (A_{(2)} + \frac{1}{2} A_{(1)}^2) X_{(1)} \right] \quad (\text{A.61})$$

and

$$\bar{V}_{L(2)}^{\beta\gamma} d_{R(3)}^{\delta\alpha} (V_L^{\alpha\beta} V_R^{\gamma\delta})_{(1)} = -\text{Tr} \left[d_{R(3)} (A_{(2)} + \frac{1}{2} A_{(1)}^2) X_{(1)} \right] + \text{Tr} \left[d_{R(3)} X_{(1)} (A_{(2)} + \frac{1}{2} A_{(1)}^2) \right] \quad (\text{A.62})$$

These two terms also cancel each other due to $d_{L(3)} = d_{R(3)}$. We therefore proved that $[Q_L^a, H_{RFT}]_{(6)} = 0$.

Thus we see that up to order $O(g^6)$ the left rotation generator Q_L^a commutes with the Hamiltonian.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] V.N. Gribov, *A reggeon diagram technique*, *Sov. Phys. JETP* **26** (1968) 414 [*Zh. Eksp. Teor. Fiz.* **53** (1967) 654] [[INSPIRE](#)].
- [2] E.A. Kuraev, L.N. Lipatov and V.S. Fadin, *The Pomeranchuk Singularity in Nonabelian Gauge Theories*, *Sov. Phys. JETP* **45** (1977) 199 [[INSPIRE](#)].
- [3] Ya.Ya. Balitsky and L.N. Lipatov, *The Pomeranchuk Singularity in Quantum Chromodynamics*, *Sov. J. Nucl. Phys.* **28** (1978) 22.
- [4] L.V. Gribov, E.M. Levin and M.G. Ryskin, *Semihard Processes in QCD*, *Phys. Rept.* **100** (1983) 1 [[INSPIRE](#)].
- [5] A.H. Mueller and J. Qiu, *Gluon Recombination and Shadowing at Small Values of x* , *Nucl. Phys. B* **268** (1986) 427 [[INSPIRE](#)].
- [6] A.H. Mueller, *Soft gluons in the infinite momentum wave function and the BFKL Pomeron*, *Nucl. Phys. B* **415** (1994) 373 [[INSPIRE](#)].
- [7] A.H. Mueller, *Unitarity and the BFKL Pomeron*, *Nucl. Phys. B* **437** (1995) 107 [[hep-ph/9408245](#)] [[INSPIRE](#)].
- [8] A.H. Mueller and B. Patel, *Single and double BFKL Pomeron exchange and a dipole picture of high-energy hard processes*, *Nucl. Phys. B* **425** (1994) 471 [[hep-ph/9403256](#)] [[INSPIRE](#)].
- [9] L.N. Lipatov, *Small x physics in perturbative QCD*, *Phys. Rept.* **286** (1997) 131 [[hep-ph/9610276](#)] [[INSPIRE](#)].
- [10] L.N. Lipatov, *High-energy scattering in QCD and in quantum gravity and two-dimensional field theories*, *Nucl. Phys. B* **365** (1991) 614 [[INSPIRE](#)].
- [11] L.N. Lipatov, *Gauge invariant effective action for high-energy processes in QCD*, *Nucl. Phys. B* **452** (1995) 369 [[hep-ph/9502308](#)] [[INSPIRE](#)].

- [12] R. Kirschner, L.N. Lipatov and L. Szymanowski, *Effective action for multi - Regge processes in QCD*, *Nucl. Phys. B* **425** (1994) 579 [[hep-th/9402010](#)] [[INSPIRE](#)].
- [13] R. Kirschner, L.N. Lipatov and L. Szymanowski, *Symmetry properties of the effective action for high-energy scattering in QCD*, *Phys. Rev. D* **51** (1995) 838 [[hep-th/9403082](#)] [[INSPIRE](#)].
- [14] J. Bartels, *Unitarity corrections to the Lipatov Pomeron and the four gluon operator in deep inelastic scattering in QCD*, *Z. Phys. C* **60** (1993) 471 [[INSPIRE](#)].
- [15] J. Bartels and M. Wusthoff, *The Triple Regge limit of diffractive dissociation in deep inelastic scattering*, *Z. Phys. C* **66** (1995) 157 [[INSPIRE](#)].
- [16] J. Bartels and C. Ewerz, *Unitarity corrections in high-energy QCD*, *JHEP* **09** (1999) 026 [[hep-ph/9908454](#)] [[INSPIRE](#)].
- [17] C. Ewerz, *Reggeization in high-energy QCD*, *JHEP* **04** (2001) 031 [[hep-ph/0103260](#)] [[INSPIRE](#)].
- [18] J. Bartels, *High-Energy Behavior in a Nonabelian Gauge Theory (II): First Corrections to $T_{n \rightarrow m}$ Beyond the Leading $\ln s$ Approximation*, *Nucl. Phys. B* **175** (1980) 365 [[INSPIRE](#)].
- [19] J. Kwiecinski and M. Praszalowicz, *Three Gluon Integral Equation and Odd c Singlet Regge Singularities in QCD*, *Phys. Lett. B* **94** (1980) 413 [[INSPIRE](#)].
- [20] A. Kovner, E. Levin, M. Li and M. Lublinsky, *The JIMWLK evolution and the s -channel unitarity*, *JHEP* **09** (2020) 199 [[arXiv:2006.15126](#)] [[INSPIRE](#)].
- [21] A. Kovner, E. Levin and M. Lublinsky, *QCD unitarity constraints on Reggeon Field Theory*, *JHEP* **08** (2016) 031 [[arXiv:1605.03251](#)] [[INSPIRE](#)].
- [22] L.D. McLerran and R. Venugopalan, *Computing quark and gluon distribution functions for very large nuclei*, *Phys. Rev. D* **49** (1994) 2233 [[hep-ph/9309289](#)] [[INSPIRE](#)].
- [23] L.D. McLerran and R. Venugopalan, *Gluon distribution functions for very large nuclei at small transverse momentum*, *Phys. Rev. D* **49** (1994) 3352 [[hep-ph/9311205](#)] [[INSPIRE](#)].
- [24] A.H. Mueller and G.P. Salam, *Large multiplicity fluctuations and saturation effects in onium collisions*, *Nucl. Phys. B* **475** (1996) 293 [[hep-ph/9605302](#)] [[INSPIRE](#)].
- [25] G.P. Salam, *Studies of unitarity at small x using the dipole formulation*, *Nucl. Phys. B* **461** (1996) 512 [[hep-ph/9509353](#)] [[INSPIRE](#)].
- [26] Y.V. Kovchegov and E. Levin, *Diffractive dissociation including multiple Pomeron exchanges in high parton density QCD*, *Nucl. Phys. B* **577** (2000) 221 [[hep-ph/9911523](#)] [[INSPIRE](#)].
- [27] M. Braun, *Structure function of the nucleus in the perturbative QCD with $N(c) \rightarrow \infty$ (BFKL Pomeron fan diagrams)*, *Eur. Phys. J. C* **16** (2000) 337 [[hep-ph/0001268](#)] [[INSPIRE](#)].
- [28] M.A. Braun and G.P. Vacca, *Triple Pomeron vertex in the limit $N(c) \rightarrow \infty$* , *Eur. Phys. J. C* **6** (1999) 147 [[hep-ph/9711486](#)] [[INSPIRE](#)].
- [29] J. Bartels, M. Braun and G.P. Vacca, *Pomeron vertices in perturbative QCD in diffractive scattering*, *Eur. Phys. J. C* **40** (2005) 419 [[hep-ph/0412218](#)] [[INSPIRE](#)].
- [30] J. Bartels, L.N. Lipatov and G.P. Vacca, *Interactions of reggeized gluons in the Mobius representation*, *Nucl. Phys. B* **706** (2005) 391 [[hep-ph/0404110](#)] [[INSPIRE](#)].
- [31] M.A. Braun, *Nucleus-nucleus scattering in perturbative QCD with $N(c) \rightarrow \infty$* , *Phys. Lett. B* **483** (2000) 115 [[hep-ph/0003004](#)] [[INSPIRE](#)].

- [32] M.A. Braun, *Nucleus nucleus interaction in the perturbative QCD*, *Eur. Phys. J. C* **33** (2004) 113 [[hep-ph/0309293](#)] [[INSPIRE](#)].
- [33] M.A. Braun, *Conformal invariant Pomeron interaction in the perturbative QCD with large N_c* , *Phys. Lett. B* **632** (2006) 297 [[hep-ph/0512057](#)] [[INSPIRE](#)].
- [34] I. Balitsky, *Factorization and high-energy effective action*, *Phys. Rev. D* **60** (1999) 014020 [[hep-ph/9812311](#)] [[INSPIRE](#)].
- [35] Y.V. Kovchegov, *Small x $F(2)$ structure function of a nucleus including multiple Pomeron exchanges*, *Phys. Rev. D* **60** (1999) 034008 [[hep-ph/9901281](#)] [[INSPIRE](#)].
- [36] T. Altinoluk, C. Contreras, A. Kovner, E. Levin, M. Lublinsky and A. Shulkin, *QCD reggeon calculus from JIMWLK Evolution*, *Int. J. Mod. Phys. Conf. Ser.* **25** (2014) 1460025 [[INSPIRE](#)].
- [37] T. Altinoluk, N. Armesto, A. Kovner, E. Levin and M. Lublinsky, *KLWMIJ Reggeon field theory beyond the large N_c limit*, *JHEP* **08** (2014) 007 [[arXiv:1402.5936](#)] [[INSPIRE](#)].
- [38] T. Altinoluk, A. Kovner, E. Levin and M. Lublinsky, *Reggeon Field Theory for Large Pomeron Loops*, *JHEP* **04** (2014) 075 [[arXiv:1401.7431](#)] [[INSPIRE](#)].
- [39] T. Altinoluk, C. Contreras, A. Kovner, E. Levin, M. Lublinsky and A. Shulkin, *QCD Reggeon Calculus From KLWMIJ/JIMWLK Evolution: Vertices, Reggeization and All*, *JHEP* **09** (2013) 115 [[arXiv:1306.2794](#)] [[INSPIRE](#)].
- [40] J. Jalilian-Marian, A. Kovner, A. Leonidov and H. Weigert, *The BFKL equation from the Wilson renormalization group*, *Nucl. Phys. B* **504** (1997) 415 [[hep-ph/9701284](#)] [[INSPIRE](#)].
- [41] J. Jalilian-Marian, A. Kovner, A. Leonidov and H. Weigert, *The Wilson renormalization group for low x physics: Towards the high density regime*, *Phys. Rev. D* **59** (1998) 014014 [[hep-ph/9706377](#)] [[INSPIRE](#)].
- [42] J. Jalilian-Marian, A. Kovner and H. Weigert, *The Wilson renormalization group for low x physics: Gluon evolution at finite parton density*, *Phys. Rev. D* **59** (1998) 014015 [[hep-ph/9709432](#)] [[INSPIRE](#)].
- [43] A. Kovner and J. Milhano, *Vector potential versus color charge density in low x evolution*, *Phys. Rev. D* **61** (2000) 014012 [[hep-ph/9904420](#)] [[INSPIRE](#)].
- [44] A. Kovner, J. Milhano and H. Weigert, *Relating different approaches to nonlinear QCD evolution at finite gluon density*, *Phys. Rev. D* **62** (2000) 114005 [[hep-ph/0004014](#)] [[INSPIRE](#)].
- [45] H. Weigert, *Unitarity at small Bjorken x* , *Nucl. Phys. A* **703** (2002) 823 [[hep-ph/0004044](#)] [[INSPIRE](#)].
- [46] A. Kovner and M. Lublinsky, *In pursuit of Pomeron loops: The JIMWLK equation and the Wess-Zumino term*, *Phys. Rev. D* **71** (2005) 085004 [[hep-ph/0501198](#)] [[INSPIRE](#)].
- [47] E. Iancu, A. Leonidov and L.D. McLerran, *Nonlinear gluon evolution in the color glass condensate. 1.*, *Nucl. Phys. A* **692** (2001) 583 [[hep-ph/0011241](#)] [[INSPIRE](#)].
- [48] E. Iancu, A. Leonidov and L.D. McLerran, *The Renormalization group equation for the color glass condensate*, *Phys. Lett. B* **510** (2001) 133 [[hep-ph/0102009](#)] [[INSPIRE](#)].
- [49] E. Ferreiro, E. Iancu, A. Leonidov and L. McLerran, *Nonlinear gluon evolution in the color glass condensate. 2.*, *Nucl. Phys. A* **703** (2002) 489 [[hep-ph/0109115](#)] [[INSPIRE](#)].

- [50] A. Kovner and M. Lublinsky, *Odderon and seven Pomerons: QCD Reggeon field theory from JIMWLK evolution*, *JHEP* **02** (2007) 058 [[hep-ph/0512316](#)] [[INSPIRE](#)].
- [51] I. Balitsky and G.A. Chirilli, *NLO evolution of color dipoles in $N = 4$ SYM*, *Nucl. Phys. B* **822** (2009) 45 [[arXiv:0903.5326](#)] [[INSPIRE](#)].
- [52] I. Balitsky and G.A. Chirilli, *Rapidity evolution of Wilson lines at the next-to-leading order*, *Phys. Rev. D* **88** (2013) 111501 [[arXiv:1309.7644](#)] [[INSPIRE](#)].
- [53] A. Kovner, M. Lublinsky and Y. Mulian, *Jalilian-Marian, Iancu, McLerran, Weigert, Leonidov, Kovner evolution at next to leading order*, *Phys. Rev. D* **89** (2014) 061704 [[arXiv:1310.0378](#)] [[INSPIRE](#)].
- [54] A. Kovner, M. Lublinsky and Y. Mulian, *NLO JIMWLK evolution unabridged*, *JHEP* **08** (2014) 114 [[arXiv:1405.0418](#)] [[INSPIRE](#)].
- [55] M. Lublinsky and Y. Mulian, *High Energy QCD at NLO: from light-cone wave function to JIMWLK evolution*, *JHEP* **05** (2017) 097 [[arXiv:1610.03453](#)] [[INSPIRE](#)].
- [56] S. Caron-Huot, *Resummation of non-global logarithms and the BFKL equation*, *JHEP* **03** (2018) 036 [[arXiv:1501.03754](#)] [[INSPIRE](#)].
- [57] S. Caron-Huot and M. Herranen, *High-energy evolution to three loops*, *JHEP* **02** (2018) 058 [[arXiv:1604.07417](#)] [[INSPIRE](#)].
- [58] A.H. Mueller and A.I. Shoshi, *Small x physics beyond the Kovchegov equation*, *Nucl. Phys. B* **692** (2004) 175 [[hep-ph/0402193](#)] [[INSPIRE](#)].
- [59] E. Iancu and A.H. Mueller, *Rare fluctuations and the high-energy limit of the S matrix in QCD*, *Nucl. Phys. A* **730** (2004) 494 [[hep-ph/0309276](#)] [[INSPIRE](#)].
- [60] E. Iancu and D.N. Triantafyllopoulos, *A Langevin equation for high energy evolution with Pomeron loops*, *Nucl. Phys. A* **756** (2005) 419 [[hep-ph/0411405](#)] [[INSPIRE](#)].
- [61] E. Iancu and D.N. Triantafyllopoulos, *Non-linear QCD evolution with improved triple-Pomeron vertices*, *Phys. Lett. B* **610** (2005) 253 [[hep-ph/0501193](#)] [[INSPIRE](#)].
- [62] A.H. Mueller, A.I. Shoshi and S.M.H. Wong, *Extension of the JIMWLK equation in the low gluon density region*, *Nucl. Phys. B* **715** (2005) 440 [[hep-ph/0501088](#)] [[INSPIRE](#)].
- [63] E. Levin and M. Lublinsky, *Towards a symmetric approach to high energy evolution: Generating functional with Pomeron loops*, *Nucl. Phys. A* **763** (2005) 172 [[hep-ph/0501173](#)] [[INSPIRE](#)].
- [64] E. Levin, J. Miller and A. Prygarin, *Summing Pomeron loops in the dipole approach*, *Nucl. Phys. A* **806** (2008) 245 [[arXiv:0706.2944](#)] [[INSPIRE](#)].
- [65] E. Iancu, G. Soyez and D.N. Triantafyllopoulos, *On the probabilistic interpretation of the evolution equations with Pomeron loops in QCD*, *Nucl. Phys. A* **768** (2006) 194 [[hep-ph/0510094](#)] [[INSPIRE](#)].
- [66] A. Kovner and M. Lublinsky, *From target to projectile and back again: Selfduality of high energy evolution*, *Phys. Rev. Lett.* **94** (2005) 181603 [[hep-ph/0502119](#)] [[INSPIRE](#)].
- [67] Y. Hatta, E. Iancu, L. McLerran, A. Stasto and D.N. Triantafyllopoulos, *Effective Hamiltonian for QCD evolution at high energy*, *Nucl. Phys. A* **764** (2006) 423 [[hep-ph/0504182](#)] [[INSPIRE](#)].
- [68] I. Balitsky, *High-energy effective action from scattering of QCD shock waves*, *Phys. Rev. D* **72** (2005) 074027 [[hep-ph/0507237](#)] [[INSPIRE](#)].

- [69] A. Kovner and M. Lublinsky, *Dense-dilute duality at work: Dipoles of the target*, *Phys. Rev. D* **72** (2005) 074023 [[hep-ph/0503155](#)] [[INSPIRE](#)].
- [70] M. Li and A. Kovner, *JIMWLK Evolution, Lindblad Equation and Quantum-Classical Correspondence*, *JHEP* **05** (2020) 036 [[arXiv:2002.02282](#)] [[INSPIRE](#)].
- [71] S. Bondarenko and L. Motyka, *Solving effective field theory of interacting QCD Pomerons in the semi-classical approximation*, *Phys. Rev. D* **75** (2007) 114015 [[hep-ph/0605185](#)] [[INSPIRE](#)].
- [72] A. Kovner and M. Lublinsky, *The Yin and Yang of high energy chromodynamics: Scattering in black and white*, *Nucl. Phys. A* **779** (2006) 220 [[hep-ph/0604085](#)] [[INSPIRE](#)].
- [73] A. Kovner and M. Lublinsky, *Remarks on high energy evolution*, *JHEP* **03** (2005) 001 [[hep-ph/0502071](#)] [[INSPIRE](#)].
- [74] A. Kovner and M. Lublinsky, *More remarks on high energy evolution*, *Nucl. Phys. A* **767** (2006) 171 [[hep-ph/0510047](#)] [[INSPIRE](#)].
- [75] O.V. Kancheli, *Parton models and frame independence of high-energy cross-sections*, [arXiv:2003.04654](#) [[INSPIRE](#)].