

RECEIVED: September 6, 2014

ACCEPTED: October 22, 2014

PUBLISHED: November 11, 2014

Lepton flavor violation in low-scale seesaw models: SUSY and non-SUSY contributions

A. Abada,^a M.E. Krauss,^b W. Porod,^b F. Staub,^c A. Vicente^d and C. Weiland^e

^a*Laboratoire de Physique Théorique, CNRS — UMR 8627, Université de Paris-Sud 11,
F-91405 Orsay Cedex, France*

^b*Institut für Theoretische Physik und Astronomie, Universität Würzburg,
Emil-Hilb-Weg 22, 97074 Würzburg, Germany*

^c*Physikalisches Institut der Universität Bonn,
Nussallee 12, 53115 Bonn, Germany*

^d*IFPA, Dep. AGO, Université de Liège,
Bat B5, Sart-Tilman B-4000 Liège 1, Belgium*

^e*Departamento de Física Teórica and Instituto de Física Teórica,
IFT-UAM/CSIC, Universidad Autónoma de Madrid,
Calle Nicolás Cabrera 13–15, Cantoblanco, 28049 Madrid, Spain*

E-mail: abada@th.u-psud.fr, manuel.krauss@physik.uni-wuerzburg.de,
porod@physik.uni-wuerzburg.de, fnstaub@th.physik.uni-bonn.de,
avelino.vicente@ulg.ac.be, cedric.weiland@uam.es

ABSTRACT: Taking the supersymmetric inverse seesaw mechanism as the explanation for neutrino oscillation data, we investigate charged lepton flavor violation in radiative and 3-body lepton decays as well as in neutrinoless $\mu - e$ conversion in muonic atoms. In contrast to former studies, we take into account all possible contributions: supersymmetric as well as non-supersymmetric. We take CMSSM-like boundary conditions for the soft supersymmetry breaking parameters. We find several regions where cancellations between various contributions exist, reducing the lepton flavor violating rates by an order of magnitude compared to the case where only the dominant contribution is taken into account. This is in particular important for the correct interpretation of existing data as well as for estimating the reach of near future experiments where the sensitivity will be improved by one to two orders of magnitude. Moreover, we demonstrate that ratios like $\text{BR}(\tau \rightarrow 3\mu)/\text{BR}(\tau \rightarrow \mu e^+ e^-)$ can be used to determine whether the supersymmetric contributions dominate over the W^\pm and H^\pm contributions or vice versa.

KEYWORDS: Rare Decays, Beyond Standard Model, Neutrino Physics, Supersymmetric Standard Model

ARXIV EPRINT: [1408.0138](https://arxiv.org/abs/1408.0138)

Contents

1	Introduction	1
2	Inverse seesaw model and its supersymmetric extension	4
3	Low energy observables	6
3.1	Effective lagrangian	6
3.2	$\ell_\alpha \rightarrow \ell_\beta \gamma$	7
3.3	$\ell_\alpha^- \rightarrow \ell_\beta^- \ell_\beta^- \ell_\beta^+$	7
3.4	$\ell_\alpha^- \rightarrow \ell_\beta^- \ell_\gamma^- \ell_\gamma^+$	8
3.5	$\ell_\alpha^- \rightarrow \ell_\beta^+ \ell_\gamma^- \ell_\gamma^-$	9
3.6	Coherent $\mu - e$ conversion in nuclei	9
4	Results	10
4.1	Numerical setup	10
4.2	Numerical results	12
5	Conclusions	18
A	Masses and vertices	20
A.1	Mass matrices	20
A.2	Vertices	24
A.2.1	Fermion-scalar vertices	24
A.2.2	Fermion-vector vertices	26
A.2.3	Scalar vertices	27
A.2.4	Scalar-vector vertices	29
A.2.5	Vector vertices	30
B	Renormalization Group Equations	30
C	Loop integrals	34
D	Photonic penguin contributions to LFV	36
D.1	Feynman diagrams	37
D.2	Neutralino contributions	38
D.3	Chargino contributions	38
D.4	W^+ and H^+ contributions	38
E	Z and Higgs penguin contributions to LFV	39
E.1	Feynman diagrams	39
E.2	Neutralino contributions	42
E.2.1	Z -penguins	42
E.2.2	Scalar penguins	43

E.3	Chargino contributions	44
E.3.1	Z-penguins	44
E.3.2	Scalar penguins	45
E.4	W^+ and H^+ contributions	48
E.4.1	Z-penguins	48
E.4.2	Scalar penguins	49
F	Box contributions to LFV	52
F.1	Four lepton boxes	52
F.1.1	Feynman diagrams	52
F.1.2	Neutralino contributions	54
F.1.3	Chargino contributions	54
F.1.4	W^+ and H^+ contributions	55
F.2	Additional boxes for $\ell_\alpha^- \rightarrow \ell_\beta^- \ell_\gamma^+ \ell_\gamma^-$	57
F.2.1	Crossed neutralino contributions	57
F.2.2	Crossed chargino contributions	57
F.2.3	Crossed W^+ and H^+ contributions	59
F.3	Two-lepton — two-quark boxes	61
F.3.1	Feynman diagrams	61
F.3.2	Down quarks	62
F.3.3	Up quarks	64
G	Form factors of the 4-fermion operators	66

1 Introduction

The recent discovery of a bosonic state at the Large Hadron Collider (LHC) [1, 2] stands as a major breakthrough in particle physics. Although further confirmation is required, all data are compatible with the long-awaited Higgs boson, thus completing the Standard Model (SM) particle content. Furthermore, the properties and decay modes of this scalar are in good agreement with the SM expectations, making the SM picture more motivated than ever.

In this context, it is crucial to keep in mind that the SM cannot be the ultimate theory. In fact, and besides theoretical arguments such as the hierarchy problem, there are very good experimental reasons to go beyond the SM (BSM). The best of these motivations is the existence of non-zero neutrino masses and mixing angles, now firmly established by neutrino oscillation experiments [3–5]. Since the SM lepton sector does not include them, one has to go beyond the SM.

A generic prediction in most of these neutrino mass models is lepton flavor violation (LFV), not only in the neutrino sector but also for the charged leptons. Depending of the exact realization of the neutrino mass model, the rates for the LFV processes can be

very different. For instance, high-scale models typically predict small branching ratios, thus making LFV hard (if not impossible) to be discovered. In contrast, one expects measurable LFV rates if the scale of new physics is not far from the electroweak (EW) scale. These low-scale mechanisms generating neutrino masses are thus more attractive from a phenomenological point of view, since they offer a window to new physics thanks to their LFV promising perspectives. Moreover, they can be directly tested at the LHC via the production of new particles if these are light enough.

On the experimental side, the field of LFV physics will live an era of unprecedented developments in the near future, with dedicated experiments in different fronts.¹ In the case of the muon radiative decay $\mu \rightarrow e\gamma$, the MEG collaboration has announced plans for future upgrades. These will allow for an improvement of the current bound, $\text{BR}(\mu \rightarrow e\gamma) < 5.7 \cdot 10^{-13}$ [7], reaching a sensitivity of about $6 \cdot 10^{-14}$ after 3 years of acquisition time [8]. Limits on τ radiative decays are less stringent, but they are expected to be improved at Belle II [9]. These will also search for lepton flavor violating B -meson decays. Moreover, the perspectives for the 3-body decays $\ell_\alpha \rightarrow 3\ell_\beta$ are good as well. The decay $\mu \rightarrow 3e$ was searched for long ago by the SINDRUM experiment [10], setting the limit $\text{Br}(\mu \rightarrow 3e) < 1.0 \cdot 10^{-12}$. The future Mu3e experiment announces a sensitivity of $\sim 10^{-16}$ [11], which would imply a 4 orders of magnitude improvement. In the case of τ decays to three charged leptons, Belle II will again be the facility where improvements are expected [12], although recently the LHCb collaboration has reported first bounds on $\tau \rightarrow 3\mu$ [13]. The LFV process where the best developments are expected in the next few years is neutrinoless $\mu - e$ conversion in muonic atoms. In the near future, many different experiments will search for a positive signal. These include Mu2e [14–16], DeeMe [17], COMET [18, 19] and PRISM/PRIME [20]. The expected sensitivities for the conversion rate range from a modest 10^{-14} in the near future to an impressive 10^{-18} . Finally, one can also search for LFV in high-energy experiments, such as the LHC. A popular process in this case is the Higgs boson LFV decay to a pair of charged leptons, $h \rightarrow \ell_\alpha \ell_\beta$, with $\alpha \neq \beta$ [21, 22], which has recently received some attention [23–34]. First bounds on $h \rightarrow \mu\tau$ have been reported by the CMS collaboration [35].² For other possibilities to search for LFV at high-energy colliders, see [36–52]. In table 1 we collect present bounds and expected near-future sensitivities for the most popular low-energy LFV observables.

With such a large variety of processes, a proper theoretical understanding of potential hierarchies or correlations in a given model becomes necessary. This goal requires detailed analytical and numerical studies of the different contributions to the LFV processes, in order to determine the dominant ones and to get a proper interpretation of the LFV bounds. Furthermore, the understanding of the LFV anatomy of several models allows one to discriminate among them by using combinations of observables which have definite predictions [58].

In this work we are interested in LFV in supersymmetric and non-supersymmetric variants of the inverse seesaw model (ISS) [59]. This low-scale neutrino mass model constitutes

¹See [6] for a recent review.

²The CMS collaboration also reports an intriguing 2.5σ excess in $h \rightarrow \mu\tau$ leading to $\text{BR}(h \rightarrow \tau\mu) \sim 1\%$.

LFV Process	Present Bound	Future Sensitivity
$\mu \rightarrow e\gamma$	5.7×10^{-13} [7]	6×10^{-14} [8]
$\tau \rightarrow e\gamma$	3.3×10^{-8} [53]	$\sim 3 \times 10^{-9}$ [9]
$\tau \rightarrow \mu\gamma$	4.4×10^{-8} [53]	$\sim 3 \times 10^{-9}$ [9]
$\mu \rightarrow eee$	1.0×10^{-12} [10]	$\sim 10^{-16}$ [11]
$\tau \rightarrow \mu\mu\mu$	2.1×10^{-8} [54]	$\sim 10^{-9}$ [9]
$\tau^- \rightarrow e^-\mu^+\mu^-$	2.7×10^{-8} [54]	$\sim 10^{-9}$ [9]
$\tau^- \rightarrow \mu^-e^+e^-$	1.8×10^{-8} [54]	$\sim 10^{-9}$ [9]
$\tau \rightarrow eee$	2.7×10^{-8} [54]	$\sim 10^{-9}$ [9]
$\mu^-, \text{Ti} \rightarrow e^-, \text{Ti}$	4.3×10^{-12} [55]	$\sim 10^{-18}$ [20]
$\mu^-, \text{Au} \rightarrow e^-, \text{Au}$	7×10^{-13} [56]	
$\mu^-, \text{Al} \rightarrow e^-, \text{Al}$		$10^{-15} - 10^{-18}$
$\mu^-, \text{SiC} \rightarrow e^-, \text{SiC}$		10^{-14} [57]

Table 1. Current experimental bounds and future sensitivities for some low-energy LFV observables.

a very interesting alternative to the usual seesaw mechanism. The suppression mechanism that guarantees the smallness of neutrino masses is the introduction of a slight breaking of lepton number in the singlet sector, in the form of a small (compared to the EW scale) Majorana mass for the X singlets. This allows for large Yukawa couplings compatible with a low (TeV or even lower) mass for the seesaw mediators. With this combination, one expects a very rich phenomenology, including sizable LFV rates and additional contributions to the radiative corrections to the Higgs mass [60–62]. In the supersymmetric (SUSY) version of the ISS, the new singlet fermions are promoted to singlet superfields. The appealing features of the ISS mechanism are kept also in the SUSY version.

LFV in models with light right-handed (RH) neutrinos has already been studied in great detail. Early studies [63–66] already pointed out the existence of large enhancements in the LFV rates with respect to those found in high-scale models. More recently, there has been a revived interest due to the expected experimental improvements in the near future. Interestingly, dominant contributions have been found in (non-SUSY) box diagrams induced by RH neutrinos. This was first shown in [67] and later confirmed in [68–70]. In this case, the future $\mu - e$ conversion experiments will play a major role in constraining light right-handed neutrino scenarios. The usual photon penguin contributions get also enhanced in the presence of light RH neutrinos, see for example [71]. Regarding the SUSY contributions, several studies have recently addressed the role of the Z -penguins. A large enhancement with respect to the usual dipole contribution was reported in [72]. Later, this result was (qualitatively) confirmed in [70] and further exploited in several phenomenological studies [73–76]. However, in [77] it was shown that the results in [72] (and the subsequent studies [73–76]) are incorrect, due to an inconsistency in [78]. While

this has a negligible impact in the case of high-scale seesaw models, this is not the case for low-scale seesaw models like the supersymmetric version of the ISS.

Given that recent studies pointed out important but partial results and the upcoming experimental improvements, we aim in this work for a complete calculation of the various LFV observables taking into account all contributions at the same time. One of our results will be that there exist several regions in parameter space where cancellations between various contributions occur, changing the interpretation of existing and future experimental results. In order to do so we have made use of `FlavorKit` [79], a tool that combines the analytical power of `SARAH` [80–84] with the numerical routines of `SPheno` [85, 86] to obtain predictions for flavor observables in a wide range of models. This setup makes use of `FeynArts/FormCalc` [87–92] to compute generic predictions for the form factors of the relevant operators and thus provides an automatic computation of the flavor observables. We use this setup to compute for the first time the Higgs penguin contributions to LFV in the inverse seesaw.³ In addition, we improve previous studies in others aspects as well: (i) we make use of the full 2-loop renormalization group equations (RGEs) including all flavor effects in the SM and SUSY sectors to obtain the parameters entering the calculation, and (ii) we include for the first time the decays $\tau^- \rightarrow \mu^+ e^- e^-$ and $\tau^- \rightarrow e^+ \mu^- \mu^-$.

The paper is organized as follows: in section 2 we present the ISS model and its supersymmetric extension. The LFV observables induced by the extended particle content and the dominant contributions are discussed in section 3, and in section 4 we present our numerical results. In section 5 we draw our conclusions. In the appendices we first introduce the formulae for the mass matrices and our convention for the loop integrals before presenting the additional contributions to the 1- and 2-loop RGEs compared to the MSSM case. More importantly, they contain the complete set of contributions to the LFV observables discussed in this paper.

2 Inverse seesaw model and its supersymmetric extension

In the inverse seesaw, the Standard Model field content is extended by n_R generations of right-handed neutrinos ν_R and n_X generations of singlet fermions X (such that $n_R + n_X = N_s$), both with lepton number $L = +1$ [59, 63, 94]. The corresponding Lagrangian before EWSB has the form

$$\mathcal{L}_{\text{ISS}} = \mathcal{L}_{\text{SM}} - Y_\nu^{ij} \overline{\nu_{Ri}} \tilde{H}^\dagger L_j - M_R^{ij} \overline{\nu_{Ri}} X_j - \frac{1}{2} \mu_X^{ij} \overline{X_i^C} X_j + h.c. , \quad (2.1)$$

where a sum over $i, j = 1, 2, 3$ is assumed.⁴ \mathcal{L}_{SM} is the SM Lagrangian, Y_ν are the neutrino Yukawa couplings and M_R is a complex mass matrix that generates a lepton number

³The Higgs penguin contributions to LFV processes were first considered in the context of the inverse seesaw in [93]. However, our paper goes beyond this reference in two ways: by doing the computation in the mass basis and by taking into account all contributions to the Higgs penguins.

⁴The ISS requires the introduction of at least two right-handed neutrinos in order to account for the active neutrino masses and mixings. The most minimal ISS realization [95–97] consists in the addition of two right-handed and two sterile neutrinos to the SM content. However, its minimal SUSY realization [98] requires only one pair of fermionic singlets.

conserving mass term for the fermion singlets. The complex symmetric mass matrix μ_X violates lepton number by two units and is naturally small, in the sense of 't Hooft [99], since in the limit $\mu_X \rightarrow 0$ lepton number is restored. This Majorana mass term also leads to a small mass splitting in the heavy neutrino sector, which then become quasi-Dirac neutrinos.

After EWSB, in the basis (ν_L, ν_R^C, X) , the 9×9 neutrino mass matrix is given by

$$M_{\text{ISS}} = \begin{pmatrix} 0 & m_D^T & 0 \\ m_D & 0 & M_R \\ 0 & M_R^T & \mu_X \end{pmatrix}. \quad (2.2)$$

where $m_D = \frac{1}{\sqrt{2}} Y_\nu v$ and $v/\sqrt{2}$ is the vacuum expectation value (vev) of the Higgs boson.

Under the assumption $\mu_X \ll m_D \ll M_R$, the mass matrix M_{ISS} can be block-diagonalized to give an effective mass matrix for the light neutrinos [100]

$$M_{\text{light}} \simeq m_D^T M_R^{T^{-1}} \mu_X M_R^{-1} m_D, \quad (2.3)$$

whereas the heavy quasi-Dirac neutrinos have masses corresponding approximately to the entries of M_R .

As usual, one can easily obtain a supersymmetric version of the model by promoting the corresponding fields to superfields $\tilde{\nu}_i^C$ and \tilde{X}_i ($i = 1, 2, 3$) and including the corresponding interactions in the superpotential. This reads

$$W = W_{\text{MSSM}} + \epsilon_{ab} Y_\nu^{ij} \tilde{\nu}_i^C \tilde{L}_j^a \tilde{H}_u^b + M_{R_{ij}} \tilde{\nu}_i^C \tilde{X}_j + \frac{1}{2} \mu_{X_{ij}} \tilde{X}_i \tilde{X}_j. \quad (2.4)$$

W_{MSSM} is the superpotential of the MSSM given by

$$W_{\text{MSSM}} = \epsilon_{ab} Y_u^{ij} \tilde{U}_i^C \tilde{Q}_j^a \tilde{H}_u^b - \epsilon_{ab} Y_d^{ij} \tilde{D}_i^C \tilde{Q}_j^a \tilde{H}_d^b - \epsilon_{ab} Y_e^{ij} \tilde{E}_i^C \tilde{L}_j^a \tilde{H}_d^b + \epsilon_{ab} \mu \tilde{H}_u^a \tilde{H}_d^b, \quad (2.5)$$

where we skipped the color indices. The corresponding soft SUSY breaking Lagrangian is given by

$$\begin{aligned} -\mathcal{L}_{\text{soft}} &= -\mathcal{L}_{\text{MSSM}}^{\text{soft}} + \tilde{\nu}_{ij}^C m_{\tilde{\nu}_i^C}^2 \tilde{\nu}_j^{C*} + \tilde{X}_i^* m_{X_{ij}}^2 \tilde{X}_j \\ &+ \left(T_\nu^{ij} \epsilon_{ab} \tilde{\nu}_i^C \tilde{L}_j^a H_u^b + B_{M_R}^{ij} \tilde{\nu}_i^C \tilde{X}_j + \frac{1}{2} B_{\mu_X}^{ij} \tilde{X}_i \tilde{X}_j + \tilde{X}_i^* m_{X_{ij}}^2 \tilde{\nu}_j^C + \text{h.c.} \right), \end{aligned} \quad (2.6)$$

where $B_{M_R}^{ij}$ and $B_{\mu_X}^{ij}$ are the new parameters involving the scalar partners of the sterile neutrino states. Notice that while the former conserves lepton number, the latter violates lepton number by two units. Finally, $\mathcal{L}_{\text{MSSM}}^{\text{soft}}$ collects the soft SUSY breaking terms of the MSSM.

$$\begin{aligned} -\mathcal{L}_{\text{MSSM}}^{\text{soft}} &= \left(\epsilon_{ab} T_u^{ij} \tilde{U}_i^C \tilde{Q}_j^a H_u^b - \epsilon_{ab} T_d^{ij} \tilde{D}_i^C \tilde{Q}_j^a H_d^b - \epsilon_{ab} T_e^{ij} \tilde{E}_i^C \tilde{L}_j^a H_d^b + \epsilon_{ab} B_\mu H_u^a H_d^b + \text{h.c.} \right) \\ &+ \frac{1}{2} \left(M_1 \lambda_B \lambda_B + \delta_{ab} M_2 \lambda_W^a \lambda_W^b + M_3 \lambda_G^\alpha \lambda_G^\beta + \text{h.c.} \right) \\ &+ \left(\delta_{ab} (\tilde{Q}_i^a)^* m_{q,ij}^2 \tilde{Q}_j^b + (\tilde{D}_i^C)^* m_{d,ij}^2 \tilde{D}_j^C + (\tilde{U}_i^C)^* m_{u,ij}^2 \tilde{U}_j^C \right) \\ &+ \tilde{E}_i^{C*} m_{e,ij}^2 \tilde{E}_j^C + \delta_{ab} (\tilde{L}_i^a)^* m_{l,ij}^2 \tilde{L}_j^b + m_{H_d}^2 |H_d|^2 + m_{H_u}^2 |H_u|^2. \end{aligned} \quad (2.7)$$

The neutrino mass matrix has the same form as in eq. (2.2), just replacing v by v_u , the vev of the up-type Higgs boson. The mass matrices of this model are the same as in the MSSM apart from the sneutrino sector. Neglecting for the moment being the soft-breaking terms which lead to a splitting between the scalar and pseudoscalar parts, the corresponding mass matrix reads

$$m_{\tilde{\nu}^i}^2 = \begin{pmatrix} m_L^2 + \frac{1}{2}v_u^2 Y_\nu^T Y_\nu^* + D_L & -\frac{1}{\sqrt{2}}(v_d \mu Y_\nu^T - v_u T_\nu^\dagger) & \frac{1}{\sqrt{2}}v_u \Re(Y_\nu^T M_R^*) \\ -\frac{1}{\sqrt{2}}(v_d \mu Y_\nu^* - v_u T_\nu) & m_{\tilde{\nu}^C}^2 + M_R M_R^\dagger + \frac{1}{2}v_u^2 Y_\nu Y_\nu^\dagger & -M_R \mu_S^* \\ \frac{1}{\sqrt{2}}v_u M_R^T Y_\nu^* & -\mu_S M_R^\dagger & M_R^T M_R^* + m_S^2 + \mu_S \mu_S^* \end{pmatrix} \quad (2.8)$$

with

$$D_L = -m_Z^2 \cos^2 \theta_W \cos 2\beta \mathbf{1}. \quad (2.9)$$

The complete mass matrices including the B -parameters as well as all other mass matrices can be found in appendix A.1.

3 Low energy observables

The fact that the LHC has not yet seen any supersymmetric particles [101, 102] implies, at least in the specific SUSY model we consider in this work, that squarks and gluinos must be heavy. However, it could well be that sleptons, charginos and neutralinos are relatively light, thus having large contributions to LFV decays. Here we will consider the processes $\ell_\alpha \rightarrow \ell_\beta \gamma$, $\ell_\alpha \rightarrow \ell_\beta \ell_\gamma \ell_\delta$ and $\mu - e$ conversion in nuclei. In this section we will present the effective low-energy lagrangian and the basic formulae for the observables. This will also serve to fix our notation (we stay close to the conventions of ref. [79]). The details for the calculations of the corresponding form factors can be found in appendices C–G.

3.1 Effective lagrangian

The interaction lagrangian relevant for LFV can be written as

$$\mathcal{L}_{\text{LFV}} = \mathcal{L}_{\ell\ell\gamma} + \mathcal{L}_{4\ell} + \mathcal{L}_{2\ell 2d} + \mathcal{L}_{2\ell 2u}. \quad (3.1)$$

with

$$\mathcal{L}_{\ell\ell\gamma} = e \bar{\ell}_\beta [\gamma^\mu (K_1^L P_L + K_1^R P_R) + i m_{\ell_\alpha} \sigma^{\mu\nu} q_\nu (K_2^L P_L + K_2^R P_R)] \ell_\alpha A_\mu + \text{h.c.} \quad (3.2)$$

$$\mathcal{L}_{4\ell} = \sum_{\substack{I=S,V,T \\ X,Y=L,R}} A_{XY}^I \bar{\ell}_\beta \Gamma_I P_X \ell_\alpha \bar{\ell}_\delta \Gamma_I P_Y \ell_\gamma + \text{h.c.} \quad (3.3)$$

$$\mathcal{L}_{2\ell 2d} = \sum_{\substack{I=S,V,T \\ X,Y=L,R}} B_{XY}^I \bar{\ell}_\beta \Gamma_I P_X \ell_\alpha \bar{d}_\gamma \Gamma_I P_Y d_\gamma + \text{h.c.} \quad (3.4)$$

$$\mathcal{L}_{2\ell 2u} = \mathcal{L}_{2\ell 2d}|_{d \rightarrow u, B \rightarrow C}. \quad (3.5)$$

Here e is the electric charge, q the 4-momenta of the photon, $P_{L,R} = \frac{1}{2}(1 \mp \gamma_5)$ are the usual chirality projectors and ℓ_α and d_α denote the lepton and d-quark flavors, respectively.

Furthermore, we have defined $\Gamma_S = 1$, $\Gamma_V = \gamma_\mu$ and $\Gamma_T = \sigma_{\mu\nu}$. We omit flavor indices in the form factors for the sake of simplicity. The underlying Feynman diagrams as well as the complete analytic results are given in appendices D–G.

Whenever possible, we have compared the explicit analytical formulae for the form factors with results already available in the literature. The supersymmetric contributions to boxes, Higgs penguins, photon penguins were found to perfectly agree with [78, 103], while the supersymmetric Z-penguins only differ from [78] via a constant term as pointed out in [77]. This constant term does not impact the result of [78] where a high-scale seesaw mechanism is considered but it can lead to non-physical results in low-scale seesaw models. We have also cross-checked our calculation of the non-SUSY boxes with [68], confirming their results. To the knowledge of the authors, this is the first calculation of the non-SUSY Higgs penguins in a two Higgs doublet Model, thus no comparison was possible.

3.2 $\ell_\alpha \rightarrow \ell_\beta \gamma$

In case of the radiative decay $\ell_\alpha \rightarrow \ell_\beta \gamma$, the corresponding decay width is given by [104]

$$\Gamma(\ell_\alpha \rightarrow \ell_\beta \gamma) = \frac{\alpha_{\text{em}} m_{\ell_\alpha}^5}{4} (|K_2^L|^2 + |K_2^R|^2), \quad (3.6)$$

where the dipole form factors $K_2^{L,R}$ are defined in eq. (3.2), α_{em} being the fine structure constant.

3.3 $\ell_\alpha^- \rightarrow \ell_\beta^- \ell_\beta^- \ell_\beta^+$

Next, we consider the $\ell_\alpha^-(p) \rightarrow \ell_\beta^-(p_1) \ell_\beta^-(p_2) \ell_\beta^+(p_3)$ 3-body decays. Using the operators in our LFV lagrangian, the decay width is given by

$$\begin{aligned} \Gamma(\ell_\alpha \rightarrow 3 \ell_\beta) = & \frac{m_{\ell_\alpha}^5}{512\pi^3} \left[e^4 \left(|K_2^L|^2 + |K_2^R|^2 \right) \left(\frac{16}{3} \log \frac{m_{\ell_\alpha}}{m_{\ell_\beta}} - \frac{22}{3} \right) \right. \\ & + \frac{1}{24} \left(|A_{LL}^S|^2 + |A_{RR}^S|^2 \right) + \frac{1}{12} \left(|A_{LR}^S|^2 + |A_{RL}^S|^2 \right) \\ & + \frac{2}{3} \left(|\hat{A}_{LL}^V|^2 + |\hat{A}_{RR}^V|^2 \right) + \frac{1}{3} \left(|\hat{A}_{LR}^V|^2 + |\hat{A}_{RL}^V|^2 \right) + 6 \left(|A_{LL}^T|^2 + |A_{RR}^T|^2 \right) \\ & + \frac{e^2}{3} (K_2^L A_{RL}^{S*} + K_2^R A_{LR}^{S*} + c.c.) - \frac{2e^2}{3} (K_2^L \hat{A}_{RL}^{V*} + K_2^R \hat{A}_{LR}^{V*} + c.c.) \\ & - \frac{4e^2}{3} (K_2^L \hat{A}_{RR}^{V*} + K_2^R \hat{A}_{LL}^{V*} + c.c.) \\ & \left. - \frac{1}{2} (A_{LL}^S A_{LL}^{T*} + A_{RR}^S A_{RR}^{T*} + c.c.) - \frac{1}{6} (A_{LR}^S \hat{A}_{LR}^{V*} + A_{RL}^S \hat{A}_{RL}^{V*} + c.c.) \right]. \end{aligned} \quad (3.7)$$

Here we have defined

$$\hat{A}_{XY}^V = A_{XY}^V + e^2 K_1^X \quad (X, Y = L, R). \quad (3.8)$$

The mass of the leptons in the final state has been neglected in this formula, with the exception of the numerical factors that multiply the $K_2^{L,R}$ contribution.⁵ Eq. (3.7) agrees

⁵We note that the correct form for the terms proportional to $K_2^{L,R}$ was first obtained in ref. [64].

This paper	[78]
$K_{1,2}^{L,R}$	$A_{1,2}^{L,R}$
A_{LL}^S	$e^2 \hat{B}_3^L$
A_{RR}^S	$e^2 \hat{B}_3^R$
A_{LL}^V	$e^2 \left(\frac{1}{2} B_1^L + F_{LL} \right)$
A_{RR}^V	$e^2 \left(\frac{1}{2} B_1^R + F_{RR} \right)$
$A_{LR}^V - \frac{1}{2} A_{RL}^S$	$e^2 \left(\hat{B}_2^L + F_{LR} \right)$
$A_{RL}^V - \frac{1}{2} A_{LR}^S$	$e^2 \left(\hat{B}_2^R + F_{RL} \right)$
A_{LL}^T	$e^2 B_4^L$
A_{RR}^T	$e^2 B_4^R$

Table 2. Relation between the form factors defined in this paper and the ones in [78]. Here $\hat{B}_2^{L,R} = B_2^{L,R} + B_{2,\text{Higgs}}^{L,R}$ and $\hat{B}_3^{L,R} = B_3^{L,R} + B_{3,\text{Higgs}}^{L,R}$, and $F_{XY} = \frac{F_X E^Y}{e^2 m_Z^2}$, with E^L and E^R the tree-level Z -boson couplings to a pair of charged leptons (see appendix A.2).

with the one in ref. [78], but includes in addition A_{LR}^S and A_{RL}^S . In [78], these contributions were absorbed in the corresponding vector form factors, A_{LR}^V and A_{RL}^V , by means of a Fierz transformation [105]. In contrast, A_{LR}^S and A_{RL}^S were explicitly added to the set of contributing form factors in [70]. The relation between our coefficients and the ones of [78] is given in table 2.

3.4 $\ell_\alpha^- \rightarrow \ell_\beta^- \ell_\gamma^- \ell_\gamma^+$

We consider the $\ell_\alpha^-(p) \rightarrow \ell_\beta^-(p_1) \ell_\gamma^-(p_2) \ell_\gamma^+(p_3)$ 3-body decays, with $\beta \neq \gamma$. The decay width is given by

$$\begin{aligned} \Gamma(\ell_\alpha^- \rightarrow \ell_\beta^- \ell_\gamma^- \ell_\gamma^+) &= \frac{m_{\ell_\alpha}^5}{512\pi^3} \left[e^4 \left(|K_2^L|^2 + |K_2^R|^2 \right) \left(\frac{16}{3} \log \frac{m_{\ell_\alpha}}{m_{\ell_\gamma}} - 8 \right) \right. \\ &\quad + \frac{1}{12} \left(|A_{LL}^S|^2 + |A_{RR}^S|^2 \right) + \frac{1}{12} \left(|A_{LR}^S|^2 + |A_{RL}^S|^2 \right) \\ &\quad + \frac{1}{3} \left(|\hat{A}_{LL}^V|^2 + |\hat{A}_{RR}^V|^2 \right) + \frac{1}{3} \left(|\hat{A}_{LR}^V|^2 + |\hat{A}_{RL}^V|^2 \right) + 4 \left(|A_{LL}^T|^2 + |A_{RR}^T|^2 \right) \\ &\quad \left. - \frac{2e^2}{3} \left(K_2^L \hat{A}_{RL}^{V*} + K_2^R \hat{A}_{LR}^{V*} + K_2^L \hat{A}_{RR}^{V*} + K_2^R \hat{A}_{LL}^{V*} + c.c. \right) \right]. \end{aligned} \quad (3.9)$$

Here we have used the same definition as in eq. (3.8). Furthermore, as for $\ell_\alpha \rightarrow 3\ell_\beta$, the mass of the leptons in the final state has been neglected in the decay width formula, with the exception of the dipole terms $K_2^{L,R}$.

Finally, we note that eqs. (3.7) and (3.9) are in perfect agreement with the expressions given in ref. [70].

3.5 $\ell_\alpha^- \rightarrow \ell_\beta^+ \ell_\gamma^- \ell_\gamma^-$

Finally, we consider the $\ell_\alpha^-(p) \rightarrow \ell_\beta^+(p_1)\ell_\gamma^-(p_2)\ell_\gamma^-(p_3)$ 3-body decays, with $\beta \neq \gamma$. The decay width is given by

$$\begin{aligned} \Gamma\left(\ell_\alpha^- \rightarrow \ell_\beta^+ \ell_\gamma^- \ell_\gamma^-\right) = & \frac{m_{\ell_\alpha}^5}{512\pi^3} \left[\frac{1}{24} \left(|A_{LL}^S|^2 + |A_{RR}^S|^2 \right) + \frac{1}{12} \left(|A_{LR}^S|^2 + |A_{RL}^S|^2 \right) \right. \\ & + \frac{2}{3} \left(|\hat{A}_{LL}^V|^2 + |\hat{A}_{RR}^V|^2 \right) + \frac{1}{3} \left(|\hat{A}_{LR}^V|^2 + |\hat{A}_{RL}^V|^2 \right) + 6 \left(|A_{LL}^T|^2 + |A_{RR}^T|^2 \right) \\ & \left. - \frac{1}{2} (A_{LL}^S A_{LL}^{T*} + A_{RR}^S A_{RR}^{T*} + c.c.) - \frac{1}{6} (A_{LR}^S \hat{A}_{LR}^{V*} + A_{RL}^S \hat{A}_{RL}^{V*} + c.c.) \right]. \end{aligned} \quad (3.10)$$

The same definitions and conventions as in the previous two observables have been used. Notice that this process does not receive contributions from penguin diagrams, but only from boxes.

3.6 Coherent $\mu - e$ conversion in nuclei

We now turn to the discussion of $\mu - e$ conversion in nuclei, which will follow the conventions and approximations described in refs. [103, 106] (see also [107–109] for detailed works regarding the effective lagrangian at the nucleon level, [68, 110] for a calculation including the effects of the atomic electric field and [111] for recent improvements on the hadronic uncertainties). The conversion rate, relative to the the muon capture rate, can be expressed as

$$\begin{aligned} \text{CR}(\mu - e, \text{Nucleus}) = & \frac{p_e E_e m_\mu^3 G_F^2 \alpha_{\text{em}}^3 Z_{\text{eff}}^4 F_p^2}{8\pi^2 Z} \\ & \times \left\{ \left| (Z+N) \left(g_{LV}^{(0)} + g_{LS}^{(0)} \right) + (Z-N) \left(g_{LV}^{(1)} + g_{LS}^{(1)} \right) \right|^2 + \right. \\ & \left. \left| (Z+N) \left(g_{RV}^{(0)} + g_{RS}^{(0)} \right) + (Z-N) \left(g_{RV}^{(1)} + g_{RS}^{(1)} \right) \right|^2 \right\} \frac{1}{\Gamma_{\text{capt}}}. \end{aligned} \quad (3.11)$$

Z and N are the number of protons and neutrons in the nucleus and Z_{eff} is the effective atomic charge [112]. Similarly, G_F is the Fermi constant, F_p is the nuclear matrix element and Γ_{capt} represents the total muon capture rate. p_e and E_e ($\simeq m_\mu$ in our numerical evaluation) are the momentum and energy of the electron and m_μ is the muon mass. In the above, $g_{XK}^{(0)}$ and $g_{XK}^{(1)}$ (with $X = L, R$ and $K = S, V$) can be written in terms of effective couplings at the quark level as

$$\begin{aligned} g_{XK}^{(0)} &= \frac{1}{2} \sum_{q=u,d,s} \left(g_{XK(q)} G_K^{(q,p)} + g_{XK(q)} G_K^{(q,n)} \right), \\ g_{XK}^{(1)} &= \frac{1}{2} \sum_{q=u,d,s} \left(g_{XK(q)} G_K^{(q,p)} - g_{XK(q)} G_K^{(q,n)} \right). \end{aligned} \quad (3.12)$$

For coherent $\mu - e$ conversion in nuclei, only scalar (S) and vector (V) couplings contribute [106]. Furthermore, sizable contributions are expected only from the u, d, s quark flavors. The numerical values of the relevant G_K factors are [106, 113]

$$G_V^{(u,p)} = G_V^{(d,p)} = 2; \quad G_V^{(d,p)} = G_V^{(u,n)} = 1;$$

α_{em}^{-1}	127.92783	G_μ	$1.11639 \cdot 10^{-5} \text{ GeV}^{-2}$
α_S	0.11720	M_Z	91.18760 GeV
$m_b(m_b)$	4.2 GeV	m_t	172.9 GeV
m_τ	1.777 GeV		

Table 3. Input values for the SM parameters taken at M_Z unless otherwise specified.

$$\begin{aligned} G_S^{(u,p)} &= G_S^{(d,n)} = 5.1; & G_S^{(d,p)} &= G_S^{(u,n)} = 4.3; \\ G_S^{(s,p)} &= G_S^{(s,n)} = 2.5. \end{aligned} \quad (3.13)$$

Finally, the $g_{XK(q)}$ coefficients can be written in terms of the form factors in eqs. (3.2), (3.4) and (3.5) as

$$g_{LV(q)} = \frac{\sqrt{2}}{G_F} \left[e^2 Q_q (K_1^L - K_2^R) - \frac{1}{2} (C_{\ell\ell qq}^{VLL} + C_{\ell\ell qq}^{VLR}) \right] \quad (3.14)$$

$$g_{RV(q)} = g_{LV(q)}|_{L \rightarrow R} \quad (3.15)$$

$$g_{LS(q)} = -\frac{\sqrt{2}}{G_F} \frac{1}{2} (C_{\ell\ell qq}^{SLL} + C_{\ell\ell qq}^{SLR}) \quad (3.16)$$

$$g_{RS(q)} = g_{LS(q)}|_{L \rightarrow R}. \quad (3.17)$$

Here Q_q is the quark electric charge ($Q_d = -1/3$, $Q_u = 2/3$) and $C_{\ell\ell qq}^{IXK} = B_{XY}^K$ (C_{XY}^K) for d-quarks (u-quarks), with $X = L, R$ and $K = S, V$.

4 Results

4.1 Numerical setup

For the numerical examples we have implemented the model in the Mathematica package **SARAH** [80–84], which creates the required modules for **SPheno** [85, 86] to calculate the masses and mixing matrices including the complete 1-loop corrections. In the Higgs sector we include in addition the known 2-loop corrections to the Higgs mass from the MSSM [114–119]. However, this does not include 2-loop corrections stemming from the extended neutrino and sneutrino sectors, where we can have sizable Yukawa couplings. Moreover, **SARAH** calculates also the full 2-loop RGEs including the entire flavor structure for the model, which we have summarized in appendix B. This will be of great importance in our numerical studies, as we use CMSSM-like boundary conditions, see below for their definition. In the flavor observables we include all possible contributions. These are calculated using the **FlavorKit** interface [79]. In the context of this project we have extended the lists of observables implemented in **FlavorKit** by $\ell_\alpha^- \rightarrow \ell_\beta^- \ell_\beta^+ \ell_\gamma^-$ and $\ell_\alpha^- \rightarrow \ell_\beta^- \ell_\beta^+ \ell_\gamma^+$.

The numerical evaluation of each parameter point is performed as follows: the Y_ν Yukawa couplings are calculated using a modified Casas-Ibarra parameterization [120], properly adapted for the inverse seesaw [121, 122] (and fixing $M_R = 2$ TeV, $\mu_X = 10^{-5}$ GeV and the lightest neutrino mass $m_{\nu_1} = 10^{-4}$ eV):

$$Y_\nu = \frac{\sqrt{2}}{v_u} V^\dagger D_{\sqrt{X}} R D_{\sqrt{m_\nu}} U_{\text{PMNS}}^\dagger. \quad (4.1)$$

Here $D_{\sqrt{m_\nu}} = \text{diag}(\sqrt{m_{\nu_i}})$, $D_{\sqrt{X}} = \text{diag}(\sqrt{\hat{X}_i})$, \hat{X}_i being the eigenvalues of $X = M_R \mu_X^{-1} M_R^T$, and V is the matrix that diagonalizes X as $V X V^T = \hat{X}$. Furthermore, we parameterize the complex orthogonal R matrix as

$$R = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos \theta_{23}^R & \sin \theta_{23}^R \\ 0 & -\sin \theta_{23}^R & \cos \theta_{23}^R \end{pmatrix} \begin{pmatrix} \cos \theta_{13}^R & 0 & \sin \theta_{13}^R \\ 0 & 1 & 0 \\ -\sin \theta_{13}^R & 0 & \cos \theta_{13}^R \end{pmatrix} \begin{pmatrix} \cos \theta_{12}^R & \sin \theta_{12}^R & 0 \\ -\sin \theta_{12}^R & \cos \theta_{12}^R & 0 \\ 0 & 0 & 1 \end{pmatrix}. \quad (4.2)$$

Below we will set R to the unit matrix except when stated otherwise. We use the best-fit values for the neutrino oscillation parameters as given in [123]:

$$\Delta m_{21}^2 = 7.60 \cdot 10^{-5} \text{ eV}^2, \quad \Delta m_{31}^2 = 2.48 \cdot 10^{-3} \text{ eV}^2, \\ \sin^2 \theta_{12} = 0.323, \quad \sin^2 \theta_{23} = 0.467, \quad \sin^2 \theta_{13} = 0.0234. \quad (4.3)$$

which are close to the ones obtained in [3–5]. We make use in our scans of the values

$$Y_\nu = f \cdot 10^{-2} \cdot \begin{pmatrix} 0.0956 & -0.0589 & 0.0348 \\ 0.616 & 0.594 & -0.687 \\ 0.404 & 1.78 & 1.91 \end{pmatrix} \quad (4.4)$$

fixed with $f = 1$ even if we vary M_R . This is because one can always adjust μ_X to fulfill neutrino oscillation data without affecting any of our observables.

SPheno derives the SM gauge and Yukawa couplings at M_Z where we take the masses and couplings given in table 3 as input. 2-loop RGEs for the dimensionless parameters are then used to evaluate these couplings at M_{GUT} , defined by the requirement $g_1 = g_2$, where g_1 and g_2 are the couplings for the $U(1)_Y$ and $SU(2)_L$ gauge groups, respectively. At M_{GUT} the CMSSM boundary conditions are applied

$$\begin{aligned} m_{\nu C}^2 &= m_X^2 = m_l^2 = m_e^2 = m_q^2 = m_d^2 = m_u^2 \equiv m_0^2 \mathbf{1} \\ m_{H_d}^2 &= m_{H_u}^2 \equiv m_0^2 \\ M_1 &= M_2 = M_3 \equiv M_{1/2} \\ T_i &\equiv A_0 Y_i \quad i = e, d, u, \nu \end{aligned}$$

The mixed soft-term $m_{X\nu C}^2$ is set to zero at the GUT scale and is not generated via RGE effects. Moreover, the phase of μ , which is an RGE invariant, is given as input. The ratio of the Higgs vevs, $\tan \beta = \frac{v_u}{v_d}$, completes the list of input parameters. Then 2-loop RGEs are used to evolve these parameters to $Q_{\text{EWSB}} = \sqrt{\tilde{t}_1 \tilde{t}_2}$. The numerical values for superpotential terms M_R and μ_X , as well as for their corresponding soft terms B_{M_R} and B_{μ_X} , are used as input at the SUSY scale. B_μ and $|\mu|$ are obtained as usual from the minimization conditions of the vacuum.⁶ At M_{SUSY} the 1-loop corrected masses are calculated before the RGEs

⁶In principle one could require that all B -parameters are proportional to each other, e.g. $B_\mu : B_{M_R} : B_{\mu_X} = \mu : M_R : \mu_X$. However, as their actual value does not have any significant impact as long as this ratio is fulfilled up to a factor 2-3 we fix for simplicity B_{M_R} and B_{μ_X} .




Figure 1. $\text{BR}(\mu \rightarrow e\gamma)$ as a function of M_{SUSY} and M_R . The other parameters are given in the text. The gray area roughly corresponds to the parameter space excluded by the LHC experiments.

m_0	1 TeV	$M_{1/2}$	1 TeV
A_0	-1.5 TeV	M_R	2 TeV
B_{μ_X}	$100 \mu_X$	B_{M_R}	$100 M_R$
$\tan \beta$	10	$\text{sign}(\mu)$	+

Table 4. Standard values for the various parameters. M_R and μ_X are taken proportional to the unit matrix.

run down to M_Z to re-calculate gauge and Yukawa couplings using the new SUSY corrections. These steps are iterated until the mass spectrum has converged with a numerical precision of 10^{-4} . Afterwards, **SPheno** runs the RGEs to $Q = 160 \text{ GeV}$ for the calculation of the operators which contribute to quark flavor violating observables and to $Q = M_Z$ for the calculation of the operators needed for lepton flavor violating observables. These operators are then combined to compute the different observables using $\alpha(0)$, which includes to a large extent the effects from running the operators between M_Z and the energy scale where the LFV processes take place (usually given by the mass of the decaying particle).

4.2 Numerical results

We will use the parameter values given in table 4 as starting point for our numerical computations unless stated otherwise. A variation of the soft SUSY parameters is denoted

by a variation of M_{SUSY} , which actually implies a variation of three parameters at the same time $M_{\text{SUSY}} = m_0 = M_{1/2} = -A_0$. For completeness, we note that fixing the ratio m_0/A_0 usually gives a Higgs boson mass, m_h , that does not agree with the ATLAS and CMS measurements. Nevertheless, we emphasize that (1) our results depend only weakly on the value of A_0 , and (2) contributions mediated by h itself are subdominant. Therefore, the actual Higgs boson mass is of little importance for our investigations here.

We start with the discussion of μ decays as the bounds are strongest in this case. In figure 1 we show the dependence of $\text{BR}(\mu \rightarrow e\gamma)$ on M_R and M_{SUSY} as well as the individual dependence of the SUSY and non-SUSY contributions. The latter consist of ν - W^\pm and the ν - H^\pm contributions. There are two particular features: (i) if $M_R = M_{\text{SUSY}}$ the SUSY contributions are more important than the non-SUSY ones and the relative importance of the SUSY contributions increases with the scale. The reason for the latter is that the mixing between light and heavy neutrinos decreases like $\sim m_D/M_R$, whereas the mixing in the sneutrino sector decreases only logarithmically with the scale. (ii) The non-SUSY contributions can flip its sign. This is due to a sign-difference between the ν - H^\pm and the ν - W^\pm contributions to the coefficients $K_2^{L,R}$. This is in contrast to the analogous decay in the quark sector, $b \rightarrow s\gamma$, where the W^\pm - and H^\pm -contributions have always the same sign. The reason for this difference can be found in eqs. (D.12)–(D.18), presented in appendix D, where the light neutrino masses appear instead of the mass of the heavy t -quark. We have checked explicitly, both numerically and analytically, that we recover the $b \rightarrow s\gamma$ result if we replace the corresponding masses and Yukawa couplings. Finally, we stress that the scalar masses are functions of M_{SUSY} , which explains why also the non-SUSY contribution actually depends on the SUSY scale. With our specific structure of the Y_ν matrices we find that M_R has to be larger than M_{SUSY} for the sign flip to occur, which is also the reason why we do not observe it in case of $M_R = M_{\text{SUSY}}$. The grey area corresponds to the part of the parameter space which is excluded in the CMSSM by the most recent ATLAS results [102]. However, we want to stress that even though the squark and gluino masses are essentially the same in our model as in the CMSSM, the cascade decays can be quite different due to (i) the enlarged sneutrino sector with additional light states and (ii) the different slepton masses. Thus, this is a rather conservative bound.

In figure 2 we display our results for the branching ratio $\text{BR}(\mu \rightarrow 3e)$ as well as the various contributions to this decay. Here we find that for the case $M_R = M_{\text{SUSY}}$ the non-SUSY boxes dominate. This fact was first noted in [67] and later confirmed by [68–70]. Note that this does not depend on the overall strength of the Y_ν couplings, which we rescale as $Y_\nu \rightarrow f Y_\nu$. This can be seen from the lower right plot: all contributions scale in the same way. However, the situation can change in principle if one allows for additional flavor violation in the soft SUSY breaking parameters. Note that the sign-flip induced by the H^\pm contributions is not as pronounced as in the case of $\mu \rightarrow e\gamma$, where it led to a change of the overall sign, as the different contributions to the off-shell photon appear with different weights. However, it is the reason for the observed kink in the non-SUSY γ -penguin. We also observe that we have negative interference between non-SUSY Z -penguins and the corresponding box contributions. In particular, for larger values of M_R this can reduce $\text{BR}(\mu \rightarrow 3e)$ by up to an order of magnitude. Since this is precisely the region which will




Figure 2. $\text{BR}(\mu \rightarrow 3e)$ as a function of M_{SUSY} , M_R and an overall scaling parameter f for Y_ν . The other parameters are given in the text. The gray area roughly corresponds to the parameter space excluded by the LHC experiments.

be probed by future experiments, the possible appearance of these cancellations has to be taken into account in order to interpret the experimental results properly.

Similar features appear in case of $\mu - e$ conversion in nuclei, as exemplified for the case of an aluminium (Al) nucleus in figure 3. The main difference is that there is a large part of parameter space where a pronounced negative interference between the non-SUSY Z -penguin and the corresponding box contributions can occur. Note that with the expected sensitivity of 10^{-18} one can probe Y_ν couplings down to a few $\times 10^{-6}$ for $M_R = M_{\text{SUSY}} = 1 \text{ TeV}$ or, equivalently, to a mass scale of about 5 TeV in case of Y_ν as given in eq. (4.4). As we found for the 3-body decays, for higher mass scales the non-SUSY Z -penguins can be as important as the corresponding box-diagrams. The overall features are essentially element independent as can be seen in figure 4 where we show all three observables discussed so far together and include also $\mu - e$ conversion in titanium (Ti). In case $M_R \simeq M_{\text{SUSY}}$, we find that $\mu - e$ conversion in nuclei is the most stringent LFV observable in our model.

Turning now to the LFV τ decays, we show in figure 5 several branching ratios for the scenario defined above. Unfortunately, they are too small to be observed in the near future. Below we will show alternative scenarios (in which the R matrix is not assumed to be the unit matrix) where this is not the case. Nevertheless, they show an interesting feature which is quite generic in this model: $\text{BR}(\tau \rightarrow \mu e^+ e^-) \simeq \text{BR}(\tau \rightarrow 3\mu)$ and $\text{BR}(\tau \rightarrow e \mu^+ \mu^-) \simeq$




Figure 3. $\mu - e$ conversion on Al as a function of M_{SUSY} , M_R and the scaling parameter f for Y_ν . The gray area roughly corresponds to the parameter space excluded by the LHC experiments.

$\text{BR}(\tau \rightarrow 3e)$. Particularly interesting is that these branching ratios are sensitive to the relative size of the non-SUSY contributions compared to the SUSY ones. We also stress that the various contributions contribute similarly as in case of $\mu \rightarrow 3e$. For completeness we note that $\text{BR}(\tau \rightarrow e\mu^+e^-)$ and $\text{BR}(\tau \rightarrow \mu e^+\mu^-)$ are strongly suppressed, at least a factor of 10^{-6} with respect to the other 3-body decays, as they require at least one additional flavor violating vertex in the dominant contributions.

It is worth stressing that the fact that the μ observables are more constraining than the τ decays is correct in large parts of the parameter space. However, there is also a substantial part where the opposite is true, as exemplified in figure 6 where we tune the parameters such that both, μ - and τ -observables can be discovered in the next generation of experiments. For this we have adjusted the diagonal entries of μ_X as well as θ_{23}^R and calculated Y_ν using eq. (4.1). Clearly, this part of the parameter space requires quite some hierarchy in μ_X to explain neutrino data correctly. Note that even in this part of parameter space the ratios $\text{BR}(\tau \rightarrow \mu e^+e^-) \simeq \text{BR}(\tau \rightarrow 3\mu)$ and $\text{BR}(\tau \rightarrow e\mu^+\mu^-) \simeq \text{BR}(\tau \rightarrow 3e)$ show the same dependence on the ratio M_R/M_{SUSY} as in the previous case.

The impact of the R matrix and the hierarchy in the μ_X entries is further illustrated for the decays $\ell_\alpha \rightarrow \ell_\beta \gamma$ in figure 7. Again, we have calculated Y_ν via eq. (4.1), such that the results from neutrino oscillation experiments are explained correctly. One finds that, depending on the region in the parameter space, either the μ decay or the τ decays




Figure 4. $\text{BR}(\mu \rightarrow e\gamma)$, $\text{BR}(\mu \rightarrow 3e)$, $\mu-e$ conversion in Ti and Al as a function of M_R and M_{SUSY} . The gray area roughly corresponds to the parameter space excluded by the LHC experiments.

are more important. As in case of the 3-body decays, one finds fine-tuned combinations of the parameters where all decays can be observed in future experiments. Note that for fixed θ_{23}^R the branching ratios scale like f_R^2/f_X where f_R and f_X denote an overall scaling of M_R and μ_X , respectively. Moreover, the branching ratios scale like $\tan^2 \beta$ if the SUSY contributions dominate. In case the non-SUSY contributions dominate we find only a slight $\tan \beta$ dependence for very large $\tan \beta$ values.

Finally, let us comment on the Higgs penguin contributions to the different LFV observables. In all our numerical scans they have been found to be completely negligible and that is why we have decided not to include them in our figures. In principle, one could look for sizable Higgs penguin contributions by going to regions in parameter space with large $\tan \beta$ and low pseudoscalar masses [93, 124]. This, however, would require dedicated parameter scans in order to overcome the constraints from flavor data, as these regions are already in strong tension after the LHCb measurement of the $B_s \rightarrow \mu^+ \mu^-$ branching ratio [125]. For this reason, we have not pursued this goal any further. Nevertheless, we have checked that the Higgs penguins contributions to $\ell_\alpha \rightarrow \ell_\beta \ell_\gamma \ell_\delta$ and $\mu-e$ conversion in nuclei have the expected decoupling behavior for large M_R and/or M_{SUSY} scales.




Figure 5. Branching ratios for τ decays as a function of M_R and M_{SUSY} . In the upper two plots the lines correspond to $\text{BR}(\mu \rightarrow 3e)$ (black solid), $\text{BR}(\tau \rightarrow 3e)$ (blue solid), $\text{BR}(\tau \rightarrow 3\mu)$ (red solid), $\text{BR}(\tau^- \rightarrow e^- \mu^+ \mu^-)$ (blue dashed) and $\text{BR}(\tau^- \rightarrow \mu^- e^+ e^-)$ (red dashed). The gray area roughly corresponds to the parameter space excluded by the LHC experiments.




Figure 6. μ - and τ -observables as a function of μ_X . The underlying parameters are given in the plots. The lines correspond to $\text{BR}(\tau \rightarrow \mu\gamma)$ (full red), $\text{BR}(\tau \rightarrow 3\mu)$ (dashed red), $\text{BR}(\tau^- \rightarrow \mu^- e^+ e^-)$ (dotted red), $\text{BR}(\mu \rightarrow e\gamma)$ (full black) and $\text{BR}(\mu \rightarrow 3e)$ (dashed black). The light gray, red, yellow and blue bands show the expected future reach of the dedicated experiments to $\tau \rightarrow \mu\gamma$, $\tau \rightarrow 3\mu$, $\mu \rightarrow e\gamma$ and $\mu \rightarrow 3e$ as given in table 1.




Figure 7. Dependence of $\ell_\alpha \rightarrow \ell_\beta \gamma$ on μ_X^{33} and θ_{23}^R . The lines correspond to $\text{BR}(\mu \rightarrow e\gamma)$ (black), $\text{BR}(\tau \rightarrow e\gamma)$ (blue dashed) and $\text{BR}(\tau \rightarrow \mu\gamma)$ (red dotted).

5 Conclusions

This paper represents the first complete computation of selected LFV observables in scenarios with light right-handed neutrinos. These include the radiative decays $\ell_\alpha \rightarrow \ell_\beta \gamma$, the 3-body decays $\ell_\alpha \rightarrow \ell_\beta \ell_\gamma \ell_\delta$ (in several variants) and neutrinoless $\mu - e$ conversion in nuclei. Our results are valid in the inverse seesaw and should also hold in low-scale type-I seesaw models with nearly conserved lepton number, the inverse seesaw being a specific realization of these models. Compared to previous studies, we have also included Higgs-penguins and considered non-supersymmetric as well as supersymmetric contributions to the corresponding LFV amplitudes simultaneously.

For the numerical examples we took a CMSSM inspired scenario where we also considered the limiting cases with either $M_R \gg M_{\text{SUSY}}$ and $M_{\text{SUSY}} \gg M_R$. Our main conclusions can be summarized as follows:

- The SUSY contributions dominate the induced photon penguins if both, M_R and M_{SUSY} , are about the same size. For $M_R \lesssim M_{\text{SUSY}}/2$ the non-SUSY contributions start to dominate the radiative decays $\ell_\alpha \rightarrow \ell_\beta \gamma$.
- For low M_R scales the LFV phenomenology is dominated by non-SUSY contributions. This holds in particular for the 3-body decays and $\mu - e$ conversion in nuclei. These are mainly given by boxes and Z -penguin diagrams containing right-handed

neutrinos in the loop. In contrast to the usual high-scale seesaw models, in which their contributions to LFV processes are tiny, the right-handed neutrinos can play a major role in low-scale seesaw scenarios. In what concerns the non-SUSY box contributions, our results confirm previous claims in the literature [67–70]. Furthermore, we have highlighted the relevance of the non-SUSY Z -penguins, previously regarded as subdominant in most studies.⁷ They are particularly relevant for larger values of M_R , where we often find a negative interference between the Z -penguins and the box contributions. This will be particularly important when the next generation of experiments start to probe this mass region.

- The proper decoupling of the different contributions has been checked explicitly, e.g. we have checked that the SUSY-contributions, the ν^C - X and the Higgs contributions decouple independently as expected.
- Currently, the radiative decay $\mu \rightarrow e\gamma$ is the most constraining LFV process. However, due to the promising experimental prospects in the near future, the situation will change. If the coming experiments perform as planned, $\mu \rightarrow 3e$ will be the most relevant LFV process in the mid term, whereas neutrinoless $\mu - e$ conversion in nuclei will set the strongest constraints in the long term.
- Ratios of τ LFV branching ratios can provide additional information about the dominant contributions. In particular, when the non-SUSY contributions dominate, one finds $\text{BR}(\tau^- \rightarrow \mu^- e^+ e^-)/\text{BR}(\tau \rightarrow 3\mu) \simeq \text{BR}(\tau^- \rightarrow e^- \mu^+ \mu^-)/\text{BR}(\tau \rightarrow 3e) \simeq 0.5 - 0.8$, whereas for a SUSY dominated scenario $\text{BR}(\tau^- \rightarrow \mu^- e^+ e^-)/\text{BR}(\tau \rightarrow 3\mu) \gg \text{BR}(\tau^- \rightarrow e^- \mu^+ \mu^-)/\text{BR}(\tau \rightarrow 3e)$. This can in turn be used to get a hint on the hierarchy between the seesaw and SUSY scales.

Acknowledgments

We thank Martin Hirsch and Ernesto Arganda for fruitful discussions. A.V. acknowledges partial support from the EXPL/FIS-NUC/0460/2013 project financed by the Portuguese FCT. M.E.K. and W.P. have been supported by the DFG, project no. PO-1337/3-1 and the DFG research training group GRK 1147. FS is supported by the BMBF PT DESY Verbundprojekt 05H2013-THEORIE ‘Vergleich von LHC-Daten mit supersymmetrischen Modellen’. C.W. receives financial support from the Spanish CICYT through the project FPA2012-31880 and a partial support from the European Union FP7 ITN INVISIBLES (Marie Curie Actions, PITN-GA-2011-289442) and the Spanish MINECO’s “Centro de Excelencia Severo Ochoa” Programme under grant SEV-2012-0249. A. A acknowledges support from the European Union FP7 ITN INVISIBLES (Marie Curie Actions, PITN-GA-2011-289442).

⁷Non-SUSY Z -penguins were also included in ref. [70], where their potentially large contributions were also shown.

A Masses and vertices

We give first our conventions for the mass matrices as well as for the corresponding rotation matrices. These matrices are then used to express in appendix A.2 all the vertices needed to calculate the LFV observables.

A.1 Mass matrices

- **Mass matrix for Neutrinos**, Basis: (ν_L, ν_R^C, X)

$$m_\nu = \begin{pmatrix} 0 & \frac{1}{\sqrt{2}}v_u Y_\nu^T & 0 \\ \frac{1}{\sqrt{2}}v_u Y_\nu & 0 & M_R \\ 0 & M_R^T & \mu_X \end{pmatrix} \quad (\text{A.1})$$

This matrix is diagonalized by U^V :

$$U^{V,*} m_\nu U^{V,\dagger} = m_\nu^{dia} \quad (\text{A.2})$$

- **Mass matrix for CP-odd Sneutrinos**, Basis: $(\sigma_L, \sigma_R, \sigma_X)$

$$m_{\nu^i}^2 = \begin{pmatrix} m_{\sigma_L \sigma_L} & m_{\sigma_L \sigma_R}^T & \frac{1}{\sqrt{2}}v_u \Re(Y_\nu^T M_R^*) \\ m_{\sigma_L \sigma_R} & m_{\sigma_R \sigma_R} & \Re(B_{M_R} - M_R \mu_X^*) \\ \frac{1}{\sqrt{2}}v_u \Re(M_R^T Y_\nu^*) & \Re(B_{M_R}^T - \mu_X M_R^\dagger) & m_{\sigma_X \sigma_X} \end{pmatrix} \quad (\text{A.3})$$

$$m_{\sigma_L \sigma_L} = \frac{1}{2}v_u^2 \Re(Y_\nu^T Y_\nu^*) + \Re(m_l^2) + \frac{1}{8}(g_1^2 + g_2^2)(v_d^2 - v_u^2) \mathbf{1} \quad (\text{A.4})$$

$$m_{\sigma_L \sigma_R} = -\frac{1}{\sqrt{2}}(v_d \Re(\mu Y_\nu^*) - v_u \Re(T_\nu)) \quad (\text{A.5})$$

$$m_{\sigma_R \sigma_R} = \Re(m_\nu^2) + \Re(M_R M_R^\dagger) + \frac{1}{2}v_u^2 \Re(Y_\nu Y_\nu^\dagger) \quad (\text{A.6})$$

$$m_{\sigma_X \sigma_X} = \Re(M_R^T M_R^*) + \Re(m_X^2) - \Re(B_{\mu_X}) + \Re(\mu_X \mu_X^*) \quad (\text{A.7})$$

This matrix is diagonalized by Z^i :

$$Z^i m_{\nu^i}^2 Z^{i,\dagger} = m_{\nu^i}^{2,dia} \quad (\text{A.8})$$

- **Mass matrix for CP-even Sneutrinos**, Basis: (ϕ_L, ϕ_R, ϕ_X)

$$m_{\nu^R}^2 = \begin{pmatrix} m_{\phi_L \phi_L} & m_{\phi_L \phi_R}^T & \frac{1}{\sqrt{2}}v_u \Re(Y_\nu^T M_R^*) \\ m_{\phi_L \phi_R} & m_{\phi_R \phi_R} & \Re(B_{M_R} + M_R \mu_X^*) \\ \frac{1}{\sqrt{2}}v_u \Re(M_R^T Y_\nu^*) & \Re(B_{M_R}^T + \mu_X M_R^\dagger) & m_{\phi_X \phi_X} \end{pmatrix} \quad (\text{A.9})$$

$$m_{\phi_L \phi_L} = \frac{1}{2}v_u^2 \Re(Y_\nu^T Y_\nu^*) + \Re(m_l^2) + \frac{1}{8}(g_1^2 + g_2^2)(v_d^2 - v_u^2) \mathbf{1} \quad (\text{A.10})$$

$$m_{\phi_L \phi_R} = -\frac{1}{\sqrt{2}} \left(v_d \Re(\mu Y_\nu^*) - v_u \Re(T_\nu) \right) \quad (\text{A.11})$$

$$m_{\phi_R \phi_R} = \Re(m_\nu^2) + \Re(M_R M_R^\dagger) + \frac{1}{2} v_u^2 \Re(Y_\nu Y_\nu^\dagger) \quad (\text{A.12})$$

$$m_{\phi_X \phi_X} = \Re(M_R^T M_R^*) + \Re(m_X^2) + \Re(B_{\mu_X}) + \Re(\mu_X \mu_X^*) \quad (\text{A.13})$$

This matrix is diagonalized by Z^R :

$$Z^R m_{\nu^R}^2 Z^{R,\dagger} = m_{\nu^R}^{2,dia} \quad (\text{A.14})$$

- **Mass matrix for Down-Squarks**, Basis: $(\tilde{d}_{L,\alpha_1}, \tilde{d}_{R,\alpha_2})$

$$m_{\tilde{d}}^2 = \begin{pmatrix} m_{\tilde{d}_L \tilde{d}_L^*} & \frac{1}{\sqrt{2}} (v_d T_d^\dagger - v_u \mu Y_d^\dagger) \delta_{\alpha_1 \beta_2} \\ \frac{1}{\sqrt{2}} \delta_{\alpha_2 \beta_1} (v_d T_d - v_u Y_d \mu^*) & m_{\tilde{d}_R \tilde{d}_R^*} \end{pmatrix} \quad (\text{A.15})$$

$$m_{\tilde{d}_L \tilde{d}_L^*} = -\frac{1}{24} (3g_2^2 + g_1^2) (v_d^2 - v_u^2) \delta_{\alpha_1 \beta_1} \mathbf{1} + \frac{1}{2} \delta_{\alpha_1 \beta_1} (2m_q^2 + v_d^2 Y_d^\dagger Y_d) \quad (\text{A.16})$$

$$m_{\tilde{d}_R \tilde{d}_R^*} = \frac{1}{12} g_1^2 (v_u^2 - v_d^2) \delta_{\alpha_2 \beta_2} \mathbf{1} + \frac{1}{2} \delta_{\alpha_2 \beta_2} (2m_d^2 + v_d^2 Y_d Y_d^\dagger) \quad (\text{A.17})$$

This matrix is diagonalized by Z^D :

$$Z^D m_{\tilde{d}}^2 Z^{D,\dagger} = m_{\tilde{d}}^{2,dia} \quad (\text{A.18})$$

- **Mass matrix for Up-Squarks**, Basis: $(\tilde{u}_{L,\alpha_1}, \tilde{u}_{R,\alpha_2})$

$$m_{\tilde{u}}^2 = \begin{pmatrix} m_{\tilde{u}_L \tilde{u}_L^*} & \frac{1}{\sqrt{2}} (-v_d \mu Y_u^\dagger + v_u T_u^\dagger) \delta_{\alpha_1 \beta_2} \\ \frac{1}{\sqrt{2}} \delta_{\alpha_2 \beta_1} (-v_d Y_u \mu^* + v_u T_u) & m_{\tilde{u}_R \tilde{u}_R^*} \end{pmatrix} \quad (\text{A.19})$$

$$m_{\tilde{u}_L \tilde{u}_L^*} = -\frac{1}{24} (-3g_2^2 + g_1^2) (v_d^2 - v_u^2) \delta_{\alpha_1 \beta_1} \mathbf{1} + \frac{1}{2} \delta_{\alpha_1 \beta_1} (2m_q^2 + v_u^2 Y_u^\dagger Y_u) \quad (\text{A.20})$$

$$m_{\tilde{u}_R \tilde{u}_R^*} = \frac{1}{2} \delta_{\alpha_2 \beta_2} (2m_u^2 + v_u^2 Y_u Y_u^\dagger) + \frac{1}{6} g_1^2 (v_d^2 - v_u^2) \delta_{\alpha_2 \beta_2} \mathbf{1} \quad (\text{A.21})$$

This matrix is diagonalized by Z^U :

$$Z^U m_{\tilde{u}}^2 Z^{U,\dagger} = m_{\tilde{u}}^{2,dia} \quad (\text{A.22})$$

- **Mass matrix for Sleptons**, Basis: $(\tilde{e}_L, \tilde{e}_R)$

$$m_{\tilde{e}}^2 = \begin{pmatrix} m_{\tilde{e}_L \tilde{e}_L^*} & \frac{1}{\sqrt{2}} (v_d T_e^\dagger - v_u \mu Y_e^\dagger) \\ \frac{1}{\sqrt{2}} (v_d T_e - v_u Y_e \mu^*) & m_{\tilde{e}_R \tilde{e}_R^*} \end{pmatrix} \quad (\text{A.23})$$

$$m_{\tilde{e}_L \tilde{e}_L^*} = \frac{1}{2} v_d^2 Y_e^\dagger Y_e + \frac{1}{8} (-g_2^2 + g_1^2) (v_d^2 - v_u^2) \mathbf{1} + m_l^2 \quad (\text{A.24})$$

$$m_{\tilde{e}_R \tilde{e}_R^*} = \frac{1}{2} v_d^2 Y_e Y_e^\dagger + \frac{1}{4} g_1^2 (v_u^2 - v_d^2) \mathbf{1} + m_e^2 \quad (\text{A.25})$$

This matrix is diagonalized by Z^E :

$$Z^E m_{\tilde{e}}^2 Z^{E,\dagger} = m_{\tilde{e}}^{2,dia} \quad (\text{A.26})$$

- **Mass matrix for CP-even Higgs**, Basis: (ϕ_d, ϕ_u)

$$m_h^2 = \begin{pmatrix} \frac{1}{8}(g_1^2 + g_2^2)(3v_d^2 - v_u^2) + m_{H_d}^2 + |\mu|^2 & -\frac{1}{4}(g_1^2 + g_2^2)v_d v_u - \Re(B_\mu) \\ -\frac{1}{4}(g_1^2 + g_2^2)v_d v_u - \Re(B_\mu) & -\frac{1}{8}(g_1^2 + g_2^2)(-3v_u^2 + v_d^2) + m_{H_u}^2 + |\mu|^2 \end{pmatrix} \quad (\text{A.27})$$

This matrix is diagonalized by Z^H :

$$Z^H m_h^2 Z^{H,\dagger} = m_h^{2,dia} \quad (\text{A.28})$$

- **Mass matrix for CP-odd Higgs**, Basis: (σ_d, σ_u)

$$m_{A^0}^2 = \begin{pmatrix} \frac{1}{8}(g_1^2 + g_2^2)(v_d^2 - v_u^2) + m_{H_d}^2 + |\mu|^2 & \Re(B_\mu) \\ \Re(B_\mu) & -\frac{1}{8}(g_1^2 + g_2^2)(v_d^2 - v_u^2) + m_{H_u}^2 + |\mu|^2 \\ + \xi_Z m^2(Z) \end{pmatrix} \quad (\text{A.29})$$

Gauge fixing contributions:

$$m^2(Z) = \begin{pmatrix} \frac{1}{4} v_d^2 (g_1 \sin \Theta_W + g_2 \cos \Theta_W)^2 & -\frac{1}{4} v_d v_u (g_1 \sin \Theta_W + g_2 \cos \Theta_W)^2 \\ -\frac{1}{4} v_d v_u (g_1 \sin \Theta_W + g_2 \cos \Theta_W)^2 & \frac{1}{4} v_u^2 (g_1 \sin \Theta_W + g_2 \cos \Theta_W)^2 \end{pmatrix} \quad (\text{A.30})$$

This matrix is diagonalized by Z^A :

$$Z^A m_{A^0}^2 Z^{A,\dagger} = m_{A^0}^{2,dia} \quad (\text{A.31})$$

- **Mass matrix for Charged Higgs**, Basis: $(H_d^-, H_u^{+,*}), (H_d^{-,*}, H_u^+)$

$$m_{H^-}^2 = \begin{pmatrix} m_{H_d^- H_d^{-,*}} & \frac{1}{4} g_2^2 v_d v_u + B_\mu^* \\ \frac{1}{4} g_2^2 v_d v_u + B_\mu & m_{H_u^{+,*} H_u^+} \end{pmatrix} + \xi_{W^-} m^2(W^-) \quad (\text{A.32})$$

$$m_{H_d^- H_d^{-,*}} = \frac{1}{8} (g_1^2 (v_d^2 - v_u^2) + g_2^2 (v_d^2 + v_u^2)) + m_{H_d}^2 + |\mu|^2 \quad (\text{A.33})$$

$$m_{H_u^{+,*} H_u^+} = \frac{1}{8} (g_1^2 (-v_d^2 + v_u^2) + g_2^2 (v_d^2 + v_u^2)) + m_{H_u}^2 + |\mu|^2 \quad (\text{A.34})$$

Gauge fixing contributions:

$$m^2(W^-) = \begin{pmatrix} \frac{1}{4} g_2^2 v_d^2 & -\frac{1}{4} g_2^2 v_d v_u \\ -\frac{1}{4} g_2^2 v_d v_u & \frac{1}{4} g_2^2 v_u^2 \end{pmatrix} \quad (\text{A.35})$$

This matrix is diagonalized by Z^+ :

$$Z^+ m_{H^-}^2 Z^{+,\dagger} = m_{H^-}^{2,dia} \quad (\text{A.36})$$

- **Mass matrix for Neutralinos**, Basis: $(\lambda_{\tilde{B}}, \tilde{W}^0, \tilde{H}_d^0, \tilde{H}_u^0)$

$$m_{\tilde{\chi}^0} = \begin{pmatrix} M_1 & 0 & -\frac{1}{2}g_1 v_d & \frac{1}{2}g_1 v_u \\ 0 & M_2 & \frac{1}{2}g_2 v_d & -\frac{1}{2}g_2 v_u \\ -\frac{1}{2}g_1 v_d & \frac{1}{2}g_2 v_d & 0 & -\mu \\ \frac{1}{2}g_1 v_u & -\frac{1}{2}g_2 v_u & -\mu & 0 \end{pmatrix} \quad (\text{A.37})$$

This matrix is diagonalized by N :

$$N^* m_{\tilde{\chi}^0} N^\dagger = m_{\tilde{\chi}^0}^{dia} \quad (\text{A.38})$$

- **Mass matrix for Charginos**, Basis: $(\tilde{W}^-, \tilde{H}_d^-), (\tilde{W}^+, \tilde{H}_u^+)$

$$m_{\tilde{\chi}^\pm} = \begin{pmatrix} M_2 & \frac{1}{\sqrt{2}}g_2 v_u \\ \frac{1}{\sqrt{2}}g_2 v_d & \mu \end{pmatrix} \quad (\text{A.39})$$

This matrix is diagonalized by U and V

$$U^* m_{\tilde{\chi}^\pm} V^\dagger = m_{\tilde{\chi}^\pm}^{dia} \quad (\text{A.40})$$

- **Mass matrix for charged Leptons**, Basis: $(e_L), (e_R^*)$

$$m_e = \left(\frac{1}{\sqrt{2}}v_d Y_e^T \right) \quad (\text{A.41})$$

This matrix is diagonalized by U_L^e and U_R^e

$$U_L^{e,*} m_e U_R^{e,\dagger} = m_e^{dia} \quad (\text{A.42})$$

- **Mass matrix for Down-Quarks**, Basis: $(d_{L,\alpha_1}), (d_{R,\beta_1}^*)$

$$m_d = \left(\frac{1}{\sqrt{2}}v_d \delta_{\alpha_1 \beta_1} Y_d^T \right) \quad (\text{A.43})$$

This matrix is diagonalized by U_L^d and U_R^d

$$U_L^{d,*} m_d U_R^{d,\dagger} = m_d^{dia} \quad (\text{A.44})$$

- **Mass matrix for Up-Quarks**, Basis: $(u_{L,\alpha_1}), (u_{R,\beta_1}^*)$

$$m_u = \left(\frac{1}{\sqrt{2}}v_u \delta_{\alpha_1 \beta_1} Y_u^T \right) \quad (\text{A.45})$$

This matrix is diagonalized by U_L^u and U_R^u

$$U_L^{u,*} m_u U_R^{u,\dagger} = m_u^{dia} \quad (\text{A.46})$$

A.2 Vertices

In this appendix we list all vertices relevant for our computations. Our conventions are as follows:

- Chiral vertices are parameterized as

$$\begin{aligned} &\Gamma_{F_a F_b S_c}^L P_L + \Gamma_{F_a F_b S_c}^R P_R \\ &\Gamma_{F_a F_b V_c^\mu}^L \gamma_\mu P_L + \Gamma_{F_a F_b V_c^\mu}^R \gamma_\mu P_R \end{aligned}$$

- The momentum flow in vector and scalar-vector vertices is

$$\begin{aligned} &\Gamma_{S_a S_b V_c^\mu} (p_{S_b}^\mu - p_{S_a}^\mu) \\ &\Gamma_{V_a^\rho V_b^\sigma V_c^\mu} (g_{\rho\mu} (-p_{V_c}^\sigma + p_{V_a}^\sigma) + g_{\rho\sigma} (-p_{V_a}^\mu + p_{V_c}^\mu) + g_{\sigma\mu} (-p_{V_a}^\rho + p_{V_c}^\rho)) \end{aligned}$$

Here we used polarization projectors $P_{L,R}$, metric $g_{\mu\nu}$ and momenta p of the external fields.

A.2.1 Fermion-scalar vertices

$$P_{ijk}^{c,L} = \Gamma_{\tilde{\chi}_i^+ \tilde{\chi}_j^- A_k^0}^L = -\frac{1}{\sqrt{2}} g_2 \left(U_{j1}^* V_{i2}^* Z_{k2}^A + U_{j2}^* V_{i1}^* Z_{k1}^A \right) \quad (\text{A.47})$$

$$P_{ijk}^{c,R} = \Gamma_{\tilde{\chi}_i^+ \tilde{\chi}_j^- A_k^0}^R = \frac{1}{\sqrt{2}} g_2 \left(U_{i1} V_{j2} Z_{k2}^A + U_{i2} V_{j1} Z_{k1}^A \right) \quad (\text{A.48})$$

$$\begin{aligned} P_{ijk}^L = \Gamma_{\tilde{\chi}_i^0 \tilde{\chi}_j^0 A_k^0}^L &= \frac{1}{2} \left(N_{i3}^* \left(g_1 N_{j1}^* - g_2 N_{j2}^* \right) Z_{k1}^A - g_2 N_{i2}^* N_{j3}^* Z_{k1}^A - g_1 N_{i4}^* N_{j1}^* Z_{k2}^A + g_2 N_{i4}^* N_{j2}^* Z_{k2}^A \right. \\ &\quad \left. + g_2 N_{i2}^* N_{j4}^* Z_{k2}^A + g_1 N_{i1}^* \left(N_{j3}^* Z_{k1}^A - N_{j4}^* Z_{k2}^A \right) \right) \end{aligned} \quad (\text{A.49})$$

$$\begin{aligned} P_{ijk}^R = \Gamma_{\tilde{\chi}_i^0 \tilde{\chi}_j^0 A_k^0}^R &= \frac{1}{2} \left(Z_{k1}^A \left(\left(-g_1 N_{i1} + g_2 N_{i2} \right) N_{j3} + N_{i3} \left(-g_1 N_{j1} + g_2 N_{j2} \right) \right) \right. \\ &\quad \left. + Z_{k2}^A \left(\left(g_1 N_{i1} - g_2 N_{i2} \right) N_{j4} + N_{i4} \left(g_1 N_{j1} - g_2 N_{j2} \right) \right) \right) \end{aligned} \quad (\text{A.50})$$

$$A_{ijk}^{d,L} = \Gamma_{\bar{d}_{i\alpha} d_{j\beta} A_k^0}^L = -\frac{i}{\sqrt{2}} \delta_{\alpha\beta} \sum_{b=1}^3 U_{L,jb}^{d,*} \sum_{a=1}^3 U_{R,ia}^{d,*} Y_{d,ab} Z_{k1}^A \quad (\text{A.51})$$

$$A_{ijk}^{d,R} = \Gamma_{\bar{d}_{i\alpha} d_{j\beta} A_k^0}^R = \frac{i}{\sqrt{2}} \delta_{\alpha\beta} \sum_{b=1}^3 \sum_{a=1}^3 Y_{d,ab}^* U_{R,ja}^d U_{L,ib}^d Z_{k1}^A \quad (\text{A.52})$$

$$A_{ijk}^L = \Gamma_{\bar{\ell}_i \ell_j A_k^0}^L = -\frac{i}{\sqrt{2}} \sum_{b=1}^3 U_{L,jb}^{e,*} \sum_{a=1}^3 U_{R,ia}^{e,*} Y_{e,ab} Z_{k1}^A \quad (\text{A.53})$$

$$A_{ijk}^R = \Gamma_{\bar{\ell}_i \ell_j A_k^0}^R = \frac{i}{\sqrt{2}} \sum_{b=1}^3 \sum_{a=1}^3 Y_{e,ab}^* U_{R,ja}^e U_{L,ib}^e Z_{k1}^A \quad (\text{A.54})$$

$$A_{ijk}^{u,L} = \Gamma_{\bar{u}_{i\alpha} u_{j\beta} A_k^0}^L = -\frac{i}{\sqrt{2}} \delta_{\alpha\beta} \sum_{b=1}^3 U_{L,jb}^{u,*} \sum_{a=1}^3 U_{R,ia}^{u,*} Y_{u,ab} Z_{k2}^A \quad (\text{A.55})$$

$$A_{ijk}^{u,R} = \Gamma_{\bar{u}_{i\alpha} u_{j\beta} A_k^0}^R = \frac{i}{\sqrt{2}} \delta_{\alpha\beta} \sum_{b=1}^3 \sum_{a=1}^3 Y_{u,ab}^* U_{R,ja}^u U_{L,ib}^u Z_{k2}^A \quad (\text{A.56})$$

$$A_{ijk}^{\nu,L} = \Gamma_{\nu_i \nu_j A_k^0}^L = -\frac{i}{\sqrt{2}} \left(\sum_{b=1}^3 U_{jb}^{V,*} \sum_{a=1}^3 U_{i3+a}^{V,*} Y_{\nu,ab} + \sum_{b=1}^3 U_{ib}^{V,*} \sum_{a=1}^3 U_{j3+a}^{V,*} Y_{\nu,ab} \right) Z_{k2}^A \quad (\text{A.57})$$

$$A_{ijk}^{\nu,R} = \Gamma_{\nu_i \nu_j A_k^0}^R = \frac{i}{\sqrt{2}} \left(\sum_{b=1}^3 \sum_{a=1}^3 Y_{\nu,ab}^* U_{j3+a}^V U_{ib}^V + \sum_{b=1}^3 \sum_{a=1}^3 Y_{\nu,ab}^* U_{i3+a}^V U_{jb}^V \right) Z_{k2}^A \quad (\text{A.58})$$

$$W_{ijk}^{u,L} = \Gamma_{\tilde{\chi}_i^- u j \beta \tilde{d}_{k\gamma}}^L = -\delta_{\beta\gamma} \left(g_2 U_{i1}^* \sum_{a=1}^3 U_{L,ja}^{u,*} Z_{ka}^D - U_{i2}^* \sum_{b=1}^3 U_{L,jb}^{u,*} \sum_{a=1}^3 Y_{d,ab} Z_{k3+a}^D \right) \quad (\text{A.59})$$

$$W_{ijk}^{u,R} = \Gamma_{\tilde{\chi}_i^- u j \beta \tilde{d}_{k\gamma}}^R = \delta_{\beta\gamma} \sum_{b=1}^3 \sum_{a=1}^3 Y_{u,ab} U_{R,ja}^u Z_{kb}^D V_{i2} \quad (\text{A.60})$$

$$S_{ijk}^{c,L} = \Gamma_{\tilde{\chi}_i^- \tilde{\chi}_j^- h_k}^L = -\frac{1}{\sqrt{2}} g_2 \left(U_{j1}^* V_{i2}^* Z_{k2}^H + U_{j2}^* V_{i1}^* Z_{k1}^H \right) \quad (\text{A.61})$$

$$S_{ijk}^{c,R} = \Gamma_{\tilde{\chi}_i^+ \tilde{\chi}_j^- h_k}^R = -\frac{1}{\sqrt{2}} g_2 \left(U_{i1} V_{j2} Z_{k2}^H + U_{i2} V_{j1} Z_{k1}^H \right) \quad (\text{A.62})$$

$$\begin{aligned} S_{ijk}^L = \Gamma_{\tilde{\chi}_i^0 \tilde{\chi}_j^0 h_k}^L &= \frac{1}{2} \left(N_{i3}^* \left(g_1 N_{j1}^* - g_2 N_{j2}^* \right) Z_{k1}^H - g_2 N_{i2}^* N_{j3}^* Z_{k1}^H - g_1 N_{i4}^* N_{j1}^* Z_{k2}^H + g_2 N_{i4}^* N_{j2}^* Z_{k2}^H \right. \\ &\quad \left. + g_2 N_{i2}^* N_{j4}^* Z_{k2}^H + g_1 N_{i1}^* \left(N_{j3}^* Z_{k1}^H - N_{j4}^* Z_{k2}^H \right) \right) \end{aligned} \quad (\text{A.63})$$

$$\begin{aligned} S_{ijk}^R = \Gamma_{\tilde{\chi}_i^0 \tilde{\chi}_j^0 h_k}^R &= \frac{1}{2} \left(Z_{k1}^H \left(\left(g_1 N_{i1} - g_2 N_{i2} \right) N_{j3} + N_{i3} \left(g_1 N_{j1} - g_2 N_{j2} \right) \right) \right. \\ &\quad \left. + Z_{k2}^H \left(\left(-g_1 N_{i1} + g_2 N_{i2} \right) N_{j4} + N_{i4} \left(-g_1 N_{j1} + g_2 N_{j2} \right) \right) \right) \end{aligned} \quad (\text{A.64})$$

$$N_{ijk}^{d,L} = \Gamma_{\tilde{\chi}_i^0 d j \beta \tilde{d}_{k\gamma}}^L = \delta_{\beta\gamma} \left(\frac{1}{\sqrt{2}} g_2 N_{i2}^* \sum_{a=1}^3 U_{L,ja}^{d,*} Z_{ka}^D - N_{i3}^* \sum_{b=1}^3 U_{L,jb}^{d,*} \sum_{a=1}^3 Y_{d,ab} Z_{k3+a}^D - \frac{1}{3\sqrt{2}} g_1 N_{i1}^* \sum_{a=1}^3 U_{L,ja}^{d,*} Z_{ka}^D \right) \quad (\text{A.65})$$

$$N_{ijk}^{d,R} = \Gamma_{\tilde{\chi}_i^0 d j \beta \tilde{d}_{k\gamma}}^R = -\delta_{\beta\gamma} \left(\sum_{b=1}^3 \sum_{a=1}^3 Y_{d,ab} U_{R,ja}^d Z_{kb}^D N_{i3} + \frac{\sqrt{2}}{3} g_1 \sum_{a=1}^3 Z_{k3+a}^D U_{R,ja}^d N_{i1} \right) \quad (\text{A.66})$$

$$N_{ijk}^L = \Gamma_{\tilde{\chi}_i^0 \ell_j \tilde{e}_k^*}^L = -N_{i3}^* \sum_{b=1}^3 U_{L,jb}^{e,*} \sum_{a=1}^3 Y_{e,ab} Z_{k3+a}^E + \frac{1}{\sqrt{2}} g_1 N_{i1}^* \sum_{a=1}^3 U_{L,ja}^{e,*} Z_{ka}^E + \frac{1}{\sqrt{2}} g_2 N_{i2}^* \sum_{a=1}^3 U_{L,ja}^{e,*} Z_{ka}^E \quad (\text{A.67})$$

$$N_{ijk}^R = \Gamma_{\tilde{\chi}_i^0 \ell_j \tilde{e}_k^*}^R = -\left(\sqrt{2} g_1 \sum_{a=1}^3 Z_{k3+a}^E U_{R,ja}^e N_{i1} + \sum_{b=1}^3 \sum_{a=1}^3 Y_{e,ab} U_{R,ja}^e Z_{kb}^E N_{i3} \right) \quad (\text{A.68})$$

$$N_{ijk}^{u,L} = \Gamma_{\tilde{\chi}_i^0 u j \beta \tilde{u}_{k\gamma}}^L = -\delta_{\beta\gamma} \left(\frac{1}{\sqrt{2}} g_2 N_{i2}^* \sum_{a=1}^3 U_{L,ja}^{u,*} Z_{ka}^U + N_{i4}^* \sum_{b=1}^3 U_{L,jb}^{u,*} \sum_{a=1}^3 Y_{u,ab} Z_{k3+a}^U + \frac{1}{3\sqrt{2}} g_1 N_{i1}^* \sum_{a=1}^3 U_{L,ja}^{u,*} Z_{ka}^U \right) \quad (\text{A.69})$$

$$N_{ijk}^{u,R} = \Gamma_{\tilde{\chi}_i^0 u j \beta \tilde{u}_{k\gamma}}^R = \delta_{\beta\gamma} \left(\frac{2\sqrt{2}}{3} g_1 \sum_{a=1}^3 Z_{k3+a}^U U_{R,ja}^u N_{i1} - \sum_{b=1}^3 \sum_{a=1}^3 Y_{u,ab} U_{R,ja}^u Z_{kb}^U N_{i4} \right) \quad (\text{A.70})$$

$$H_{ijk}^{d,L} = \Gamma_{d_{i\alpha} d_{j\beta} h_k}^L = -\frac{1}{\sqrt{2}} \delta_{\alpha\beta} \sum_{b=1}^3 U_{L,jb}^{d,*} \sum_{a=1}^3 U_{R,ia}^{d,*} Y_{d,ab} Z_{k1}^H \quad (\text{A.71})$$

$$H_{ijk}^{d,R} = \Gamma_{d_{i\alpha} d_{j\beta} h_k}^R = -\frac{1}{\sqrt{2}} \delta_{\alpha\beta} \sum_{b=1}^3 \sum_{a=1}^3 Y_{d,ab} U_{R,ja}^d U_{L,ib}^d Z_{k1}^H \quad (\text{A.72})$$

$$W_{ijk}^{d,L} = \Gamma_{\tilde{\chi}_i^0 d_{j\beta} \tilde{u}_{k\gamma}}^L = -\delta_{\beta\gamma} \left(g_2 V_{i1}^* \sum_{a=1}^3 U_{L,ja}^{d,*} Z_{ka}^U - V_{i2}^* \sum_{b=1}^3 U_{L,jb}^{d,*} \sum_{a=1}^3 Y_{u,ab} Z_{k3+a}^U \right) \quad (\text{A.73})$$

$$W_{ijk}^{d,R} = \Gamma_{\tilde{\chi}_i^0 d_{j\beta} \tilde{u}_{k\gamma}}^R = \delta_{\beta\gamma} \sum_{b=1}^3 \sum_{a=1}^3 Y_{d,ab} U_{R,ja}^d Z_{kb}^U U_{i2} \quad (\text{A.74})$$

$$V_{ijk}^{u,L} = \Gamma_{\bar{u}_{i\alpha} d_{j\beta} H_k^+}^L = \delta_{\alpha\beta} \sum_{b=1}^3 U_{L,jb}^{d,*} \sum_{a=1}^3 U_{R,ia}^{u,*} Y_{u,ab} Z_{k2}^+ \quad (\text{A.75})$$

$$V_{ijk}^{u,R} = \Gamma_{\bar{u}_{i\alpha} d_{j\beta} H_k^+}^R = \delta_{\alpha\beta} \sum_{b=1}^3 \sum_{a=1}^3 Y_{d,ab} U_{R,ja}^d U_{L,ib}^u Z_{k1}^+ \quad (\text{A.76})$$

$$V_{ijk}^{+,L} = \Gamma_{\nu_i \ell_j H_k^+}^L = \sum_{b=1}^3 U_{L,jb}^{e,*} \sum_{a=1}^3 U_{i3+a}^{V,*} Y_{\nu,ab} Z_{k2}^+ \quad (\text{A.77})$$

$$V_{ijk}^{+,R} = \Gamma_{\nu_i \ell_j h_k}^R = \sum_{b=1}^3 \sum_{a=1}^3 Y_{e,ab}^* U_{R,ja}^e U_{ib}^V Z_{k1}^+ \quad (\text{A.78})$$

$$H_{ijk}^L = \Gamma_{\tilde{\ell}_i \ell_j h_k}^L = -\frac{1}{\sqrt{2}} \sum_{b=1}^3 U_{L,jb}^{e,*} \sum_{a=1}^3 U_{R,ia}^{e,*} Y_{e,ab} Z_{k1}^H \quad (\text{A.79})$$

$$H_{ijk}^R = \Gamma_{\tilde{\ell}_i \ell_j h_k}^R = -\frac{1}{\sqrt{2}} \sum_{b=1}^3 \sum_{a=1}^3 Y_{e,ab}^* U_{R,ja}^e U_{L,ib}^e Z_{k1}^H \quad (\text{A.80})$$

$$X_{ijk}^L = \Gamma_{\tilde{\chi}_i^+ \ell_j \nu_k^i}^L = -\frac{i}{\sqrt{2}} \left(-g_2 V_{i1}^* \sum_{a=1}^3 U_{L,ja}^{e,*} Z_{ka}^{i,*} + V_{i2}^* \sum_{b=1}^3 U_{L,jb}^{e,*} \sum_{a=1}^3 Z_{k3+a}^{i,*} Y_{\nu,ab} \right) \quad (\text{A.81})$$

$$X_{ijk}^R = \Gamma_{\tilde{\chi}_i^+ \ell_j \nu_k^i}^R = -\frac{i}{\sqrt{2}} \sum_{b=1}^3 Z_{kb}^{i,*} \sum_{a=1}^3 Y_{e,ab}^* U_{R,ja}^e U_{i2} \quad (\text{A.82})$$

$$\hat{X}_{ijk}^L = \Gamma_{\tilde{\chi}_i^+ \ell_j \nu_k^R}^L = -\frac{1}{\sqrt{2}} \left(g_2 V_{i1}^* \sum_{a=1}^3 U_{L,ja}^{e,*} Z_{ka}^{R,*} - V_{i2}^* \sum_{b=1}^3 U_{L,jb}^{e,*} \sum_{a=1}^3 Z_{k3+a}^{R,*} Y_{\nu,ab} \right) \quad (\text{A.83})$$

$$\hat{X}_{ijk}^R = \Gamma_{\tilde{\chi}_i^+ \ell_j \nu_k^R}^R = \frac{1}{\sqrt{2}} \sum_{b=1}^3 Z_{kb}^{R,*} \sum_{a=1}^3 Y_{e,ab}^* U_{R,ja}^e U_{i2} \quad (\text{A.84})$$

$$H_{ijk}^{u,L} = \Gamma_{\bar{u}_{i\alpha} u_{j\beta} h_k}^L = -\frac{1}{\sqrt{2}} \delta_{\alpha\beta} \sum_{b=1}^3 U_{L,jb}^{u,*} \sum_{a=1}^3 U_{R,ia}^{u,*} Y_{u,ab} Z_{k2}^H \quad (\text{A.85})$$

$$H_{ijk}^{u,R} = \Gamma_{\bar{u}_{i\alpha} u_{j\beta} h_k}^R = -\frac{1}{\sqrt{2}} \delta_{\alpha\beta} \sum_{b=1}^3 \sum_{a=1}^3 Y_{u,ab}^* U_{R,ja}^u U_{L,ib}^u Z_{k2}^H \quad (\text{A.86})$$

$$H_{ijk}^{\nu,L} = \Gamma_{\nu_i \nu_j h_k}^L = -\frac{1}{\sqrt{2}} \left(\sum_{b=1}^3 U_{jb}^{V,*} \sum_{a=1}^3 U_{i3+a}^{V,*} Y_{\nu,ab} + \sum_{b=1}^3 U_{ib}^{V,*} \sum_{a=1}^3 U_{j3+a}^{V,*} Y_{\nu,ab} \right) Z_{k2}^H \quad (\text{A.87})$$

$$H_{ijk}^{\nu,R} = \Gamma_{\nu_i \nu_j h_k}^R = -\frac{1}{\sqrt{2}} \left(\sum_{b=1}^3 \sum_{a=1}^3 Y_{\nu,ab}^* U_{j3+a}^V U_{ib}^V + \sum_{b=1}^3 \sum_{a=1}^3 Y_{\nu,ab}^* U_{i3+a}^V U_{jb}^V \right) Z_{k2}^H \quad (\text{A.88})$$

$$F_{ijk}^{c,L} = \Gamma_{\tilde{\chi}_i^+ \nu_j \tilde{\epsilon}_k}^L = V_{i2}^* \sum_{b=1}^3 Z_{kb}^{E,*} \sum_{a=1}^3 U_{j3+a}^{V,*} Y_{\nu,ab} \quad (\text{A.89})$$

$$F_{ijk}^{c,R} = \Gamma_{\tilde{\chi}_i^+ \nu_j \tilde{\epsilon}_k}^R = - \left(g_2 \sum_{a=1}^3 Z_{ka}^{E,*} U_{ja}^V U_{i1} - \sum_{b=1}^3 \sum_{a=1}^3 Y_{e,ab}^* Z_{k3+a}^{E,*} U_{jb}^V U_{i2} \right) \quad (\text{A.90})$$

In addition, we introduce

$$\bar{W}_{ijk}^{d,L} = (W_{jik}^{d,R})^* \quad \bar{W}_{ijk}^{d,R} = (W_{jik}^{d,L})^* \quad \bar{W}_{ijk}^{u,L} = (W_{jik}^{u,R})^* \quad \bar{W}_{ijk}^{u,R} = (W_{jik}^{u,L})^* \quad (\text{A.91})$$

$$\bar{X}_{ijk}^L = (X_{jik}^R)^* \quad \bar{X}_{ijk}^R = (X_{jik}^L)^* \quad \hat{\bar{X}}_{ijk}^L = (\hat{X}_{jik}^R)^* \quad \hat{\bar{X}}_{ijk}^R = (\hat{X}_{jik}^L)^* \quad (\text{A.92})$$

$$\bar{N}_{ijk}^{d,L} = (N_{jik}^{d,R})^* \quad \bar{N}_{ijk}^{d,R} = (N_{jik}^{d,L})^* \quad \bar{N}_{ijk}^{u,L} = (N_{jik}^{u,R})^* \quad \bar{N}_{ijk}^{u,R} = (N_{jik}^{u,L})^* \quad (\text{A.93})$$

$$\bar{N}_{ijk}^L = (N_{jik}^R)^* \quad \bar{N}_{ijk}^R = (N_{jik}^L)^* \quad \bar{V}_{ijk}^{d,L} = (V_{jik}^{u,R})^* \quad \bar{V}_{ijk}^{d,R} = (V_{jik}^{u,L})^* \quad (\text{A.94})$$

$$\bar{V}_{ijk}^{+,L} = (V_{jik}^{+,R})^* \quad \bar{V}_{ijk}^{+,R} = (V_{jik}^{+,L})^* \quad (\text{A.95})$$

A.2.2 Fermion-vector vertices

$$F_{ij}^{c,L} = \Gamma_{\tilde{\chi}_i^+ \tilde{\chi}_j^- \gamma_\mu}^L = F_{ij}^{c,R} = \Gamma_{\tilde{\chi}_i^+ \tilde{\chi}_j^- \gamma_\mu}^R = e \delta_{ij} \quad (\text{A.96})$$

$$C_{ij}^L = \Gamma_{\tilde{\chi}_i^+ \tilde{\chi}_j^- Z_\mu}^L = g_2 U_{j1}^* \cos \Theta_W U_{i1} + \frac{1}{2} U_{j2}^* \left(-g_1 \sin \Theta_W + g_2 \cos \Theta_W \right) U_{i2} \quad (\text{A.97})$$

$$C_{ij}^R = \Gamma_{\tilde{\chi}_i^+ \tilde{\chi}_j^- Z_\mu}^R = g_2 V_{i1}^* \cos \Theta_W V_{j1} + \frac{1}{2} V_{i2}^* \left(-g_1 \sin \Theta_W + g_2 \cos \Theta_W \right) V_{j2} \quad (\text{A.98})$$

$$M_{ij}^L = \Gamma_{\tilde{\chi}_i^0 \tilde{\chi}_j^0 Z_\mu}^L = -\frac{1}{2} \left(g_1 \sin \Theta_W + g_2 \cos \Theta_W \right) \left(N_{j3}^* N_{i3} - N_{j4}^* N_{i4} \right) \quad (\text{A.99})$$

$$M_{ij}^R = \Gamma_{\tilde{\chi}_i^0 \tilde{\chi}_j^0 Z_\mu}^R = \frac{1}{2} \left(g_1 \sin \Theta_W + g_2 \cos \Theta_W \right) \left(N_{i3}^* N_{j3} - N_{i4}^* N_{j4} \right) \quad (\text{A.100})$$

$$D_{ij}^L = \Gamma_{\bar{d}_{i\alpha} d_{j\beta} Z_\mu}^L = \frac{1}{6} \delta_{\alpha\beta} \delta_{ij} \left(3g_2 \cos \Theta_W + g_1 \sin \Theta_W \right) \quad (\text{A.101})$$

$$D_{ij}^R = \Gamma_{\bar{d}_{i\alpha} d_{j\beta} Z_\mu}^R = -\frac{1}{3} g_1 \delta_{\alpha\beta} \delta_{ij} \sin \Theta_W \quad (\text{A.102})$$

$$\hat{V}_{ij}^{u,L} = \Gamma_{\bar{u}_{i\alpha} d_{j\beta} W_\mu^+}^L = -\frac{1}{\sqrt{2}} g_2 \delta_{\alpha\beta} \sum_{a=1}^3 U_{L,ja}^{d,*} U_{L,ia}^u \quad (\text{A.103})$$

$$\hat{V}_{ij}^{u,R} = \Gamma_{\bar{u}_{i\alpha} d_{j\beta} W_\mu^+}^R = 0 \quad (\text{A.104})$$

$$\hat{V}_{ij}^{+,L} = \Gamma_{\nu_i \ell_j W_\mu^+}^L = -\frac{1}{\sqrt{2}} g_2 \sum_{a=1}^3 U_{L,ja}^{e,*} U_{ia}^V \quad (\text{A.105})$$

$$\hat{V}_{ij}^{+,R} = \Gamma_{\nu_i \ell_j W_\mu^+}^R = 0 \quad (\text{A.106})$$

$$E_{ij}^L = \Gamma_{\bar{\ell}_i \ell_j Z_\mu}^L = \frac{1}{2} \delta_{ij} \left(-g_1 \sin \Theta_W + g_2 \cos \Theta_W \right) \quad (\text{A.107})$$

$$E_{ij}^R = \Gamma_{\bar{\ell}_i \ell_j Z_\mu}^R = -g_1 \delta_{ij} \sin \Theta_W \quad (\text{A.108})$$

$$U_{ij}^L = \Gamma_{\bar{u}_{i\alpha} u_{j\beta} Z_\mu}^L = -\frac{1}{6} \delta_{\alpha\beta} \delta_{ij} \left(3g_2 \cos \Theta_W - g_1 \sin \Theta_W \right) \quad (\text{A.109})$$

$$U_{ij}^R = \Gamma_{\bar{u}_{i\alpha} u_{j\beta} Z_\mu}^R = \frac{2}{3} g_1 \delta_{\alpha\beta} \delta_{ij} \sin \Theta_W \quad (\text{A.110})$$

$$V_{ij}^L = \Gamma_{\nu_i \nu_j Z_\mu}^L = -\frac{1}{2} \left(g_1 \sin \Theta_W + g_2 \cos \Theta_W \right) \sum_{a=1}^3 U_{ja}^{V,*} U_{ia}^V \quad (\text{A.111})$$

$$V_{ij}^R = \Gamma_{\nu_i \nu_j Z_\mu}^R = \frac{1}{2} \left(g_1 \sin \Theta_W + g_2 \cos \Theta_W \right) \sum_{a=1}^3 U_{ia}^{V,*} U_{ja}^V \quad (\text{A.112})$$

In addition, we introduce

$$\hat{\hat{V}}_{ij}^{d,L} = (\hat{V}_{ji}^{u,L})^* \quad \hat{\hat{V}}_{ij}^{d,R} = (\hat{V}_{ji}^{u,R})^* \quad \hat{\hat{V}}_{ij}^{+,L} = (\hat{V}_{ij}^{+,L})^* \quad \hat{\hat{V}}_{ij}^{+,R} = (\hat{V}_{ij}^{+,R})^* \quad (\text{A.113})$$

A.2.3 Scalar vertices

$$A_{ijk}^{hh} = \Gamma_{A_i^0 H_j^- H_k^+}^{hh} = -\frac{i}{4} g_2^2 \left(v_d Z_{i2}^A + v_u Z_{i1}^A \right) \left(-Z_{j1}^{+,*} Z_{k2}^+ + Z_{j2}^{+,*} Z_{k1}^+ \right) \quad (\text{A.114})$$

$$\begin{aligned} \tilde{A}_{ijk} &= \Gamma_{A_i^0 \bar{e}_j \bar{e}_k^*} = -\frac{i}{\sqrt{2}} \left(\sum_{b=1}^3 Z_{jb}^{E,*} \sum_{a=1}^3 Z_{k3+a}^E T_{e,ab} Z_{i1}^A - \sum_{b=1}^3 \sum_{a=1}^3 Z_{j3+a}^{E,*} T_{e,ab}^* Z_{kb}^E Z_{i1}^A \right. \\ &\quad \left. + \left(-\mu \sum_{b=1}^3 \sum_{a=1}^3 Y_{e,ab}^* Z_{j3+a}^{E,*} Z_{kb}^E + \mu^* \sum_{b=1}^3 Z_{jb}^{E,*} \sum_{a=1}^3 Y_{e,ab} Z_{k3+a}^E \right) Z_{i2}^A \right) \end{aligned} \quad (\text{A.115})$$

$$\tilde{A}_{ijk}^{ii} = \Gamma_{A_i^0 \nu_j^i \nu_k^i}^{ii} = 0 \quad (\text{A.116})$$

$$\begin{aligned} \tilde{A}_{ijk}^{ir} &= \Gamma_{A_i^0 \nu_j^i \nu_k^r}^{ir} = -\frac{1}{2\sqrt{2}} \left(\mu \sum_{b=1}^3 Z_{kb}^{R,*} \sum_{a=1}^3 Y_{\nu,ab}^* Z_{j3+a}^{i,*} Z_{i1}^A - \mu \sum_{b=1}^3 Z_{jb}^{i,*} \sum_{a=1}^3 Y_{\nu,ab}^* Z_{k3+a}^{R,*} Z_{i1}^A \right. \\ &\quad \left. + \mu^* \sum_{b=1}^3 Z_{kb}^{R,*} \sum_{a=1}^3 Z_{j3+a}^{i,*} Y_{\nu,ab} Z_{i1}^A - \mu^* \sum_{b=1}^3 Z_{jb}^{i,*} \sum_{a=1}^3 Z_{k3+a}^{R,*} Y_{\nu,ab} Z_{i1}^A \right) \end{aligned}$$

$$\begin{aligned}
& + \sum_{b=1}^3 Z_{kb}^{R,*} \sum_{a=1}^3 Z_{j3+a}^{i,*} T_{\nu,ab}^* Z_{i2}^A - \sum_{b=1}^3 Z_{jb}^{i,*} \sum_{a=1}^3 Z_{k3+a}^{R,*} T_{\nu,ab}^* Z_{i2}^A \\
& + \sum_{b=1}^3 Z_{kb}^{R,*} \sum_{a=1}^3 Z_{j3+a}^{i,*} T_{\nu,ab}^* Z_{i2}^A - \sum_{b=1}^3 Z_{jb}^{i,*} \sum_{a=1}^3 Z_{k3+a}^{R,*} T_{\nu,ab}^* Z_{i2}^A \\
& + \sum_{c=1}^3 Z_{kc}^{R,*} \sum_{b=1}^3 Z_{j6+b}^{i,*} \sum_{a=1}^3 Y_{\nu,ac}^* M_{R,ab} Z_{i2}^A - \sum_{c=1}^3 Z_{jc}^{i,*} \sum_{b=1}^3 Z_{k6+b}^{R,*} \sum_{a=1}^3 Y_{\nu,ac}^* M_{R,ab} Z_{i2}^A \\
& - \sum_{c=1}^3 Z_{k6+c}^{R,*} \sum_{b=1}^3 Z_{jb}^{i,*} \sum_{a=1}^3 M_{\nu,ac}^* Y_{\nu,ab} Z_{i2}^A + \sum_{c=1}^3 Z_{j6+c}^{i,*} \sum_{b=1}^3 Z_{kb}^{R,*} \sum_{a=1}^3 M_{\nu,ac}^* Y_{\nu,ab} Z_{i2}^A \Big)
\end{aligned} \tag{A.117}$$

$$\tilde{A}_{ijk}^{rr} = \Gamma_{A_i^0 \nu_j^R \nu_k^R} = 0 \tag{A.118}$$

$$\begin{aligned}
H_{ijk}^{hh} = \Gamma_{h_i H_j^- H_k^+} &= -\frac{1}{4} \left(Z_{j1}^{+,*} \left(Z_{i1}^H \left((g_1^2 + g_2^2) v_d Z_{k1}^+ + g_2^2 v_u Z_{k2}^+ \right) + Z_{i2}^H \left((-g_1^2 + g_2^2) v_u Z_{k1}^+ + g_2^2 v_d Z_{k2}^+ \right) \right) \right. \\
&\quad \left. + Z_{j2}^{+,*} \left(Z_{i1}^H \left((-g_1^2 + g_2^2) v_d Z_{k2}^+ + g_2^2 v_u Z_{k1}^+ \right) + Z_{i2}^H \left((g_1^2 + g_2^2) v_u Z_{k2}^+ + g_2^2 v_d Z_{k1}^+ \right) \right) \right)
\end{aligned} \tag{A.119}$$

$$\begin{aligned}
\tilde{H}_{ijk} = \Gamma_{h_i \tilde{e}_j \tilde{e}_k^*} &= -\frac{1}{4} \left((-g_2^2 + g_1^2) \sum_{a=1}^3 Z_{ja}^{E,*} Z_{ka}^E \left(v_d Z_{i1}^H - v_u Z_{i2}^H \right) \right. \\
&\quad + 2 \left(\sqrt{2} \sum_{b=1}^3 Z_{jb}^{E,*} \sum_{a=1}^3 Z_{k3+a}^E T_{e,ab} Z_{i1}^H + \sqrt{2} \sum_{b=1}^3 \sum_{a=1}^3 Z_{j3+a}^{E,*} T_{e,ab}^* Z_{kb}^E Z_{i1}^H \right. \\
&\quad + 2v_d \sum_{c=1}^3 Z_{j3+c}^{E,*} \sum_{b=1}^3 \sum_{a=1}^3 Y_{e,ca}^* Y_{e,ba} Z_{k3+b}^E Z_{i1}^H + 2v_d \sum_{c=1}^3 \sum_{b=1}^3 Z_{jb}^{E,*} \sum_{a=1}^3 Y_{e,ac}^* Y_{e,ab} Z_{kc}^E Z_{i1}^H \\
&\quad - \sqrt{2} \mu^* \sum_{b=1}^3 Z_{jb}^{E,*} \sum_{a=1}^3 Y_{e,ab} Z_{k3+a}^E Z_{i2}^H - \sqrt{2} \mu \sum_{b=1}^3 \sum_{a=1}^3 Y_{e,ab}^* Z_{j3+a}^{E,*} Z_{kb}^E Z_{i2}^H \\
&\quad \left. \left. + g_1^2 \sum_{a=1}^3 Z_{j3+a}^{E,*} Z_{k3+a}^E \left(-v_d Z_{i1}^H + v_u Z_{i2}^H \right) \right) \right)
\end{aligned} \tag{A.120}$$

$$\begin{aligned}
\tilde{H}_{ijk}^{ii} = \Gamma_{h_i \nu_j^i \nu_k^i} &= -\frac{1}{4} \left(-\sqrt{2} \mu \sum_{b=1}^3 Z_{kb}^{i,*} \sum_{a=1}^3 Y_{\nu,ab}^* Z_{j3+a}^{i,*} Z_{i1}^H - \sqrt{2} \mu \sum_{b=1}^3 Z_{jb}^{i,*} \sum_{a=1}^3 Y_{\nu,ab}^* Z_{k3+a}^{i,*} Z_{i1}^H \right. \\
&\quad - \sqrt{2} \mu^* \sum_{b=1}^3 Z_{kb}^{i,*} \sum_{a=1}^3 Z_{j3+a}^{i,*} Y_{\nu,ab} Z_{i1}^H - \sqrt{2} \mu^* \sum_{b=1}^3 Z_{jb}^{i,*} \sum_{a=1}^3 Z_{k3+a}^{i,*} Y_{\nu,ab} Z_{i1}^H \\
&\quad + \sqrt{2} \sum_{b=1}^3 Z_{kb}^{i,*} \sum_{a=1}^3 Z_{j3+a}^{i,*} T_{\nu,ab}^* Z_{i2}^H + \sqrt{2} \sum_{b=1}^3 Z_{jb}^{i,*} \sum_{a=1}^3 Z_{k3+a}^{i,*} T_{\nu,ab}^* Z_{i2}^H \\
&\quad + \sqrt{2} \sum_{b=1}^3 Z_{kb}^{i,*} \sum_{a=1}^3 Z_{j3+a}^{i,*} T_{\nu,ab} Z_{i2}^H + \sqrt{2} \sum_{b=1}^3 Z_{jb}^{i,*} \sum_{a=1}^3 Z_{k3+a}^{i,*} T_{\nu,ab} Z_{i2}^H \\
&\quad + \sqrt{2} \sum_{c=1}^3 Z_{kc}^{i,*} \sum_{b=1}^3 Z_{j6+b}^{i,*} \sum_{a=1}^3 Y_{\nu,ac}^* M_{R,ab} Z_{i2}^H + \sqrt{2} \sum_{c=1}^3 Z_{jc}^{i,*} \sum_{b=1}^3 Z_{k6+b}^{i,*} \sum_{a=1}^3 Y_{\nu,ac}^* M_{R,ab} Z_{i2}^H \\
&\quad + \sqrt{2} \sum_{c=1}^3 Z_{k6+c}^{i,*} \sum_{b=1}^3 Z_{jb}^{i,*} \sum_{a=1}^3 M_{\nu,ac}^* Y_{\nu,ab} Z_{i2}^H + \sqrt{2} \sum_{c=1}^3 Z_{j6+c}^{i,*} \sum_{b=1}^3 Z_{kb}^{i,*} \sum_{a=1}^3 M_{\nu,ac}^* Y_{\nu,ab} Z_{i2}^H \\
&\quad + 2v_u \sum_{c=1}^3 Z_{kc}^{i,*} \sum_{b=1}^3 Z_{jb}^{i,*} \sum_{a=1}^3 Y_{\nu,ac}^* Y_{\nu,ab} Z_{i2}^H + 2v_u \sum_{c=1}^3 Z_{jc}^{i,*} \sum_{b=1}^3 Z_{kb}^{i,*} \sum_{a=1}^3 Y_{\nu,ac}^* Y_{\nu,ab} Z_{i2}^H \\
&\quad + 2v_u \sum_{c=1}^3 Z_{k3+c}^{i,*} \sum_{b=1}^3 Z_{j3+b}^{i,*} \sum_{a=1}^3 Y_{\nu,ca}^* Y_{\nu,ba} Z_{i2}^H
\end{aligned}$$

$$\begin{aligned}
& + 2v_u \sum_{c=1}^3 Z_{j3+c}^{i,*} \sum_{b=1}^3 Z_{k3+b}^{i,*} \sum_{a=1}^3 Y_{\nu,ca}^* Y_{\nu,ba} Z_{i2}^H \\
& + \left(g_1^2 + g_2^2 \right) \sum_{a=1}^3 Z_{ja}^{i,*} Z_{ka}^{i,*} \left(v_d Z_{i1}^H - v_u Z_{i2}^H \right)
\end{aligned} \tag{A.121}$$

$$\tilde{H}_{ijk}^{ir} = \Gamma_{h_i \nu_j^i \nu_k^R} = 0 \tag{A.122}$$

$$\begin{aligned}
\tilde{H}_{ijk}^{rr} = \Gamma_{h_i \nu_j^R \nu_k^R} = & -\frac{1}{4} \left(-\sqrt{2}\mu \sum_{b=1}^3 Z_{kb}^{R,*} \sum_{a=1}^3 Y_{\nu,ab}^* Z_{j3+a}^R Z_{i1}^H - \sqrt{2}\mu \sum_{b=1}^3 Z_{jb}^{R,*} \sum_{a=1}^3 Y_{\nu,ab}^* Z_{k3+a}^R Z_{i1}^H \right. \\
& - \sqrt{2}\mu^* \sum_{b=1}^3 Z_{kb}^{R,*} \sum_{a=1}^3 Z_{j3+a}^R Y_{\nu,ab} Z_{i1}^H - \sqrt{2}\mu^* \sum_{b=1}^3 Z_{jb}^{R,*} \sum_{a=1}^3 Z_{k3+a}^R Y_{\nu,ab} Z_{i1}^H \\
& + \sqrt{2} \sum_{b=1}^3 Z_{kb}^{R,*} \sum_{a=1}^3 Z_{j3+a}^R T_{\nu,ab}^* Z_{i2}^H + \sqrt{2} \sum_{b=1}^3 Z_{jb}^{R,*} \sum_{a=1}^3 Z_{k3+a}^R T_{\nu,ab}^* Z_{i2}^H \\
& + \sqrt{2} \sum_{b=1}^3 Z_{kb}^{R,*} \sum_{a=1}^3 Z_{j3+a}^R T_{\nu,ab} Z_{i2}^H + \sqrt{2} \sum_{b=1}^3 Z_{jb}^{R,*} \sum_{a=1}^3 Z_{k3+a}^R T_{\nu,ab} Z_{i2}^H \\
& + \sqrt{2} \sum_{c=1}^3 Z_{kc}^{R,*} \sum_{b=1}^3 Z_{j6+b}^{R,*} \sum_{a=1}^3 Y_{\nu,ac}^* M_{R,ab} Z_{i2}^H + \sqrt{2} \sum_{c=1}^3 Z_{jc}^{R,*} \sum_{b=1}^3 Z_{k6+b}^{R,*} \sum_{a=1}^3 Y_{\nu,ac}^* M_{R,ab} Z_{i2}^H \\
& + \sqrt{2} \sum_{c=1}^3 Z_{k6+c}^{R,*} \sum_{b=1}^3 Z_{jb}^{R,*} \sum_{a=1}^3 M_{\nu,ac}^* Y_{\nu,ab} Z_{i2}^H + \sqrt{2} \sum_{c=1}^3 Z_{j6+c}^{R,*} \sum_{b=1}^3 Z_{kb}^{R,*} \sum_{a=1}^3 M_{\nu,ac}^* Y_{\nu,ab} Z_{i2}^H \\
& + 2v_u \sum_{c=1}^3 Z_{kc}^{R,*} \sum_{b=1}^3 Z_{jb}^{R,*} \sum_{a=1}^3 Y_{\nu,ac}^* Y_{\nu,ab} Z_{i2}^H + 2v_u \sum_{c=1}^3 Z_{jc}^{R,*} \sum_{b=1}^3 Z_{kb}^{R,*} \sum_{a=1}^3 Y_{\nu,ac}^* Y_{\nu,ab} Z_{i2}^H \\
& + 2v_u \sum_{c=1}^3 Z_{k3+c}^{R,*} \sum_{b=1}^3 Z_{j3+b}^{R,*} \sum_{a=1}^3 Y_{\nu,ca}^* Y_{\nu,ba} Z_{i2}^H \\
& + 2v_u \sum_{c=1}^3 Z_{j3+c}^{R,*} \sum_{b=1}^3 Z_{k3+b}^{R,*} \sum_{a=1}^3 Y_{\nu,ca}^* Y_{\nu,ba} Z_{i2}^H \\
& \left. + \left(g_1^2 + g_2^2 \right) \sum_{a=1}^3 Z_{ja}^{R,*} Z_{ka}^{R,*} \left(v_d Z_{i1}^H - v_u Z_{i2}^H \right) \right)
\end{aligned} \tag{A.123}$$

A.2.4 Scalar-vector vertices

$$A_{ij}^{hw} = \Gamma_{A_i^0 H_j^- W_\mu^+} = \frac{i}{2} g_2 \left(Z_{j1}^{+,*} Z_{i1}^A + Z_{j2}^{+,*} Z_{i2}^A \right) \tag{A.124}$$

$$H_{ij}^{hw} = \Gamma_{h_i H_j^- W_\mu^+} = \frac{1}{2} g_2 \left(Z_{j2}^{+,*} Z_{i2}^H - Z_{j1}^{+,*} Z_{i1}^H \right) \tag{A.125}$$

$$F_{ij}^h = \Gamma_{H_i^- H_j^+ \gamma_\mu} = -e \delta_{ij} \tag{A.126}$$

$$Z_{ij}^{hh} = \Gamma_{H_i^- H_j^+ Z_\mu} = \frac{1}{2} \left(g_1 \sin \Theta_W - g_2 \cos \Theta_W \right) \delta_{ij} \tag{A.127}$$

$$F_{ij}^{\tilde{e}} = \Gamma_{\tilde{e}_i \tilde{e}_j^* \gamma_\mu} = -e \delta_{ij} \tag{A.128}$$

$$\begin{aligned}
\tilde{E}_{ij} = \Gamma_{\tilde{e}_i \tilde{e}_j^* Z_\mu} = & g_1 \sin \Theta_W \sum_{a=1}^3 Z_{i3+a}^{E,*} Z_{j3+a}^E + \frac{1}{2} \left(g_1 \sin \Theta_W - g_2 \cos \Theta_W \right) \sum_{a=1}^3 Z_{ia}^{E,*} Z_{ja}^E \\
\end{aligned} \tag{A.129}$$

$$\tilde{V}_{ij} = \Gamma_{\nu_i^i \nu_j^R Z_\mu} = \frac{i}{2} \left(g_1 \sin \Theta_W + g_2 \cos \Theta_W \right) \sum_{a=1}^3 Z_{ia}^{i,*} Z_{ja}^{R,*} \tag{A.130}$$

$$H_i^{ww} = \Gamma_{h_i W_\sigma^+ W_\mu^-} = -\frac{1}{2} g_2^2 (v_d Z_{i1}^H + v_u Z_{i2}^H) \quad (\text{A.131})$$

$$F_i^{hw} = \Gamma_{H_i^- W_\sigma^+ \gamma_\mu} = -\frac{i}{2 \sin \Theta_W} e^2 (v_d Z_{i1}^{+,*} - v_u Z_{i2}^{+,*}) \quad (\text{A.132})$$

$$Z_i^{hw} = \Gamma_{H_i^- W_\sigma^+ Z_\mu} = \frac{1}{2} g_1 g_2 (v_d Z_{i1}^{+,*} - v_u Z_{i2}^{+,*}) \sin \Theta_W \quad (\text{A.133})$$

(A.134)

In addition, we introduce

$$\bar{A}_{ij}^{hw} = (A_{ij}^{hw})^* \quad \bar{H}_{ij}^{hw} = (H_{ij}^{hw})^* \quad \bar{F}_i^{hw} = (F_i^{hw})^* \quad \bar{Z}_i^{hw} = (Z_i^{hw})^* \quad (\text{A.135})$$

A.2.5 Vector vertices

$$F^w = \Gamma_{W_\rho^+ \gamma_\sigma W_\mu^-} = g_2 \sin \Theta_W \quad (\text{A.136})$$

$$Z^{ww} = \Gamma_{W_\rho^+ W_\sigma^- Z_\mu} = -g_2 \cos \Theta_W \quad (\text{A.137})$$

B Renormalization Group Equations

We give in the following the 2-loop RGEs for the considered model. For parameters present in the MSSM we show only the difference with respect to the MSSM RGEs. In general, the RGEs for a parameter X are defined by

$$\frac{d}{dt} X = \frac{1}{16\pi^2} \beta_X^{(1)} + \frac{1}{(16\pi^2)^2} \beta_X^{(2)} \quad (\text{B.1})$$

Here, $t = \log(Q/M)$, with Q the renormalization scale and M a reference scale.

Gauge couplings

$$\Delta \beta_{g_1}^{(2)} = -\frac{6}{5} g_1^3 \text{Tr}(Y_\nu Y_\nu^\dagger) \quad (\text{B.2})$$

$$\Delta \beta_{g_2}^{(2)} = -2g_2^3 \text{Tr}(Y_\nu Y_\nu^\dagger) \quad (\text{B.3})$$

Gaugino mass parameters

$$\Delta \beta_{M_1}^{(2)} = -\frac{12}{5} g_1^2 (M_1 \text{Tr}(Y_\nu Y_\nu^\dagger) - \text{Tr}(Y_\nu^\dagger T_\nu)) \quad (\text{B.4})$$

$$\Delta \beta_{M_2}^{(2)} = 4g_2^2 (-M_2 \text{Tr}(Y_\nu Y_\nu^\dagger) + \text{Tr}(Y_\nu^\dagger T_\nu)) \quad (\text{B.5})$$

Trilinear superpotential parameters

$$\Delta \beta_{Y_d}^{(2)} = -Y_d \text{Tr}(Y_e Y_\nu^\dagger Y_\nu Y_e^\dagger) - Y_d Y_u^\dagger Y_u \text{Tr}(Y_\nu Y_\nu^\dagger) \quad (\text{B.6})$$

$$\Delta \beta_{Y_e}^{(1)} = Y_e Y_\nu^\dagger Y_\nu \quad (\text{B.7})$$

$$\Delta \beta_{Y_e}^{(2)} = -2Y_e Y_\nu^\dagger Y_\nu Y_e^\dagger Y_e - 2Y_e Y_\nu^\dagger Y_\nu Y_\nu^\dagger Y_\nu - 3Y_e Y_\nu^\dagger Y_\nu \text{Tr}(Y_u Y_u^\dagger)$$

$$- Y_e Y_\nu^\dagger Y_\nu \text{Tr} \left(Y_\nu Y_\nu^\dagger \right) - Y_e \text{Tr} \left(Y_e Y_\nu^\dagger Y_\nu Y_e^\dagger \right) \quad (\text{B.8})$$

$$\Delta \beta_{Y_u}^{(1)} = Y_u \text{Tr} \left(Y_u Y_\nu^\dagger \right) \quad (\text{B.9})$$

$$\Delta \beta_{Y_u}^{(2)} = -3 Y_u \text{Tr} \left(Y_\nu Y_\nu^\dagger Y_\nu Y_\nu^\dagger \right) - 3 Y_u Y_u^\dagger Y_u \text{Tr} \left(Y_\nu Y_\nu^\dagger \right) - Y_u \text{Tr} \left(Y_e Y_\nu^\dagger Y_\nu Y_e^\dagger \right) \quad (\text{B.10})$$

$$\beta_{Y_\nu}^{(1)} = 3 Y_\nu Y_\nu^\dagger Y_\nu + Y_\nu \left(-3 g_2^2 + 3 \text{Tr} \left(Y_u Y_u^\dagger \right) - \frac{3}{5} g_1^2 + \text{Tr} \left(Y_\nu Y_\nu^\dagger \right) \right) + Y_\nu Y_e^\dagger Y_e \quad (\text{B.11})$$

$$\begin{aligned} \beta_{Y_\nu}^{(2)} = & +\frac{6}{5} g_1^2 Y_\nu Y_\nu^\dagger Y_\nu + 6 g_2^2 Y_\nu Y_\nu^\dagger Y_\nu - 2 Y_\nu Y_e^\dagger Y_e Y_e^\dagger Y_e - 2 Y_\nu Y_e^\dagger Y_e Y_\nu^\dagger Y_\nu \\ & - 4 Y_\nu Y_\nu^\dagger Y_\nu Y_\nu^\dagger Y_\nu + Y_\nu Y_e^\dagger Y_e \left(-3 \text{Tr} \left(Y_d Y_d^\dagger \right) + \frac{6}{5} g_1^2 - \text{Tr} \left(Y_e Y_e^\dagger \right) \right) - 9 Y_\nu Y_\nu^\dagger Y_\nu \text{Tr} \left(Y_u Y_u^\dagger \right) \\ & - 3 Y_\nu Y_\nu^\dagger Y_\nu \text{Tr} \left(Y_\nu Y_\nu^\dagger \right) \\ & + Y_\nu \left(\frac{207}{50} g_1^4 + \frac{9}{5} g_1^2 g_2^2 + \frac{15}{2} g_2^4 + \frac{4}{5} \left(20 g_3^2 + g_1^2 \right) \text{Tr} \left(Y_u Y_u^\dagger \right) - 3 \text{Tr} \left(Y_d Y_u^\dagger Y_u Y_d^\dagger \right) \right. \\ & \left. - \text{Tr} \left(Y_e Y_\nu^\dagger Y_\nu Y_e^\dagger \right) - 9 \text{Tr} \left(Y_u Y_u^\dagger Y_u Y_u^\dagger \right) - 3 \text{Tr} \left(Y_\nu Y_\nu^\dagger Y_\nu Y_\nu^\dagger \right) \right) \end{aligned} \quad (\text{B.12})$$

Bilinear superpotential parameters

$$\Delta \beta_\mu^{(1)} = \mu \text{Tr} \left(Y_\nu Y_\nu^\dagger \right) \quad (\text{B.13})$$

$$\Delta \beta_\mu^{(2)} = -\mu \left(2 \text{Tr} \left(Y_e Y_\nu^\dagger Y_\nu Y_e^\dagger \right) + 3 \text{Tr} \left(Y_\nu Y_\nu^\dagger Y_\nu Y_\nu^\dagger \right) \right) \quad (\text{B.14})$$

$$\beta_{\mu X}^{(1)} = \beta_{\mu X}^{(2)} = 0 \quad (\text{B.15})$$

$$\beta_{M_R}^{(1)} = 2 Y_\nu Y_\nu^\dagger M_R \quad (\text{B.16})$$

$$\begin{aligned} \beta_{M_R}^{(2)} = & -2 \left(Y_\nu Y_e^\dagger Y_e Y_\nu^\dagger M_R + Y_\nu Y_\nu^\dagger Y_\nu Y_\nu^\dagger M_R \right) \\ & + Y_\nu Y_\nu^\dagger M_R \left(-2 \text{Tr} \left(Y_\nu Y_\nu^\dagger \right) + 6 g_2^2 - 6 \text{Tr} \left(Y_u Y_u^\dagger \right) + \frac{6}{5} g_1^2 \right) \end{aligned} \quad (\text{B.17})$$

Trilinear soft-breaking parameters

$$\begin{aligned} \Delta \beta_{T_d}^{(2)} = & -2 Y_d Y_u^\dagger T_u \text{Tr} \left(Y_\nu Y_\nu^\dagger \right) - T_d Y_u^\dagger Y_u \text{Tr} \left(Y_\nu Y_\nu^\dagger \right) - 2 Y_d Y_u^\dagger Y_u \text{Tr} \left(Y_\nu^\dagger T_\nu \right) \\ & - T_d \text{Tr} \left(Y_e Y_\nu^\dagger Y_\nu Y_e^\dagger \right) - 2 Y_d \text{Tr} \left(Y_e Y_\nu^\dagger T_\nu Y_e^\dagger \right) - 2 Y_d \text{Tr} \left(Y_\nu Y_e^\dagger T_e Y_\nu^\dagger \right) \end{aligned} \quad (\text{B.18})$$

$$\Delta \beta_{T_e}^{(1)} = 2 Y_e Y_\nu^\dagger T_\nu + T_e Y_\nu^\dagger Y_\nu \quad (\text{B.19})$$

$$\begin{aligned} \Delta \beta_{T_e}^{(2)} = & -2 Y_e Y_\nu^\dagger Y_\nu Y_e^\dagger T_e - 4 Y_e Y_\nu^\dagger Y_\nu Y_\nu^\dagger T_\nu - 4 Y_e Y_\nu^\dagger T_\nu Y_e^\dagger Y_e - 4 Y_e Y_\nu^\dagger T_\nu Y_\nu^\dagger Y_\nu \\ & - 4 T_e Y_\nu^\dagger Y_\nu Y_e^\dagger Y_e - 2 T_e Y_\nu^\dagger Y_\nu Y_\nu^\dagger Y_\nu - 6 Y_e Y_\nu^\dagger T_\nu \text{Tr} \left(Y_u Y_u^\dagger \right) \\ & - 3 T_e Y_\nu^\dagger Y_\nu \text{Tr} \left(Y_u Y_u^\dagger \right) - 2 Y_e Y_\nu^\dagger T_\nu \text{Tr} \left(Y_\nu Y_\nu^\dagger \right) - T_e Y_\nu^\dagger Y_\nu \text{Tr} \left(Y_\nu Y_\nu^\dagger \right) \\ & - 6 Y_e Y_\nu^\dagger Y_\nu \text{Tr} \left(Y_u^\dagger T_u \right) - 2 Y_e Y_\nu^\dagger Y_\nu \text{Tr} \left(Y_\nu^\dagger T_\nu \right) - T_e \text{Tr} \left(Y_e Y_\nu^\dagger Y_\nu Y_e^\dagger \right) \\ & - 2 Y_e \text{Tr} \left(Y_e Y_\nu^\dagger T_\nu Y_e^\dagger \right) - 2 Y_e \text{Tr} \left(Y_\nu Y_e^\dagger T_e Y_\nu^\dagger \right) \end{aligned} \quad (\text{B.20})$$

$$\Delta \beta_{T_u}^{(1)} = 2 Y_u \text{Tr} \left(Y_\nu^\dagger T_\nu \right) + T_u \text{Tr} \left(Y_\nu Y_\nu^\dagger \right) \quad (\text{B.21})$$

$$\Delta \beta_{T_u}^{(2)} = -4 Y_u Y_u^\dagger T_u \text{Tr} \left(Y_\nu Y_\nu^\dagger \right) - 5 T_u Y_u^\dagger Y_u \text{Tr} \left(Y_\nu Y_\nu^\dagger \right) - 6 Y_u Y_u^\dagger Y_u \text{Tr} \left(Y_\nu^\dagger T_\nu \right)$$

$$\begin{aligned}
& - T_u \text{Tr} \left(Y_e Y_\nu^\dagger Y_\nu Y_e^\dagger \right) - 2Y_u \text{Tr} \left(Y_e Y_\nu^\dagger T_\nu Y_e^\dagger \right) - 2Y_u \text{Tr} \left(Y_\nu Y_e^\dagger T_e Y_\nu^\dagger \right) \\
& - 3T_u \text{Tr} \left(Y_\nu Y_\nu^\dagger Y_\nu Y_\nu^\dagger \right) - 12Y_u \text{Tr} \left(Y_\nu Y_\nu^\dagger T_\nu Y_\nu^\dagger \right)
\end{aligned} \tag{B.22}$$

$$\begin{aligned}
\beta_{T_\nu}^{(1)} &= +2Y_\nu Y_e^\dagger T_e + 4Y_\nu Y_\nu^\dagger T_\nu + T_\nu Y_e^\dagger Y_e + 5T_\nu Y_\nu^\dagger Y_\nu - \frac{3}{5}g_1^2 T_\nu - 3g_2^2 T_\nu \\
& + 3T_\nu \text{Tr} \left(Y_u Y_u^\dagger \right) + T_\nu \text{Tr} \left(Y_\nu Y_\nu^\dagger \right) + Y_\nu \left(2\text{Tr} \left(Y_\nu^\dagger T_\nu \right) + 6g_2^2 M_2 + 6\text{Tr} \left(Y_u^\dagger T_u \right) + \frac{6}{5}g_1^2 M_1 \right)
\end{aligned} \tag{B.23}$$

$$\begin{aligned}
\beta_{T_\nu}^{(2)} &= +\frac{12}{5}g_1^2 Y_\nu Y_e^\dagger T_e - \frac{12}{5}g_1^2 M_1 Y_\nu Y_\nu^\dagger Y_\nu - 12g_2^2 M_2 Y_\nu Y_\nu^\dagger Y_\nu + \frac{6}{5}g_1^2 Y_\nu Y_\nu^\dagger T_\nu \\
& + 6g_2^2 Y_\nu Y_\nu^\dagger T_\nu + \frac{6}{5}g_1^2 T_\nu Y_e^\dagger Y_e + \frac{12}{5}g_1^2 T_\nu Y_\nu^\dagger Y_\nu + 12g_2^2 T_\nu Y_\nu^\dagger Y_\nu \\
& - 4Y_\nu Y_e^\dagger Y_e Y_e^\dagger T_e - 2Y_\nu Y_e^\dagger Y_e Y_\nu^\dagger T_\nu - 4Y_\nu Y_e^\dagger T_e Y_e^\dagger Y_e - 4Y_\nu Y_e^\dagger T_e Y_\nu^\dagger Y_\nu \\
& - 6Y_\nu Y_\nu^\dagger Y_\nu Y_\nu^\dagger T_\nu - 8Y_\nu Y_\nu^\dagger T_\nu Y_\nu^\dagger Y_\nu - 2T_\nu Y_e^\dagger Y_e Y_e^\dagger Y_e - 4T_\nu Y_e^\dagger Y_e Y_\nu^\dagger Y_\nu \\
& - 6T_\nu Y_\nu^\dagger Y_\nu Y_\nu^\dagger Y_\nu + \frac{207}{50}g_1^4 T_\nu + \frac{9}{5}g_2^2 g_2^2 T_\nu + \frac{15}{2}g_2^4 T_\nu - 6Y_\nu Y_e^\dagger T_e \text{Tr} \left(Y_d Y_d^\dagger \right) \\
& - 3T_\nu Y_e^\dagger Y_e \text{Tr} \left(Y_d Y_d^\dagger \right) - 2Y_\nu Y_e^\dagger T_e \text{Tr} \left(Y_e Y_e^\dagger \right) - T_\nu Y_e^\dagger Y_e \text{Tr} \left(Y_e Y_e^\dagger \right) \\
& - 12Y_\nu Y_\nu^\dagger T_\nu \text{Tr} \left(Y_u Y_u^\dagger \right) - 15T_\nu Y_\nu^\dagger Y_\nu \text{Tr} \left(Y_u Y_u^\dagger \right) + \frac{4}{5}g_1^2 T_\nu \text{Tr} \left(Y_u Y_u^\dagger \right) \\
& + 16g_3^2 T_\nu \text{Tr} \left(Y_u Y_u^\dagger \right) - 4Y_\nu Y_\nu^\dagger T_\nu \text{Tr} \left(Y_\nu Y_\nu^\dagger \right) - 5T_\nu Y_\nu^\dagger Y_\nu \text{Tr} \left(Y_\nu Y_\nu^\dagger \right) \\
& - \frac{2}{5}Y_\nu Y_e^\dagger Y_e \left(15\text{Tr} \left(Y_d^\dagger T_d \right) + 5\text{Tr} \left(Y_e^\dagger T_e \right) + 6g_1^2 M_1 \right) - 18Y_\nu Y_\nu^\dagger Y_\nu \text{Tr} \left(Y_u^\dagger T_u \right) \\
& - 6Y_\nu Y_\nu^\dagger Y_\nu \text{Tr} \left(Y_\nu^\dagger T_\nu \right) - 3T_\nu \text{Tr} \left(Y_d Y_u^\dagger Y_u Y_d^\dagger \right) - T_\nu \text{Tr} \left(Y_e Y_\nu^\dagger Y_\nu Y_e^\dagger \right) \\
& - 9T_\nu \text{Tr} \left(Y_u Y_u^\dagger Y_u Y_u^\dagger \right) - 3T_\nu \text{Tr} \left(Y_\nu Y_\nu^\dagger Y_\nu Y_\nu^\dagger \right) \\
& - \frac{2}{25}Y_\nu \left(207g_1^4 M_1 + 45g_1^2 g_2^2 M_1 + 45g_1^2 g_2^2 M_2 + 375g_2^4 M_2 + 20 \left(20g_3^2 M_3 + g_1^2 M_1 \right) \text{Tr} \left(Y_u Y_u^\dagger \right) \right. \\
& \left. - 20 \left(20g_3^2 + g_1^2 \right) \text{Tr} \left(Y_u^\dagger T_u \right) + 75 \text{Tr} \left(Y_d Y_u^\dagger T_u Y_d^\dagger \right) + 25 \text{Tr} \left(Y_e Y_\nu^\dagger T_\nu Y_e^\dagger \right) \right. \\
& \left. + 75 \text{Tr} \left(Y_u Y_d^\dagger T_d Y_u^\dagger \right) + 450 \text{Tr} \left(Y_u Y_u^\dagger T_u Y_u^\dagger \right) + 25 \text{Tr} \left(Y_\nu Y_e^\dagger T_e Y_\nu^\dagger \right) + 150 \text{Tr} \left(Y_\nu Y_\nu^\dagger T_\nu Y_\nu^\dagger \right) \right)
\end{aligned} \tag{B.24}$$

Bilinear soft-breaking parameters

$$\Delta\beta_{B_\mu}^{(1)} = 2\mu \text{Tr} \left(Y_\nu^\dagger T_\nu \right) + B_\mu \text{Tr} \left(Y_\nu Y_\nu^\dagger \right) \tag{B.25}$$

$$\begin{aligned}
\Delta\beta_{B_\mu}^{(2)} &= -B_\mu \left(2\text{Tr} \left(Y_e Y_\nu^\dagger Y_\nu Y_e^\dagger \right) + 3\text{Tr} \left(Y_\nu Y_\nu^\dagger Y_\nu Y_\nu^\dagger \right) \right) \\
& - 4\mu \left(3\text{Tr} \left(Y_\nu Y_\nu^\dagger T_\nu Y_\nu^\dagger \right) + \text{Tr} \left(Y_e Y_\nu^\dagger T_\nu Y_e^\dagger \right) + \text{Tr} \left(Y_\nu Y_e^\dagger T_e Y_\nu^\dagger \right) \right)
\end{aligned} \tag{B.26}$$

$$\beta_{B_{\mu_X}}^{(1)} = \beta_{B_{\mu_X}}^{(2)} = 0 \tag{B.27}$$

$$\beta_{B_{M_R}}^{(1)} = 2 \left(2T_\nu Y_\nu^\dagger M_R + Y_\nu Y_\nu^\dagger B_{M_R} \right) \tag{B.28}$$

$$\begin{aligned}
\beta_{B_{M_R}}^{(2)} &= -\frac{2}{5} \left(-6g_1^2 T_\nu Y_\nu^\dagger M_R - 30g_2^2 T_\nu Y_\nu^\dagger M_R + 5Y_\nu Y_e^\dagger Y_e Y_\nu^\dagger B_{M_R} + 10Y_\nu Y_e^\dagger T_e Y_\nu^\dagger M_R \right. \\
& + 5Y_\nu Y_\nu^\dagger Y_\nu Y_\nu^\dagger B_{M_R} + 10Y_\nu Y_\nu^\dagger T_\nu Y_\nu^\dagger M_R + 10T_\nu Y_\nu^\dagger Y_e Y_\nu^\dagger M_R + 10T_\nu Y_\nu^\dagger Y_\nu Y_\nu^\dagger M_R \\
& + 30T_\nu Y_\nu^\dagger M_R \text{Tr} \left(Y_u Y_u^\dagger \right) + 10T_\nu Y_\nu^\dagger M_R \text{Tr} \left(Y_\nu Y_\nu^\dagger \right) \\
& + Y_\nu Y_\nu^\dagger B_{M_R} \left(15\text{Tr} \left(Y_u Y_u^\dagger \right) - 3 \left(5g_2^2 + g_1^2 \right) + 5\text{Tr} \left(Y_\nu Y_\nu^\dagger \right) \right) \\
& \left. + 2Y_\nu Y_\nu^\dagger M_R \left(15g_2^2 M_2 + 15\text{Tr} \left(Y_u^\dagger T_u \right) + 3g_1^2 M_1 + 5\text{Tr} \left(Y_\nu^\dagger T_\nu \right) \right) \right)
\end{aligned} \tag{B.29}$$

Soft-breaking scalar masses

The RGEs of the soft SUSY breaking masses are usually written in terms of a set of traces (see e.g. [126]). In the model considered here, only one changes with respect to the MSSM:

$$\Delta\sigma_{3,1} = \frac{1}{20}\frac{1}{\sqrt{15}}g_1\left(-30m_{H_u}^2\text{Tr}\left(Y_\nu Y_\nu^\dagger\right) + 30\text{Tr}\left(Y_\nu m_l^{2*} Y_\nu^\dagger\right)\right) \quad (\text{B.30})$$

The resulting RGEs are:

$$\begin{aligned} \Delta\beta_{m_q^2}^{(2)} &= -\left(2T_u^\dagger T_u + 2Y_u^\dagger m_u^2 Y_u + 4m_{H_u}^2 Y_u^\dagger Y_u + m_q^2 Y_u^\dagger Y_u + Y_u^\dagger Y_u m_q^2\right)\text{Tr}\left(Y_\nu Y_\nu^\dagger\right) \\ &\quad - 2\left(T_u^\dagger Y_u \text{Tr}\left(Y_\nu^\dagger T_\nu\right) + Y_u^\dagger T_u \text{Tr}\left(T_\nu^* Y_\nu^T\right) + Y_u^\dagger Y_u \text{Tr}\left(T_\nu^* T_\nu^T\right)\right. \\ &\quad \left.+ Y_u^\dagger Y_u \text{Tr}\left(m_l^2 Y_\nu^\dagger Y_\nu\right) + Y_u^\dagger Y_u \text{Tr}\left(m_\nu^2 Y_\nu Y_\nu^\dagger\right)\right) \end{aligned} \quad (\text{B.31})$$

$$\Delta\beta_{m_l^2}^{(1)} = 2m_{H_u}^2 Y_\nu^\dagger Y_\nu + 2T_\nu^\dagger T_\nu + 2Y_\nu^\dagger m_\nu^2 Y_\nu + m_l^2 Y_\nu^\dagger Y_\nu + Y_\nu^\dagger Y_\nu m_l^2 \quad (\text{B.32})$$

$$\begin{aligned} \Delta\beta_{m_l^2}^{(2)} &= -3\left(2T_\nu^\dagger T_\nu + 2Y_\nu^\dagger m_\nu^2 Y_\nu + 4m_{H_u}^2 Y_\nu^\dagger Y_\nu + m_l^2 Y_\nu^\dagger Y_\nu + Y_\nu^\dagger Y_\nu m_l^2\right)\text{Tr}\left(Y_u Y_u^\dagger\right) \\ &\quad - \left(2T_\nu^\dagger T_\nu + 2Y_\nu^\dagger m_\nu^2 Y_\nu + 4m_{H_u}^2 Y_\nu^\dagger Y_\nu + m_l^2 Y_\nu^\dagger Y_\nu + Y_\nu^\dagger Y_\nu m_l^2\right)\text{Tr}\left(Y_\nu Y_\nu^\dagger\right) \\ &\quad - 2\left(4m_{H_u}^2 Y_\nu^\dagger Y_\nu Y_\nu^\dagger Y_\nu + 2Y_\nu^\dagger Y_\nu T_\nu^\dagger T_\nu + 2Y_\nu^\dagger T_\nu T_\nu^\dagger Y_\nu + 2T_\nu^\dagger Y_\nu Y_\nu^\dagger T_\nu\right. \\ &\quad \left.+ 2T_\nu^\dagger T_\nu Y_\nu^\dagger Y_\nu + m_l^2 Y_\nu^\dagger Y_\nu Y_\nu^\dagger Y_\nu + 2Y_\nu^\dagger m_\nu^2 Y_\nu Y_\nu^\dagger Y_\nu + 2Y_\nu^\dagger Y_\nu m_l^2 Y_\nu^\dagger Y_\nu\right. \\ &\quad \left.+ 2Y_\nu^\dagger Y_\nu Y_\nu^\dagger m_\nu^2 Y_\nu + Y_\nu^\dagger Y_\nu Y_\nu^\dagger Y_\nu m_l^2 + 3T_\nu^\dagger Y_\nu \text{Tr}\left(Y_\nu^\dagger T_u\right) + T_\nu^\dagger Y_\nu \text{Tr}\left(Y_\nu^\dagger T_\nu\right)\right. \\ &\quad \left.+ 3Y_\nu^\dagger T_\nu \text{Tr}\left(T_u^* Y_u^T\right) + 3Y_\nu^\dagger Y_\nu \text{Tr}\left(T_u^* T_u^T\right) + Y_\nu^\dagger T_\nu \text{Tr}\left(T_\nu^* Y_\nu^T\right)\right. \\ &\quad \left.+ Y_\nu^\dagger Y_\nu \text{Tr}\left(T_\nu^* T_\nu^T\right) + Y_\nu^\dagger Y_\nu \text{Tr}\left(m_l^2 Y_\nu^\dagger Y_\nu\right) + 3Y_\nu^\dagger Y_\nu \text{Tr}\left(m_q^2 Y_u^\dagger Y_u\right)\right. \\ &\quad \left.+ 3Y_\nu^\dagger Y_\nu \text{Tr}\left(m_u^2 Y_u Y_u^\dagger\right) + Y_\nu^\dagger Y_\nu \text{Tr}\left(m_\nu^2 Y_\nu Y_\nu^\dagger\right)\right) \end{aligned} \quad (\text{B.33})$$

$$\begin{aligned} \Delta\beta_{m_{H_d}^2}^{(2)} &= -2\left(\left(m_{H_d}^2 + m_{H_u}^2\right)\text{Tr}\left(Y_e Y_\nu^\dagger Y_\nu Y_e^\dagger\right) + \text{Tr}\left(Y_e Y_\nu^\dagger T_\nu T_e^\dagger\right) + \text{Tr}\left(Y_e T_\nu^\dagger T_\nu Y_e^\dagger\right) + \text{Tr}\left(Y_\nu Y_e^\dagger T_e T_\nu^\dagger\right)\right. \\ &\quad \left.+ \text{Tr}\left(Y_\nu T_e^\dagger T_e Y_\nu^\dagger\right) + \text{Tr}\left(m_e^2 Y_e Y_\nu^\dagger Y_\nu Y_e^\dagger\right) + \text{Tr}\left(m_l^2 Y_e^\dagger Y_e Y_\nu^\dagger Y_\nu\right) + \text{Tr}\left(m_l^2 Y_\nu^\dagger Y_\nu Y_e^\dagger Y_e\right)\right. \\ &\quad \left.+ \text{Tr}\left(m_\nu^2 Y_\nu Y_e^\dagger Y_e Y_\nu^\dagger\right)\right) \end{aligned} \quad (\text{B.34})$$

$$\Delta\beta_{m_{H_u}^2}^{(1)} = 2\left(m_{H_u}^2 \text{Tr}\left(Y_\nu Y_\nu^\dagger\right) + \text{Tr}\left(T_\nu^* T_\nu^T\right) + \text{Tr}\left(m_l^2 Y_\nu^\dagger Y_\nu\right) + \text{Tr}\left(m_\nu^2 Y_\nu Y_\nu^\dagger\right)\right) \quad (\text{B.35})$$

$$\begin{aligned} \Delta\beta_{m_{H_u}^2}^{(2)} &= -2\left(\left(m_{H_d}^2 + m_{H_u}^2\right)\text{Tr}\left(Y_e Y_\nu^\dagger Y_\nu Y_e^\dagger\right) + \text{Tr}\left(Y_e Y_\nu^\dagger T_\nu T_e^\dagger\right) + \text{Tr}\left(Y_e T_\nu^\dagger T_\nu Y_e^\dagger\right) + \text{Tr}\left(Y_\nu Y_e^\dagger T_e T_\nu^\dagger\right)\right. \\ &\quad \left.+ 6m_{H_u}^2 \text{Tr}\left(Y_\nu Y_\nu^\dagger Y_\nu Y_\nu^\dagger\right) + 6\text{Tr}\left(Y_\nu Y_\nu^\dagger T_\nu T_\nu^\dagger\right) + \text{Tr}\left(Y_\nu T_e^\dagger T_e Y_\nu^\dagger\right) + 6\text{Tr}\left(Y_\nu T_\nu^\dagger T_\nu Y_\nu^\dagger\right)\right. \\ &\quad \left.+ \text{Tr}\left(m_e^2 Y_e Y_\nu^\dagger Y_\nu Y_e^\dagger\right) + \text{Tr}\left(m_l^2 Y_e^\dagger Y_e Y_\nu^\dagger Y_\nu\right) + \text{Tr}\left(m_l^2 Y_\nu^\dagger Y_\nu Y_e^\dagger Y_e\right) + 6\text{Tr}\left(m_l^2 Y_\nu^\dagger Y_\nu Y_\nu^\dagger Y_\nu\right)\right. \\ &\quad \left.+ \text{Tr}\left(m_\nu^2 Y_\nu Y_e^\dagger Y_e Y_\nu^\dagger\right) + 6\text{Tr}\left(m_\nu^2 Y_\nu Y_\nu^\dagger Y_\nu Y_\nu^\dagger\right)\right) \end{aligned} \quad (\text{B.36})$$

$$\begin{aligned} \Delta\beta_{m_u^2}^{(2)} &= -2\left(\left(2T_u^\dagger T_u + 2Y_u m_q^2 Y_u^\dagger + 4m_{H_u}^2 Y_u^\dagger Y_u + m_u^2 Y_u^\dagger Y_u + Y_u^\dagger Y_u m_u^2\right)\text{Tr}\left(Y_\nu Y_\nu^\dagger\right)\right. \\ &\quad \left.+ 2\left(Y_u^\dagger T_u \text{Tr}\left(Y_\nu^\dagger T_\nu\right) + T_u^\dagger Y_u \text{Tr}\left(T_\nu^* Y_\nu^T\right) + Y_u^\dagger Y_u \text{Tr}\left(T_\nu^* T_\nu^T\right)\right.\right. \\ &\quad \left.\left.+ Y_u^\dagger Y_u \text{Tr}\left(m_l^2 Y_\nu^\dagger Y_\nu\right) + Y_u^\dagger Y_u \text{Tr}\left(m_\nu^2 Y_\nu Y_\nu^\dagger\right)\right)\right) \end{aligned} \quad (\text{B.37})$$

$$\begin{aligned} \Delta\beta_{m_e^2}^{(2)} &= -2\left(2\left(m_{H_d}^2 + m_{H_u}^2\right)Y_e Y_\nu^\dagger Y_\nu Y_e^\dagger + 2Y_e Y_\nu^\dagger T_\nu T_e^\dagger + 2Y_e T_\nu^\dagger T_\nu Y_e^\dagger + 2T_e Y_\nu^\dagger Y_\nu T_e^\dagger\right. \\ &\quad \left.+ 2T_e T_\nu^\dagger Y_\nu Y_e^\dagger + m_e^2 Y_e Y_\nu^\dagger Y_\nu Y_e^\dagger + 2Y_e m_l^2 Y_\nu^\dagger Y_\nu Y_e^\dagger\right. \\ &\quad \left.+ 2Y_e Y_\nu^\dagger m_\nu^2 Y_\nu Y_e^\dagger + 2Y_e Y_\nu^\dagger Y_\nu m_l^2 Y_e^\dagger + Y_e Y_\nu^\dagger Y_\nu Y_e^\dagger m_e^2\right) \end{aligned} \quad (\text{B.38})$$

$$\beta_{m_\nu^2}^{(1)} = 2 \left(2m_{H_u}^2 Y_\nu Y_\nu^\dagger + 2T_\nu T_\nu^\dagger + 2Y_\nu m_l^2 Y_\nu^\dagger + m_\nu^2 Y_\nu Y_\nu^\dagger + Y_\nu Y_\nu^\dagger m_\nu^2 \right) \quad (\text{B.39})$$

$$\begin{aligned} \beta_{m_\nu^2}^{(2)} = & -\frac{2}{5} \left(6g_1^2 M_1^* T_\nu Y_\nu^\dagger + 30g_2^2 M_2^* T_\nu Y_\nu^\dagger - 6g_1^2 T_\nu T_\nu^\dagger - 30g_2^2 T_\nu T_\nu^\dagger \right. \\ & - 3g_1^2 m_\nu^2 Y_\nu Y_\nu^\dagger - 15g_2^2 m_\nu^2 Y_\nu Y_\nu^\dagger - 6g_1^2 Y_\nu m_l^2 Y_\nu^\dagger - 30g_2^2 Y_\nu m_l^2 Y_\nu^\dagger \\ & - 3g_1^2 Y_\nu Y_\nu^\dagger m_\nu^2 - 15g_2^2 Y_\nu Y_\nu^\dagger m_\nu^2 + 10m_{H_d}^2 Y_\nu Y_e^\dagger Y_e Y_\nu^\dagger \\ & + 10m_{H_u}^2 Y_\nu Y_e^\dagger Y_e Y_\nu^\dagger + 10Y_\nu Y_e^\dagger T_e T_\nu^\dagger + 20m_{H_u}^2 Y_\nu Y_\nu^\dagger Y_\nu Y_\nu^\dagger + 10Y_\nu Y_\nu^\dagger T_\nu T_\nu^\dagger \\ & + 10Y_\nu T_e^\dagger T_e Y_\nu^\dagger + 10Y_\nu T_\nu^\dagger T_\nu Y_\nu^\dagger + 10T_\nu Y_e^\dagger Y_e T_\nu^\dagger + 10T_\nu Y_\nu^\dagger Y_\nu T_\nu^\dagger \\ & + 10T_\nu T_e^\dagger Y_e Y_\nu^\dagger + 10T_\nu T_\nu^\dagger Y_\nu Y_\nu^\dagger + 5m_\nu^2 Y_\nu Y_e^\dagger Y_e Y_\nu^\dagger + 5m_\nu^2 Y_\nu Y_\nu^\dagger Y_\nu Y_\nu^\dagger \\ & + 10Y_\nu m_l^2 Y_e^\dagger Y_e Y_\nu^\dagger + 10Y_\nu m_l^2 Y_\nu^\dagger Y_\nu Y_\nu^\dagger + 10Y_\nu Y_e^\dagger m_e^2 Y_e Y_\nu^\dagger \\ & + 10Y_\nu Y_e^\dagger Y_e m_l^2 Y_\nu^\dagger + 5Y_\nu Y_e^\dagger Y_e Y_\nu^\dagger m_\nu^2 + 10Y_\nu Y_\nu^\dagger m_\nu^2 Y_\nu Y_\nu^\dagger + 10Y_\nu Y_\nu^\dagger Y_\nu m_l^2 Y_\nu^\dagger \\ & + 5Y_\nu Y_\nu^\dagger Y_\nu Y_\nu^\dagger m_\nu^2 + 30T_\nu T_\nu^\dagger \text{Tr}(Y_u Y_u^\dagger) + 15m_\nu^2 Y_\nu Y_\nu^\dagger \text{Tr}(Y_u Y_u^\dagger) \\ & + 30Y_\nu m_l^2 Y_\nu^\dagger \text{Tr}(Y_u Y_u^\dagger) + 15Y_\nu Y_\nu^\dagger m_\nu^2 \text{Tr}(Y_u Y_u^\dagger) + 10T_\nu T_\nu^\dagger \text{Tr}(Y_\nu Y_\nu^\dagger) \\ & + 5m_\nu^2 Y_\nu Y_\nu^\dagger \text{Tr}(Y_\nu Y_\nu^\dagger) + 10Y_\nu m_l^2 Y_\nu^\dagger \text{Tr}(Y_\nu Y_\nu^\dagger) + 5Y_\nu Y_\nu^\dagger m_\nu^2 \text{Tr}(Y_\nu Y_\nu^\dagger) \\ & + 2Y_\nu T_\nu^\dagger \left(15g_2^2 M_2 + 15\text{Tr}(Y_u^\dagger T_u) + 3g_1^2 M_1 + 5\text{Tr}(Y_\nu^\dagger T_\nu) \right) + 30T_\nu Y_\nu^\dagger \text{Tr}(T_u^* Y_u^T) \\ & + 10T_\nu Y_\nu^\dagger \text{Tr}(T_\nu^* Y_\nu^T) \\ & + \frac{4}{5} Y_\nu Y_\nu^\dagger \left(3g_1^2 m_{H_u}^2 + 15g_2^2 m_{H_u}^2 + 6g_1^2 |M_1|^2 + 30g_2^2 |M_2|^2 - 30m_{H_u}^2 \text{Tr}(Y_u Y_u^\dagger) - 10m_{H_u}^2 \text{Tr}(Y_\nu Y_\nu^\dagger) \right. \\ & \left. - 15\text{Tr}(T_u^* T_u^T) - 5\text{Tr}(T_\nu^* T_\nu^T) - 5\text{Tr}(m_l^2 Y_\nu^\dagger Y_\nu) - 15\text{Tr}(m_q^2 Y_u^\dagger Y_u) - 15\text{Tr}(m_u^2 Y_u Y_u^\dagger) \right. \\ & \left. - 5\text{Tr}(m_\nu^2 Y_\nu Y_\nu^\dagger) \right) \end{aligned} \quad (\text{B.40})$$

$$\beta_{m_X^2}^{(1)} = \beta_{m_X^2}^{(2)} = 0 \quad (\text{B.41})$$

Vacuum expectation values

$$\Delta\beta_{v_d}^{(2)} = v_d \text{Tr}(Y_e Y_\nu^\dagger Y_\nu Y_e^\dagger) \quad (\text{B.42})$$

$$\Delta\beta_{v_u}^{(1)} = -v_u \text{Tr}(Y_\nu Y_\nu^\dagger) \quad (\text{B.43})$$

$$\Delta\beta_{v_u}^{(2)} = 3v_u \text{Tr}(Y_\nu Y_\nu^\dagger Y_\nu Y_\nu^\dagger) - \frac{3}{10} \left(5g_2^2 + g_1^2 \right) v_u \xi \text{Tr}(Y_\nu Y_\nu^\dagger) + v_u \text{Tr}(Y_e Y_\nu^\dagger Y_\nu Y_e^\dagger) \quad (\text{B.44})$$

C Loop integrals

The B -functions with vanishing external momenta and the arguments (a, b) are given by

$$B_0 = 1 - \log\left(\frac{b}{Q^2}\right) + \frac{1}{b-a} \left[a \log\left(\frac{a}{b}\right) \right], \quad (\text{C.1})$$

$$B_1 = -\frac{1}{2} + \frac{1}{2} \log\left(\frac{b}{Q^2}\right) - \frac{1}{4(a-b)^2} \left[a^2 - b^2 + 2a^2 \log\left(\frac{b}{a}\right) \right], \quad (\text{C.2})$$

The C -functions with vanishing external momenta and the arguments (a, b, c) read

$$C_0 = -\frac{1}{(a-b)(a-c)(b-c)} \times$$

$$\times \left[b(c-a) \log\left(\frac{b}{a}\right) + c(a-b) \log\left(\frac{c}{a}\right) \right] \quad (\text{C.3})$$

$$C_{00} = \frac{1}{8(a-b)(a-c)(b-c)} \times \\ \times \left[(c-a) \left((a-b)(2 \log\left(\frac{a}{Q^2}\right) - 3(b-c) - 2b^2 \log\left(\frac{b}{a}\right)) + 2c^2(b-a) \log\left(\frac{c}{a}\right) \right) \right] \quad (\text{C.4})$$

$$C_1 = -\frac{1}{2(a-b)^2(a-c)(b-c)^2} \times \left[c^2(a-b)^2 \log\left(\frac{c}{a}\right) \right. \\ \left. + b(c-a) \left((b-a)(b-c) - (a(b-2c) + bc) \log\left(\frac{b}{a}\right) \right) \right] \quad (\text{C.5})$$

$$C_2 = -\frac{1}{2(a-b)(a-c)^2(b-c)^2} \times \left[a^2(b-c)^2 \log\left(\frac{b}{a}\right) \right. \quad (\text{C.6})$$

$$\left. + c(b-a) \left((a-c)(b-c) + (c(a+b) - 2ab) \log\left(\frac{b}{c}\right) \right) \right] \quad (\text{C.7})$$

$$C_{11} = -\frac{1}{6(a-b)^3(a-c)(b-c)^3} \times \\ \times \left[b(a-c) \left(-2(a^2(b^2 - 3bc + 3c^2) + abc(b-3c) + b^2c^2) \log\left(\frac{b}{a}\right) \right. \right. \\ \left. \left. - (b-a)(b-c) (-3ab + 5ac + b^2 - 3bc) \right) + 2c^3(a-b)^3 \log\left(\frac{c}{a}\right) \right] \quad (\text{C.8})$$

$$C_{12} = \frac{1}{6(a-b)^2(a-c)^2(b-c)^3} \times \left[(a-b) \left((a-c)(b-c)(a(b^2 + c^2) \right. \right. \\ \left. \left. - bc(b+c) \right) + c^2(a-b)(3ab - c(a+2b)) \log\left(\frac{c}{a}\right) \right. \\ \left. + b^2(a-c)^2(a(b-3c) + 2bc) \log\left(\frac{b}{a}\right) \right] \quad (\text{C.9})$$

$$C_{22} = \frac{1}{6(a-b)(a-c)^3(b-c)^3} \times \left[2a^3(b-c)^3 \log\left(\frac{b}{a}\right) \right. \\ \left. + c(a-b) \left(2(c^2(a^2 + ab + b^2) + 3a^2b^2 - 3abc(a+b)) \log\left(\frac{b}{c}\right) \right. \right. \\ \left. \left. - (a-c)(b-c) (-3c(a+b) + 5ab + c^2) \right) \right] \quad (\text{C.10})$$

In the case of external photons, often the same combinations of C -functions appear. If the arguments are (a, b, b) , these can be expressed as

$$C_2 + C_{12} + C_{22} = C_1 + C_{12} + C_{11} = \frac{1}{12b(x-1)^4} ((x-1)(x(2x+5)-1) - 6x^2 \log(x)) \quad (\text{C.11})$$

$$C_0 + C_1 + C_2 = \frac{1}{2b(x-1)^3} (-x^2 + 2x \log(x) + 1) \quad (\text{C.12})$$

$$2C_{12} + 2C_{11} - C_2 = 2C_{12} + 2C_{22} - C_1 = \frac{1}{12b(x-1)^4} (6(1-3x)x^2 \log(x) + (x-1)(x(31x-26)+7)) \quad (\text{C.13})$$

$$C_1 + C_2 = \frac{1}{2b(x-1)^3} (2x^2 \log(x) + (4-3x)x - 1) \quad (\text{C.14})$$

and for (a, a, b) we get

$$C_2 + C_{12} + C_{22} = -C_{12} = \frac{1}{12b(x-1)^4} (x((x-6)x+3) + 6x\log(x) + 2) \quad (\text{C.15})$$

$$C_0 + C_1 + C_2 = -C_1 = -\frac{1}{4b(x-1)^3} (x^2 - 4x + 2\log(x) + 3) \quad (\text{C.16})$$

In the previous expressions we used $x = a/b$.

For the photonic monopole operators we define special loop functions

$$M_{SFF}(a, b) = \frac{((a-b)(16a^2 - 29ab + 7b^2) + 6a^2(2a-3b)\log(\frac{b}{a}))}{36(a-b)^4} \quad (\text{C.17})$$

$$M_{FSS}(a, b) = \frac{6a^3\log(\frac{b}{a}) + 11a^3 - 18a^2b + 9ab^2 - 2b^3}{36(a-b)^4} \quad (\text{C.18})$$

$$M_{FSV}(a, b) = \frac{\sqrt{a}(2a^3 + 3a^2b + 6a^2b\log(\frac{b}{a}) - 6ab^2 + b^3)}{12b(a-b)^4} \quad (\text{C.19})$$

$$M_{FVS}(a, b) = \frac{\sqrt{a}(2a^3 + 3a^2b + 6a^2b\log(\frac{b}{a}) - 6ab^2 + b^3)}{12b(a-b)^4} \quad (\text{C.20})$$

$$M_{FVV}(a, b) = \frac{6a^2(a-3b)\log(\frac{a}{b}) - (a-b)(5a^2 - 22ab + 5b^2)}{9(a-b)^4} \quad (\text{C.21})$$

The necessary box functions with the arguments (a, b, c, d) read, in the limit of vanishing external momenta,

$$D_0 = - \left[\frac{b\log\frac{b}{a}}{(b-a)(b-c)(b-d)} + \frac{c\log\frac{c}{a}}{(c-a)(c-b)(c-d)} \right. \\ \left. + \frac{d\log\frac{d}{a}}{(d-a)(d-b)(d-c)} \right] \quad (\text{C.22})$$

$$D_{27} = -\frac{1}{4} \left[\frac{b^2\log\frac{b}{a}}{(b-a)(b-c)(b-d)} + \frac{c^2\log\frac{c}{a}}{(c-a)(c-b)(c-d)} \right. \\ \left. + \frac{d^2\log\frac{d}{a}}{(d-a)(d-b)(d-c)} \right] \quad (\text{C.23})$$


In addition, we define

$$I_{C_0 D_0}(a, b, c, d) = C_0(a, b, c) + dD_0(a, b, c, d) \quad (\text{C.24})$$

D Photonic penguin contributions to LFV

In the following appendices we present our results for the form factors of the operators involved in our computation, done in the mass basis. The flavor of the external fermions will be denoted with Greek characters ($\alpha, \beta, \gamma, \delta$), whereas the mass eigenstates of the particles in the loops will be denoted with Latin characters (a, b, c, d). A sum over repeated indices will be assumed.

D.1 Feynman diagrams



We give in the following the contribution of each diagram to the different operators. We indicate the diagram by the corresponding index (a_i) with $i = 1, \dots, 7$.

D.2 Neutralino contributions

$$C_i = C_i(m_{\tilde{\chi}_a^0}^2, m_{\tilde{e}_c}^2, m_{\tilde{e}_b}^2) \quad (\text{D.1})$$

$$\begin{aligned} A_{(a_1)}^{2L} &= 2F_{c,b}^{\tilde{e}}(\bar{N}_{\alpha,a,b}^L N_{a,\beta,c}^R(C_{12} + C_{22} + C_2)m_{\ell_\alpha} + \bar{N}_{\alpha,a,b}^R(N_{a,\beta,c}^L(C_{11} \\ &\quad + C_{12} + C_1)m_{\ell_\beta} - N_{a,\beta,c}^R(C_0 + C_1 + C_2)m_{\tilde{\chi}_a^0})) \end{aligned} \quad (\text{D.2})$$

$$A_{(a_1)}^{1L} = -\bar{N}_{\alpha,a,b}^R N_{a,\beta,c}^L F_{c,b}^{\tilde{e}} M_{FSS}(m_{\tilde{\chi}_a^0}^2, m_{\tilde{e}_b}^2) \quad (\text{D.3})$$

D.3 Chargino contributions

$$C_i = C_i(m_{\tilde{\chi}_c^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^i}^2) \quad (\text{D.4})$$

$$\begin{aligned} A_{(a_2)}^{2L} &= -2\left(\bar{X}_{\alpha,b,a}^L X_{c,\beta,a}^R(F_{b,c}^{c,L})^* C_{12} m_{\ell_\alpha} - \bar{X}_{\alpha,b,a}^R(X_{c,\beta,a}^L(F_{b,c}^{c,R})^*(C_{12} + C_{22} + C_2)m_{\ell_\beta} \right. \\ &\quad \left. + X_{c,\beta,a}^R((F_{b,c}^{c,L})^* C_1 m_{\tilde{\chi}_b^-} - (F_{b,c}^{c,R})^*(C_0 + C_1 + C_2)m_{\tilde{\chi}_c^-}))\right) \end{aligned} \quad (\text{D.5})$$

$$A_{(a_2)}^{1L} = -\bar{X}_{\alpha,b,a}^R X_{c,\beta,a}^L(F_{b,c}^{c,R})^* M_{SFF}(m_{\nu_a^i}^2, m_{\tilde{\chi}_b^-}^2) \quad (\text{D.6})$$

$$C_i = C_i(m_{\tilde{\chi}_c^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^R}^2) \quad (\text{D.7})$$

$$\begin{aligned} A_{(a_3)}^{2L} &= -2\left(\hat{X}_{\alpha,b,a}^L \hat{X}_{c,\beta,a}^R(F_{b,c}^{c,L})^* C_{12} m_{\ell_\alpha} - \hat{X}_{\alpha,b,a}^R(\hat{X}_{c,\beta,a}^L(F_{b,c}^{c,R})^*(C_{12} + C_{22} + C_2)m_{\ell_\beta} \right. \\ &\quad \left. + \hat{X}_{c,\beta,a}^R((F_{b,c}^{c,L})^* C_1 m_{\tilde{\chi}_b^-} - (F_{b,c}^{c,R})^*(C_0 + C_1 + C_2)m_{\tilde{\chi}_c^-}))\right) \end{aligned} \quad (\text{D.8})$$

$$A_{(a_3)}^{1L} = -\hat{X}_{\alpha,b,a}^R \hat{X}_{c,\beta,a}^L(F_{b,c}^{c,R})^* M_{SFF}(m_{\nu_a^R}^2, m_{\tilde{\chi}_b^-}^2) \quad (\text{D.9})$$

D.4 W^+ and H^+ contributions

$$C_i = C_i(m_{\nu_a}^2, m_{H_c^-}^2, m_{H_b^-}^2) \quad (\text{D.10})$$

$$\begin{aligned} A_{(a_4)}^{2L} &= 2F_{c,b}^h(\bar{V}_{\alpha,a,b}^{+,L} V_{a,\beta,c}^{+,R}(C_{12} + C_{22} + C_2)m_{\ell_\alpha} + \bar{V}_{\alpha,a,b}^{+,R}(V_{a,\beta,c}^{+,L}(C_{11} + C_{12} + C_1)m_{\ell_\beta} \\ &\quad - V_{a,\beta,c}^{+,R}(C_0 + C_1 + C_2)m_{\nu_a})) \end{aligned} \quad (\text{D.11})$$

$$A_{(a_4)}^{1L} = -\bar{V}_{\alpha,a,b}^{+,R} V_{a,\beta,c}^{+,L} F_{c,b}^h M_{FSS}(m_{\nu_a}^2, m_{H_b^-}^2) \quad (\text{D.12})$$

$$A_{(a_5)}^{2L} = 2(\hat{V}_{\alpha,a}^{+,L})^* V_{a,\beta,c}^{+,R} F_c^{hw} C_2(m_{\nu_a}^2, m_{H_c^-}^2, m_{W^-}^2) \quad (\text{D.13})$$

$$A_{(a_5)}^{1L} = (\hat{V}_{\alpha,a}^{+,R})^* V_{a,\beta,c}^{+,L} F_c^{hw} M_{FVS}(m_{\nu_a}^2, m_{W^-}^2) \quad (\text{D.14})$$

$$A_{(a_6)}^{2L} = 2\bar{V}_{\alpha,a,b}^{+,R}(\hat{V}_{a,\beta}^{+,R})^* \bar{F}_b^{hw} C_1(m_{\nu_a}^2, m_{W^-}^2, m_{H_b^-}^2) \quad (\text{D.15})$$

$$A_{(a_6)}^{1L} = \bar{V}_{\alpha,a,b}^{+,L}(\hat{V}_{a,\beta}^{+,L})^* \bar{F}_b^{hw} M_{FSV}(m_{\nu_a}^2, m_{H_b^-}^2) \quad (\text{D.16})$$

$$C_i = C_i(m_{\nu_a}^2, m_{W^-}^2, m_{W^-}^2) \quad (\text{D.17})$$

$$A_{(a_7)}^{2L} = -2F^w\left((\hat{V}_{\alpha,a}^{+,R})^* (\hat{V}_{a,\beta}^{+,R})^*(2C_{12} - C_1 + 2C_{22})m_{\ell_\alpha}\right)$$

$$+ (\hat{V}_{\alpha,a}^{+,L})^* ((\hat{V}_{a,\beta}^{+,L})^* (2C_{11} + 2C_{12} - C_2)m_{\ell_\beta} + 3(\hat{V}_{a,\beta}^{+,R})^* (C_1 + C_2)m_{\nu_a}) \Big) \quad (\text{D.18})$$

$$A_{(a_7)}^{1L} = (\hat{V}_{\alpha,a}^{+,R})^* (\hat{V}_{a,\beta}^{+,R})^* F^w M_{FVV}(m_{\nu_a}^2, m_{W^-}^2) \quad (\text{D.19})$$

These coefficients are related to the ones used in the calculation of the flavor observables by

$$K_L^1 = \frac{1}{e} \sum_p A_{(a_p)}^{1L} \quad (\text{D.20})$$

$$K_L^2 = -\frac{1}{2e m_{\ell_\alpha}} \sum_p A_{(a_p)}^{2L} \quad (\text{D.21})$$


E Z and Higgs penguin contributions to LFV

E.1 Feynman diagrams


In the following $B = Z, h_p, A_p^0$ is used.

Neutralino diagrams

Self energy corrections




Vertex corrections





Chargino diagrams

Self energy corrections




Vertex corrections






W^+ and H^+ diagrams

Self energy corrections



Vertex corrections



E.2 Neutralino contributions

E.2.1 Z-penguins

Self-energy corrections.

$$I_1 = B_0(m_{\tilde{\chi}_a^0}^2, m_{\tilde{e}_b}^2) \quad (\text{E.1})$$

$$I_2 = B_1(m_{\tilde{\chi}_a^0}^2, m_{\tilde{e}_b}^2) \quad (\text{E.2})$$

$$\begin{aligned} V_{Z,(n_1,n_2)}^{LL} = & (E_{\beta,c}^L(N_{a,\alpha,b}^L \bar{N}_{c,a,b}^R I_2 m_{\ell_\alpha}^2 - N_{a,\alpha,b}^R \bar{N}_{c,a,b}^R I_1 m_{\ell_\alpha} m_{\tilde{\chi}_a^0} + N_{a,\alpha,b}^R \bar{N}_{c,a,b}^L I_2 m_{\ell_\alpha} m_{\ell_c} \\ & - N_{a,\alpha,b}^L \bar{N}_{c,a,b}^L I_1 m_{\tilde{\chi}_a^0} m_{\ell_c})) / (m_{\ell_\alpha}^2 - m_{\ell_c}^2) \quad + (\alpha \leftrightarrow \beta) \end{aligned} \quad (\text{E.3})$$

$$\begin{aligned} V_{Z,(n_1,n_2)}^{LR} = & (E_{\beta,c}^L(N_{a,\alpha,b}^L \bar{N}_{c,a,b}^R I_2 m_{\ell_\alpha}^2 - N_{a,\alpha,b}^R \bar{N}_{c,a,b}^R I_1 m_{\ell_\alpha} m_{\tilde{\chi}_a^0} + N_{a,\alpha,b}^R \bar{N}_{c,a,b}^L I_2 m_{\ell_\alpha} m_{\ell_c} \\ & - N_{a,\alpha,b}^L \bar{N}_{c,a,b}^L I_1 m_{\tilde{\chi}_a^0} m_{\ell_c})) / (m_{\ell_\alpha}^2 - m_{\ell_c}^2) \quad + (\alpha \leftrightarrow \beta) \end{aligned} \quad (\text{E.4})$$

Vertex corrections.

$$V_{Z,(n_3)}^{LL} = -2N_{a,\alpha,b}^L \bar{N}_{\beta,a,c}^R \tilde{E}_{b,c} C_{00}(m_{\tilde{\chi}_a^0}^2, m_{\tilde{e}_c}^2, m_{\tilde{e}_b}^2) \quad (\text{E.5})$$

$$V_{Z,(n_3)}^{LR} = -2N_{a,\alpha,b}^L \bar{N}_{\beta,a,c}^R \tilde{E}_{b,c} C_{00}(m_{\tilde{\chi}_a^0}^2, m_{\tilde{e}_c}^2, m_{\tilde{e}_b}^2) \quad (\text{E.6})$$

$$I_1 = B_0(m_{\tilde{\chi}_b^0}^2, m_{\tilde{\chi}_c^0}^2) \quad (\text{E.7})$$

$$I_2 = C_{00}(m_{\tilde{\chi}_c^0}^2, m_{\tilde{\chi}_b^0}^2, m_{\tilde{e}_a}^2) \quad (\text{E.8})$$

$$I_3 = C_0(m_{\tilde{\chi}_c^0}^2, m_{\tilde{\chi}_b^0}^2, m_{\tilde{e}_a}^2) \quad (\text{E.9})$$

$$V_{Z,(n_4)}^{LL} = N_{b,\alpha,a}^L \bar{N}_{\beta,c,a}^R (-(M_{c,b}^L I_3 m_{\tilde{\chi}_b^0} m_{\tilde{\chi}_c^0}) + M_{c,b}^R (I_1 - 2I_2 + I_3 m_{\tilde{e}_a}^2)) \quad (\text{E.10})$$

$$V_{Z,(n_4)}^{LR} = N_{b,\alpha,a}^L \bar{N}_{\beta,c,a}^R (-(M_{c,b}^L I_3 m_{\tilde{\chi}_b^0} m_{\tilde{\chi}_c^0}) + M_{c,b}^R (I_1 - 2I_2 + I_3 m_{\tilde{e}_a}^2)) \quad (\text{E.11})$$

E.2.2 Scalar penguins

CP even scalars.

Self-energy corrections.

$$I_1 = B_0(m_{\tilde{\chi}_a^0}^2, m_{\tilde{e}_b}^2) \quad (\text{E.12})$$

$$I_2 = B_1(m_{\tilde{\chi}_a^0}^2, m_{\tilde{e}_b}^2) \quad (\text{E.13})$$

$$\begin{aligned} S_{h_p,(n_1,n_2)}^{LL} = & (H_{\beta,c,p}^L (-(N_{a,\alpha,b}^L \bar{N}_{c,a,b}^R I_2 m_{\ell_\alpha}^2) + N_{a,\alpha,b}^R \bar{N}_{c,a,b}^R I_1 m_{\ell_\alpha} m_{\tilde{\chi}_a^0} - N_{a,\alpha,b}^R \bar{N}_{c,a,b}^L I_2 m_{\ell_\alpha} m_{\ell_c} \\ & + N_{a,\alpha,b}^L \bar{N}_{c,a,b}^L I_1 m_{\tilde{\chi}_a^0} m_{\ell_c})) / (m_{\ell_\alpha}^2 - m_{\ell_c}^2) \quad + (\alpha \leftrightarrow \beta) \end{aligned} \quad (\text{E.14})$$

$$\begin{aligned} S_{h_p,(n_1,n_2)}^{LR} = & (H_{\beta,c,p}^L (-(N_{a,\alpha,b}^L \bar{N}_{c,a,b}^R I_2 m_{\ell_\alpha}^2) + N_{a,\alpha,b}^R \bar{N}_{c,a,b}^R I_1 m_{\ell_\alpha} m_{\tilde{\chi}_a^0} - N_{a,\alpha,b}^R \bar{N}_{c,a,b}^L I_2 m_{\ell_\alpha} m_{\ell_c} \\ & + N_{a,\alpha,b}^L \bar{N}_{c,a,b}^L I_1 m_{\tilde{\chi}_a^0} m_{\ell_c})) / (m_{\ell_\alpha}^2 - m_{\ell_c}^2) \quad + (\alpha \leftrightarrow \beta) \end{aligned} \quad (\text{E.15})$$

Vertex corrections.

$$S_{h_p,(n_3)}^{LL} = N_{a,\alpha,b}^L \bar{N}_{\beta,a,c}^L \tilde{H}_{p,b,c} C_0(m_{\tilde{\chi}_a^0}^2, m_{\tilde{e}_c}^2, m_{\tilde{e}_b}^2) m_{\tilde{\chi}_a^0} \quad (\text{E.16})$$

$$S_{h_p,(n_3)}^{LR} = N_{a,\alpha,b}^L \bar{N}_{\beta,a,c}^L \tilde{H}_{p,b,c} C_0(m_{\tilde{\chi}_a^0}^2, m_{\tilde{e}_c}^2, m_{\tilde{e}_b}^2) m_{\tilde{\chi}_a^0} \quad (\text{E.17})$$

$$I_1 = B_0(m_{\tilde{\chi}_b^0}^2, m_{\tilde{\chi}_c^0}^2) \quad (\text{E.18})$$

$$I_2 = C_0(m_{\tilde{\chi}_c^0}^2, m_{\tilde{\chi}_b^0}^2, m_{\tilde{e}_a}^2) \quad (\text{E.19})$$

$$S_{h_p,(n_4)}^{LL} = N_{b,\alpha,a}^L \bar{N}_{\beta,c,a}^L (S_{c,b,p}^L I_2 m_{\tilde{\chi}_b^0} m_{\tilde{\chi}_c^0} + S_{c,b,p}^R (I_1 + I_2 m_{\tilde{e}_a}^2)) \quad (\text{E.20})$$

$$S_{h_p,(n_4)}^{LR} = N_{b,\alpha,a}^L \bar{N}_{\beta,c,a}^L (S_{c,b,p}^L I_2 m_{\tilde{\chi}_b^0} m_{\tilde{\chi}_c^0} + S_{c,b,p}^R (I_1 + I_2 m_{\tilde{e}_a}^2)) \quad (\text{E.21})$$

CP odd scalars.

Self-energy corrections.

$$I_1 = B_0(m_{\tilde{\chi}_a^0}^2, m_{\tilde{e}_b}^2) \quad (\text{E.22})$$

$$I_2 = B_1(m_{\tilde{\chi}_a^0}^2, m_{\tilde{e}_b}^2) \quad (\text{E.23})$$

$$S_{A_p^0, (n_1, n_2)}^{LL} = (A_{\beta, c, p}^L(-(N_{a, \alpha, b}^L \bar{N}_{c, a, b}^R I_2 m_{\ell_\alpha}^2) + N_{a, \alpha, b}^R \bar{N}_{c, a, b}^R I_1 m_{\ell_\alpha} m_{\tilde{\chi}_a^0} - N_{a, \alpha, b}^R \bar{N}_{c, a, b}^L I_2 m_{\ell_\alpha} m_{\ell_c} + N_{a, \alpha, b}^L \bar{N}_{c, a, b}^L I_1 m_{\tilde{\chi}_a^0} m_{\ell_c})/(m_{\ell_\alpha}^2 - m_{\ell_c}^2)) + (\alpha \leftrightarrow \beta) \quad (\text{E.24})$$

$$S_{A_p^0, (n_1, n_2)}^{LR} = (A_{\beta, c, p}^L(-(N_{a, \alpha, b}^L \bar{N}_{c, a, b}^R I_2 m_{\ell_\alpha}^2) + N_{a, \alpha, b}^R \bar{N}_{c, a, b}^R I_1 m_{\ell_\alpha} m_{\tilde{\chi}_a^0} - N_{a, \alpha, b}^R \bar{N}_{c, a, b}^L I_2 m_{\ell_\alpha} m_{\ell_c} + N_{a, \alpha, b}^L \bar{N}_{c, a, b}^L I_1 m_{\tilde{\chi}_a^0} m_{\ell_c})/(m_{\ell_\alpha}^2 - m_{\ell_c}^2)) + (\alpha \leftrightarrow \beta) \quad (\text{E.25})$$

Vertex corrections.

$$S_{A_p^0, (n_3)}^{LL} = N_{a, \alpha, b}^L \bar{N}_{\beta, a, c}^L \tilde{A}_{p, b, c} C_0(m_{\tilde{\chi}_a^0}^2, m_{\tilde{e}_c}^2, m_{\tilde{e}_b}^2) m_{\tilde{\chi}_a^0} \quad (\text{E.26})$$

$$S_{A_p^0, (n_3)}^{LR} = N_{a, \alpha, b}^L \bar{N}_{\beta, a, c}^L \tilde{A}_{p, b, c} C_0(m_{\tilde{\chi}_a^0}^2, m_{\tilde{e}_c}^2, m_{\tilde{e}_b}^2) m_{\tilde{\chi}_a^0} \quad (\text{E.27})$$

$$I_1 = B_0(m_{\tilde{\chi}_b^0}^2, m_{\tilde{\chi}_c^0}^2) \quad (\text{E.28})$$

$$I_2 = C_0(m_{\tilde{\chi}_c^0}^2, m_{\tilde{\chi}_b^0}^2, m_{\tilde{e}_a}^2) \quad (\text{E.29})$$

$$S_{A_p^0, (n_4)}^{LL} = N_{b, \alpha, a}^L \bar{N}_{\beta, c, a}^L (P_{c, b, p}^L I_2 m_{\tilde{\chi}_b^0} m_{\tilde{\chi}_c^0} + P_{c, b, p}^R (I_1 + I_2 m_{\tilde{e}_a}^2)) \quad (\text{E.30})$$

$$S_{A_p^0, (n_4)}^{LR} = N_{b, \alpha, a}^L \bar{N}_{\beta, c, a}^L (P_{c, b, p}^L I_2 m_{\tilde{\chi}_b^0} m_{\tilde{\chi}_c^0} + P_{c, b, p}^R (I_1 + I_2 m_{\tilde{e}_a}^2)) \quad (\text{E.31})$$

E.3 Chargino contributions

E.3.1 Z-penguins

Self-energy corrections.

$$I_1 = B_0(m_{\tilde{\chi}_b^-}^2, m_{\nu_a^i}^2) \quad (\text{E.32})$$

$$I_2 = B_1(m_{\tilde{\chi}_b^-}^2, m_{\nu_a^i}^2) \quad (\text{E.33})$$

$$V_{Z, (c_1, c_3)}^{LL} = (E_{\beta, c}^L (W_{b, \alpha, a}^L \bar{X}_{c, b, a}^R I_2 m_{\ell_\alpha}^2 - W_{b, \alpha, a}^R \bar{X}_{c, b, a}^R I_1 m_{\ell_\alpha} m_{\tilde{\chi}_b^-} + W_{b, \alpha, a}^R \bar{X}_{c, b, a}^L I_2 m_{\ell_\alpha} m_{\ell_c} - W_{b, \alpha, a}^L \bar{X}_{c, b, a}^L I_1 m_{\tilde{\chi}_b^-} m_{\ell_c})/(m_{\ell_\alpha}^2 - m_{\ell_c}^2)) + (\alpha \leftrightarrow \beta) \quad (\text{E.34})$$

$$V_{Z, (c_1, c_3)}^{LR} = (E_{\beta, c}^L (W_{b, \alpha, a}^L \bar{X}_{c, b, a}^R I_2 m_{\ell_\alpha}^2 - W_{b, \alpha, a}^R \bar{X}_{c, b, a}^R I_1 m_{\ell_\alpha} m_{\tilde{\chi}_b^-} + W_{b, \alpha, a}^R \bar{X}_{c, b, a}^L I_2 m_{\ell_\alpha} m_{\ell_c} - W_{b, \alpha, a}^L \bar{X}_{c, b, a}^L I_1 m_{\tilde{\chi}_b^-} m_{\ell_c})/(m_{\ell_\alpha}^2 - m_{\ell_c}^2)) + (\alpha \leftrightarrow \beta) \quad (\text{E.35})$$

$$I_1 = B_0(m_{\tilde{\chi}_b^-}^2, m_{\nu_a^R}^2) \quad (\text{E.36})$$

$$I_2 = B_1(m_{\tilde{\chi}_b^-}^2, m_{\nu_a^R}^2) \quad (\text{E.37})$$

$$V_{Z, (c_2, c_4)}^{LL} = (E_{\beta, c}^L (\hat{X}_{b, \alpha, a}^L \hat{\bar{X}}_{c, b, a}^R I_2 m_{\ell_\alpha}^2 - \hat{X}_{b, \alpha, a}^R \hat{\bar{X}}_{c, b, a}^R I_1 m_{\ell_\alpha} m_{\tilde{\chi}_b^-} + \hat{X}_{b, \alpha, a}^R \hat{\bar{X}}_{c, b, a}^L I_2 m_{\ell_\alpha} m_{\ell_c} - \hat{X}_{b, \alpha, a}^L \hat{\bar{X}}_{c, b, a}^L I_1 m_{\tilde{\chi}_b^-} m_{\ell_c})/(m_{\ell_\alpha}^2 - m_{\ell_c}^2)) + (\alpha \leftrightarrow \beta) \quad (\text{E.38})$$

$$V_{Z, (c_2, c_4)}^{LR} = (E_{\beta, c}^L (\hat{X}_{b, \alpha, a}^L \hat{\bar{X}}_{c, b, a}^R I_2 m_{\ell_\alpha}^2 - \hat{X}_{b, \alpha, a}^R \hat{\bar{X}}_{c, b, a}^R I_1 m_{\ell_\alpha} m_{\tilde{\chi}_b^-} + \hat{X}_{b, \alpha, a}^R \hat{\bar{X}}_{c, b, a}^L I_2 m_{\ell_\alpha} m_{\ell_c} - \hat{X}_{b, \alpha, a}^L \hat{\bar{X}}_{c, b, a}^L I_1 m_{\tilde{\chi}_b^-} m_{\ell_c})/(m_{\ell_\alpha}^2 - m_{\ell_c}^2)) + (\alpha \leftrightarrow \beta) \quad (\text{E.39})$$

Vertex corrections.

$$I_1 = B_0(m_{\tilde{\chi}_b^-}^2, m_{\tilde{\chi}_c^-}^2) \quad (\text{E.40})$$

$$I_2 = C_{00}(m_{\tilde{\chi}_c^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^i}^2) \quad (\text{E.41})$$

$$I_3 = C_0(m_{\tilde{\chi}_c^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^i}^2) \quad (\text{E.42})$$

$$V_{Z,(c_5)}^{LL} = W_{b,\alpha,a}^L \bar{X}_{\beta,c,a}^R (-C_{c,b}^L I_3 m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_c^-}) + C_{c,b}^R (I_1 - 2I_2 + I_3 m_{\nu_a^i}^2) \quad (\text{E.43})$$

$$V_{Z,(c_5)}^{LR} = W_{b,\alpha,a}^L \bar{X}_{\beta,c,a}^R (-C_{c,b}^L I_3 m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_c^-}) + C_{c,b}^R (I_1 - 2I_2 + I_3 m_{\nu_a^i}^2) \quad (\text{E.44})$$

$$I_1 = B_0(m_{\tilde{\chi}_b^-}^2, m_{\tilde{\chi}_c^-}^2) \quad (\text{E.45})$$

$$I_2 = C_{00}(m_{\tilde{\chi}_c^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^R}^2) \quad (\text{E.46})$$

$$I_3 = C_0(m_{\tilde{\chi}_c^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^R}^2) \quad (\text{E.47})$$

$$V_{Z,(c_6)}^{LL} = \hat{X}_{b,\alpha,a}^L \hat{\bar{X}}_{\beta,c,a}^R (-C_{c,b}^L I_3 m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_c^-}) + C_{c,b}^R (I_1 - 2I_2 + I_3 m_{\nu_a^R}^2) \quad (\text{E.48})$$

$$V_{Z,(c_6)}^{LR} = \hat{X}_{b,\alpha,a}^L \hat{\bar{X}}_{\beta,c,a}^R (-C_{c,b}^L I_3 m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_c^-}) + C_{c,b}^R (I_1 - 2I_2 + I_3 m_{\nu_a^R}^2) \quad (\text{E.49})$$

$$V_{Z,(c_7)}^{LL} = -2 \hat{X}_{a,\alpha,b}^L \bar{X}_{\beta,a,c}^R \tilde{V}_{c,b} C_{00}(m_{\tilde{\chi}_a^-}^2, m_{\nu_c^i}^2, m_{\nu_b^R}^2) \quad (\text{E.50})$$

$$V_{Z,(c_7)}^{LR} = -2 \hat{X}_{a,\alpha,b}^L \bar{X}_{\beta,a,c}^R \tilde{V}_{c,b} C_{00}(m_{\tilde{\chi}_a^-}^2, m_{\nu_c^i}^2, m_{\nu_b^R}^2) \quad (\text{E.51})$$

$$V_{Z,(c_8)}^{LL} = 2 W_{a,\alpha,b}^L \hat{\bar{X}}_{\beta,a,c}^R \tilde{V}_{b,c} C_{00}(m_{\tilde{\chi}_a^-}^2, m_{\nu_c^R}^2, m_{\nu_b^i}^2) \quad (\text{E.52})$$

$$V_{Z,(c_8)}^{LR} = 2 W_{a,\alpha,b}^L \hat{\bar{X}}_{\beta,a,c}^R \tilde{V}_{b,c} C_{00}(m_{\tilde{\chi}_a^-}^2, m_{\nu_c^R}^2, m_{\nu_b^i}^2) \quad (\text{E.53})$$

E.3.2 Scalar penguins

CP even scalars.

Self-energy corrections.

$$I_1 = B_0(m_{\tilde{\chi}_b^-}^2, m_{\nu_a^i}^2) \quad (\text{E.54})$$

$$I_2 = B_1(m_{\tilde{\chi}_b^-}^2, m_{\nu_a^i}^2) \quad (\text{E.55})$$

$$\begin{aligned} S_{h_p,(c_1,c_3)}^{LL} &= (H_{\beta,c,p}^L (-(W_{b,\alpha,a}^L \bar{X}_{c,b,a}^R I_2 m_{\ell_\alpha}^2) + W_{b,\alpha,a}^R \bar{X}_{c,b,a}^R I_1 m_{\ell_\alpha} m_{\tilde{\chi}_b^-}) - W_{b,\alpha,a}^R \bar{X}_{c,b,a}^L I_2 m_{\ell_\alpha} m_{\ell_c} \\ &\quad + W_{b,\alpha,a}^L \bar{X}_{c,b,a}^L I_1 m_{\tilde{\chi}_b^-} m_{\ell_c})) / (m_{\ell_\alpha}^2 - m_{\ell_c}^2) \quad + (\alpha \leftrightarrow \beta) \end{aligned} \quad (\text{E.56})$$

$$\begin{aligned} S_{h_p,(c_1,c_3)}^{LR} &= (H_{\beta,c,p}^L (-(W_{b,\alpha,a}^L \bar{X}_{c,b,a}^R I_2 m_{\ell_\alpha}^2) + W_{b,\alpha,a}^R \bar{X}_{c,b,a}^R I_1 m_{\ell_\alpha} m_{\tilde{\chi}_b^-}) - W_{b,\alpha,a}^R \bar{X}_{c,b,a}^L I_2 m_{\ell_\alpha} m_{\ell_c} \\ &\quad + W_{b,\alpha,a}^L \bar{X}_{c,b,a}^L I_1 m_{\tilde{\chi}_b^-} m_{\ell_c})) / (m_{\ell_\alpha}^2 - m_{\ell_c}^2) \quad + (\alpha \leftrightarrow \beta) \end{aligned}$$

$$I_1 = B_0(m_{\tilde{\chi}_b^-}^2, m_{\nu_a^R}^2) \quad (\text{E.57})$$

$$I_2 = B_1(m_{\tilde{\chi}_b^-}^2, m_{\nu_a^R}^2) \quad (\text{E.58})$$

$$S_{h_p,(c_2,c_4)}^{LL} = (H_{\beta,c,p}^L (-(\hat{X}_{b,\alpha,a}^L \hat{\bar{X}}_{c,b,a}^R I_2 m_{\ell_\alpha}^2) + \hat{X}_{b,\alpha,a}^R \hat{\bar{X}}_{c,b,a}^R I_1 m_{\ell_\alpha} m_{\tilde{\chi}_b^-}) - \hat{X}_{b,\alpha,a}^R \hat{\bar{X}}_{c,b,a}^L I_2 m_{\ell_\alpha} m_{\ell_c})$$

$$+ \hat{X}_{b,\alpha,a}^L \hat{\bar{X}}_{c,b,a}^L I_1 m_{\tilde{\chi}_b^-} m_{\ell_c}) / (m_{\ell_\alpha}^2 - m_{\ell_c}^2) \quad + (\alpha \leftrightarrow \beta) \quad (\text{E.59})$$

$$\begin{aligned} S_{h_p,(c_2,c_4)}^{LR} = & (H_{\beta,c,p}^L (-(\hat{X}_{b,\alpha,a}^L \hat{\bar{X}}_{c,b,a}^R I_2 m_{\ell_\alpha}^2) + \hat{X}_{b,\alpha,a}^R \hat{\bar{X}}_{c,b,a}^R I_1 m_{\ell_\alpha} m_{\tilde{\chi}_b^-} - \hat{X}_{b,\alpha,a}^R \hat{\bar{X}}_{c,b,a}^L I_2 m_{\ell_\alpha} m_{\ell_c} \\ & + \hat{X}_{b,\alpha,a}^L \hat{\bar{X}}_{c,b,a}^L I_1 m_{\tilde{\chi}_b^-} m_{\ell_c})) / (m_{\ell_\alpha}^2 - m_{\ell_c}^2) \quad + (\alpha \leftrightarrow \beta) \end{aligned} \quad (\text{E.60})$$

Vertex corrections.

$$I_1 = B_0(m_{\tilde{\chi}_b^-}^2, m_{\tilde{\chi}_c^-}^2) \quad (\text{E.61})$$

$$I_2 = C_0(m_{\tilde{\chi}_c^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^i}^2) \quad (\text{E.62})$$

$$S_{h_p,(c_5)}^{LL} = W_{b,\alpha,a}^L \bar{X}_{\beta,c,a}^L (S_{c,b,p}^{c,L} I_2 m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_c^-} + S_{c,b,p}^{c,R} (I_1 + I_2 m_{\nu_a^i}^2)) \quad (\text{E.63})$$

$$S_{h_p,(c_5)}^{LR} = W_{b,\alpha,a}^L \bar{X}_{\beta,c,a}^L (S_{c,b,p}^{c,L} I_2 m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_c^-} + S_{c,b,p}^{c,R} (I_1 + I_2 m_{\nu_a^i}^2)) \quad (\text{E.64})$$

$$I_1 = B_0(m_{\tilde{\chi}_b^-}^2, m_{\tilde{\chi}_c^-}^2) \quad (\text{E.65})$$

$$I_2 = C_0(m_{\tilde{\chi}_c^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^R}^2) \quad (\text{E.66})$$

$$S_{h_p,(c_6)}^{LL} = \hat{X}_{b,\alpha,a}^L \hat{\bar{X}}_{\beta,c,a}^L (S_{c,b,p}^{c,L} I_2 m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_c^-} + S_{c,b,p}^{c,R} (I_1 + I_2 m_{\nu_a^R}^2)) \quad (\text{E.67})$$

$$S_{h_p,(c_6)}^{LR} = \hat{X}_{b,\alpha,a}^L \hat{\bar{X}}_{\beta,c,a}^L (S_{c,b,p}^{c,L} I_2 m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_c^-} + S_{c,b,p}^{c,R} (I_1 + I_2 m_{\nu_a^R}^2)) \quad (\text{E.68})$$

$$S_{h_p,(c_{10})}^{LL} = W_{a,\alpha,b}^L \bar{X}_{\beta,a,c}^L \tilde{H}_{p,c,b}^{ii} C_0(m_{\tilde{\chi}_a^-}^2, m_{\nu_c^i}^2, m_{\nu_b^i}^2) m_{\tilde{\chi}_a^-} \quad (\text{E.69})$$

$$S_{h_p,(c_{10})}^{LR} = W_{a,\alpha,b}^L \bar{X}_{\beta,a,c}^L \tilde{H}_{p,c,b}^{ii} C_0(m_{\tilde{\chi}_a^-}^2, m_{\nu_c^i}^2, m_{\nu_b^i}^2) m_{\tilde{\chi}_a^-} \quad (\text{E.70})$$

$$S_{h_p,(c_7)}^{LL} = \hat{X}_{a,\alpha,b}^L \bar{X}_{\beta,a,c}^L \tilde{H}_{p,c,b}^{ir} C_0(m_{\tilde{\chi}_a^-}^2, m_{\nu_c^i}^2, m_{\nu_b^R}^2) m_{\tilde{\chi}_a^-} \quad (\text{E.71})$$

$$S_{h_p,(c_7)}^{LR} = \hat{X}_{a,\alpha,b}^L \bar{X}_{\beta,a,c}^L \tilde{H}_{p,c,b}^{ir} C_0(m_{\tilde{\chi}_a^-}^2, m_{\nu_c^i}^2, m_{\nu_b^R}^2) m_{\tilde{\chi}_a^-} \quad (\text{E.72})$$

$$S_{h_p,(c_8)}^{LL} = W_{a,\alpha,b}^L \hat{\bar{X}}_{\beta,a,c}^L \tilde{H}_{p,b,c}^{ir} C_0(m_{\tilde{\chi}_a^-}^2, m_{\nu_c^R}^2, m_{\nu_b^i}^2) m_{\tilde{\chi}_a^-} \quad (\text{E.73})$$

$$S_{h_p,(c_8)}^{LR} = W_{a,\alpha,b}^L \hat{\bar{X}}_{\beta,a,c}^L \tilde{H}_{p,b,c}^{ir} C_0(m_{\tilde{\chi}_a^-}^2, m_{\nu_c^R}^2, m_{\nu_b^i}^2) m_{\tilde{\chi}_a^-} \quad (\text{E.74})$$

$$S_{h_p,(c_9)}^{LL} = \hat{X}_{a,\alpha,b}^L \hat{\bar{X}}_{\beta,a,c}^L \tilde{H}_{p,c,b}^{rr} C_0(m_{\tilde{\chi}_a^-}^2, m_{\nu_c^R}^2, m_{\nu_b^R}^2) m_{\tilde{\chi}_a^-} \quad (\text{E.75})$$

$$S_{h_p,(c_9)}^{LR} = \hat{X}_{a,\alpha,b}^L \hat{\bar{X}}_{\beta,a,c}^L \tilde{H}_{p,c,b}^{rr} C_0(m_{\tilde{\chi}_a^-}^2, m_{\nu_c^R}^2, m_{\nu_b^R}^2) m_{\tilde{\chi}_a^-} \quad (\text{E.76})$$

$$(\text{E.77})$$

CP odd scalars.

Self-energy corrections.

$$I_1 = B_0(m_{\tilde{\chi}_b^-}^2, m_{\nu_a^i}^2) \quad (\text{E.78})$$

$$I_2 = B_1(m_{\tilde{\chi}_b^-}^2, m_{\nu_a^i}^2) \quad (\text{E.79})$$

$$S_{A_p^0, (c_1, c_3)}^{LL} = (A_{\beta, c, p}^L(-(W_{b, \alpha, a}^L \bar{X}_{c, b, a}^R I_2 m_{\ell_\alpha}^2) + W_{b, \alpha, a}^R \bar{X}_{c, b, a}^R I_1 m_{\ell_\alpha} m_{\tilde{\chi}_b^-} - W_{b, \alpha, a}^R \bar{X}_{c, b, a}^L I_2 m_{\ell_\alpha} m_{\ell_c} + W_{b, \alpha, a}^L \bar{X}_{c, b, a}^L I_1 m_{\tilde{\chi}_b^-} m_{\ell_c})) / (m_{\ell_\alpha}^2 - m_{\ell_c}^2) + (\alpha \leftrightarrow \beta) \quad (\text{E.80})$$

$$S_{A_p^0, (c_1, c_3)}^{LR} = (A_{\beta, c, p}^L(-(W_{b, \alpha, a}^L \bar{X}_{c, b, a}^R I_2 m_{\ell_\alpha}^2) + W_{b, \alpha, a}^R \bar{X}_{c, b, a}^R I_1 m_{\ell_\alpha} m_{\tilde{\chi}_b^-} - W_{b, \alpha, a}^R \bar{X}_{c, b, a}^L I_2 m_{\ell_\alpha} m_{\ell_c} + W_{b, \alpha, a}^L \bar{X}_{c, b, a}^L I_1 m_{\tilde{\chi}_b^-} m_{\ell_c})) / (m_{\ell_\alpha}^2 - m_{\ell_c}^2) + (\alpha \leftrightarrow \beta) \quad (\text{E.80})$$

$$I_1 = B_0(m_{\tilde{\chi}_b^-}^2, m_{\nu_a^R}^2) \quad (\text{E.81})$$

$$I_2 = B_1(m_{\tilde{\chi}_b^-}^2, m_{\nu_a^R}^2) \quad (\text{E.82})$$

$$S_{A_p^0, (c_2, c_4)}^{LL} = (A_{\beta, c, p}^L(-(\hat{X}_{b, \alpha, a}^L \hat{\bar{X}}_{c, b, a}^R I_2 m_{\ell_\alpha}^2) + \hat{X}_{b, \alpha, a}^R \hat{\bar{X}}_{c, b, a}^R I_1 m_{\ell_\alpha} m_{\tilde{\chi}_b^-} - \hat{X}_{b, \alpha, a}^R \hat{\bar{X}}_{c, b, a}^L I_2 m_{\ell_\alpha} m_{\ell_c} + \hat{X}_{b, \alpha, a}^L \hat{\bar{X}}_{c, b, a}^L I_1 m_{\tilde{\chi}_b^-} m_{\ell_c})) / (m_{\ell_\alpha}^2 - m_{\ell_c}^2) + (\alpha \leftrightarrow \beta) \quad (\text{E.83})$$

$$S_{A_p^0, (c_2, c_4)}^{LR} = (A_{\beta, c, p}^L(-(\hat{X}_{b, \alpha, a}^L \hat{\bar{X}}_{c, b, a}^R I_2 m_{\ell_\alpha}^2) + \hat{X}_{b, \alpha, a}^R \hat{\bar{X}}_{c, b, a}^R I_1 m_{\ell_\alpha} m_{\tilde{\chi}_b^-} - \hat{X}_{b, \alpha, a}^R \hat{\bar{X}}_{c, b, a}^L I_2 m_{\ell_\alpha} m_{\ell_c} + \hat{X}_{b, \alpha, a}^L \hat{\bar{X}}_{c, b, a}^L I_1 m_{\tilde{\chi}_b^-} m_{\ell_c})) / (m_{\ell_\alpha}^2 - m_{\ell_c}^2) + (\alpha \leftrightarrow \beta) \quad (\text{E.84})$$

Vertex corrections.

$$I_1 = B_0(m_{\tilde{\chi}_b^-}^2, m_{\tilde{\chi}_c^-}^2) \quad (\text{E.85})$$

$$I_2 = C_0(m_{\tilde{\chi}_c^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^i}^2) \quad (\text{E.86})$$

$$S_{A_p^0, (c_5)}^{LL} = W_{b, \alpha, a}^L \bar{X}_{\beta, c, a}^L (P_{c, b, p}^{c, L} I_2 m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_c^-} + P_{c, b, p}^{c, R} (I_1 + I_2 m_{\nu_a^i}^2)) \quad (\text{E.87})$$

$$S_{A_p^0, (c_5)}^{LR} = W_{b, \alpha, a}^L \bar{X}_{\beta, c, a}^L (P_{c, b, p}^{c, L} I_2 m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_c^-} + P_{c, b, p}^{c, R} (I_1 + I_2 m_{\nu_a^i}^2)) \quad (\text{E.88})$$

$$I_1 = B_0(m_{\tilde{\chi}_b^-}^2, m_{\tilde{\chi}_c^-}^2) \quad (\text{E.89})$$

$$I_2 = C_0(m_{\tilde{\chi}_c^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^R}^2) \quad (\text{E.90})$$

$$S_{A_p^0, (c_6)}^{LL} = \hat{X}_{b, \alpha, a}^L \hat{\bar{X}}_{\beta, c, a}^L (P_{c, b, p}^{c, L} I_2 m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_c^-} + P_{c, b, p}^{c, R} (I_1 + I_2 m_{\nu_a^R}^2)) \quad (\text{E.91})$$

$$S_{A_p^0, (c_6)}^{LR} = \hat{X}_{b, \alpha, a}^L \hat{\bar{X}}_{\beta, c, a}^L (P_{c, b, p}^{c, L} I_2 m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_c^-} + P_{c, b, p}^{c, R} (I_1 + I_2 m_{\nu_a^R}^2)) \quad (\text{E.92})$$

$$S_{A_p^0, (c_{10})}^{LL} = W_{a, \alpha, b}^L \bar{X}_{\beta, a, c}^L \tilde{A}_{p, c, b}^{ii} C_0(m_{\tilde{\chi}_a^-}^2, m_{\nu_c^i}^2, m_{\nu_b^i}^2) m_{\tilde{\chi}_a^-} \quad (\text{E.93})$$

$$S_{A_p^0, (c_{10})}^{LR} = W_{a, \alpha, b}^L \bar{X}_{\beta, a, c}^L \tilde{A}_{p, c, b}^{ii} C_0(m_{\tilde{\chi}_a^-}^2, m_{\nu_c^i}^2, m_{\nu_b^R}^2) m_{\tilde{\chi}_a^-} \quad (\text{E.94})$$

$$S_{A_p^0, (c_7)}^{LL} = \hat{X}_{a, \alpha, b}^L \hat{\bar{X}}_{\beta, a, c}^L \tilde{A}_{p, c, b}^{ir} C_0(m_{\tilde{\chi}_a^-}^2, m_{\nu_c^i}^2, m_{\nu_b^R}^2) m_{\tilde{\chi}_a^-} \quad (\text{E.95})$$

$$S_{A_p^0, (c_7)}^{LR} = \hat{X}_{a, \alpha, b}^L \hat{\bar{X}}_{\beta, a, c}^L \tilde{A}_{p, c, b}^{ir} C_0(m_{\tilde{\chi}_a^-}^2, m_{\nu_c^i}^2, m_{\nu_b^R}^2) m_{\tilde{\chi}_a^-} \quad (\text{E.96})$$

$$S_{A_p^0, (c_8)}^{LL} = W_{a, \alpha, b}^L \hat{\bar{X}}_{\beta, a, c}^L \tilde{A}_{p, b, c}^{ir} C_0(m_{\tilde{\chi}_a^-}^2, m_{\nu_c^R}^2, m_{\nu_b^2}^2) m_{\tilde{\chi}_a^-} \quad (\text{E.97})$$

$$S_{A_p^0, (c_8)}^{LR} = W_{a, \alpha, b}^L \hat{\bar{X}}_{\beta, a, c}^L \tilde{A}_{p, b, c}^{ir} C_0(m_{\tilde{\chi}_a^-}^2, m_{\nu_c^R}^2, m_{\nu_b^i}^2) m_{\tilde{\chi}_a^-} \quad (\text{E.98})$$

$$S_{A_p^0, (c_9)}^{LL} = \hat{X}_{a,\alpha,b}^L \hat{\bar{X}}_{\beta,a,c}^L \tilde{A}_{p,c,b}^{rr} C_0(m_{\tilde{\chi}_a^-}^2, m_{\nu_c^R}^2, m_{\nu_b^R}^2) m_{\tilde{\chi}_a^-} \quad (\text{E.99})$$

$$S_{A_p^0, (c_9)}^{LR} = \hat{X}_{a,\alpha,b}^L \hat{\bar{X}}_{\beta,a,c}^L \tilde{A}_{p,c,b}^{rr} C_0(m_{\tilde{\chi}_a^-}^2, m_{\nu_c^R}^2, m_{\nu_b^R}^2) m_{\tilde{\chi}_a^-} \quad (\text{E.100})$$

E.4 W^+ and H^+ contributions

E.4.1 Z-penguins

Self-energy corrections.

$$I_1 = B_0(m_{\nu_a}^2, m_{H_b^-}^2) \quad (\text{E.101})$$

$$I_2 = B_1(m_{\nu_a}^2, m_{H_b^-}^2) \quad (\text{E.102})$$

$$\begin{aligned} V_{Z,(w_1,w_3)}^{LL} = & (E_{\beta,c}^L (V_{a,\alpha,b}^{+,L} \bar{V}_{c,a,b}^{+,R} I_2 m_{\ell_\alpha}^2 - V_{a,\alpha,b}^{+,R} \bar{V}_{c,a,b}^{+,R} I_1 m_{\ell_\alpha} m_{\nu_a} + V_{a,\alpha,b}^{+,R} \bar{V}_{c,a,b}^{+,L} I_2 m_{\ell_\alpha} m_{\ell_c} \\ & - V_{a,\alpha,b}^{+,L} \bar{V}_{c,a,b}^{+,L} I_1 m_{\nu_a} m_{\ell_c})) / (m_{\ell_\alpha}^2 - m_{\ell_c}^2) \quad + (\alpha \leftrightarrow \beta) \end{aligned} \quad (\text{E.103})$$

$$\begin{aligned} V_{Z,(w_1,w_3)}^{LR} = & (E_{\beta,c}^L (V_{a,\alpha,b}^{+,L} \bar{V}_{c,a,b}^{+,R} I_2 m_{\ell_\alpha}^2 - V_{a,\alpha,b}^{+,R} \bar{V}_{c,a,b}^{+,R} I_1 m_{\ell_\alpha} m_{\nu_a} + V_{a,\alpha,b}^{+,R} \bar{V}_{c,a,b}^{+,L} I_2 m_{\ell_\alpha} m_{\ell_c} \\ & - V_{a,\alpha,b}^{+,L} \bar{V}_{c,a,b}^{+,L} I_1 m_{\nu_a} m_{\ell_c})) / (m_{\ell_\alpha}^2 - m_{\ell_c}^2) \quad + (\alpha \leftrightarrow \beta) \end{aligned}$$

$$I_1 = B_0(m_{\nu_a}^2, m_{W_-}^2) \quad (\text{E.104})$$

$$I_2 = B_1(m_{\nu_a}^2, m_{W_-}^2) \quad (\text{E.105})$$

$$\begin{aligned} V_{Z,(w_2,w_4)}^{LL} = & (E_{\beta,c}^L (\hat{V}_{a,i}^{+,R} m_{\ell_\alpha} (-2 \hat{\bar{V}}_{c,a}^{+,L} (1 - 2 I_1) m_{\nu_a} + \hat{\bar{V}}_{c,a}^{+,R} (1 + 2 I_2) m_{\ell_c}) \\ & + \hat{V}_{a,i}^{+,L} (\hat{\bar{V}}_{c,a}^{+,L} (1 + 2 I_2) m_{\ell_\alpha}^2 - 2 \hat{\bar{V}}_{c,a}^{+,R} (1 - 2 I_1) m_{\nu_a} m_{\ell_c})) / (m_{\ell_\alpha}^2 - m_{\ell_c}^2) \quad + (\alpha \leftrightarrow \beta) \end{aligned} \quad (\text{E.106})$$

$$\begin{aligned} V_{Z,(w_2,w_4)}^{LR} = & (E_{\beta,c}^L (\hat{V}_{a,i}^{+,R} m_{\ell_\alpha} (-2 \hat{\bar{V}}_{c,a}^{+,L} (1 - 2 I_1) m_{\nu_a} + \hat{\bar{V}}_{c,a}^{+,R} (1 + 2 I_2) m_{\ell_c}) \\ & + \hat{V}_{a,i}^{+,L} (\hat{\bar{V}}_{c,a}^{+,L} (1 + 2 I_2) m_{\ell_\alpha}^2 - 2 \hat{\bar{V}}_{c,a}^{+,R} (1 - 2 I_1) m_{\nu_a} m_{\ell_c})) / (m_{\ell_\alpha}^2 - m_{\ell_c}^2) \quad + (\alpha \leftrightarrow \beta) \end{aligned} \quad (\text{E.107})$$

Vertex corrections.

$$V_{Z,(w_5)}^{LL} = -2 V_{a,\alpha,b}^{+,L} \bar{V}_{\beta,a,c}^{+,R} Z_{b,c}^{hh} C_{00}(m_{\nu_a}^2, m_{H_c^-}^2, m_{H_b^-}^2) \quad (\text{E.108})$$

$$V_{Z,(w_5)}^{LR} = -2 V_{a,\alpha,b}^{+,L} \bar{V}_{\beta,a,c}^{+,R} Z_{b,c}^{hh} C_{00}(m_{\nu_a}^2, m_{H_c^-}^2, m_{H_b^-}^2) \quad (\text{E.109})$$

$$V_{Z,(w_6)}^{LL} = \hat{V}_{a,i}^{+,L} \bar{V}_{\beta,a,c}^{+,R} \bar{Z}_c^{hw} C_0(m_{\nu_a}^2, m_{H_c^-}^2, m_{W_-}^2) m_{\nu_a} \quad (\text{E.110})$$

$$V_{Z,(w_6)}^{LR} = \hat{V}_{a,i}^{+,L} \bar{V}_{\beta,a,c}^{+,R} \bar{Z}_c^{hw} C_0(m_{\nu_a}^2, m_{H_c^-}^2, m_{W_-}^2) m_{\nu_a} \quad (\text{E.111})$$

$$V_{Z,(w_7)}^{LL} = V_{a,\alpha,b}^{+,L} \hat{\bar{V}}_{\beta,a}^{+,L} Z_b^{hw} C_0(m_{\nu_a}^2, m_{W_-}^2, m_{H_b^-}^2) m_{\nu_a} \quad (\text{E.112})$$

$$V_{Z,(w_7)}^{LR} = V_{a,\alpha,b}^{+,L} \hat{\bar{V}}_{\beta,a}^{+,L} Z_b^{hw} C_0(m_{\nu_a}^2, m_{W_-}^2, m_{H_b^-}^2) m_{\nu_a} \quad (\text{E.113})$$

$$I_1 = B_0(m_{W_-}^2, m_{W_-}^2) \quad (\text{E.114})$$

$$I_2 = C_{00}(m_{\nu_a}^2, m_{W_-}^2, m_{W_-}^2) \quad (\text{E.115})$$

$$I_3 = C_0(m_{\nu_a}^2, m_{W^-}^2, m_{W^-}^2) \quad (\text{E.116})$$

$$V_{Z,(w_8)}^{LL} = -\hat{V}_{a,i}^{+,L} \hat{\bar{V}}_{\beta,a}^{+,L} Z^{ww} (-1 + 2(I_1 + 2I_2 + I_3 m_{\nu_a}^2)) \quad (\text{E.117})$$

$$V_{Z,(w_8)}^{LR} = -\hat{V}_{a,i}^{+,L} \hat{\bar{V}}_{\beta,a}^{+,L} Z^{ww} (-1 + 2(I_1 + 2I_2 + I_3 m_{\nu_a}^2)) \quad (\text{E.118})$$

$$I_1 = B_0(m_{\nu_b}^2, m_{\nu_c}^2) \quad (\text{E.119})$$

$$I_2 = C_{00}(m_{\nu_c}^2, m_{\nu_b}^2, m_{H_a^-}^2) \quad (\text{E.120})$$

$$I_3 = C_0(m_{\nu_c}^2, m_{\nu_b}^2, m_{H_a^-}^2) \quad (\text{E.121})$$

$$V_{Z,(w_9)}^{LL} = V_{b,\alpha,a}^{+,L} \bar{V}_{\beta,c,a}^{+,R} (- (V_{c,b}^L I_3 m_{\nu_b} m_{\nu_c}) + V_{c,b}^R (I_1 - 2I_2 + I_3 m_{H_a^-}^2)) \quad (\text{E.122})$$

$$V_{Z,(w_9)}^{LR} = V_{b,\alpha,a}^{+,L} \bar{V}_{\beta,c,a}^{+,R} (- (V_{c,b}^L I_3 m_{\nu_b} m_{\nu_c}) + V_{c,b}^R (I_1 - 2I_2 + I_3 m_{H_a^-}^2)) \quad (\text{E.123})$$

$$I_1 = B_0(m_{\nu_b}^2, m_{\nu_c}^2) \quad (\text{E.124})$$

$$I_2 = C_{00}(m_{\nu_c}^2, m_{\nu_b}^2, m_{W^-}^2) \quad (\text{E.125})$$

$$I_3 = C_0(m_{\nu_c}^2, m_{\nu_b}^2, m_{W^-}^2) \quad (\text{E.126})$$

$$V_{Z,(w_{10})}^{LL} = -(\hat{V}_{b,i}^{+,L} \hat{\bar{V}}_{\beta,c}^{+,L} (2V_{c,b}^R I_3 m_{\nu_b} m_{\nu_c} + V_{c,b}^L (1 - 2(I_1 - 2I_2 + I_3 m_{W^-}^2)))) \quad (\text{E.127})$$

$$V_{Z,(w_{10})}^{LR} = -(\hat{V}_{b,i}^{+,L} \hat{\bar{V}}_{\beta,c}^{+,L} (2V_{c,b}^R I_3 m_{\nu_b} m_{\nu_c} + V_{c,b}^L (1 - 2(I_1 - 2I_2 + I_3 m_{W^-}^2)))) \quad (\text{E.128})$$

E.4.2 Scalar penguins

CP even scalars.

Self-energy corrections.

$$I_1 = B_0(m_{\nu_a}^2, m_{H_b^-}^2) \quad (\text{E.129})$$

$$I_2 = B_1(m_{\nu_a}^2, m_{H_b^-}^2) \quad (\text{E.130})$$

$$\begin{aligned} S_{h_p,(w_1,w_3)}^{LL} &= (H_{\beta,c,p}^L (- (V_{a,\alpha,b}^{+,L} \bar{V}_{c,a,b}^{+,R} I_2 m_{\ell_\alpha}^2) + V_{a,\alpha,b}^{+,R} \bar{V}_{c,a,b}^{+,R} I_1 m_{\ell_\alpha} m_{\nu_a} - V_{a,\alpha,b}^{+,R} \bar{V}_{c,a,b}^{+,L} I_2 m_{\ell_\alpha} m_{\ell_c} \\ &\quad + V_{a,\alpha,b}^{+,L} \bar{V}_{c,a,b}^{+,L} I_1 m_{\nu_a} m_{\ell_c})) / (m_{\ell_\alpha}^2 - m_{\ell_c}^2) \quad + (\alpha \leftrightarrow \beta) \end{aligned} \quad (\text{E.131})$$

$$\begin{aligned} S_{h_p,(w_1,w_3)}^{LR} &= (H_{\beta,c,p}^L (- (V_{a,\alpha,b}^{+,L} \bar{V}_{c,a,b}^{+,R} I_2 m_{\ell_\alpha}^2) + V_{a,\alpha,b}^{+,R} \bar{V}_{c,a,b}^{+,R} I_1 m_{\ell_\alpha} m_{\nu_a} - V_{a,\alpha,b}^{+,R} \bar{V}_{c,a,b}^{+,L} I_2 m_{\ell_\alpha} m_{\ell_c} \\ &\quad + V_{a,\alpha,b}^{+,L} \bar{V}_{c,a,b}^{+,L} I_1 m_{\nu_a} m_{\ell_c})) / (m_{\ell_\alpha}^2 - m_{\ell_c}^2) \quad + (\alpha \leftrightarrow \beta) \end{aligned}$$

$$I_1 = B_0(m_{\nu_a}^2, m_{W^-}^2) \quad (\text{E.132})$$

$$I_2 = B_1(m_{\nu_a}^2, m_{W^-}^2) \quad (\text{E.133})$$

$$\begin{aligned} S_{h_p,(w_2,w_4)}^{LL} &= -((H_{\beta,c,p}^L (\hat{V}_{a,i}^{+,R} m_{\ell_\alpha} (-2\hat{V}_{c,a}^{+,L} (1 - 2I_1) m_{\nu_a} + \hat{V}_{c,a}^{+,R} (1 + 2I_2) m_{\ell_c}) + \hat{V}_{a,i}^{+,L} (\hat{V}_{c,a}^{+,L} (1 + 2I_2) m_{\ell_\alpha}^2 \\ &\quad - 2\hat{V}_{c,a}^{+,R} (1 - 2I_1) m_{\nu_a} m_{\ell_c}))) / (m_{\ell_\alpha}^2 - m_{\ell_c}^2)) \quad + (\alpha \leftrightarrow \beta) \end{aligned} \quad (\text{E.134})$$

$$\begin{aligned} S_{h_p,(w_2,w_4)}^{LR} &= -((H_{\beta,c,p}^L (\hat{V}_{a,i}^{+,R} m_{\ell_\alpha} (-2\hat{V}_{c,a}^{+,L} (1 - 2I_1) m_{\nu_a} + \hat{V}_{c,a}^{+,R} (1 + 2I_2) m_{\ell_c}) + \hat{V}_{a,i}^{+,L} (\hat{V}_{c,a}^{+,L} (1 + 2I_2) m_{\ell_\alpha}^2 \\ &\quad - 2\hat{V}_{c,a}^{+,R} (1 - 2I_1) m_{\nu_a} m_{\ell_c}))) / (m_{\ell_\alpha}^2 - m_{\ell_c}^2)) \quad + (\alpha \leftrightarrow \beta) \end{aligned} \quad (\text{E.135})$$

Vertex corrections.

$$S_{h_p,(w_5)}^{LL} = V_{a,\alpha,b}^{+,L} \bar{V}_{\beta,a,c}^{+,L} H_{p,b,c}^{hh} C_0(m_{\nu_a}^2, m_{H_c^-}^2, m_{H_b^-}^2) m_{\nu_a} \quad (\text{E.136})$$

$$S_{h_p, (w_5)}^{LR} = V_{a,\alpha,b}^{+,L} \bar{V}_{\beta,a,c}^{+,L} H_{p,b,c}^{hh} C_0(m_{\nu_a}^2, m_{H_c^-}^2, m_{H_b^-}^2) m_{\nu_a} \quad (\text{E.137})$$

$$I_1 = B_0(m_{H_c^-}^2, m_{W^-}^2) \quad (\text{E.138})$$

$$I_2 = C_0(m_{\nu_a}^2, m_{H_c^-}^2, m_{W^-}^2) \quad (\text{E.139})$$

$$S_{h_p, (w_6)}^{LL} = -\hat{V}_{a,i}^{+,L} \bar{V}_{\beta,a,c}^{+,L} \bar{H}_{p,c}^{hw} (I_1 + I_2 m_{\nu_a}^2) \quad (\text{E.140})$$

$$S_{h_p, (w_6)}^{LR} = -\hat{V}_{a,i}^{+,L} \bar{V}_{\beta,a,c}^{+,L} \bar{H}_{p,c}^{hw} (I_1 + I_2 m_{\nu_a}^2) \quad (\text{E.141})$$

$$I_1 = B_0(m_{H_b^-}^2, m_{W^-}^2) \quad (\text{E.142})$$

$$I_2 = C_0(m_{\nu_a}^2, m_{W^-}^2, m_{H_b^-}^2) \quad (\text{E.143})$$

$$S_{h_p, (w_7)}^{LL} = V_{a,\alpha,b}^{+,L} \hat{\bar{V}}_{\beta,a}^{+,R} H_{p,b}^{hw} (I_1 + I_2 m_{\nu_a}^2) \quad (\text{E.144})$$

$$S_{h_p, (w_7)}^{LR} = V_{a,\alpha,b}^{+,L} \hat{\bar{V}}_{\beta,a}^{+,R} H_{p,b}^{hw} (I_1 + I_2 m_{\nu_a}^2) \quad (\text{E.145})$$

$$S_{h_p, (w_8)}^{LL} = 4 \hat{V}_{a,i}^{+,L} \hat{\bar{V}}_{\beta,a}^{+,R} H_p^{ww} C_0(m_{\nu_a}^2, m_{W^-}^2, m_{W^-}^2) m_{\nu_a} \quad (\text{E.146})$$

$$S_{h_p, (w_8)}^{LR} = 4 \hat{V}_{a,i}^{+,L} \hat{\bar{V}}_{\beta,a}^{+,R} H_p^{ww} C_0(m_{\nu_a}^2, m_{W^-}^2, m_{W^-}^2) m_{\nu_a} \quad (\text{E.147})$$

$$I_1 = B_0(m_{\nu_b}^2, m_{\nu_c}^2) \quad (\text{E.148})$$

$$I_2 = C_0(m_{\nu_c}^2, m_{\nu_b}^2, m_{H_a^-}^2) \quad (\text{E.149})$$

$$S_{h_p, (w_9)}^{LL} = V_{b,\alpha,a}^{+,L} \bar{V}_{\beta,c,a}^{+,L} (H_{c,b,p}^{\nu,L} I_2 m_{\nu_b} m_{\nu_c} + H_{c,b,p}^{\nu,R} (I_1 + I_2 m_{H_a^-}^2)) \quad (\text{E.150})$$

$$S_{h_p, (w_9)}^{LR} = V_{b,\alpha,a}^{+,L} \bar{V}_{\beta,c,a}^{+,L} (H_{c,b,p}^{\nu,L} I_2 m_{\nu_b} m_{\nu_c} + H_{c,b,p}^{\nu,R} (I_1 + I_2 m_{H_a^-}^2)) \quad (\text{E.151})$$

$$I_1 = B_0(m_{\nu_b}^2, m_{\nu_c}^2) \quad (\text{E.152})$$

$$I_2 = C_0(m_{\nu_c}^2, m_{\nu_b}^2, m_{W^-}^2) \quad (\text{E.153})$$

$$S_{h_p, (w_{10})}^{LL} = 2 \hat{V}_{b,i}^{+,L} \hat{\bar{V}}_{\beta,c}^{+,R} (-2 H_{c,b,p}^{\nu,R} I_2 m_{\nu_b} m_{\nu_c} + H_{c,b,p}^{\nu,L} (1 - 2(I_1 + I_2 m_{W^-}^2))) \quad (\text{E.154})$$

$$S_{h_p, (w_{10})}^{LR} = 2 \hat{V}_{b,i}^{+,L} \hat{\bar{V}}_{\beta,c}^{+,R} (-2 H_{c,b,p}^{\nu,R} I_2 m_{\nu_b} m_{\nu_c} + H_{c,b,p}^{\nu,L} (1 - 2(I_1 + I_2 m_{W^-}^2))) \quad (\text{E.155})$$

CP odd scalars.

Self-energy corrections.

$$I_1 = B_0(m_{\nu_a}^2, m_{H_b^-}^2) \quad (\text{E.156})$$

$$I_2 = B_1(m_{\nu_a}^2, m_{H_b^-}^2) \quad (\text{E.157})$$

$$\begin{aligned} S_{A_p^0, (w_1, w_3)}^{LL} = & (A_{\beta,c,p}^L (-(V_{a,\alpha,b}^{+,L} \bar{V}_{c,a,b}^{+,R} I_2 m_{\ell_\alpha}^2) + V_{a,\alpha,b}^{+,R} \bar{V}_{c,a,b}^{+,R} I_1 m_{\ell_\alpha} m_{\nu_a} - V_{a,\alpha,b}^{+,R} \bar{V}_{c,a,b}^{+,L} I_2 m_{\ell_\alpha} m_{\ell_c} \\ & + V_{a,\alpha,b}^{+,L} \bar{V}_{c,a,b}^{+,L} I_1 m_{\nu_a} m_{\ell_c})) / (m_{\ell_\alpha}^2 - m_{\ell_c}^2) \quad + (\alpha \leftrightarrow \beta) \end{aligned} \quad (\text{E.158})$$

$$\begin{aligned} S_{A_p^0, (w_1, w_3)}^{LR} = & (A_{\beta,c,p}^L (-(V_{a,\alpha,b}^{+,L} \bar{V}_{c,a,b}^{+,R} I_2 m_{\ell_\alpha}^2) + V_{a,\alpha,b}^{+,R} \bar{V}_{c,a,b}^{+,R} I_1 m_{\ell_\alpha} m_{\nu_a} - V_{a,\alpha,b}^{+,R} \bar{V}_{c,a,b}^{+,L} I_2 m_{\ell_\alpha} m_{\ell_c} \\ & + V_{a,\alpha,b}^{+,L} \bar{V}_{c,a,b}^{+,L} I_1 m_{\nu_a} m_{\ell_c})) / (m_{\ell_\alpha}^2 - m_{\ell_c}^2) \quad + (\alpha \leftrightarrow \beta) \end{aligned}$$

$$I_1 = B_0(m_{\nu_a}^2, m_{W^-}^2) \quad (\text{E.159})$$

$$I_2 = B_1(m_{\nu_a}^2, m_{W^-}^2) \quad (\text{E.160})$$

$$\begin{aligned} S_{A_p^0, (w_2, w_4)}^{LL} = & - (A_{\beta, c, p}^L (\hat{V}_{a, i}^{+, R} m_{\ell_\alpha} (-2 \hat{V}_{c, a}^{+, L} (1 - 2I_1) m_{\nu_a} + \hat{V}_{c, a}^{+, R} (1 + 2I_2) m_{\ell_c}) + \hat{V}_{a, i}^{+, L} (\hat{V}_{c, a}^{+, L} (1 + 2I_2) m_{\ell_\alpha}^2 \\ & - 2 \hat{V}_{c, a}^{+, R} (1 - 2I_1) m_{\nu_a} m_{\ell_c})))/(m_{\ell_\alpha}^2 - m_{\ell_c}^2) \quad + (\alpha \leftrightarrow \beta) \end{aligned} \quad (\text{E.161})$$

$$\begin{aligned} S_{A_p^0, (w_2, w_4)}^{LR} = & - (A_{\beta, c, p}^L (\hat{V}_{a, i}^{+, R} m_{\ell_\alpha} (-2 \hat{V}_{c, a}^{+, L} (1 - 2I_1) m_{\nu_a} + \hat{V}_{c, a}^{+, R} (1 + 2I_2) m_{\ell_c}) + \hat{V}_{a, i}^{+, L} (\hat{V}_{c, a}^{+, L} (1 + 2I_2) m_{\ell_\alpha}^2 \\ & - 2 \hat{V}_{c, a}^{+, R} (1 - 2I_1) m_{\nu_a} m_{\ell_c})))/(m_{\ell_\alpha}^2 - m_{\ell_c}^2) \quad + (\alpha \leftrightarrow \beta) \end{aligned} \quad (\text{E.162})$$

Vertex corrections.

$$S_{A_p^0, (w_5)}^{LL} = V_{a, \alpha, b}^{+, L} \bar{V}_{\beta, a, c}^{+, L} A_{p, b, c}^{hh} C_0(m_{\nu_a}^2, m_{H_b^-}^2, m_{H_a^-}^2) m_{\nu_a} \quad (\text{E.163})$$

$$S_{A_p^0, (w_5)}^{LR} = V_{a, \alpha, b}^{+, L} \bar{V}_{\beta, a, c}^{+, L} A_{p, b, c}^{hh} C_0(m_{\nu_a}^2, m_{H_b^-}^2, m_{H_a^-}^2) m_{\nu_a} \quad (\text{E.164})$$

$$I_1 = B_0(m_{H_a^-}^2, m_{W^-}^2) \quad (\text{E.165})$$

$$I_2 = C_0(m_{\nu_a}^2, m_{H_a^-}^2, m_{W^-}^2) \quad (\text{E.166})$$

$$S_{A_p^0, (w_6)}^{LL} = - \hat{V}_{a, i}^{+, L} \bar{V}_{\beta, a, c}^{+, L} \bar{A}_{p, c}^{hw} (I_1 + I_2 m_{\nu_a}^2) \quad (\text{E.167})$$

$$S_{A_p^0, (w_6)}^{LR} = - \hat{V}_{a, i}^{+, L} \bar{V}_{\beta, a, c}^{+, L} \bar{A}_{p, c}^{hw} (I_1 + I_2 m_{\nu_a}^2) \quad (\text{E.168})$$

$$I_1 = B_0(m_{H_b^-}^2, m_{W^-}^2) \quad (\text{E.169})$$

$$I_2 = C_0(m_{\nu_a}^2, m_{W^-}^2, m_{H_b^-}^2) \quad (\text{E.170})$$

$$S_{A_p^0, (w_7)}^{LL} = V_{a, \alpha, b}^{+, L} \hat{V}_{\beta, a}^{+, R} A_{p, b}^{hw} (I_1 + I_2 m_{\nu_a}^2) \quad (\text{E.171})$$

$$S_{A_p^0, (w_7)}^{LR} = V_{a, \alpha, b}^{+, L} \hat{V}_{\beta, a}^{+, R} A_{p, b}^{hw} (I_1 + I_2 m_{\nu_a}^2) \quad (\text{E.172})$$

$$I_1 = B_0(m_{\nu_b}^2, m_{\nu_c}^2) \quad (\text{E.173})$$

$$I_2 = C_0(m_{\nu_c}^2, m_{\nu_b}^2, m_{H_a^-}^2) \quad (\text{E.174})$$

$$S_{A_p^0, (w_9)}^{LL} = V_{b, \alpha, a}^{+, L} \bar{V}_{\beta, c, a}^{+, L} (A_{c, b, p}^{\nu, L} I_2 m_{\nu_b} m_{\nu_c} + A_{c, b, p}^{\nu, R} (I_1 + I_2 m_{H_a^-}^2)) \quad (\text{E.175})$$

$$S_{A_p^0, (w_9)}^{LR} = V_{b, \alpha, a}^{+, L} \bar{V}_{\beta, c, a}^{+, L} (A_{c, b, p}^{\nu, L} I_2 m_{\nu_b} m_{\nu_c} + A_{c, b, p}^{\nu, R} (I_1 + I_2 m_{H_a^-}^2)) \quad (\text{E.176})$$

$$I_1 = B_0(m_{\nu_b}^2, m_{\nu_c}^2) \quad (\text{E.177})$$

$$I_2 = C_0(m_{\nu_c}^2, m_{\nu_b}^2, m_{W^-}^2) \quad (\text{E.178})$$

$$S_{A_p^0, (w_{10})}^{LL} = 2 \hat{V}_{b, i}^{+, L} \hat{V}_{\beta, c}^{+, R} (-2 A_{c, b, p}^{\nu, R} I_2 m_{\nu_b} m_{\nu_c} + A_{c, b, p}^{\nu, L} (1 - 2(I_1 + I_2 m_{W^-}^2))) \quad (\text{E.179})$$

$$S_{A_p^0, (w_{10})}^{LR} = 2 \hat{V}_{b, i}^{+, L} \hat{V}_{\beta, c}^{+, R} (-2 A_{c, b, p}^{\nu, R} I_2 m_{\nu_b} m_{\nu_c} + A_{c, b, p}^{\nu, L} (1 - 2(I_1 + I_2 m_{W^-}^2))) \quad (\text{E.180})$$


$$(\text{E.181})$$

F Box contributions to LFV


F.1 Four lepton boxes


F.1.1 Feynman diagrams

Neutralino diagrams



Chargino diagrams





F.1.2 Neutralino contributions

$$S_{(n_1^l)}^{LL} = -N_{a,\alpha,d}^L \bar{N}_{\beta,a,b}^L N_{c,\gamma,b}^L \bar{N}_{\delta,c,d}^L m_{\tilde{\chi}_a^0} m_{\tilde{\chi}_c^0} D_0(m_{\tilde{\chi}_a^0}^2, m_{\tilde{\chi}_c^0}^2, m_{\tilde{e}_d}^2, m_{\tilde{e}_b}^2) \quad (\text{F.1})$$

$$S_{(n_1^l)}^{LR} = -N_{a,\alpha,d}^L \bar{N}_{\beta,a,b}^L N_{c,\gamma,b}^R \bar{N}_{\delta,c,d}^R m_{\tilde{\chi}_a^0} m_{\tilde{\chi}_c^0} D_0(m_{\tilde{\chi}_a^0}^2, m_{\tilde{\chi}_c^0}^2, m_{\tilde{e}_d}^2, m_{\tilde{e}_b}^2) \quad (\text{F.2})$$

$$V_{(n_1^l)}^{LL} = -N_{a,\alpha,d}^L \bar{N}_{\beta,a,b}^R N_{c,\gamma,b}^L \bar{N}_{\delta,c,d}^R D_{27}(m_{\tilde{\chi}_a^0}^2, m_{\tilde{\chi}_c^0}^2, m_{\tilde{e}_d}^2, m_{\tilde{e}_b}^2) \quad (\text{F.3})$$

$$V_{(n_1^l)}^{LR} = -N_{a,\alpha,d}^L \bar{N}_{\beta,a,b}^R N_{c,\gamma,b}^R \bar{N}_{\delta,c,d}^L D_{27}(m_{\tilde{\chi}_a^0}^2, m_{\tilde{\chi}_c^0}^2, m_{\tilde{e}_d}^2, m_{\tilde{e}_b}^2) \quad (\text{F.4})$$

$$S_{(n_2^l)}^{LL} = \frac{1}{2} N_{a,\alpha,d}^L N_{a,\gamma,b}^L \bar{N}_{\beta,c,b}^L \bar{N}_{\delta,c,d}^L m_{\tilde{\chi}_a^0} m_{\tilde{\chi}_c^0} D_0(m_{\tilde{\chi}_a^0}^2, m_{\tilde{\chi}_c^0}^2, m_{\tilde{e}_d}^2, m_{\tilde{e}_b}^2) \quad (\text{F.5})$$

$$S_{(n_2^l)}^{LR} = -2 N_{a,\alpha,d}^L N_{a,\gamma,b}^R \bar{N}_{\beta,c,b}^L \bar{N}_{\delta,c,d}^R D_{27}(m_{\tilde{\chi}_a^0}^2, m_{\tilde{\chi}_c^0}^2, m_{\tilde{e}_d}^2, m_{\tilde{e}_b}^2) \quad (\text{F.6})$$

$$V_{(n_2^l)}^{LL} = -\frac{1}{2} N_{a,\alpha,d}^L N_{a,\gamma,b}^L \bar{N}_{\beta,c,b}^R \bar{N}_{\delta,c,d}^R m_{\tilde{\chi}_a^0} m_{\tilde{\chi}_c^0} D_0(m_{\tilde{\chi}_a^0}^2, m_{\tilde{\chi}_c^0}^2, m_{\tilde{e}_d}^2, m_{\tilde{e}_b}^2) \quad (\text{F.7})$$

$$V_{(n_2^l)}^{LR} = -N_{a,\alpha,d}^L N_{a,\gamma,b}^R \bar{N}_{\beta,c,b}^R \bar{N}_{\delta,c,d}^L D_{27}(m_{\tilde{\chi}_a^0}^2, m_{\tilde{\chi}_c^0}^2, m_{\tilde{e}_d}^2, m_{\tilde{e}_b}^2) \quad (\text{F.8})$$

$$T_{(n_2^l)}^{LL} = \frac{1}{8} N_{a,\alpha,d}^L N_{a,\gamma,b}^L \bar{N}_{\beta,c,b}^L \bar{N}_{\delta,c,d}^L m_{\tilde{\chi}_a^0} m_{\tilde{\chi}_c^0} D_0(m_{\tilde{\chi}_a^0}^2, m_{\tilde{\chi}_c^0}^2, m_{\tilde{e}_d}^2, m_{\tilde{e}_b}^2) \quad (\text{F.9})$$

F.1.3 Chargino contributions

$$S_{(c_1^l)}^{LL} = -W_{a,\alpha,d}^L \bar{X}_{\beta,a,b}^L W_{c,\gamma,b}^L \bar{X}_{\delta,c,d}^L m_{\tilde{\chi}_a^-} m_{\tilde{\chi}_c^-} D_0(m_{\tilde{\chi}_a^-}^2, m_{\tilde{\chi}_c^-}^2, m_{\nu_d^i}^2, m_{\nu_b^i}^2) \quad (\text{F.10})$$

$$S_{(c_1^l)}^{LR} = -W_{a,\alpha,d}^L \bar{X}_{\beta,a,b}^L W_{c,\gamma,b}^R \bar{X}_{\delta,c,d}^R m_{\tilde{\chi}_a^-} m_{\tilde{\chi}_c^-} D_0(m_{\tilde{\chi}_a^-}^2, m_{\tilde{\chi}_c^-}^2, m_{\nu_d^i}^2, m_{\nu_b^i}^2) \quad (\text{F.11})$$

$$V_{(c_1^l)}^{LL} = -W_{a,\alpha,d}^L \bar{X}_{\beta,a,b}^R W_{c,\gamma,b}^L \bar{X}_{\delta,c,d}^R D_{27}(m_{\tilde{\chi}_a^-}^2, m_{\tilde{\chi}_c^-}^2, m_{\nu_d^i}^2, m_{\nu_b^i}^2) \quad (\text{F.12})$$

$$V_{(c_1^l)}^{LR} = -W_{a,\alpha,d}^L \bar{X}_{\beta,a,b}^R W_{c,\gamma,b}^R \bar{X}_{\delta,c,d}^L D_{27}(m_{\tilde{\chi}_a^-}^2, m_{\tilde{\chi}_c^-}^2, m_{\nu_d^i}^2, m_{\nu_b^i}^2) \quad (\text{F.13})$$

$$S_{(c_2^l)}^{LL} = -W_{a,\alpha,d}^L \hat{\bar{X}}_{\beta,a,b}^L \hat{X}_{c,\gamma,b}^L \bar{X}_{\delta,c,d}^L m_{\tilde{\chi}_a^-} m_{\tilde{\chi}_c^-} D_0(m_{\tilde{\chi}_a^-}^2, m_{\tilde{\chi}_c^-}^2, m_{\nu_d^i}^2, m_{\nu_b^R}^2) \quad (\text{F.14})$$

$$S_{(c_2^l)}^{LR} = -W_{a,\alpha,d}^L \hat{\bar{X}}_{\beta,a,b}^L \hat{X}_{c,\gamma,b}^R \bar{X}_{\delta,c,d}^R m_{\tilde{\chi}_a^-} m_{\tilde{\chi}_c^-} D_0(m_{\tilde{\chi}_a^-}^2, m_{\tilde{\chi}_c^-}^2, m_{\nu_d^i}^2, m_{\nu_b^R}^2) \quad (\text{F.15})$$

$$V_{(c_2^l)}^{LL} = -W_{a,\alpha,d}^L \hat{\bar{X}}_{\beta,a,b}^R \hat{X}_{c,\gamma,b}^L \bar{X}_{\delta,c,d}^R D_{27}(m_{\tilde{\chi}_a^-}^2, m_{\tilde{\chi}_c^-}^2, m_{\nu_d^i}^2, m_{\nu_b^R}^2) \quad (\text{F.16})$$

$$V_{(c_2^l)}^{LR} = -W_{a,\alpha,d}^L \hat{\bar{X}}_{\beta,a,b}^R \hat{X}_{c,\gamma,b}^R \bar{X}_{\delta,c,d}^L D_{27}(m_{\tilde{\chi}_a^-}^2, m_{\tilde{\chi}_c^-}^2, m_{\nu_d^i}^2, m_{\nu_b^R}^2) \quad (\text{F.17})$$

$$S_{(c_3^l)}^{LL} = -\hat{X}_{a,\alpha,d}^L \bar{X}_{\beta,a,b}^L W_{c,\gamma,b}^L \hat{\bar{X}}_{\delta,c,d}^L m_{\tilde{\chi}_a^-} m_{\tilde{\chi}_c^-} D_0(m_{\tilde{\chi}_a^-}^2, m_{\tilde{\chi}_c^-}^2, m_{\nu_d^R}^2, m_{\nu_b^i}^2) \quad (\text{F.18})$$

$$S_{(c_3^l)}^{LR} = -\hat{X}_{a,\alpha,d}^L \bar{X}_{\beta,a,b}^L W_{c,\gamma,b}^R \hat{\bar{X}}_{\delta,c,d}^R m_{\tilde{\chi}_a^-} m_{\tilde{\chi}_c^-} D_0(m_{\tilde{\chi}_a^-}^2, m_{\tilde{\chi}_c^-}^2, m_{\nu_d^R}^2, m_{\nu_b^i}^2) \quad (\text{F.19})$$

$$V_{(c_3^l)}^{LL} = -\hat{X}_{a,\alpha,d}^L \bar{X}_{\beta,a,b}^R W_{c,\gamma,b}^L \hat{\bar{X}}_{\delta,c,d}^R D_{27}(m_{\tilde{\chi}_a^-}^2, m_{\tilde{\chi}_c^-}^2, m_{\nu_d^R}^2, m_{\nu_b^i}^2) \quad (\text{F.20})$$

$$V_{(c_3^l)}^{LR} = -\hat{X}_{a,\alpha,d}^L \bar{X}_{\beta,a,b}^R W_{c,\gamma,b}^R \hat{\bar{X}}_{\delta,c,d}^L D_{27}(m_{\tilde{\chi}_a^-}^2, m_{\tilde{\chi}_c^-}^2, m_{\nu_d^R}^2, m_{\nu_b^i}^2) \quad (\text{F.21})$$

$$S_{(c_4^l)}^{LL} = -\hat{X}_{a,\alpha,d}^L \hat{\bar{X}}_{\beta,a,b}^L \hat{X}_{c,\gamma,b}^L \hat{\bar{X}}_{\delta,c,d}^L m_{\tilde{\chi}_a^-} m_{\tilde{\chi}_c^-} D_0(m_{\tilde{\chi}_a^-}^2, m_{\tilde{\chi}_c^-}^2, m_{\nu_d^R}^2, m_{\nu_b^R}^2) \quad (\text{F.22})$$

$$S_{(c_4^l)}^{LR} = -\hat{X}_{a,\alpha,d}^L \hat{\bar{X}}_{\beta,a,b}^L \hat{X}_{c,\gamma,b}^R \hat{\bar{X}}_{\delta,c,d}^R m_{\tilde{\chi}_a^-} m_{\tilde{\chi}_c^-} D_0(m_{\tilde{\chi}_a^-}^2, m_{\tilde{\chi}_c^-}^2, m_{\nu_d^R}^2, m_{\nu_b^R}^2) \quad (\text{F.23})$$

$$V_{(c_4^l)}^{LL} = -\hat{X}_{a,\alpha,d}^L \hat{\bar{X}}_{\beta,a,b}^R \hat{X}_{c,\gamma,b}^L \hat{\bar{X}}_{\delta,c,d}^R D_{27}(m_{\tilde{\chi}_a^-}^2, m_{\tilde{\chi}_c^-}^2, m_{\nu_d^R}^2, m_{\nu_b^R}^2) \quad (\text{F.24})$$

$$V_{(c_4^l)}^{LR} = -\hat{X}_{a,\alpha,d}^L \hat{\bar{X}}_{\beta,a,b}^R \hat{X}_{c,\gamma,b}^R \hat{\bar{X}}_{\delta,c,d}^L D_{27}(m_{\tilde{\chi}_a^-}^2, m_{\tilde{\chi}_c^-}^2, m_{\nu_d^R}^2, m_{\nu_b^R}^2) \quad (\text{F.25})$$

$$S_{(c_5^l)}^{LL} = -W_{a,\alpha,d}^L \bar{X}_{\beta,a,b}^L \bar{X}_{\delta,c,b}^L W_{c,\gamma,d}^L m_{\tilde{\chi}_a^-} m_{\tilde{\chi}_c^-} D_0(m_{\tilde{\chi}_a^-}^2, m_{\tilde{\chi}_c^-}^2, m_{\nu_d^i}^2, m_{\nu_b^i}^2) \quad (\text{F.26})$$

$$S_{(c_5^l)}^{LR} = -W_{a,\alpha,d}^L \bar{X}_{\beta,a,b}^L \bar{X}_{\delta,c,b}^R W_{c,\gamma,d}^R m_{\tilde{\chi}_a^-} m_{\tilde{\chi}_c^-} D_0(m_{\tilde{\chi}_a^-}^2, m_{\tilde{\chi}_c^-}^2, m_{\nu_d^i}^2, m_{\nu_b^i}^2) \quad (\text{F.27})$$

$$V_{(c_5^l)}^{LL} = W_{a,\alpha,d}^L \bar{X}_{\beta,a,b}^R \bar{X}_{\delta,c,b}^L W_{c,\gamma,d}^L D_{27}(m_{\tilde{\chi}_a^-}^2, m_{\tilde{\chi}_c^-}^2, m_{\nu_d^i}^2, m_{\nu_b^i}^2) \quad (\text{F.28})$$

$$V_{(c_5^l)}^{LR} = W_{a,\alpha,d}^L \bar{X}_{\beta,a,b}^R \bar{X}_{\delta,c,b}^L W_{c,\gamma,d}^R D_{27}(m_{\tilde{\chi}_a^-}^2, m_{\tilde{\chi}_c^-}^2, m_{\nu_d^i}^2, m_{\nu_b^i}^2) \quad (\text{F.29})$$

$$S_{(c_6^l)}^{LL} = -W_{a,\alpha,d}^L \hat{\bar{X}}_{\beta,a,b}^L \hat{\bar{X}}_{\delta,c,b}^L W_{c,\gamma,d}^L m_{\tilde{\chi}_a^-} m_{\tilde{\chi}_c^-} D_0(m_{\tilde{\chi}_a^-}^2, m_{\tilde{\chi}_c^-}^2, m_{\nu_d^i}^2, m_{\nu_b^i}^2) \quad (\text{F.30})$$

$$S_{(c_6^l)}^{LR} = -W_{a,\alpha,d}^L \hat{\bar{X}}_{\beta,a,b}^L \hat{\bar{X}}_{\delta,c,b}^R W_{c,\gamma,d}^R m_{\tilde{\chi}_a^-} m_{\tilde{\chi}_c^-} D_0(m_{\tilde{\chi}_a^-}^2, m_{\tilde{\chi}_c^-}^2, m_{\nu_d^i}^2, m_{\nu_b^i}^2) \quad (\text{F.31})$$

$$V_{(c_6^l)}^{LL} = W_{a,\alpha,d}^L \hat{\bar{X}}_{\beta,a,b}^R \hat{\bar{X}}_{\delta,c,b}^R W_{c,\gamma,d}^L D_{27}(m_{\tilde{\chi}_a^-}^2, m_{\tilde{\chi}_c^-}^2, m_{\nu_d^i}^2, m_{\nu_b^i}^2) \quad (\text{F.32})$$

$$V_{(c_6^l)}^{LR} = W_{a,\alpha,d}^L \hat{\bar{X}}_{\beta,a,b}^R \hat{\bar{X}}_{\delta,c,b}^L W_{c,\gamma,d}^R D_{27}(m_{\tilde{\chi}_a^-}^2, m_{\tilde{\chi}_c^-}^2, m_{\nu_d^i}^2, m_{\nu_b^i}^2) \quad (\text{F.33})$$

$$S_{(c_7^l)}^{LL} = -\hat{X}_{a,\alpha,d}^L \bar{X}_{\beta,a,b}^L \bar{X}_{\delta,c,b}^L \hat{X}_{c,\gamma,d}^L m_{\tilde{\chi}_a^-} m_{\tilde{\chi}_c^-} D_0(m_{\tilde{\chi}_a^-}^2, m_{\tilde{\chi}_c^-}^2, m_{\nu_d^R}^2, m_{\nu_b^R}^2) \quad (\text{F.34})$$

$$S_{(c_7^l)}^{LR} = -\hat{X}_{a,\alpha,d}^L \bar{X}_{\beta,a,b}^L \bar{X}_{\delta,c,b}^R \hat{X}_{c,\gamma,d}^R m_{\tilde{\chi}_a^-} m_{\tilde{\chi}_c^-} D_0(m_{\tilde{\chi}_a^-}^2, m_{\tilde{\chi}_c^-}^2, m_{\nu_d^R}^2, m_{\nu_b^R}^2) \quad (\text{F.35})$$

$$V_{(c_7^l)}^{LL} = \hat{X}_{a,\alpha,d}^L \bar{X}_{\beta,a,b}^R \bar{X}_{\delta,c,b}^R \hat{X}_{c,\gamma,d}^L D_{27}(m_{\tilde{\chi}_a^-}^2, m_{\tilde{\chi}_c^-}^2, m_{\nu_d^R}^2, m_{\nu_b^R}^2) \quad (\text{F.36})$$

$$V_{(c_7^l)}^{LR} = \hat{X}_{a,\alpha,d}^L \bar{X}_{\beta,a,b}^R \bar{X}_{\delta,c,b}^L \hat{X}_{c,\gamma,d}^R D_{27}(m_{\tilde{\chi}_a^-}^2, m_{\tilde{\chi}_c^-}^2, m_{\nu_d^R}^2, m_{\nu_b^R}^2) \quad (\text{F.37})$$

$$S_{(c_8^l)}^{LL} = -\hat{X}_{a,\alpha,d}^L \hat{\bar{X}}_{\beta,a,b}^L \hat{\bar{X}}_{\delta,c,b}^L \hat{X}_{c,\gamma,d}^L m_{\tilde{\chi}_a^-} m_{\tilde{\chi}_c^-} D_0(m_{\tilde{\chi}_a^-}^2, m_{\tilde{\chi}_c^-}^2, m_{\nu_d^R}^2, m_{\nu_b^R}^2) \quad (\text{F.38})$$

$$S_{(c_8^l)}^{LR} = -\hat{X}_{a,\alpha,d}^L \hat{\bar{X}}_{\beta,a,b}^L \hat{\bar{X}}_{\delta,c,b}^R \hat{X}_{c,\gamma,d}^R m_{\tilde{\chi}_a^-} m_{\tilde{\chi}_c^-} D_0(m_{\tilde{\chi}_a^-}^2, m_{\tilde{\chi}_c^-}^2, m_{\nu_d^R}^2, m_{\nu_b^R}^2) \quad (\text{F.39})$$

$$V_{(c_8^l)}^{LL} = \hat{X}_{a,\alpha,d}^L \hat{\bar{X}}_{\beta,a,b}^R \hat{\bar{X}}_{\delta,c,b}^R \hat{X}_{c,\gamma,d}^L D_{27}(m_{\tilde{\chi}_a^-}^2, m_{\tilde{\chi}_c^-}^2, m_{\nu_d^R}^2, m_{\nu_b^R}^2) \quad (\text{F.40})$$

$$V_{(c_8^l)}^{LR} = \hat{X}_{a,\alpha,d}^L \hat{\bar{X}}_{\beta,a,b}^R \hat{\bar{X}}_{\delta,c,b}^L \hat{X}_{c,\gamma,d}^R D_{27}(m_{\tilde{\chi}_a^-}^2, m_{\tilde{\chi}_c^-}^2, m_{\nu_d^R}^2, m_{\nu_b^R}^2) \quad (\text{F.41})$$

F.1.4 W^+ and H^+ contributions

$$S_{(w_1^l)}^{LL} = -V_{a,\alpha,d}^{+,L} \bar{V}_{\beta,a,b}^{+,L} V_{c,\gamma,d}^{+,L} \bar{V}_{\delta,c,d}^{+,L} m_{\nu_a} m_{\nu_c} D_0(m_{\nu_a}^2, m_{\nu_c}^2, m_{H_d^-}^2, m_{H_b^-}^2) \quad (\text{F.42})$$

$$S_{(w_1^l)}^{LR} = -V_{a,\alpha,d}^{+,L} \bar{V}_{\beta,a,b}^{+,L} V_{c,\gamma,b}^{+,R} \bar{V}_{\delta,c,d}^{+,R} m_{\nu_a} m_{\nu_c} D_0(m_{\nu_a}^2, m_{\nu_c}^2, m_{H_d^-}^2, m_{H_b^-}^2) \quad (\text{F.43})$$

$$V_{(w_1^l)}^{LL} = -V_{a,\alpha,d}^{+,L} \bar{V}_{\beta,a,b}^{+,R} V_{c,\gamma,b}^{+,L} \bar{V}_{\delta,c,d}^{+,R} D_{27}(m_{\nu_a}^2, m_{\nu_c}^2, m_{H_d^-}^2, m_{H_b^-}^2) \quad (\text{F.44})$$

$$V_{(w_1^l)}^{LR} = -V_{a,\alpha,d}^{+,L} \bar{V}_{\beta,a,b}^{+,R} V_{c,\gamma,b}^{+,R} \bar{V}_{\delta,c,d}^{+,L} D_{27}(m_{\nu_a}^2, m_{\nu_c}^2, m_{H_d^-}^2, m_{H_b^-}^2) \quad (\text{F.45})$$

$$S_{(w_2^l)}^{LL} = 2 V_{a,\alpha,d}^{+,L} \hat{\bar{V}}_{\beta,a}^{+,R} \hat{V}_{c,\gamma}^{+,L} \bar{V}_{\delta,c,d}^{+,L} (I_{C_0 D_0}(m_{\nu_c}^2, m_{W^-}^2, m_{H_d^-}^2, m_{\nu_a}^2) - 2D_{27}(m_{\nu_a}^2, m_{\nu_c}^2, m_{W^-}^2, m_{H_d^-}^2)) \quad (\text{F.46})$$

$$S_{(w_2^l)}^{LR} = 2 V_{a,\alpha,d}^{+,L} \hat{\bar{V}}_{\beta,a}^{+,R} \hat{V}_{c,\gamma}^{+,R} \bar{V}_{\delta,c,d}^{+,R} (I_{C_0 D_0}(m_{\nu_c}^2, m_{W^-}^2, m_{H_d^-}^2, m_{\nu_a}^2) - 2D_{27}(m_{\nu_a}^2, m_{\nu_c}^2, m_{W^-}^2, m_{H_d^-}^2)) \quad (\text{F.47})$$

$$V_{(w_2^l)}^{LL} = V_{a,\alpha,d}^{+,L} \hat{\bar{V}}_{\beta,a}^{+,L} \hat{V}_{c,\gamma}^{+,L} \bar{V}_{\delta,c,d}^{+,R} m_{\nu_a} m_{\nu_c} D_0(m_{\nu_a}^2, m_{\nu_c}^2, m_{W^-}^2, m_{H_d^-}^2) \quad (\text{F.48})$$

$$V_{(w_2^l)}^{LR} = V_{a,\alpha,d}^{+,L} \hat{V}_{\beta,a}^{+,L} \hat{V}_{c,\gamma}^{+,R} \bar{V}_{\delta,c,d}^{+,L} m_{\nu_a} m_{\nu_c} D_0(m_{\nu_a}^2, m_{\nu_c}^2, m_{W-}^2, m_{H_d^-}^2) \quad (\text{F.49})$$

$$S_{(w_3^l)}^{LL} = 2 \hat{V}_{a,i}^{+,L} \bar{V}_{\beta,a,b}^{+,L} V_{c,\gamma,b}^{+,L} \hat{V}_{\delta,c}^{+,R} (I_{C_0 D_0}(m_{\nu_c}^2, m_{H_b^-}^2, m_{W-}^2, m_{\nu_a}^2) - 2D_{27}(m_{\nu_a}^2, m_{\nu_c}^2, m_{H_b^-}^2, m_{W-}^2)) \quad (\text{F.50})$$

$$S_{(w_3^l)}^{LR} = 2 \hat{V}_{a,i}^{+,L} \bar{V}_{\beta,a,b}^{+,L} V_{c,\gamma,b}^{+,R} \hat{V}_{\delta,c}^{+,L} (I_{C_0 D_0}(m_{\nu_c}^2, m_{H_b^-}^2, m_{W-}^2, m_{\nu_a}^2) - 2D_{27}(m_{\nu_a}^2, m_{\nu_c}^2, m_{H_b^-}^2, m_{W-}^2)) \quad (\text{F.51})$$

$$V_{(w_3^l)}^{LL} = \hat{V}_{a,i}^{+,L} \bar{V}_{\beta,a,b}^{+,R} V_{c,\gamma,b}^{+,L} \hat{V}_{\delta,c}^{+,L} m_{\nu_a} m_{\nu_c} D_0(m_{\nu_a}^2, m_{\nu_c}^2, m_{H_b^-}^2, m_{W-}^2) \quad (\text{F.52})$$

$$V_{(w_3^l)}^{LR} = \hat{V}_{a,i}^{+,L} \bar{V}_{\beta,a,b}^{+,R} V_{c,\gamma,b}^{+,R} \hat{V}_{\delta,c}^{+,R} m_{\nu_a} m_{\nu_c} D_0(m_{\nu_a}^2, m_{\nu_c}^2, m_{H_b^-}^2, m_{W-}^2) \quad (\text{F.53})$$

$$T_{(w_3^l)}^{LL} = -\hat{V}_{a,i}^{+,L} \bar{V}_{\beta,a,b}^{+,L} V_{c,\gamma,b}^{+,L} \hat{V}_{\delta,c}^{+,R} D_{27}(m_{\nu_a}^2, m_{\nu_c}^2, m_{H_b^-}^2, m_{W-}^2) \quad (\text{F.54})$$

$$S_{(w_4^l)}^{LL} = -4 \hat{V}_{a,i}^{+,L} \hat{V}_{\beta,a}^{+,R} \hat{V}_{c,\gamma}^{+,L} \hat{V}_{\delta,c}^{+,R} m_{\nu_a} m_{\nu_c} D_0(m_{\nu_a}^2, m_{\nu_c}^2, m_{W-}^2, m_{W-}^2) \quad (\text{F.55})$$

$$S_{(w_4^l)}^{LR} = -4 \hat{V}_{a,i}^{+,L} \hat{V}_{\beta,a}^{+,R} \hat{V}_{c,\gamma}^{+,L} \hat{V}_{\delta,c}^{+,R} m_{\nu_a} m_{\nu_c} D_0(m_{\nu_a}^2, m_{\nu_c}^2, m_{W-}^2, m_{W-}^2) \quad (\text{F.56})$$

$$V_{(w_4^l)}^{LL} = -4 \hat{V}_{a,i}^{+,L} \hat{V}_{\beta,a}^{+,L} \hat{V}_{c,\gamma}^{+,L} \hat{V}_{\delta,c}^{+,L} (I_{C_0 D_0}(m_{\nu_c}^2, m_{W-}^2, m_{W-}^2, m_{\nu_a}^2) - 3D_{27}(m_{\nu_a}^2, m_{\nu_c}^2, m_{W-}^2, m_{W-}^2)) \quad (\text{F.57})$$

$$V_{(w_4^l)}^{LR} = -4 \hat{V}_{a,i}^{+,L} \hat{V}_{\beta,a}^{+,L} \hat{V}_{c,\gamma}^{+,L} \hat{V}_{\delta,c}^{+,R} I_{C_0 D_0}(m_{\nu_c}^2, m_{W-}^2, m_{\nu_a}^2) \quad (\text{F.58})$$

$$T_{(w_4^l)}^{LL} = \hat{V}_{a,i}^{+,L} \hat{V}_{\beta,a}^{+,R} \hat{V}_{c,\gamma}^{+,L} \hat{V}_{\delta,c}^{+,R} m_{\nu_a} m_{\nu_c} D_0(m_{\nu_a}^2, m_{\nu_c}^2, m_{W-}^2, m_{W-}^2) \quad (\text{F.59})$$

$$S_{(w_5^l)}^{LL} = \frac{1}{2} V_{a,\alpha,d}^{+,L} \bar{V}_{\delta,c,d}^{+,L} V_{a,\gamma,b}^{+,L} \bar{V}_{\beta,c,b}^{+,L} m_{\nu_a} m_{\nu_c} D_0(m_{\nu_a}^2, m_{\nu_c}^2, m_{H_d^-}^2, m_{H_b^-}^2) \quad (\text{F.60})$$

$$S_{(w_5^l)}^{LR} = -2 V_{a,\alpha,d}^{+,L} \bar{V}_{\delta,c,d}^{+,R} V_{a,\gamma,b}^{+,R} \bar{V}_{\beta,c,b}^{+,L} D_{27}(m_{\nu_a}^2, m_{\nu_c}^2, m_{H_d^-}^2, m_{H_b^-}^2) \quad (\text{F.61})$$

$$V_{(w_5^l)}^{LL} = -\frac{1}{2} V_{a,\alpha,d}^{+,L} \bar{V}_{\delta,c,d}^{+,R} V_{a,\gamma,b}^{+,L} \bar{V}_{\beta,c,b}^{+,R} m_{\nu_a} m_{\nu_c} D_0(m_{\nu_a}^2, m_{\nu_c}^2, m_{H_d^-}^2, m_{H_b^-}^2) \quad (\text{F.62})$$

$$V_{(w_5^l)}^{LR} = -V_{a,\alpha,d}^{+,L} \bar{V}_{\delta,c,d}^{+,L} V_{a,\gamma,b}^{+,R} \bar{V}_{\beta,c,b}^{+,R} D_{27}(m_{\nu_a}^2, m_{\nu_c}^2, m_{H_d^-}^2, m_{H_b^-}^2) \quad (\text{F.63})$$

$$T_{(w_5^l)}^{LL} = \frac{1}{8} V_{a,\alpha,d}^{+,L} \bar{V}_{\delta,c,d}^{+,L} V_{a,\gamma,b}^{+,L} \bar{V}_{\beta,c,b}^{+,L} m_{\nu_a} m_{\nu_c} D_0(m_{\nu_a}^2, m_{\nu_c}^2, m_{H_d^-}^2, m_{H_b^-}^2) \quad (\text{F.64})$$

$$S_{(w_6^l)}^{LL} = -V_{a,\alpha,d}^{+,L} \bar{V}_{\delta,c,d}^{+,L} \hat{V}_{a,\gamma}^{+,L} \hat{V}_{\beta,c}^{+,R} (I_{C_0 D_0}(m_{\nu_c}^2, m_{W-}^2, m_{H_d^-}^2, m_{\nu_a}^2) - 8D_{27}(m_{\nu_a}^2, m_{\nu_c}^2, m_{W-}^2, m_{H_d^-}^2)) \quad (\text{F.65})$$

$$S_{(w_6^l)}^{LR} = 2 V_{a,\alpha,d}^{+,L} \bar{V}_{\delta,c,d}^{+,R} \hat{V}_{a,\gamma}^{+,R} \hat{V}_{\beta,c}^{+,R} m_{\nu_a} m_{\nu_c} D_0(m_{\nu_a}^2, m_{\nu_c}^2, m_{W-}^2, m_{H_d^-}^2) \quad (\text{F.66})$$

$$V_{(w_6^l)}^{LL} = V_{a,\alpha,d}^{+,L} \bar{V}_{\delta,c,d}^{+,R} \hat{V}_{a,\gamma}^{+,R} \hat{V}_{\beta,c}^{+,L} (I_{C_0 D_0}(m_{\nu_c}^2, m_{W-}^2, m_{H_d^-}^2, m_{\nu_a}^2) - 2D_{27}(m_{\nu_a}^2, m_{\nu_c}^2, m_{W-}^2, m_{H_d^-}^2)) \quad (\text{F.67})$$

$$V_{(w_6^l)}^{LR} l = V_{a,\alpha,d}^{+,L} \bar{V}_{\delta,c,d}^{+,R} \hat{V}_{a,\gamma}^{+,R} \hat{V}_{\beta,c}^{+,L} m_{\nu_a} m_{\nu_c} D_0(m_{\nu_a}^2, m_{\nu_c}^2, m_{W-}^2, m_{H_d^-}^2) \quad (\text{F.68})$$

$$T_{(w_6^l)}^{LL} = -\frac{1}{4} V_{a,\alpha,d}^{+,L} \bar{V}_{\delta,c,d}^{+,L} \hat{V}_{a,\gamma}^{+,L} \hat{V}_{\beta,c}^{+,R} I_{C_0 D_0}(m_{\nu_c}^2, m_{W-}^2, m_{H_d^-}^2, m_{\nu_a}^2) \quad (\text{F.69})$$

$$S_{(w_7^l)}^{LL} = -\hat{V}_{a,i}^{+,L} \bar{V}_{\delta,c}^{+,R} V_{a,\gamma,b}^{+,L} \bar{V}_{\beta,c,b}^{+,L} (I_{C_0 D_0}(m_{\nu_c}^2, m_{H_b^-}^2, m_{W-}^2, m_{\nu_a}^2) - 8D_{27}(m_{\nu_a}^2, m_{\nu_c}^2, m_{H_b^-}^2, m_{W-}^2)) \quad (\text{F.70})$$

$$S_{(w_7^l)}^{LR} = 2 \hat{V}_{a,i}^{+,L} \bar{V}_{\delta,c}^{+,R} V_{a,\gamma,b}^{+,R} \bar{V}_{\beta,c,b}^{+,L} m_{\nu_a} m_{\nu_c} D_0(m_{\nu_a}^2, m_{\nu_c}^2, m_{H_b^-}^2, m_{W-}^2) \quad (\text{F.71})$$

$$V_{(w_7^l)}^{LL} = \hat{V}_{a,i}^{+,L} \bar{V}_{\delta,c}^{+,L} V_{a,\gamma,b}^{+,L} \bar{V}_{\beta,c,b}^{+,R} (I_{C_0 D_0}(m_{\nu_c}^2, m_{H_b^-}^2, m_{W-}^2, m_{\nu_a}^2) - 2D_{27}(m_{\nu_a}^2, m_{\nu_c}^2, m_{H_b^-}^2, m_{W-}^2)) \quad (\text{F.72})$$

$$V_{(w_7^l)}^{LR} = \hat{V}_{a,i}^{+,L} \bar{V}_{\delta,c}^{+,R} V_{a,\gamma,b}^{+,R} \bar{V}_{\beta,c,b}^{+,R} m_{\nu_a} m_{\nu_c} D_0(m_{\nu_a}^2, m_{\nu_c}^2, m_{H_b^-}^2, m_{W-}^2) \quad (\text{F.73})$$

$$T_{(w_7^l)}^{LL} = -\frac{1}{4} \hat{V}_{a,i}^{+,L} \bar{V}_{\delta,c}^{+,R} V_{a,\gamma,b}^{+,L} \bar{V}_{\beta,c,b}^{+,L} I_{C_0 D_0}(m_{\nu_c}^2, m_{H_b^-}^2, m_{W-}^2, m_{\nu_a}^2) \quad (\text{F.74})$$

$$S_{(w_8^l)}^{LL} = -4 \hat{V}_{a,i}^{+,L} \bar{V}_{\delta,c}^{+,R} \hat{V}_{a,\gamma}^{+,L} \bar{V}_{\beta,c}^{+,R} m_{\nu_a} m_{\nu_c} D_0(m_{\nu_a}^2, m_{\nu_c}^2, m_{W-}^2, m_{W-}^2) \quad (\text{F.75})$$

$$S_{(w_8^l)}^{LR} = -8 \hat{V}_{a,i}^{+,L} \bar{V}_{\delta,c}^{+,L} \hat{V}_{a,\gamma}^{+,R} \bar{V}_{\beta,c}^{+,R} (I_{C_0 D_0}(m_{\nu_c}^2, m_{W-}^2, m_{W-}^2, m_{\nu_a}^2) - 3D_{27}(m_{\nu_a}^2, m_{\nu_c}^2, m_{W-}^2, m_{W-}^2)) \quad (\text{F.76})$$

$$V_{(w_8^l)}^{LL} = -2 \hat{V}_{a,i}^{+,L} \bar{V}_{\delta,c}^{+,R} \hat{V}_{a,\gamma}^{+,L} \bar{V}_{\beta,c}^{+,L} m_{\nu_a} m_{\nu_c} D_0(m_{\nu_a}^2, m_{\nu_c}^2, m_{W-}^2, m_{W-}^2) \quad (\text{F.77})$$

$$V_{(w_8^l)}^{LR} = -4\hat{V}_{a,i}^{+,L}\hat{V}_{\delta,c}^{+,R}\hat{V}_{a,\gamma}^{+,R}\hat{V}_{\beta,c}^{+,L}I_{C_0D_0}(m_{\nu_c}^2, m_{W^-}^2, m_{W^-}^2, m_{\nu_a}^2) \quad (\text{F.78})$$

$$T_{(w_8^l)}^{LL} = \hat{V}_{a,i}^{+,L}\hat{V}_{\delta,c}^{+,R}\hat{V}_{a,\gamma}^{+,L}\hat{V}_{\beta,c}^{+,R}m_{\nu_a}m_{\nu_c}D_0(m_{\nu_a}^2, m_{\nu_c}^2, m_{W^-}^2, m_{W^-}^2) \quad (\text{F.79})$$

F.2 Additional boxes for $\ell_\alpha^- \rightarrow \ell_\beta^- \ell_\gamma^+ \ell_\gamma^-$

In the case of $\ell_\alpha^- \rightarrow \ell_\beta^- \ell_\gamma^+ \ell_\gamma^-$ it is necessary to calculate the crossed diagrams with exchanged indices $\beta \leftrightarrow \gamma$ explicitly.

F.2.1 Crossed neutralino contributions

$$S_{(n_1^l)}^{LL} = \frac{1}{2}N_{d,\alpha,a}^L\bar{N}_{\beta,b,a}^L N_{b,\gamma,c}^L\bar{N}_{\delta,d,c}^L m_{\tilde{\chi}_b^0}m_{\tilde{\chi}_d^0}D_0(m_{\tilde{\chi}_d^0}^2, m_{\tilde{\chi}_b^0}^2, m_{\tilde{e}_a}^2, m_{\tilde{e}_c}^2) \quad (\text{F.80})$$

$$S_{(n_1^l)}^{LR} = 2N_{d,\alpha,a}^L\bar{N}_{\beta,b,a}^L N_{b,\gamma,c}^R\bar{N}_{\delta,d,c}^R D_{27}(m_{\tilde{\chi}_d^0}^2, m_{\tilde{\chi}_b^0}^2, m_{\tilde{e}_a}^2, m_{\tilde{e}_c}^2) \quad (\text{F.81})$$

$$V_{(n_1^l)}^{LL} = -N_{d,\alpha,a}^L\bar{N}_{\beta,b,a}^R N_{b,\gamma,c}^L\bar{N}_{\delta,d,c}^R D_{27}(m_{\tilde{\chi}_d^0}^2, m_{\tilde{\chi}_b^0}^2, m_{\tilde{e}_a}^2, m_{\tilde{e}_c}^2) \quad (\text{F.82})$$

$$V_{(n_1^l)}^{LR} = \frac{1}{2}N_{d,\alpha,a}^L\bar{N}_{\beta,b,a}^R N_{b,\gamma,c}^R\bar{N}_{\delta,d,c}^L m_{\tilde{\chi}_b^0}m_{\tilde{\chi}_d^0}D_0(m_{\tilde{\chi}_d^0}^2, m_{\tilde{\chi}_b^0}^2, m_{\tilde{e}_a}^2, m_{\tilde{e}_c}^2) \quad (\text{F.83})$$

$$T_{(n_1^l)}^{LL} = -\frac{1}{8}N_{d,\alpha,a}^L\bar{N}_{\beta,b,a}^L N_{b,\gamma,c}^L\bar{N}_{\delta,d,c}^L m_{\tilde{\chi}_b^0}m_{\tilde{\chi}_d^0}D_0(m_{\tilde{\chi}_d^0}^2, m_{\tilde{\chi}_b^0}^2, m_{\tilde{e}_a}^2, m_{\tilde{e}_c}^2) \quad (\text{F.84})$$

$$S_{(n_2^l)}^{LL} = \frac{1}{2}N_{d,\alpha,a}^L\bar{N}_{\beta,b,a}^L N_{\delta,b,c}^L\bar{N}_{d,\gamma,c}^L m_{\tilde{\chi}_b^0}m_{\tilde{\chi}_d^0}D_0(m_{\tilde{\chi}_b^0}^2, m_{\tilde{\chi}_d^0}^2, m_{\tilde{e}_a}^2, m_{\tilde{e}_c}^2) \quad (\text{F.85})$$

$$S_{(n_2^l)}^{LR} = 2N_{d,\alpha,a}^L\bar{N}_{\beta,b,a}^L N_{\delta,b,c}^R\bar{N}_{d,\gamma,c}^R D_{27}(m_{\tilde{\chi}_b^0}^2, m_{\tilde{\chi}_d^0}^2, m_{\tilde{e}_a}^2, m_{\tilde{e}_c}^2) \quad (\text{F.86})$$

$$V_{(n_2^l)}^{LL} = -\frac{1}{2}N_{d,\alpha,a}^L\bar{N}_{\beta,b,a}^R N_{\delta,b,c}^L\bar{N}_{d,\gamma,c}^L m_{\tilde{\chi}_b^0}m_{\tilde{\chi}_d^0}D_0(m_{\tilde{\chi}_b^0}^2, m_{\tilde{\chi}_d^0}^2, m_{\tilde{e}_a}^2, m_{\tilde{e}_c}^2) \quad (\text{F.87})$$

$$V_{(n_2^l)}^{LR} = N_{d,\alpha,a}^L\bar{N}_{\beta,b,a}^R\bar{N}_{\delta,b,c}^L N_{d,\gamma,c}^R D_{27}(m_{\tilde{\chi}_b^0}^2, m_{\tilde{\chi}_d^0}^2, m_{\tilde{e}_a}^2, m_{\tilde{e}_c}^2) \quad (\text{F.88})$$

$$T_{(n_2^l)}^{LL} = \frac{1}{8}N_{d,\alpha,a}^L\bar{N}_{\beta,b,a}^L\bar{N}_{\delta,b,c}^L N_{d,\gamma,c}^L m_{\tilde{\chi}_b^0}m_{\tilde{\chi}_d^0}D_0(m_{\tilde{\chi}_b^0}^2, m_{\tilde{\chi}_d^0}^2, m_{\tilde{e}_a}^2, m_{\tilde{e}_c}^2) \quad (\text{F.89})$$

$$(F.90)$$

F.2.2 Crossed chargino contributions

$$S_{(c_1^l)}^{LL} = \frac{1}{2}X_{d,\alpha,a}^L\bar{X}_{\delta,d,c}^L X_{b,\gamma,a}^L\bar{X}_{\beta,b,c}^L m_{\tilde{\chi}_b^-}m_{\tilde{\chi}_d^-}D_0(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^i}^2, m_{\nu_c^i}^2) \quad (\text{F.91})$$

$$S_{(c_1^l)}^{LR} = -2X_{d,\alpha,a}^L\bar{X}_{\delta,d,c}^R X_{b,\gamma,a}^R\bar{X}_{\beta,b,c}^L D_{27}(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^i}^2, m_{\nu_c^i}^2) \quad (\text{F.92})$$

$$V_{(c_1^l)}^{LL} = X_{d,\alpha,a}^L\bar{X}_{\delta,d,c}^R X_{b,\gamma,a}^L\bar{X}_{\beta,b,c}^R D_{27}(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^i}^2, m_{\nu_c^i}^2) \quad (\text{F.93})$$

$$V_{(c_1^l)}^{LR} = \frac{1}{2}X_{d,\alpha,a}^L\bar{X}_{\delta,d,c}^L X_{b,\gamma,a}^R\bar{X}_{\beta,b,c}^R m_{\tilde{\chi}_b^-}m_{\tilde{\chi}_d^-}D_0(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^i}^2, m_{\nu_c^i}^2) \quad (\text{F.94})$$

$$T_{(c_1^l)}^{LL} = -\frac{1}{8}X_{d,\alpha,a}^L\bar{X}_{\delta,d,c}^L X_{b,\gamma,a}^L\bar{X}_{\beta,b,c}^L m_{\tilde{\chi}_d^-}D_0(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^i}^2, m_{\nu_c^i}^2) \quad (\text{F.95})$$

$$S_{(c_2^l)}^{LL} = \frac{1}{2}X_{d,\alpha,a}^L\bar{X}_{\beta,b,a}^L X_{b,\gamma,c}^L\bar{X}_{\delta,d,c}^L m_{\tilde{\chi}_b^-}m_{\tilde{\chi}_d^-}D_0(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^i}^2, m_{\nu_c^R}^2) \quad (\text{F.96})$$

$$S_{(c_2')}^{LR} = 2 X_{d,\alpha,a}^L \bar{X}_{\beta,b,a}^L \hat{X}_{b,\gamma,c}^R \hat{\bar{X}}_{\delta,d,c}^R D_{27}(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^i}^2, m_{\nu_c^R}^2) \quad (\text{F.97})$$

$$V_{(c_2')}^{LL} = -X_{d,\alpha,a}^L \bar{X}_{\beta,b,a}^R \hat{X}_{b,\gamma,c}^L \hat{\bar{X}}_{\delta,d,c}^R D_{27}(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^i}^2, m_{\nu_c^R}^2) \quad (\text{F.98})$$

$$V_{(c_2')}^{LR} = \frac{1}{2} X_{d,\alpha,a}^L \bar{X}_{\beta,b,a}^R \hat{X}_{b,\gamma,c}^R \hat{\bar{X}}_{\delta,d,c}^L m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_d^-} D_0(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^i}^2, m_{\nu_c^R}^2) \quad (\text{F.99})$$

$$T_{(c_2')}^{LL} = -\frac{1}{8} X_{d,\alpha,a}^L \bar{X}_{\beta,b,a}^L \hat{X}_{b,\gamma,c}^L m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_d^-} D_0(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^i}^2, m_{\nu_c^R}^2) \quad (\text{F.100})$$

$$S_{(c_3')}^{LL} = \frac{1}{2} \hat{X}_{d,\alpha,a}^L \hat{\bar{X}}_{\beta,b,a}^L X_{b,\gamma,c}^L \bar{X}_{\delta,d,c}^L m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_d^-} D_0(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^R}^2, m_{\nu_c^i}^2) \quad (\text{F.101})$$

$$S_{(c_3')}^{LR} = 2 \hat{X}_{d,\alpha,a}^L \hat{\bar{X}}_{\beta,b,a}^L X_{b,\gamma,c}^R \bar{X}_{\delta,d,c}^R D_{27}(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^R}^2, m_{\nu_c^i}^2) \quad (\text{F.102})$$

$$V_{(c_3')}^{LL} = -\hat{X}_{d,\alpha,a}^L \hat{\bar{X}}_{\beta,b,a}^R X_{b,\gamma,c}^L \bar{X}_{\delta,d,c}^R D_{27}(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^R}^2, m_{\nu_c^i}^2) \quad (\text{F.103})$$

$$V_{(c_3')}^{LR} = \frac{1}{2} \hat{X}_{d,\alpha,a}^L \hat{\bar{X}}_{\beta,b,a}^R X_{b,\gamma,c}^R \bar{X}_{\delta,d,c}^L m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_d^-} D_0(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^R}^2, m_{\nu_c^i}^2) \quad (\text{F.104})$$

$$T_{(c_3')}^{LL} = -\frac{1}{8} \hat{X}_{d,\alpha,a}^L \hat{\bar{X}}_{\beta,b,a}^L X_{b,\gamma,c}^L \bar{X}_{\delta,d,c}^L m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_d^-} D_0(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^R}^2, m_{\nu_c^i}^2) \quad (\text{F.105})$$

$$S_{(c_4')}^{LL} = \frac{1}{2} \hat{X}_{d,\alpha,a}^L \hat{\bar{X}}_{\beta,b,a}^L \hat{X}_{b,\gamma,c}^L \bar{X}_{\delta,d,c}^L m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_d^-} D_0(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^R}^2, m_{\nu_c^R}^2) \quad (\text{F.106})$$

$$S_{(c_4')}^{LR} = 2 \hat{X}_{d,\alpha,a}^L \hat{\bar{X}}_{\beta,b,a}^L \hat{X}_{b,\gamma,c}^R \bar{X}_{\delta,d,c}^R D_{27}(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^R}^2, m_{\nu_c^R}^2) \quad (\text{F.107})$$

$$V_{(c_4')}^{LL} = -\hat{X}_{d,\alpha,a}^L \hat{\bar{X}}_{\beta,b,a}^R \hat{X}_{b,\gamma,c}^L \bar{X}_{\delta,d,c}^R D_{27}(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^R}^2, m_{\nu_c^R}^2) \quad (\text{F.108})$$

$$V_{(c_4')}^{LR} = \frac{1}{2} \hat{X}_{d,\alpha,a}^L \hat{\bar{X}}_{\beta,b,a}^R \hat{X}_{b,\gamma,c}^R \bar{X}_{\delta,d,c}^L m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_d^-} D_0(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^R}^2, m_{\nu_c^R}^2) \quad (\text{F.109})$$

$$T_{(c_4')}^{LL} = -\frac{1}{8} \hat{X}_{d,\alpha,a}^L \hat{\bar{X}}_{\beta,b,a}^L \hat{X}_{b,\gamma,c}^L \bar{X}_{\delta,d,c}^L m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_d^-} D_0(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^R}^2, m_{\nu_c^R}^2) \quad (\text{F.110})$$

$$S_{(c_5')}^{LL} = \frac{1}{2} X_{d,\alpha,a}^L \bar{X}_{\delta,d,c}^L X_{b,\gamma,a}^L \bar{X}_{\beta,b,c}^L m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_d^-} D_0(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^i}^2, m_{\nu_c^i}^2) \quad (\text{F.111})$$

$$S_{(c_5')}^{LR} = -2 X_{d,\alpha,a}^L \bar{X}_{\delta,d,c}^R X_{b,\gamma,a}^R \bar{X}_{\beta,b,c}^L D_{27}(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^i}^2, m_{\nu_c^i}^2) \quad (\text{F.112})$$

$$V_{(c_5')}^{LL} = X_{d,\alpha,a}^L \bar{X}_{\delta,d,c}^R X_{b,\gamma,a}^L \bar{X}_{\beta,b,c}^R D_{27}(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^i}^2, m_{\nu_c^i}^2) \quad (\text{F.113})$$

$$V_{(c_5')}^{LR} = \frac{1}{2} X_{d,\alpha,a}^L \bar{X}_{\delta,d,c}^L X_{b,\gamma,a}^R \bar{X}_{\beta,b,c}^R m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_d^-} D_0(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^i}^2, m_{\nu_c^i}^2) \quad (\text{F.114})$$

$$T_{(c_5')}^{LL} = -\frac{1}{8} X_{d,\alpha,a}^L \bar{X}_{\delta,d,c}^L X_{b,\gamma,a}^R \bar{X}_{\beta,b,c}^L m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_d^-} D_0(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^i}^2, m_{\nu_c^i}^2) \quad (\text{F.115})$$

$$S_{(c_6')}^{LL} = \frac{1}{2} X_{d,\alpha,a}^L \hat{\bar{X}}_{\delta,d,c}^L X_{b,\gamma,a}^L \hat{\bar{X}}_{\beta,b,c}^R m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_d^-} D_0(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^i}^2, m_{\nu_c^R}^2) \quad (\text{F.116})$$

$$S_{(c_6')}^{LR} = -2 X_{d,\alpha,a}^L \hat{\bar{X}}_{\delta,d,c}^R X_{b,\gamma,a}^R \hat{\bar{X}}_{\beta,b,c}^L D_{27}(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^i}^2, m_{\nu_c^R}^2) \quad (\text{F.117})$$

$$V_{(c_6')}^{LL} = X_{d,\alpha,a}^L \hat{\bar{X}}_{\delta,d,c}^R X_{b,\gamma,a}^L \hat{\bar{X}}_{\beta,b,c}^R D_{27}(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^i}^2, m_{\nu_c^R}^2) \quad (\text{F.118})$$

$$V_{(c_6^l)}^{LR} = \frac{1}{2} X_{d,\alpha,a}^L \hat{X}_{\delta,d,c}^L X_{b,\gamma,a}^R \hat{X}_{\beta,b,c}^R m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_d^-} D_0(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^i}^2, m_{\nu_c^R}^2) \quad (\text{F.119})$$

$$T_{(c_6^l)}^{LL} = -\frac{1}{8} X_{d,\alpha,a}^L \hat{X}_{\delta,d,c}^L X_{b,\gamma,a}^L \hat{X}_{\beta,b,c}^L m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_d^-} D_0(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^i}^2, m_{\nu_c^R}^2) \quad (\text{F.120})$$

$$S_{(c_7^l)}^{LL} = \frac{1}{2} \hat{X}_{d,\alpha,a}^L \bar{X}_{\delta,d,c}^L \hat{X}_{b,\gamma,a}^L \bar{X}_{\beta,b,c}^L m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_d^-} D_0(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^R}^2, m_{\nu_c^i}^2) \quad (\text{F.121})$$

$$S_{(c_7^l)}^{LR} = -2 \hat{X}_{d,\alpha,a}^L \bar{X}_{\delta,d,c}^R \hat{X}_{b,\gamma,a}^R \bar{X}_{\beta,b,c}^L D_{27}(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^R}^2, m_{\nu_c^i}^2) \quad (\text{F.122})$$

$$V_{(c_7^l)}^{LL} = \hat{X}_{d,\alpha,a}^L \bar{X}_{\delta,d,c}^R \hat{X}_{b,\gamma,a}^L \bar{X}_{\beta,b,c}^R D_{27}(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^R}^2, m_{\nu_c^i}^2) \quad (\text{F.123})$$

$$V_{(c_7^l)}^{LR} = \frac{1}{2} \hat{X}_{d,\alpha,a}^L \bar{X}_{\delta,d,c}^R \hat{X}_{b,\gamma,a}^R \bar{X}_{\beta,b,c}^R m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_d^-} D_0(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^R}^2, m_{\nu_c^i}^2) \quad (\text{F.124})$$

$$T_{(c_7^l)}^{LL} = -\frac{1}{8} \hat{X}_{d,\alpha,a}^L \bar{X}_{\delta,d,c}^R \hat{X}_{b,\gamma,a}^L \bar{X}_{\beta,b,c}^L m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_d^-} D_0(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^R}^2, m_{\nu_c^i}^2) \quad (\text{F.125})$$

$$S_{(c_8^l)}^{LL} = \frac{1}{2} \hat{X}_{d,\alpha,a}^L \hat{X}_{\delta,d,c}^L \hat{X}_{b,\gamma,a}^L \hat{X}_{\beta,b,c}^L m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_d^-} D_0(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^R}^2, m_{\nu_c^i}^2) \quad (\text{F.126})$$

$$S_{(c_8^l)}^{LR} = -2 \hat{X}_{d,\alpha,a}^L \hat{X}_{\delta,d,c}^R \hat{X}_{b,\gamma,a}^R \hat{X}_{\beta,b,c}^L D_{27}(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^R}^2, m_{\nu_c^R}^2) \quad (\text{F.127})$$

$$V_{(c_8^l)}^{LL} = \hat{X}_{d,\alpha,a}^L \hat{X}_{\delta,d,c}^R \hat{X}_{b,\gamma,a}^L \hat{X}_{\beta,b,c}^R D_{27}(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^R}^2, m_{\nu_c^R}^2) \quad (\text{F.128})$$

$$V_{(c_8^l)}^{LR} = \frac{1}{2} \hat{X}_{d,\alpha,a}^L \hat{X}_{\delta,d,c}^R \hat{X}_{b,\gamma,a}^R \hat{X}_{\beta,b,c}^L m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_d^-} D_0(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^R}^2, m_{\nu_c^R}^2) \quad (\text{F.129})$$

$$T_{(c_8^l)}^{LL} = -\frac{1}{8} \hat{X}_{d,\alpha,a}^L \hat{X}_{\delta,d,c}^R \hat{X}_{b,\gamma,a}^L \hat{X}_{\beta,b,c}^L m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_d^-} D_0(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^R}^2, m_{\nu_c^R}^2) \quad (\text{F.130})$$

F.2.3 Crossed W^+ and H^+ contributions

$$S_{(w_1^l)}^{LL} = \frac{1}{2} V_{d,\alpha,a}^{+,L} \bar{V}_{\beta,b,a}^{+,L} V_{b,\gamma,c}^{+,L} \bar{V}_{\delta,d,c}^{+,L} m_{\nu_b} m_{\nu_d} D_0(m_{\nu_d}^2, m_{\nu_b}^2, m_{H_a^-}^2, m_{H_c^-}^2) \quad (\text{F.131})$$

$$S_{(w_1^l)}^{LR} = 2 V_{d,\alpha,a}^{+,L} \bar{V}_{\beta,b,a}^{+,L} V_{b,\gamma,c}^{+,R} \bar{V}_{\delta,d,c}^{+,R} D_{27}(m_{\nu_d}^2, m_{\nu_b}^2, m_{H_a^-}^2, m_{H_c^-}^2) \quad (\text{F.132})$$

$$V_{(w_1^l)}^{LL} = -V_{d,\alpha,a}^{+,L} \bar{V}_{\beta,b,a}^{+,R} V_{b,\gamma,c}^{+,L} \bar{V}_{\delta,d,c}^{+,R} D_{27}(m_{\nu_d}^2, m_{\nu_b}^2, m_{H_a^-}^2, m_{H_c^-}^2) \quad (\text{F.133})$$

$$V_{(w_1^l)}^{LR} = \frac{1}{2} V_{d,\alpha,a}^{+,L} \bar{V}_{\beta,b,a}^{+,R} V_{b,\gamma,c}^{+,R} \bar{V}_{\delta,d,c}^{+,L} m_{\nu_b} m_{\nu_d} D_0(m_{\nu_d}^2, m_{\nu_b}^2, m_{H_a^-}^2, m_{H_c^-}^2) \quad (\text{F.134})$$

$$T_{(w_1^l)}^{LL} = -\frac{1}{8} V_{d,\alpha,a}^{+,L} \bar{V}_{\beta,b,a}^{+,L} V_{b,\gamma,c}^{+,L} \bar{V}_{\delta,d,c}^{+,L} m_{\nu_b} m_{\nu_d} D_0(m_{\nu_d}^2, m_{\nu_b}^2, m_{H_a^-}^2, m_{H_c^-}^2) \quad (\text{F.135})$$

$$I_1 = I_{C_0 D_0}(m_{\nu_d}^2, m_{\nu_b}^2, m_{W^-}^2, m_{H_a^-}^2) \quad (\text{F.136})$$

$$I_2 = D_{27}(m_{\nu_d}^2, m_{\nu_b}^2, m_{H_a^-}^2, m_{W^-}^2) \quad (\text{F.137})$$

$$I_3 = m_{\nu_b} m_{\nu_d} D_0(m_{\nu_d}^2, m_{\nu_b}^2, m_{H_a^-}^2, m_{W^-}^2) \quad (\text{F.138})$$

$$S_{(w_2^l)}^{LL} = \frac{1}{4} (-3 V_{d,\alpha,a}^{+,R} \bar{V}_{\beta,b,a}^{+,R} \hat{V}_{b,\gamma}^{+,L} \bar{V}_{\delta,d}^{+,L} I_2 - V_{d,\alpha,a}^{+,L} \bar{V}_{\beta,b,a}^{+,L} \hat{V}_{b,\gamma}^{+,L} \bar{V}_{\delta,d}^{+,R} (4I_1 + 13I_2)) \quad (\text{F.139})$$

$$S_{(w_2^l)}^{LR} = \frac{3}{4} (- (V_{d,\alpha,a}^{+,R} \bar{V}_{\beta,b,a}^{+,R} \hat{V}_{b,\gamma}^{+,L} \bar{V}_{\delta,d}^{+,L}) + V_{d,\alpha,a}^{+,L} \bar{V}_{\beta,b,a}^{+,L} \hat{V}_{b,\gamma}^{+,L} \bar{V}_{\delta,d}^{+,R}) I_2 - 2 V_{d,\alpha,a}^{+,L} \bar{V}_{\beta,b,a}^{+,L} \hat{V}_{b,\gamma}^{+,R} \bar{V}_{\delta,d}^{+,L} I_3 \quad (\text{F.140})$$

$$V_{(w_2^l)}^{LL} = V_{d,\alpha,a}^{+,L} \bar{V}_{\beta,b,a}^{+,R} \hat{V}_{b,\gamma}^{+,L} \bar{V}_{\delta,d}^{+,L} I_3 \quad (\text{F.141})$$

$$V_{(w_2^l)}^{LR} = -V_{d,\alpha,a}^{+,L} \bar{V}_{\beta,b,a}^{+,R} \hat{V}_{b,\gamma}^{+,R} \bar{V}_{\delta,d}^{+,R} (I_1 - 2I_2) \quad (\text{F.142})$$

$$T_{(w_2^l)}^{LL} = \frac{1}{16} (V_{d,\alpha,a}^{+,L} \bar{V}_{\beta,b,a}^{+,L} \hat{V}_{b,\gamma}^{+,L} \bar{V}_{\delta,d}^{+,R} (4I_1 - 21I_2) + 5 V_{d,\alpha,a}^{+,R} \bar{V}_{\beta,b,a}^{+,R} \hat{V}_{b,\gamma}^{+,R} \bar{V}_{\delta,d}^{+,L} I_2) \quad (\text{F.143})$$

$$I_1 = I_{C_0 D_0}(m_{\nu_d}^2, m_{\nu_b}^2, m_{H_c^-}^2, m_{W^-}^2) \quad (\text{F.144})$$

$$I_2 = D_{27}(m_{\nu_d}^2, m_{\nu_b}^2, m_{W^-}^2, m_{H_c^-}^2) \quad (\text{F.145})$$

$$I_3 = m_{\nu_b} m_{\nu_d} D_0(m_{\nu_d}^2, m_{\nu_b}^2, m_{W^-}^2, m_{H_c^-}^2) \quad (\text{F.146})$$

$$S_{(w_3')}^{LL} = \frac{1}{4}(-3\hat{V}_{d,i}^{+,L}\hat{\bar{V}}_{\beta,b}^{+,L}V_{b,\gamma,c}^{+,R}\bar{V}_{\delta,d,c}^{+,R}I_2 - \hat{V}_{d,i}^{+,L}\hat{\bar{V}}_{\beta,b}^{+,R}V_{b,\gamma,c}^{+,L}\bar{V}_{\delta,d,c}^{+,L}(4I_1 + 13I_2)) \quad (\text{F.147})$$

$$S_{(w_3')}^{LR} = \frac{3}{4}(\hat{V}_{d,i}^{+,L}\hat{\bar{V}}_{\beta,b}^{+,R}V_{b,\gamma,c}^{+,L}\bar{V}_{\delta,d,c}^{+,L} - \hat{V}_{d,i}^{+,R}\hat{\bar{V}}_{\beta,b}^{+,L}V_{b,\gamma,c}^{+,R}\bar{V}_{\delta,d,c}^{+,R})I_2 - 2\hat{V}_{d,i}^{+,L}\hat{\bar{V}}_{\beta,b}^{+,R}V_{b,\gamma,c}^{+,R}\bar{V}_{\delta,d,c}^{+,R}I_3 \quad (\text{F.148})$$

$$V_{(w_3')}^{LL} = \hat{V}_{d,i}^{+,L}\hat{\bar{V}}_{\beta,b}^{+,L}V_{b,\gamma,c}^{+,L}\bar{V}_{\delta,d,c}^{+,R}I_3 \quad (\text{F.149})$$

$$V_{(w_3')}^{LR} = -\hat{V}_{d,i}^{+,L}\hat{\bar{V}}_{\beta,b}^{+,L}V_{b,\gamma,c}^{+,R}\bar{V}_{\delta,d,c}^{+,L}(I_1 - 2I_2) \quad (\text{F.150})$$

$$T_{(w_3')}^{LL} = \frac{1}{16}(\hat{V}_{d,i}^{+,L}\hat{\bar{V}}_{\beta,b}^{+,R}V_{b,\gamma,c}^{+,L}\bar{V}_{\delta,d,c}^{+,L}(4I_1 - 21I_2) + 5\hat{V}_{d,i}^{+,R}\hat{\bar{V}}_{\beta,b}^{+,L}V_{b,\gamma,c}^{+,R}\bar{V}_{\delta,d,c}^{+,R}I_2) \quad (\text{F.151})$$

$$S_{(w_4')}^{LL} = 8\hat{V}_{d,i}^{+,L}\hat{\bar{V}}_{\beta,b}^{+,R}\hat{V}_{b,\gamma}^{+,L}\bar{V}_{\delta,d}^{+,R}m_{\nu_b} m_{\nu_d} D_0(m_{\nu_d}^2, m_{\nu_b}^2, m_{W^-}^2, m_{W^-}^2) \quad (\text{F.152})$$

$$S_{(w_4')}^{LR} = 8\hat{V}_{d,i}^{+,L}\hat{\bar{V}}_{\beta,b}^{+,R}\hat{V}_{b,\gamma}^{+,R}\bar{V}_{\delta,d}^{+,L}I_{C_0 D_0}(m_{\nu_d}^2, m_{\nu_b}^2, m_{W^-}^2, m_{W^-}^2) \quad (\text{F.153})$$

$$V_{(w_4')}^{LL} = -4\hat{V}_{d,i}^{+,L}\hat{\bar{V}}_{\beta,b}^{+,L}V_{b,\gamma}^{+,L}\bar{V}_{\delta,d}^{+,L}(I_{C_0 D_0}(m_{\nu_d}^2, m_{\nu_b}^2, m_{W^-}^2, m_{W^-}^2) - 3D_{27}(m_{\nu_d}^2, m_{\nu_b}^2, m_{W^-}^2, m_{W^-}^2)) \quad (\text{F.154})$$

$$V_{(w_4')}^{LR} = 2\hat{V}_{d,i}^{+,L}\hat{\bar{V}}_{\beta,b}^{+,L}V_{b,\gamma}^{+,R}\bar{V}_{\delta,d}^{+,R}m_{\nu_b} m_{\nu_d} D_0(m_{\nu_d}^2, m_{\nu_b}^2, m_{W^-}^2, m_{W^-}^2) \quad (\text{F.155})$$

$$S_{(w_5')}^{LL} = \frac{1}{2}V_{d,\alpha,a}^{+,L}\bar{V}_{\beta,b,a}^{+,L}\bar{V}_{\delta,b,c}^{+,L}V_{d,\gamma,c}^{+,L}m_{\nu_b} m_{\nu_d} D_0(m_{\nu_b}^2, m_{\nu_d}^2, m_{H_a^-}^2, m_{H_c^-}^2) \quad (\text{F.156})$$

$$S_{(w_5')}^{LR} = 2V_{d,\alpha,a}^{+,L}\bar{V}_{\beta,b,a}^{+,L}\bar{V}_{\delta,b,c}^{+,R}V_{d,\gamma,c}^{+,R}D_{27}(m_{\nu_b}^2, m_{\nu_d}^2, m_{H_a^-}^2, m_{H_c^-}^2) \quad (\text{F.157})$$

$$V_{(w_5')}^{LL} = -\frac{1}{2}V_{d,\alpha,a}^{+,L}\bar{V}_{\beta,b,a}^{+,R}\bar{V}_{\delta,b,c}^{+,R}V_{d,\gamma,c}^{+,L}m_{\nu_b} m_{\nu_d} D_0(m_{\nu_b}^2, m_{\nu_d}^2, m_{H_a^-}^2, m_{H_c^-}^2) \quad (\text{F.158})$$

$$V_{(w_5')}^{LR} = V_{d,\alpha,a}^{+,L}\bar{V}_{\beta,b,a}^{+,R}\bar{V}_{\delta,b,c}^{+,L}V_{d,\gamma,c}^{+,R}D_{27}(m_{\nu_b}^2, m_{\nu_d}^2, m_{H_a^-}^2, m_{H_c^-}^2) \quad (\text{F.159})$$

$$T_{(w_5')}^{LL} = \frac{1}{8}V_{d,\alpha,a}^{+,L}\bar{V}_{\beta,b,a}^{+,L}\bar{V}_{\delta,b,c}^{+,L}V_{d,\gamma,c}^{+,L}m_{\nu_b} m_{\nu_d} D_0(m_{\nu_b}^2, m_{\nu_d}^2, m_{H_a^-}^2, m_{H_c^-}^2) \quad (\text{F.160})$$

$$S_{(w_6')}^{LL} = -8V_{d,\alpha,a}^{+,L}\bar{V}_{\beta,b,a}^{+,L}\hat{\bar{V}}_{\delta,b}^{+,R}\hat{V}_{d,\gamma}^{+,L}D_{27}(m_{\nu_b}^2, m_{\nu_d}^2, m_{H_a^-}^2, m_{W^-}^2) \quad (\text{F.161})$$

$$S_{(w_6')}^{LR} = -2V_{d,\alpha,a}^{+,L}\bar{V}_{\beta,b,a}^{+,L}\hat{\bar{V}}_{\delta,b}^{+,L}\hat{V}_{d,\gamma}^{+,R}m_{\nu_b} m_{\nu_d} D_0(m_{\nu_b}^2, m_{\nu_d}^2, m_{H_a^-}^2, m_{W^-}^2) \quad (\text{F.162})$$

$$V_{(w_6')}^{LL} = 2V_{d,\alpha,a}^{+,L}\bar{V}_{\beta,b,a}^{+,L}\hat{\bar{V}}_{\delta,b}^{+,R}\hat{V}_{d,\gamma}^{+,L}D_{27}(m_{\nu_b}^2, m_{\nu_d}^2, m_{H_a^-}^2, m_{W^-}^2) \quad (\text{F.163})$$

$$V_{(w_6')}^{LR} = -V_{d,\alpha,a}^{+,L}\bar{V}_{\beta,b,a}^{+,R}\hat{\bar{V}}_{\delta,b}^{+,R}\hat{V}_{d,\gamma}^{+,R}m_{\nu_b} m_{\nu_d} D_0(m_{\nu_b}^2, m_{\nu_d}^2, m_{H_a^-}^2, m_{W^-}^2) \quad (\text{F.164})$$

$$S_{(w_7')}^{LL} = -8\hat{V}_{d,i}^{+,L}\hat{\bar{V}}_{\beta,b}^{+,R}\bar{V}_{\delta,b,c}^{+,L}V_{d,\gamma,c}^{+,L}D_{27}(m_{\nu_b}^2, m_{\nu_d}^2, m_{W^-}^2, m_{H_c^-}^2) \quad (\text{F.165})$$

$$S_{(w_7')}^{LR} = -2\hat{V}_{d,i}^{+,L}\hat{\bar{V}}_{\beta,b}^{+,R}\bar{V}_{\delta,b,c}^{+,R}V_{d,\gamma,c}^{+,R}m_{\nu_b} m_{\nu_d} D_0(m_{\nu_b}^2, m_{\nu_d}^2, m_{W^-}^2, m_{H_c^-}^2) \quad (\text{F.166})$$

$$V_{(w_7')}^{LL} = 2\hat{V}_{d,i}^{+,L}\hat{\bar{V}}_{\beta,b}^{+,L}\bar{V}_{\delta,b,c}^{+,R}V_{d,\gamma,c}^{+,L}D_{27}(m_{\nu_b}^2, m_{\nu_d}^2, m_{W^-}^2, m_{H_c^-}^2) \quad (\text{F.167})$$

$$V_{(w_7')}^{LR} = -\hat{V}_{d,i}^{+,L}\hat{\bar{V}}_{\beta,b}^{+,L}\bar{V}_{\delta,b,c}^{+,L}V_{d,\gamma,c}^{+,R}m_{\nu_b} m_{\nu_d} D_0(m_{\nu_b}^2, m_{\nu_d}^2, m_{W^-}^2, m_{H_c^-}^2) \quad (\text{F.168})$$

$$S_{(w_8')}^{LL} = 8\hat{V}_{d,i}^{+,L}\hat{\bar{V}}_{\beta,b}^{+,R}\hat{\bar{V}}_{\delta,b}^{+,R}\hat{V}_{d,\gamma}^{+,L}m_{\nu_b} m_{\nu_d} D_0(m_{\nu_b}^2, m_{\nu_d}^2, m_{W^-}^2, m_{W^-}^2) \quad (\text{F.169})$$

$$S_{(w_8')}^{LR} = 32\hat{V}_{d,i}^{+,L}\hat{\bar{V}}_{\beta,b}^{+,R}\hat{\bar{V}}_{\delta,b}^{+,L}\hat{V}_{d,\gamma}^{+,R}D_{27}(m_{\nu_b}^2, m_{\nu_d}^2, m_{W^-}^2, m_{W^-}^2) \quad (\text{F.170})$$


$$V_{(w_8^l)}^{LL} = -2\hat{V}_{d,i}^{+,L}\hat{V}_{\beta,b}^{+,L}\hat{V}_{\delta,b}^{+,L}\hat{V}_{d,\gamma}^{+,L}m_{\nu_b}m_{\nu_d}D_0(m_{\nu_b}^2, m_{\nu_d}^2, m_{W^-}^2, m_{W^-}^2) \quad (\text{F.171})$$

$$V_{(w_8^l)}^{LR} = 4\hat{V}_{d,i}^{+,L}\hat{V}_{\beta,b}^{+,L}\hat{V}_{\delta,b}^{+,R}\hat{V}_{d,\gamma}^{+,R}D_{27}(m_{\nu_b}^2, m_{\nu_d}^2, m_{W^-}^2, m_{W^-}^2) \quad (\text{F.172})$$


F.3 Two-lepton — two-quark boxes


F.3.1 Feynman diagrams

Neutralino diagrams



Chargino diagrams



W^+ and H^+ diagrams

F.3.2 Down quarks
Neutralino.

$$S_{(n_1^d)}^{LL} = \frac{1}{2} N_{d,\alpha,a}^L \bar{N}_{\beta,b,a}^L N_{b,\gamma,c}^{d,L} \bar{N}_{\delta,d,c}^{d,L} m_{\tilde{\chi}_b^0} m_{\tilde{\chi}_d^0} D_0(m_{\tilde{\chi}_d^0}^2, m_{\tilde{\chi}_b^0}^2, m_{\tilde{e}_a}^2, m_{\tilde{d}_c}^2) \quad (\text{F.173})$$

$$S_{(n_1^d)}^{LR} = 2 N_{d,\alpha,a}^L \bar{N}_{\beta,b,a}^L N_{b,\gamma,c}^{d,R} \bar{N}_{\delta,d,c}^{d,R} D_{27}(m_{\tilde{\chi}_d^0}^2, m_{\tilde{\chi}_b^0}^2, m_{\tilde{e}_a}^2, m_{\tilde{d}_c}^2) \quad (\text{F.174})$$

$$V_{(n_1^d)}^{LL} = -N_{d,\alpha,a}^L \bar{N}_{\beta,b,a}^R N_{b,\gamma,c}^{d,L} \bar{N}_{\delta,d,c}^{d,R} D_{27}(m_{\tilde{\chi}_d^0}^2, m_{\tilde{\chi}_b^0}^2, m_{\tilde{e}_a}^2, m_{\tilde{d}_c}^2) \quad (\text{F.175})$$

$$V_{(n_1^d)}^{LR} = \frac{1}{2} N_{d,\alpha,a}^L \bar{N}_{\beta,b,a}^R N_{b,\gamma,c}^{d,R} \bar{N}_{\delta,d,c}^{d,L} m_{\tilde{\chi}_b^0} m_{\tilde{\chi}_d^0} D_0(m_{\tilde{\chi}_d^0}^2, m_{\tilde{\chi}_b^0}^2, m_{\tilde{e}_a}^2, m_{\tilde{d}_c}^2) \quad (\text{F.176})$$

$$T_{(n_1^d)}^{LL} = -\frac{1}{8} N_{d,\alpha,a}^L \bar{N}_{\beta,b,a}^L N_{b,\gamma,c}^{d,L} \bar{N}_{\delta,d,c}^{d,L} m_{\tilde{\chi}_b^0} m_{\tilde{\chi}_d^0} D_0(m_{\tilde{\chi}_d^0}^2, m_{\tilde{\chi}_b^0}^2, m_{\tilde{e}_a}^2, m_{\tilde{d}_c}^2) \quad (\text{F.177})$$

$$S_{(n_2^d)}^{LL} = \frac{1}{2} N_{d,\alpha,a}^L \bar{N}_{\beta,b,a}^L \bar{N}_{\delta,b,c}^{d,L} N_{d,\gamma,c}^{d,L} m_{\tilde{\chi}_b^0} m_{\tilde{\chi}_d^0} D_0(m_{\tilde{\chi}_b^0}^2, m_{\tilde{\chi}_d^0}^2, m_{\tilde{e}_a}^2, m_{\tilde{d}_c}^2) \quad (\text{F.178})$$

$$S_{(n_2^d)}^{LR} = 2 N_{d,\alpha,a}^L \bar{N}_{\beta,b,a}^L \bar{N}_{\delta,b,c}^{d,R} N_{d,\gamma,c}^{d,R} D_{27}(m_{\tilde{\chi}_b^0}^2, m_{\tilde{\chi}_d^0}^2, m_{\tilde{e}_a}^2, m_{\tilde{d}_c}^2) \quad (\text{F.179})$$

$$V_{(n_2^d)}^{LL} = -\frac{1}{2} N_{d,\alpha,a}^L \bar{N}_{\beta,b,a}^R \bar{N}_{\delta,b,c}^{d,R} N_{d,\gamma,c}^{d,L} m_{\tilde{\chi}_b^0} m_{\tilde{\chi}_d^0} D_0(m_{\tilde{\chi}_d^0}^2, m_{\tilde{\chi}_b^0}^2, m_{\tilde{e}_a}^2, m_{\tilde{d}_c}^2) \quad (\text{F.180})$$

$$V_{(n_2^d)}^{LR} = N_{d,\alpha,a}^L \bar{N}_{\beta,b,a}^R \bar{N}_{\delta,b,c}^{d,L} N_{d,\gamma,c}^{d,R} D_{27}(m_{\tilde{\chi}_b^0}^2, m_{\tilde{\chi}_d^0}^2, m_{\tilde{e}_a}^2, m_{\tilde{d}_c}^2) \quad (\text{F.181})$$

$$T_{(n_2^d)}^{LL} = \frac{1}{8} N_{d,\alpha,a}^L \bar{N}_{\beta,b,a}^L \bar{N}_{\delta,b,c}^{d,L} N_{d,\gamma,c}^{d,L} m_{\tilde{\chi}_b^0} m_{\tilde{\chi}_d^0} D_0(m_{\tilde{\chi}_d^0}^2, m_{\tilde{\chi}_b^0}^2, m_{\tilde{e}_a}^2, m_{\tilde{d}_c}^2) \quad (\text{F.182})$$

Chargino.

$$S_{(c_1^d)}^{LL} = \frac{1}{2} X_{d,\alpha,a}^L \bar{X}_{\beta,b,a}^L W_{b,\gamma,c}^{d,L} \bar{W}_{\delta,d,c}^{d,L} m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_d^-} D_0(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^i}^2, m_{\tilde{u}_c}^2) \quad (\text{F.183})$$

$$S_{(c_1^d)}^{LR} = 2 X_{d,\alpha,a}^L \bar{X}_{\beta,b,a}^L W_{b,\gamma,c}^{d,R} \bar{W}_{\delta,d,c}^{d,R} D_{27}(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^i}^2, m_{\tilde{u}_c}^2) \quad (\text{F.184})$$

$$V_{(c_1^d)}^{LL} = -X_{d,\alpha,a}^L \bar{X}_{\beta,b,a}^R W_{b,\gamma,c}^{d,L} \bar{W}_{\delta,d,c}^{d,R} D_{27}(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^i}^2, m_{\tilde{u}_c}^2) \quad (\text{F.185})$$

$$V_{(c_1^d)}^{LR} = \frac{1}{2} X_{d,\alpha,a}^L \bar{X}_{\beta,b,a}^R W_{b,\gamma,c}^{d,R} \bar{W}_{\delta,d,c}^{d,L} m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_d^-} D_0(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^i}^2, m_{\tilde{u}_c}^2) \quad (\text{F.186})$$

$$T_{(c_1^d)}^{LL} = -\frac{1}{8} X_{d,\alpha,a}^L \bar{X}_{\beta,b,a}^L W_{b,\gamma,c}^{d,L} \bar{W}_{\delta,d,c}^{d,L} m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_d^-} D_0(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^i}^2, m_{\tilde{u}_c}^2) \quad (\text{F.187})$$

$$S_{(c_2^d)}^{LL} = \frac{1}{2} \hat{X}_{d,\alpha,a}^L \hat{\bar{X}}_{\beta,b,a}^L W_{b,\gamma,c}^{d,L} \bar{W}_{\delta,d,c}^{d,L} m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_d^-} D_0(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^R}^2, m_{\tilde{u}_c}^2) \quad (\text{F.188})$$

$$S_{(c_2^d)}^{LR} = 2 \hat{X}_{d,\alpha,a}^L \hat{\bar{X}}_{\beta,b,a}^L W_{b,\gamma,c}^{d,R} \bar{W}_{\delta,d,c}^{d,R} D_{27}(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^R}^2, m_{\tilde{u}_c}^2) \quad (\text{F.189})$$

$$V_{(c_2^d)}^{LL} = -\hat{X}_{d,\alpha,a}^L \hat{\bar{X}}_{\beta,b,a}^R W_{b,\gamma,c}^{d,L} \bar{W}_{\delta,d,c}^{d,R} D_{27}(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^R}^2, m_{\tilde{u}_c}^2) \quad (\text{F.190})$$

$$V_{(c_2^d)}^{LR} = \frac{1}{2} \hat{X}_{d,\alpha,a}^L \hat{\bar{X}}_{\beta,b,a}^R W_{b,\gamma,c}^{d,R} \bar{W}_{\delta,d,c}^{d,L} m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_d^-} D_0(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^R}^2, m_{\tilde{u}_c}^2) \quad (\text{F.191})$$

$$T_{(c_2^d)}^{LL} = -\frac{1}{8} \hat{X}_{d,\alpha,a}^L \hat{\bar{X}}_{\beta,b,a}^L W_{b,\gamma,c}^{d,L} \bar{W}_{\delta,d,c}^{d,L} m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_d^-} D_0(m_{\tilde{\chi}_d^-}^2, m_{\tilde{\chi}_b^-}^2, m_{\nu_a^R}^2, m_{\tilde{u}_c}^2) \quad (\text{F.192})$$

W^+ and H^+ .

$$S_{(w_1^d)}^{LL} = -V_{a,\alpha,d}^{+,L} \bar{V}_{\beta,a,b}^{+,L} V_{c,\gamma,b}^{u,L} \bar{V}_{\delta,c,d}^{d,L} m_{\nu_a} m_{u_c} D_0(m_{\nu_a}^2, m_{u_c}^2, m_{H_d^-}^2, m_{H_b^-}^2) \quad (\text{F.193})$$

$$S_{(w_1^d)}^{LR} = -V_{a,\alpha,d}^{+,L} \bar{V}_{\beta,a,b}^{+,L} V_{c,\gamma,b}^{u,R} \bar{V}_{\delta,c,d}^{d,R} m_{\nu_a} m_{u_c} D_0(m_{\nu_a}^2, m_{u_c}^2, m_{H_d^-}^2, m_{H_b^-}^2) \quad (\text{F.194})$$

$$V_{(w_1^d)}^{LL} = -V_{a,\alpha,d}^{+,L} \bar{V}_{\beta,a,b}^{+,R} V_{c,\gamma,b}^{u,L} \bar{V}_{\delta,c,d}^{d,R} D_{27}(m_{\nu_a}^2, m_{u_c}^2, m_{H_d^-}^2, m_{H_b^-}^2) \quad (\text{F.195})$$

$$V_{(w_1^d)}^{LR} = -V_{a,\alpha,d}^{+,L} \bar{V}_{\beta,a,b}^{+,R} V_{c,\gamma,b}^{u,R} \bar{V}_{\delta,c,d}^{d,L} D_{27}(m_{\nu_a}^2, m_{u_c}^2, m_{H_d^-}^2, m_{H_b^-}^2) \quad (\text{F.196})$$

$$S_{(w_2^d)}^{LL} = 2 V_{a,\alpha,d}^{+,L} \hat{\bar{V}}_{\beta,a}^{+,R} \hat{V}_{c,\gamma}^{u,L} \bar{V}_{\delta,c,d}^{d,L} (I_{C_0 D_0}(m_{u_c}^2, m_{W^-}^2, m_{H_d^-}^2, m_{\nu_a}^2) - 2 D_{27}(m_{\nu_a}^2, m_{u_c}^2, m_{W^-}^2, m_{H_d^-}^2)) \quad (\text{F.197})$$

$$S_{(w_2^d)}^{LR} = 2 V_{a,\alpha,d}^{+,L} \hat{\bar{V}}_{\beta,a}^{+,R} \hat{V}_{c,\gamma}^{u,R} \bar{V}_{\delta,c,d}^{d,R} (I_{C_0 D_0}(m_{u_c}^2, m_{W^-}^2, m_{H_d^-}^2, m_{\nu_a}^2) - 2 D_{27}(m_{\nu_a}^2, m_{u_c}^2, m_{W^-}^2, m_{H_d^-}^2)) \quad (\text{F.198})$$

$$V_{(w_2^d)}^{LL} = V_{a,\alpha,d}^{+,L} \hat{V}_{\beta,a}^{+,L} \hat{V}_{c,\gamma}^{u,L} \bar{V}_{\delta,c,d}^{d,R} m_{\nu_a} m_{u_c} D_0(m_{\nu_a}^2, m_{u_c}^2, m_{W^-}^2, m_{H_d^-}^2) \quad (\text{F.199})$$

$$V_{(w_2^d)}^{LR} = V_{a,\alpha,d}^{+,L} \hat{V}_{\beta,a}^{+,L} \hat{V}_{c,\gamma}^{u,R} \bar{V}_{\delta,c,d}^{d,L} m_{\nu_a} m_{u_c} D_0(m_{\nu_a}^2, m_{u_c}^2, m_{W^-}^2, m_{H_d^-}^2) \quad (\text{F.200})$$

$$T_{(w_2^d)}^{LL} = -V_{a,\alpha,d}^{+,L} \hat{V}_{\beta,a}^{+,R} \hat{V}_{c,\gamma}^{u,L} \bar{V}_{\delta,c,d}^{d,L} D_{27}(m_{\nu_a}^2, m_{u_c}^2, m_{W^-}^2, m_{H_d^-}^2) \quad (\text{F.201})$$

$$S_{(w_3^d)}^{LL} = 2 \hat{V}_{a,i}^{+,L} \bar{V}_{\beta,a,b}^{+,L} V_{c,\gamma,b}^{u,L} \hat{V}_{\delta,c}^{d,R} (I_{C_0 D_0}(m_{u_c}^2, m_{H_b^-}^2, m_{W^-}^2, m_{\nu_a}^2) - 2D_{27}(m_{\nu_a}^2, m_{u_c}^2, m_{S22}, m_{W^-}^2)) \quad (\text{F.202})$$

$$S_{(w_3^d)}^{LR} = 2 \hat{V}_{a,i}^{+,L} \bar{V}_{\beta,a,b}^{+,L} V_{c,\gamma,b}^{u,R} \hat{V}_{\delta,c}^{d,L} (I_{C_0 D_0}(m_{u_c}^2, m_{H_b^-}^2, m_{W^-}^2, m_{\nu_a}^2) - 2D_{27}(m_{\nu_a}^2, m_{u_c}^2, m_{S22}, m_{W^-}^2)) \quad (\text{F.203})$$

$$V_{(w_3^d)}^{LL} = \hat{V}_{a,i}^{+,L} \bar{V}_{\beta,a,b}^{+,R} V_{c,\gamma,b}^{u,L} \hat{V}_{\delta,c}^{d,L} m_{\nu_a} m_{u_c} D_0(m_{\nu_a}^2, m_{u_c}^2, m_{H_b^-}^2, m_{W^-}^2) \quad (\text{F.204})$$

$$V_{(w_3^d)}^{LR} = \hat{V}_{a,i}^{+,L} \bar{V}_{\beta,a,b}^{+,R} V_{c,\gamma,b}^{u,R} \hat{V}_{\delta,c}^{d,R} m_{\nu_a} m_{u_c} D_0(m_{\nu_a}^2, m_{u_c}^2, m_{H_b^-}^2, m_{W^-}^2) \quad (\text{F.205})$$

$$T_{(w_3^d)}^{LL} d = -\hat{V}_{a,i}^{+,L} \bar{V}_{\beta,a,b}^{+,L} V_{c,\gamma,b}^{u,L} \hat{V}_{\delta,c}^{d,R} D_{27}(m_{\nu_a}^2, m_{u_c}^2, m_{H_b^-}^2, m_{W^-}^2) \quad (\text{F.206})$$

$$S_{(w_4^d)}^{LL} = -4 \hat{V}_{a,i}^{+,L} \bar{V}_{\beta,a}^{+,R} \hat{V}_{c,\gamma}^{u,L} \hat{V}_{\delta,c}^{d,R} m_{\nu_a} m_{u_c} D_0(m_{\nu_a}^2, m_{u_c}^2, m_{W^-}^2, m_{W^-}^2) \quad (\text{F.207})$$

$$S_{(w_4^d)}^{LR} = -4 \hat{V}_{a,i}^{+,L} \bar{V}_{\beta,a}^{+,R} \hat{V}_{c,\gamma}^{u,R} \hat{V}_{\delta,c}^{d,L} m_{\nu_a} m_{u_c} D_0(m_{\nu_a}^2, m_{u_c}^2, m_{W^-}^2, m_{W^-}^2) \quad (\text{F.208})$$

$$V_{(w_4^d)}^{LL} = -4 \hat{V}_{a,i}^{+,L} \bar{V}_{\beta,a}^{+,L} \hat{V}_{c,\gamma}^{u,L} \hat{V}_{\delta,c}^{d,L} (I_{C_0 D_0}(m_{u_c}^2, m_{W^-}^2, m_{W^-}^2, m_{\nu_a}^2) - 3D_{27}(m_{\nu_a}^2, m_{u_c}^2, m_{W^-}^2, m_{W^-}^2)) \quad (\text{F.209})$$

$$V_{(w_4^d)}^{LR} = -4 \hat{V}_{a,i}^{+,L} \bar{V}_{\beta,a}^{+,L} \hat{V}_{c,\gamma}^{u,R} \hat{V}_{\delta,c}^{d,R} I_{C_0 D_0}(m_{u_c}^2, m_{W^-}^2, m_{W^-}^2, m_{\nu_a}^2) \quad (\text{F.210})$$

$$T_{(w_4^d)}^{LL} d = \hat{V}_{a,i}^{+,L} \bar{V}_{\beta,a}^{+,R} \hat{V}_{c,\gamma}^{u,L} \hat{V}_{\delta,c}^{d,R} m_{\nu_a} m_{u_c} D_0(m_{\nu_a}^2, m_{u_c}^2, m_{W^-}^2, m_{W^-}^2) \quad (\text{F.211})$$

F.3.3 Up quarks

Neutralino.

$$S_{(n_1^u)}^{LL} = \frac{1}{2} N_{d,\alpha,a}^L \bar{N}_{\beta,b,a}^L N_{b,\gamma,c}^{u,L} \bar{N}_{\delta,d,c}^{u,L} D_{27}(m_{\tilde{\chi}_d^0}^2, m_{\tilde{\chi}_b^0}^2, m_{\tilde{e}_a}^2, m_{\tilde{u}_c}^2) \quad (\text{F.212})$$

$$S_{(n_1^u)}^{LR} = 2 N_{d,\alpha,a}^L \bar{N}_{\beta,b,a}^L N_{b,\gamma,c}^{u,R} \bar{N}_{\delta,d,c}^{u,R} D_{27}(m_{\tilde{\chi}_d^0}^2, m_{\tilde{\chi}_b^0}^2, m_{\tilde{e}_a}^2, m_{\tilde{u}_c}^2) \quad (\text{F.213})$$

$$V_{(n_1^u)}^{LL} = -N_{d,\alpha,a}^L \bar{N}_{\beta,b,a}^R N_{b,\gamma,c}^{u,L} \bar{N}_{\delta,d,c}^{u,R} D_{27}(m_{\tilde{\chi}_d^0}^2, m_{\tilde{\chi}_b^0}^2, m_{\tilde{e}_a}^2, m_{\tilde{u}_c}^2) \quad (\text{F.214})$$

$$V_{(n_1^u)}^{LR} = \frac{1}{2} N_{d,\alpha,a}^L \bar{N}_{\beta,b,a}^R N_{b,\gamma,c}^{u,R} \bar{N}_{\delta,d,c}^{u,L} D_{27}(m_{\tilde{\chi}_d^0}^2, m_{\tilde{\chi}_b^0}^2, m_{\tilde{e}_a}^2, m_{\tilde{u}_c}^2) \quad (\text{F.215})$$

$$S_{(n_2^u)}^{LL} = \frac{1}{2} N_{d,\alpha,a}^L \bar{N}_{\beta,b,a}^L \bar{N}_{\delta,b,c}^{u,L} N_{d,\gamma,c}^{u,L} m_{\tilde{\chi}_b^0} m_{\tilde{\chi}_d^0} D_0(m_{\tilde{\chi}_b^0}^2, m_{\tilde{\chi}_d^0}^2, m_{\tilde{e}_a}^2, m_{\tilde{u}_c}^2) \quad (\text{F.216})$$

$$S_{(n_2^u)}^{LR} = 2 N_{d,\alpha,a}^L \bar{N}_{\beta,b,a}^L \bar{N}_{\delta,b,c}^{u,R} N_{d,\gamma,c}^{u,R} D_{27}(m_{\tilde{\chi}_b^0}^2, m_{\tilde{\chi}_d^0}^2, m_{\tilde{e}_a}^2, m_{\tilde{u}_c}^2) \quad (\text{F.217})$$

$$V_{(n_2^u)}^{LL} = -\frac{1}{2} N_{d,\alpha,a}^L \bar{N}_{\beta,b,a}^R \bar{N}_{\delta,b,c}^{u,R} N_{d,\gamma,c}^{u,L} m_{\tilde{\chi}_b^0} m_{\tilde{\chi}_d^0} D_0(m_{\tilde{\chi}_b^0}^2, m_{\tilde{\chi}_d^0}^2, m_{\tilde{e}_a}^2, m_{\tilde{u}_c}^2) \quad (\text{F.218})$$

$$V_{(n_2^u)}^{LR} = N_{d,\alpha,a}^L \bar{N}_{\beta,b,a}^R \bar{N}_{\delta,b,c}^{u,L} N_{d,\gamma,c}^{u,R} D_{27}(m_{\tilde{\chi}_b^0}^2, m_{\tilde{\chi}_d^0}^2, m_{\tilde{e}_a}^2, m_{\tilde{u}_c}^2) \quad (\text{F.219})$$

$$T_{(n_2^u)}^{LL} = \frac{1}{8} N_{d,\alpha,a}^L \bar{N}_{\beta,b,a}^L \bar{N}_{\delta,b,c}^{u,L} N_{d,\gamma,c}^{u,L} m_{\tilde{\chi}_b^0} m_{\tilde{\chi}_d^0} D_0(m_{\tilde{\chi}_b^0}^2, m_{\tilde{\chi}_d^0}^2, m_{\tilde{e}_a}^2, m_{\tilde{u}_c}^2) \quad (\text{F.220})$$

Chargino.

$$S_{(c_1^u)}^{LL} = \frac{1}{2} X_{d,\alpha,a}^L \bar{X}_{\beta,b,a}^L W_{b,\delta,c}^{u,L} \bar{W}_{d,\gamma,c}^{u,L} m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_d^-} D_0(m_{\tilde{\chi}_b^-}^2, m_{\tilde{\chi}_d^-}^2, m_{\nu_a^i}^2, m_{d_c^2}^2) \quad (\text{F.221})$$

$$S_{(c_1^u)}^{LR} = 2 X_{d,\alpha,a}^L \bar{X}_{\beta,b,a}^L W_{b,\delta,c}^{u,R} \bar{W}_{d,\gamma,c}^{u,R} D_{27}(m_{\tilde{\chi}_b^-}^2, m_{\tilde{\chi}_d^-}^2, m_{\nu_a^i}^2, m_{\tilde{d}_c}^2) \quad (\text{F.222})$$

$$V_{(c_1^u)}^{LL} = -\frac{1}{2} X_{d,\alpha,a}^L \bar{X}_{\beta,b,a}^R W_{b,\delta,c}^{u,R} \bar{W}_{d,\gamma,c}^{u,L} m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_d^-} D_0(m_{\tilde{\chi}_b^-}^2, m_{\tilde{\chi}_d^-}^2, m_{\nu_a^i}^2, m_{\tilde{d}_c}^2) \quad (\text{F.223})$$

$$V_{(c_1^u)}^{LR} = X_{d,\alpha,a}^L \bar{X}_{\beta,b,a}^R W_{b,\delta,c}^{u,L} \bar{W}_{d,\gamma,c}^{u,R} D_{27}(m_{\tilde{\chi}_b^-}^2, m_{\tilde{\chi}_d^-}^2, m_{\nu_a^i}^2, m_{\tilde{d}_c}^2) \quad (\text{F.224})$$

$$T_{(c_1^u)}^{LL} = \frac{1}{8} X_{d,\alpha,a}^L \bar{X}_{\beta,b,a}^L W_{b,\delta,c}^{u,L} \bar{W}_{d,\gamma,c}^{u,L} m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_d^-} D_0(m_{\tilde{\chi}_b^-}^2, m_{\tilde{\chi}_d^-}^2, m_{\nu_a^i}^2, m_{\tilde{d}_c}^2) \quad (\text{F.225})$$

$$S_{(c_2^u)}^{LL} = \frac{1}{2} \hat{X}_{d,\alpha,a}^L \hat{\bar{X}}_{\beta,b,a}^L W_{b,\delta,c}^{u,L} \bar{W}_{d,\gamma,c}^{u,L} m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_d^-} D_0(m_{\tilde{\chi}_b^-}^2, m_{\tilde{\chi}_d^-}^2, m_{\nu_a^R}^2, m_{\tilde{d}_c}^2) \quad (\text{F.226})$$

$$S_{(c_2^u)}^{LR} = 2 \hat{X}_{d,\alpha,a}^L \hat{\bar{X}}_{\beta,b,a}^L W_{b,\delta,c}^{u,R} \bar{W}_{d,\gamma,c}^{u,R} D_{27}(m_{\tilde{\chi}_b^-}^2, m_{\tilde{\chi}_d^-}^2, m_{\nu_a^R}^2, m_{\tilde{d}_c}^2) \quad (\text{F.227})$$

$$V_{(c_2^u)}^{LL} = -\frac{1}{2} \hat{X}_{d,\alpha,a}^L \hat{\bar{X}}_{\beta,b,a}^R W_{b,\delta,c}^{u,R} \bar{W}_{d,\gamma,c}^{u,L} m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_d^-} D_0(m_{\tilde{\chi}_b^-}^2, m_{\tilde{\chi}_d^-}^2, m_{\nu_a^R}^2, m_{\tilde{d}_c}^2) \quad (\text{F.228})$$

$$V_{(c_2^u)}^{LR} = \hat{X}_{d,\alpha,a}^L \hat{\bar{X}}_{\beta,b,a}^R W_{b,\delta,c}^{u,L} \bar{W}_{d,\gamma,c}^{u,R} D_{27}(m_{\tilde{\chi}_b^-}^2, m_{\tilde{\chi}_d^-}^2, m_{\nu_a^R}^2, m_{\tilde{d}_c}^2) \quad (\text{F.229})$$

$$T_{(c_2^u)}^{LL} = \frac{1}{8} \hat{X}_{d,\alpha,a}^L \hat{\bar{X}}_{\beta,b,a}^L W_{b,\delta,c}^{u,L} \bar{W}_{d,\gamma,c}^{u,L} m_{\tilde{\chi}_b^-} m_{\tilde{\chi}_d^-} D_0(m_{\tilde{\chi}_b^-}^2, m_{\tilde{\chi}_d^-}^2, m_{\nu_a^R}^2, m_{\tilde{d}_c}^2) \quad (\text{F.230})$$

W^+ and H^+ .

$$S_{(w_1^u)}^{LL} = -V_{a,\alpha,d}^{+,L} \bar{V}_{\beta,a,b}^{+,L} V_{\delta,c,b}^{u,L} \bar{V}_{c,\gamma,d}^{d,L} m_{\nu_a} m_{d_c} D_0(m_{\nu_a}^2, m_{d_c}^2, m_{H_d^-}^2, m_{H_b^-}^2) \quad (\text{F.231})$$

$$S_{(w_1^u)}^{LR} = -V_{a,\alpha,d}^{+,L} \bar{V}_{\beta,a,b}^{+,L} V_{\delta,c,b}^{u,R} \bar{V}_{c,\gamma,d}^{d,R} m_{\nu_a} m_{d_c} D_0(m_{\nu_a}^2, m_{d_c}^2, m_{H_d^-}^2, m_{H_b^-}^2) \quad (\text{F.232})$$

$$V_{(w_1^u)}^{LL} = V_{a,\alpha,d}^{+,L} \bar{V}_{\beta,a,b}^{+,R} V_{\delta,c,b}^{u,R} \bar{V}_{c,\gamma,d}^{d,L} D_{27}(m_{\nu_a}^2, m_{d_c}^2, m_{H_d^-}^2, m_{H_b^-}^2) \quad (\text{F.233})$$

$$V_{(w_1^u)}^{LR} = V_{a,\alpha,d}^{+,L} \bar{V}_{\beta,a,b}^{+,R} V_{\delta,c,b}^{u,L} \bar{V}_{c,\gamma,d}^{d,R} D_{27}(m_{\nu_a}^2, m_{d_c}^2, m_{H_d^-}^2, m_{H_b^-}^2) \quad (\text{F.234})$$

$$S_{(w_2^u)}^{LL} = -4 V_{a,\alpha,d}^{+,L} \hat{\bar{V}}_{\beta,a}^{+,R} \hat{V}_{\delta,c}^{u,R} \bar{V}_{c,\gamma,d}^{d,L} D_{27}(m_{\nu_a}^2, m_{d_c}^2, m_{W^-}^2, m_{H_d^-}^2) \quad (\text{F.235})$$

$$S_{(w_2^u)}^{LR} = -4 V_{a,\alpha,d}^{+,L} \hat{\bar{V}}_{\beta,a}^{+,R} \hat{V}_{\delta,c}^{u,L} \bar{V}_{c,\gamma,d}^{d,R} D_{27}(m_{\nu_a}^2, m_{d_c}^2, m_{W^-}^2, m_{H_d^-}^2) \quad (\text{F.236})$$

$$V_{(w_2^u)}^{LL} = V_{a,\alpha,d}^{+,L} \hat{\bar{V}}_{\beta,a}^{+,L} \hat{V}_{\delta,c}^{u,L} \bar{V}_{c,\gamma,d}^{d,L} m_{\nu_a} m_{d_c} D_0(m_{\nu_a}^2, m_{d_c}^2, m_{W^-}^2, m_{H_d^-}^2) \quad (\text{F.237})$$

$$V_{(w_2^u)}^{LR} = V_{a,\alpha,d}^{+,L} \hat{\bar{V}}_{\beta,a}^{+,L} \hat{V}_{\delta,c}^{u,R} \bar{V}_{c,\gamma,d}^{d,R} m_{\nu_a} m_{d_c} D_0(m_{\nu_a}^2, m_{d_c}^2, m_{W^-}^2, m_{H_d^-}^2) \quad (\text{F.238})$$

$$T_{(w_2^u)}^{LL} = -V_{a,\alpha,d}^{+,L} \hat{\bar{V}}_{\beta,a}^{+,R} \hat{V}_{\delta,c}^{u,R} \bar{V}_{c,\gamma,d}^{d,L} D_{27}(m_{\nu_a}^2, m_{d_c}^2, m_{W^-}^2, m_{H_d^-}^2) \quad (\text{F.239})$$

$$S_{(w_3^u)}^{LL} = -4 \hat{V}_{a,i}^{+,L} \bar{V}_{\beta,a,b}^{+,L} V_{\delta,c,b}^{u,L} \hat{\bar{V}}_{c,\gamma}^{d,L} D_{27}(m_{\nu_a}^2, m_{d_c}^2, m_{H_b^-}^2, m_{W^-}^2) \quad (\text{F.240})$$

$$S_{(w_3^u)}^{LR} = -4 \hat{V}_{a,i}^{+,L} \bar{V}_{\beta,a,b}^{+,L} V_{\delta,c,b}^{u,R} \hat{\bar{V}}_{c,\gamma}^{d,R} D_{27}(m_{\nu_a}^2, m_{d_c}^2, m_{H_b^-}^2, m_{W^-}^2) \quad (\text{F.241})$$

$$V_{(w_3^u)}^{LL} = \hat{V}_{a,i}^{+,L} \bar{V}_{\beta,a,b}^{+,R} V_{\delta,c,b}^{u,R} \hat{\bar{V}}_{c,\gamma}^{d,L} m_{\nu_a} m_{d_c} D_0(m_{\nu_a}^2, m_{d_c}^2, m_{H_b^-}^2, m_{W^-}^2) \quad (\text{F.242})$$

$$V_{(w_3^u)}^{LR} = \hat{V}_{a,i}^{+,L} \bar{V}_{\beta,a,b}^{+,R} V_{\delta,c,b}^{u,L} \hat{\bar{V}}_{c,\gamma}^{d,R} m_{\nu_a} m_{d_c} D_0(m_{\nu_a}^2, m_{d_c}^2, m_{H_b^-}^2, m_{W^-}^2) \quad (\text{F.243})$$

$$T_{(w_3^u)}^{LL} = -\hat{V}_{a,i}^{+,L} \bar{V}_{\beta,a,b}^{+,L} V_{\delta,c,b}^{u,L} \hat{\bar{V}}_{c,\gamma}^{d,L} D_{27}(m_{\nu_a}^2, m_{d_c}^2, m_{H_b^-}^2, m_{W^-}^2) \quad (\text{F.244})$$

$$S_{(w_4^u)}^{LL} = -4\hat{V}_{a,i}^{+,L}\hat{\bar{V}}_{\beta,a}^{+,R}\hat{V}_{\delta,c}^{u,R}\hat{\bar{V}}_{c,\gamma}^{d,L}m_{\nu_a}m_{d_c}D_0(m_{\nu_a}^2, m_{d_c}^2, m_{W^-}^2, m_{W^-}^2) \quad (\text{F.245})$$

$$S_{(w_4^u)}^{LR} = -4\hat{V}_{a,i}^{+,L}\hat{\bar{V}}_{\beta,a}^{+,R}\hat{V}_{\delta,c}^{u,L}\hat{\bar{V}}_{c,\gamma}^{d,R}m_{\nu_a}m_{d_c}D_0(m_{\nu_a}^2, m_{d_c}^2, m_{W^-}^2, m_{W^-}^2) \quad (\text{F.246})$$

$$V_{(w_4^u)}^{LL} = 16\hat{V}_{a,i}^{+,L}\hat{\bar{V}}_{\beta,a}^{+,L}\hat{V}_{\delta,c}^{u,L}\hat{\bar{V}}_{c,\gamma}^{d,L}D_{27}(m_{\nu_a}^2, m_{d_c}^2, m_{W^-}^2, m_{W^-}^2) \quad (\text{F.247})$$

$$V_{(w_4^u)}^{LR} = 4\hat{V}_{a,i}^{+,L}\hat{\bar{V}}_{\beta,a}^{+,L}\hat{V}_{\delta,c}^{u,R}\hat{\bar{V}}_{c,\gamma}^{d,R}D_{27}(m_{\nu_a}^2, m_{d_c}^2, m_{W^-}^2, m_{W^-}^2) \quad (\text{F.248})$$

$$T_{(w_4^u)}^{LL} = -\hat{V}_{a,i}^{+,L}\hat{\bar{V}}_{\beta,a}^{+,R}\hat{V}_{\delta,c}^{u,R}\hat{\bar{V}}_{c,\gamma}^{d,L}m_{\nu_a}m_{d_c}D_0(m_{\nu_a}^2, m_{d_c}^2, m_{W^-}^2, m_{W^-}^2) \quad (\text{F.249})$$

G Form factors of the 4-fermion operators

We define the sum over all penguin diagrams as

$$V_{Z,sum}^{LL} = \frac{1}{16\pi^2} \left(\sum_a V_{Z,(n_a)}^{LL} + \sum_a V_{Z,(c_a)}^{LL} + \sum_a V_{Z,(w_a)}^{LL} \right) \quad (\text{G.1})$$

$$V_{Z,sum}^{LR} = \frac{1}{16\pi^2} \left(\sum_a V_{Z,(n_a)}^{LR} + \sum_a V_{Z,(c_a)}^{LR} + \sum_a V_{Z,(w_a)}^{LR} \right) \quad (\text{G.2})$$

$$S_{h_p,sum}^{LL} = \frac{1}{16\pi^2} \left(\sum_a S_{h_p,(n_a)}^{LL} + \sum_a S_{h_p,(c_a)}^{LL} + \sum_a S_{h_p,(w_a)}^{LL} \right) \quad (\text{G.3})$$

$$S_{h_p,sum}^{LR} = \frac{1}{16\pi^2} \left(\sum_a S_{h_p,(n_a)}^{LR} + \sum_a S_{h_p,(c_a)}^{LR} + \sum_a S_{h_p,(w_a)}^{LR} \right) \quad (\text{G.4})$$

$$S_{A_p^0,sum}^{LL} = \frac{1}{16\pi^2} \left(\sum_a S_{A_p^0,(n_a)}^{LL} + \sum_a S_{A_p^0,(c_a)}^{LL} + \sum_a S_{A_p^0,(w_a)}^{LL} \right) \quad (\text{G.5})$$

$$S_{A_p^0,sum}^{LR} = \frac{1}{16\pi^2} \left(\sum_a S_{A_p^0,(n_a)}^{LR} + \sum_a S_{A_p^0,(c_a)}^{LR} + \sum_a S_{A_p^0,(w_a)}^{LR} \right) \quad (\text{G.6})$$

and the sum over all boxes as

$$X_{(sum^x)}^{LL} = \frac{1}{16\pi^2} \left(\sum_a X_{(n_a^x)}^{LL} + \sum_a X_{(c_a^x)}^{LL} + \sum_a X_{(w_a^x)}^{LL} \right) \quad (\text{G.7})$$

$$X_{(sum^x)}^{LR} = \frac{1}{16\pi^2} \left(\sum_a X_{(n_a^x)}^{LR} + \sum_a X_{(c_a^x)}^{LR} + \sum_a X_{(w_a^x)}^{LR} \right) \quad (\text{G.8})$$

with $X = V, S, T$ and $x = \ell, d, u$. With these, we can finally obtain the form factors of the 4-lepton operators as follows:

$$A_{LL}^V = V_{Z,sum}^{LL}E_{\gamma,\delta}^L\frac{1}{m_Z^2} + V_{(sum^\ell)}^{LL} \quad (\text{G.9})$$

$$A_{LR}^V = V_{Z,sum}^{LR}E_{\gamma,\delta}^R\frac{1}{m_Z^2} + V_{(sum^\ell)}^{LR} \quad (\text{G.10})$$

$$A_{LL}^S = \sum_p S_{h_p,sum}^{LL}H_{\gamma,\delta,p}^L\frac{1}{m_{h_p}^2} + \sum_p S_{A_p^0,sum}^{LL}A_{\gamma,\delta,p}^L\frac{1}{m_{A_p^0}^2} + S_{(sum^\ell)}^{LL} \quad (\text{G.11})$$

$$A_{LR}^S = \sum_p S_{h_p,sum}^{LR}H_{\gamma,\delta,p}^R\frac{1}{m_{h_p}^2} + \sum_p S_{A_p^0,sum}^{LR}A_{\gamma,\delta,p}^R\frac{1}{m_{A_p^0}^2} + S_{(sum^\ell)}^{LR} \quad (\text{G.12})$$

$$A_{LL}^T = T_{(sum^\ell)}^{LL} \quad (G.13)$$

$$A_{LR}^T = T_{(sum^\ell)}^{LR} \quad (G.14)$$

$$B_{LL}^V = V_{Z,sum}^{LL} D_{\gamma,\delta}^L \frac{1}{m_Z^2} + V_{(sum^\ell)}^{LL} \quad (G.15)$$

$$B_{LR}^V = V_{Z,sum}^{LR} D_{\gamma,\delta}^R \frac{1}{m_Z^2} + V_{(sum^\ell)}^{LR} \quad (G.16)$$

$$B_{LL}^S = \sum_p S_{h_p,sum}^{LL} H_{\gamma,\delta,p}^{d,L} \frac{1}{m_{h_p}^2} + \sum_p S_{A_p^0,sum}^{LL} A_{\gamma,\delta,p}^{d,L} \frac{1}{m_{A_p^0}^2} + S_{(sum^d)}^{LL} \quad (G.17)$$

$$B_{LR}^S = \sum_p S_{h_p,sum}^{LR} H_{\gamma,\delta,p}^{d,R} \frac{1}{m_{h_p}^2} + \sum_p S_{A_p^0,sum}^{LR} A_{\gamma,\delta,p}^{d,R} \frac{1}{m_{A_p^0}^2} + S_{(sum^d)}^{LR} \quad (G.18)$$

$$B_{LL}^T = T_{(sum^d)}^{LL} \quad (G.19)$$

$$B_{LR}^T = T_{(sum^d)}^{LR} \quad (G.20)$$

$$C_{LL}^V = V_{Z,sum}^{LL} U_{\gamma,\delta}^L \frac{1}{m_Z^2} + V_{(sum^u)}^{LL} \quad (G.21)$$

$$C_{LR}^V = V_{Z,sum}^{LR} U_{\gamma,\delta}^R \frac{1}{m_Z^2} + V_{(sum^u)}^{LR} \quad (G.22)$$

$$C_{LL}^S = \sum_p S_{h_p,sum}^{LL} H_{\gamma,\delta,p}^{u,L} \frac{1}{m_{h_p}^2} + \sum_p S_{A_p^0,sum}^{LL} A_{\gamma,\delta,p}^{u,L} \frac{1}{m_{A_p^0}^2} + S_{(sum^u)}^{LL} \quad (G.23)$$

$$C_{LR}^S = \sum_p S_{h_p,sum}^{LR} H_{\gamma,\delta,p}^{u,R} \frac{1}{m_{h_p}^2} + \sum_p S_{A_p^0,sum}^{LR} A_{\gamma,\delta,p}^{u,R} \frac{1}{m_{A_p^0}^2} + S_{(sum^u)}^{LR} \quad (G.24)$$

$$C_{LL}^T = T_{(sum^u)}^{LL} \quad (G.25)$$

$$C_{LR}^T = T_{(sum^u)}^{LR} \quad (G.26)$$

and the other chiralities are given by $X_{RL}^W = X_{LR}^W(R \leftrightarrow L)$ and $X_{RR}^W = X_{LL}^W(R \leftrightarrow L)$ ($X = A, B, C; W = S, T, V$).

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](#)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] ATLAS collaboration, *Observation of a new particle in the search for the standard model Higgs boson with the ATLAS detector at the LHC*, *Phys. Lett. B* **716** (2012) 1 [[arXiv:1207.7214](#)] [[INSPIRE](#)].
- [2] CMS collaboration, *Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC*, *Phys. Lett. B* **716** (2012) 30 [[arXiv:1207.7235](#)] [[INSPIRE](#)].
- [3] D.V. Forero, M. Tortola and J.W.F. Valle, *Global status of neutrino oscillation parameters after Neutrino-2012*, *Phys. Rev. D* **86** (2012) 073012 [[arXiv:1205.4018](#)] [[INSPIRE](#)].

- [4] M.C. Gonzalez-Garcia, M. Maltoni, J. Salvado and T. Schwetz, *Global fit to three neutrino mixing: critical look at present precision*, *JHEP* **12** (2012) 123 [[arXiv:1209.3023](#)] [[INSPIRE](#)].
- [5] F. Capozzi et al., *Status of three-neutrino oscillation parameters, circa 2013*, *Phys. Rev. D* **89** (2014) 093018 [[arXiv:1312.2878](#)] [[INSPIRE](#)].
- [6] S. Mihara, J.P. Miller, P. Paradisi and G. Piredda, *Charged lepton flavor-violation experiments*, *Ann. Rev. Nucl. Part. Sci.* **63** (2013) 531 [[INSPIRE](#)].
- [7] MEG collaboration, J. Adam et al., *New constraint on the existence of the $\mu^+ \rightarrow e^+\gamma$ decay*, *Phys. Rev. Lett.* **110** (2013) 201801 [[arXiv:1303.0754](#)] [[INSPIRE](#)].
- [8] A.M. Baldini et al., *MEG upgrade proposal*, [arXiv:1301.7225](#) [[INSPIRE](#)].
- [9] T. Aushev et al., *Physics at super B factory*, [arXiv:1002.5012](#) [[INSPIRE](#)].
- [10] SINDRUM collaboration, U. Bellgardt et al., *Search for the decay $\mu^+ \rightarrow e^+e^+e^-$* , *Nucl. Phys. B* **299** (1988) 1 [[INSPIRE](#)].
- [11] A. Blondel et al., *Research proposal for an experiment to search for the decay $\mu \rightarrow eee$* , [arXiv:1301.6113](#) [[INSPIRE](#)].
- [12] BABAR and BELLE collaborations, A.J. Bevan et al., *The physics of the B factories*, [arXiv:1406.6311](#) [[INSPIRE](#)].
- [13] LHCb collaboration, *Searches for violation of lepton flavour and baryon number in τ lepton decays at LHCb*, *Phys. Lett. B* **724** (2013) 36 [[arXiv:1304.4518](#)] [[INSPIRE](#)].
- [14] MU2E collaboration, R.M. Carey et al., *Proposal to search for $\mu^-N \rightarrow e^-N$ with a single event sensitivity below 10^{-16}* , FERMILAB-PROPOSAL-0973, Fermilab, Batavia U.S.A. (2008) [[INSPIRE](#)].
- [15] MU2E collaboration, D. Glenzinski, *The Mu2e experiment at Fermilab*, *AIP Conf. Proc.* **1222** (2010) 383 [[INSPIRE](#)].
- [16] MU2E collaboration, R.J. Abrams et al., *Mu2e conceptual design report*, [arXiv:1211.7019](#) [[INSPIRE](#)].
- [17] DEEME collaboration, M. Aoki, *A new idea for an experimental search for ν -e conversion*, *PoS(ICHEP 2010)279* [[INSPIRE](#)].
- [18] COMET collaboration, Y.G. Cui et al., *Conceptual design report for experimental search for lepton flavor violating μ^-e^- conversion at sensitivity of 10^{-16} with a slow-extracted bunched proton beam (COMET)*, KEK-2009-10, Japan (2009) [[INSPIRE](#)].
- [19] COMET collaboration, Y. Kuno, *A search for muon-to-electron conversion at J-PARC: the COMET experiment*, *Prog. Theor. Exp. Phys.* **2013** (2013) 022C01 [[INSPIRE](#)].
- [20] PRISM/PRIME WORKING GROUP collaboration, *An experimental search for a μ^-e^- conversion at sensitivity of the order of 10^{-18} with a highly intense muon source: PRISM*, http://j-parc.jp/researcher/Hadron/en/pac_0606/pdf/p20-Kuno.pdf.
- [21] A. Pilaftsis, *Lepton flavor nonconservation in H^0 decays*, *Phys. Lett. B* **285** (1992) 68 [[INSPIRE](#)].
- [22] J.L. Diaz-Cruz and J.J. Toscano, *Lepton flavor violating decays of Higgs bosons beyond the standard model*, *Phys. Rev. D* **62** (2000) 116005 [[hep-ph/9910233](#)] [[INSPIRE](#)].

- [23] G. Blankenburg, J. Ellis and G. Isidori, *Flavour-changing decays of a 125 GeV Higgs-like particle*, *Phys. Lett. B* **712** (2012) 386 [[arXiv:1202.5704](#)] [[INSPIRE](#)].
- [24] R. Harnik, J. Kopp and J. Zupan, *Flavor violating Higgs decays*, *JHEP* **03** (2013) 026 [[arXiv:1209.1397](#)] [[INSPIRE](#)].
- [25] S. Davidson and P. Verdier, *LHC sensitivity to the decay $h \rightarrow \tau^\pm mu^\mp$* , *Phys. Rev. D* **86** (2012) 111701 [[arXiv:1211.1248](#)] [[INSPIRE](#)].
- [26] A. Arhrib, Y. Cheng and O.C.W. Kong, *Higgs to $\mu^\mp\tau^+$ decay in supersymmetry without R-parity*, *Europhys. Lett.* **101** (2013) 31003 [[arXiv:1208.4669](#)] [[INSPIRE](#)].
- [27] A. Arhrib, Y. Cheng and O.C.W. Kong, *A comprehensive analysis on lepton flavor violating Higgs to $\mu\bar{\tau} + \tau\bar{n}u$ decay in supersymmetry without R-parity*, *Phys. Rev. D* **87** (2013) 015025 [[arXiv:1210.8241](#)] [[INSPIRE](#)].
- [28] P.S. Bhupal Dev, R. Franceschini and R.N. Mohapatra, *Bounds on TeV seesaw models from LHC Higgs data*, *Phys. Rev. D* **86** (2012) 093010 [[arXiv:1207.2756](#)] [[INSPIRE](#)].
- [29] M. Arana-Catania, E. Arganda and M.J. Herrero, *Non-decoupling SUSY in LFV Higgs decays: a window to new physics at the LHC*, *JHEP* **09** (2013) 160 [[arXiv:1304.3371](#)] [[INSPIRE](#)].
- [30] A. Falkowski, D.M. Straub and A. Vicente, *Vector-like leptons: Higgs decays and collider phenomenology*, *JHEP* **05** (2014) 092 [[arXiv:1312.5329](#)] [[INSPIRE](#)].
- [31] E. Arganda, M.J. Herrero, X. Marcano and C. Weiland, *Imprints of massive ISS neutrinos in LFV Higgs decays*, [arXiv:1405.4300](#) [[INSPIRE](#)].
- [32] S. Bressler, A. Dery and A. Efrati, *Asymmetric lepton-flavor violating Higgs decays*, *Phys. Rev. D* **90** (2014) 015025 [[arXiv:1405.4545](#)] [[INSPIRE](#)].
- [33] J. Kopp and M. Nardecchia, *Flavor and CP-violation in Higgs decays*, *JHEP* **10** (2014) 156 [[arXiv:1406.5303](#)] [[INSPIRE](#)].
- [34] D.A. Sierra and A. Vicente, *Explaining the CMS Higgs flavor violating decay excess*, [arXiv:1409.7690](#) [[INSPIRE](#)].
- [35] CMS collaboration, *Search for lepton flavour violating decays of the Higgs boson*, CMS-PAS-HIG-14-005, CERN, Geneva Switzerland (2014).
- [36] W. Porod and W. Majerotto, *Large lepton flavor violating signals in supersymmetric particle decays at future e^+e^- colliders*, *Phys. Rev. D* **66** (2002) 015003 [[hep-ph/0201284](#)] [[INSPIRE](#)].
- [37] A. Bartl et al., *Test of lepton flavor violation at LHC*, *Eur. Phys. J. C* **46** (2006) 783 [[hep-ph/0510074](#)] [[INSPIRE](#)].
- [38] M. Hirsch, J.W.F. Valle, W. Porod, J.C. Romao and A. Villanova del Moral, *Probing minimal supergravity in type-I seesaw with lepton flavour violation at the LHC*, *Phys. Rev. D* **78** (2008) 013006 [[arXiv:0804.4072](#)] [[INSPIRE](#)].
- [39] M. Hirsch, S. Kaneko and W. Porod, *Supersymmetric seesaw type. II. LHC and lepton flavour violating phenomenology*, *Phys. Rev. D* **78** (2008) 093004 [[arXiv:0806.3361](#)] [[INSPIRE](#)].
- [40] S. Kaneko, J. Sato, T. Shimomura, O. Vives and M. Yamanaka, *Measuring lepton flavour violation at LHC with long-lived slepton in the coannihilation region*, *Phys. Rev. D* **78** (2008) 116013 [[arXiv:0811.0703](#)] [[INSPIRE](#)].

- [41] F. del Aguila and J.A. Aguilar-Saavedra, *Distinguishing seesaw models at LHC with multi-lepton signals*, *Nucl. Phys. B* **813** (2009) 22 [[arXiv:0808.2468](#)] [[INSPIRE](#)].
- [42] A. Atre, T. Han, S. Pascoli and B. Zhang, *The search for heavy Majorana neutrinos*, *JHEP* **05** (2009) 030 [[arXiv:0901.3589](#)] [[INSPIRE](#)].
- [43] A. Abada, A.J.R. Figueiredo, J.C. Romao and A.M. Teixeira, *Interplay of LFV and slepton mass splittings at the LHC as a probe of the SUSY seesaw*, *JHEP* **10** (2010) 104 [[arXiv:1007.4833](#)] [[INSPIRE](#)].
- [44] J.N. Esteves et al., *LHC and lepton flavour violation phenomenology of a left-right extension of the MSSM*, *JHEP* **12** (2010) 077 [[arXiv:1011.0348](#)] [[INSPIRE](#)].
- [45] J.N. Esteves, J.C. Romao, M. Hirsch, F. Staub and W. Porod, *Supersymmetric type-III seesaw: lepton flavour violating decays and dark matter*, *Phys. Rev. D* **83** (2011) 013003 [[arXiv:1010.6000](#)] [[INSPIRE](#)].
- [46] A. Abada, A.J.R. Figueiredo, J.C. Romao and A.M. Teixeira, *Probing the supersymmetric type-III seesaw: LFV at low-energies and at the LHC*, *JHEP* **08** (2011) 099 [[arXiv:1104.3962](#)] [[INSPIRE](#)].
- [47] A. Abada, A.J.R. Figueiredo, J.C. Romao and A.M. Teixeira, *Lepton flavour violation: physics potential of a linear collider*, *JHEP* **08** (2012) 138 [[arXiv:1206.2306](#)] [[INSPIRE](#)].
- [48] P. Bandyopadhyay, E.J. Chun, H. Okada and J.-C. Park, *Higgs signatures in inverse seesaw model at the LHC*, *JHEP* **01** (2013) 079 [[arXiv:1209.4803](#)] [[INSPIRE](#)].
- [49] S. Mondal, S. Biswas, P. Ghosh and S. Roy, *Exploring novel correlations in trilepton channels at the LHC for the minimal supersymmetric inverse seesaw model*, *JHEP* **05** (2012) 134 [[arXiv:1201.1556](#)] [[INSPIRE](#)].
- [50] A. Das and N. Okada, *Inverse seesaw neutrino signatures at the LHC and ILC*, *Phys. Rev. D* **88** (2013) 113001 [[arXiv:1207.3734](#)] [[INSPIRE](#)].
- [51] A.M. Teixeira, A. Abada, A.J.R. Figueiredo and J.C. Romao, *Lepton flavour violation at high energies: the LHC and a linear collider*, *Nuovo Cim. C* **037** (2014) 19 [[arXiv:1402.1426](#)] [[INSPIRE](#)].
- [52] P.S. Bhupal Dev, S. Mondal, B. Mukhopadhyaya and S. Roy, *Phenomenology of light sneutrino dark matter in cmSSM/mSUGRA with inverse seesaw*, *JHEP* **09** (2012) 110 [[arXiv:1207.6542](#)] [[INSPIRE](#)].
- [53] BABAR collaboration, B. Aubert et al., *Searches for lepton flavor violation in the decays $\tau^\pm \rightarrow e^\pm \gamma$ and $\tau^\pm \rightarrow \mu^\pm \gamma$* , *Phys. Rev. Lett.* **104** (2010) 021802 [[arXiv:0908.2381](#)] [[INSPIRE](#)].
- [54] K. Hayasaka et al., *Search for lepton flavor violating τ decays into three leptons with 719 million produced $\tau^+\tau^-$ pairs*, *Phys. Lett. B* **687** (2010) 139 [[arXiv:1001.3221](#)] [[INSPIRE](#)].
- [55] SINDRUM II collaboration, C. Dohmen et al., *Test of lepton flavor conservation in $\mu \rightarrow e$ conversion on titanium*, *Phys. Lett. B* **317** (1993) 631 [[INSPIRE](#)].
- [56] SINDRUM II collaboration, W.H. Bertl et al., *A search for muon to electron conversion in muonic gold*, *Eur. Phys. J. C* **47** (2006) 337 [[INSPIRE](#)].
- [57] DEEME collaboration, M. Aoki, *An experimental search for muon-electron conversion in nuclear field at sensitivity of 10^{-14} with a pulsed proton beam*, *AIP Conf. Proc.* **1441** (2012) 599 [[INSPIRE](#)].

- [58] A.J. Buras, B. Duling, T. Feldmann, T. Heidsieck and C. Promberger, *Lepton flavour violation in the presence of a fourth generation of quarks and leptons*, *JHEP* **09** (2010) 104 [[arXiv:1006.5356](#)] [[INSPIRE](#)].
- [59] R.N. Mohapatra and J.W.F. Valle, *Neutrino mass and baryon number nonconservation in superstring models*, *Phys. Rev. D* **34** (1986) 1642 [[INSPIRE](#)].
- [60] A. Elsayed, S. Khalil and S. Moretti, *Higgs mass corrections in the SUSY B-L model with inverse seesaw*, *Phys. Lett. B* **715** (2012) 208 [[arXiv:1106.2130](#)] [[INSPIRE](#)].
- [61] M. Hirsch, M. Malinsky, W. Porod, L. Reichert and F. Staub, *Hefty MSSM-like light Higgs in extended gauge models*, *JHEP* **02** (2012) 084 [[arXiv:1110.3037](#)] [[INSPIRE](#)].
- [62] E.J. Chun, V.S. Mummidri and S.K. Vempati, *Anatomy of Higgs mass in supersymmetric inverse seesaw models*, *Phys. Lett. B* **736** (2014) 470 [[arXiv:1405.5478](#)] [[INSPIRE](#)].
- [63] J. Bernabeu, A. Santamaria, J. Vidal, A. Mendez and J.W.F. Valle, *Lepton flavor nonconservation at high-energies in a superstring inspired standard model*, *Phys. Lett. B* **187** (1987) 303 [[INSPIRE](#)].
- [64] A. Ilakovac and A. Pilaftsis, *Flavor violating charged lepton decays in seesaw-type models*, *Nucl. Phys. B* **437** (1995) 491 [[hep-ph/9403398](#)] [[INSPIRE](#)].
- [65] F. Deppisch and J.W.F. Valle, *Enhanced lepton flavor violation in the supersymmetric inverse seesaw model*, *Phys. Rev. D* **72** (2005) 036001 [[hep-ph/0406040](#)] [[INSPIRE](#)].
- [66] F. Deppisch, T.S. Kosmas and J.W.F. Valle, *Enhanced μ^- - e^- conversion in nuclei in the inverse seesaw model*, *Nucl. Phys. B* **752** (2006) 80 [[hep-ph/0512360](#)] [[INSPIRE](#)].
- [67] A. Ilakovac and A. Pilaftsis, *Supersymmetric lepton flavour violation in low-scale seesaw models*, *Phys. Rev. D* **80** (2009) 091902 [[arXiv:0904.2381](#)] [[INSPIRE](#)].
- [68] R. Alonso, M. Dhen, M.B. Gavela and T. Hambye, *Muon conversion to electron in nuclei in type-I seesaw models*, *JHEP* **01** (2013) 118 [[arXiv:1209.2679](#)] [[INSPIRE](#)].
- [69] D.N. Dinh, A. Ibarra, E. Molinaro and S.T. Petcov, *The μ - e conversion in nuclei, $\mu \rightarrow e\gamma$, $\mu \rightarrow 3e$ decays and TeV scale see-saw scenarios of neutrino mass generation*, *JHEP* **08** (2012) 125 [Erratum *ibid.* **09** (2013) 023] [[arXiv:1205.4671](#)] [[INSPIRE](#)].
- [70] A. Ilakovac, A. Pilaftsis and L. Popov, *Charged lepton flavor violation in supersymmetric low-scale seesaw models*, *Phys. Rev. D* **87** (2013) 053014 [[arXiv:1212.5939](#)] [[INSPIRE](#)].
- [71] C.-H. Lee, P.S. Bhupal Dev and R.N. Mohapatra, *Natural TeV-scale left-right seesaw mechanism for neutrinos and experimental tests*, *Phys. Rev. D* **88** (2013) 093010 [[arXiv:1309.0774](#)] [[INSPIRE](#)].
- [72] M. Hirsch, F. Staub and A. Vicente, *Enhancing $\ell_i \rightarrow 3\ell_j$ with the Z^0 -penguin*, *Phys. Rev. D* **85** (2012) 113013 [[arXiv:1202.1825](#)] [[INSPIRE](#)].
- [73] H.K. Dreiner, K. Nickel, F. Staub and A. Vicente, *New bounds on trilinear R-parity violation from lepton flavor violating observables*, *Phys. Rev. D* **86** (2012) 015003 [[arXiv:1204.5925](#)] [[INSPIRE](#)].
- [74] M. Hirsch, W. Porod, L. Reichert and F. Staub, *Phenomenology of the minimal supersymmetric $U(1)_{B-L} \times U(1)_R$ extension of the standard model*, *Phys. Rev. D* **86** (2012) 093018 [[arXiv:1206.3516](#)] [[INSPIRE](#)].

- [75] A. Abada, D. Das, A. Vicente and C. Weiland, *Enhancing lepton flavour violation in the supersymmetric inverse seesaw beyond the dipole contribution*, *JHEP* **09** (2012) 015 [[arXiv:1206.6497](#)] [[INSPIRE](#)].
- [76] M.E. Krauss, W. Porod and F. Staub, SO(10) *inspired gauge-mediated supersymmetry breaking*, *Phys. Rev. D* **88** (2013) 015014 [[arXiv:1304.0769](#)] [[INSPIRE](#)].
- [77] M.E. Krauss et al., *Decoupling of heavy sneutrinos in low-scale seesaw models*, *Phys. Rev. D* **90** (2014) 013008 [[arXiv:1312.5318](#)] [[INSPIRE](#)].
- [78] E. Arganda and M.J. Herrero, *Testing supersymmetry with lepton flavor violating τ and μ decays*, *Phys. Rev. D* **73** (2006) 055003 [[hep-ph/0510405](#)] [[INSPIRE](#)].
- [79] W. Porod, F. Staub and A. Vicente, *A flavor kit for BSM models*, *Eur. Phys. J. C* **74** (2014) 2992 [[arXiv:1405.1434](#)] [[INSPIRE](#)].
- [80] F. Staub, *SARAH*, [arXiv:0806.0538](#) [[INSPIRE](#)].
- [81] F. Staub, *From superpotential to model files for FeynArts and CalcHep/CompHEP*, *Comput. Phys. Commun.* **181** (2010) 1077 [[arXiv:0909.2863](#)] [[INSPIRE](#)].
- [82] F. Staub, *Automatic calculation of supersymmetric renormalization group equations and self energies*, *Comput. Phys. Commun.* **182** (2011) 808 [[arXiv:1002.0840](#)] [[INSPIRE](#)].
- [83] F. Staub, *SARAH 3.2: Dirac gauginos, UFO output and more*, *Comput. Phys. Commun.* **184** (2013) 1792 [[arXiv:1207.0906](#)] [[INSPIRE](#)].
- [84] F. Staub, *SARAH 4: a tool for (not only SUSY) model builders*, *Comput. Phys. Commun.* **185** (2014) 1773 [[arXiv:1309.7223](#)] [[INSPIRE](#)].
- [85] W. Porod, *SPheno, a program for calculating supersymmetric spectra, SUSY particle decays and SUSY particle production at e^+e^- colliders*, *Comput. Phys. Commun.* **153** (2003) 275 [[hep-ph/0301101](#)] [[INSPIRE](#)].
- [86] W. Porod and F. Staub, *SPheno 3.1: extensions including flavour, CP-phases and models beyond the MSSM*, *Comput. Phys. Commun.* **183** (2012) 2458 [[arXiv:1104.1573](#)] [[INSPIRE](#)].
- [87] T. Hahn and M. Pérez-Victoria, *Automatized one loop calculations in four-dimensions and D-dimensions*, *Comput. Phys. Commun.* **118** (1999) 153 [[hep-ph/9807565](#)] [[INSPIRE](#)].
- [88] T. Hahn, *Generating Feynman diagrams and amplitudes with FeynArts 3*, *Comput. Phys. Commun.* **140** (2001) 418 [[hep-ph/0012260](#)] [[INSPIRE](#)].
- [89] T. Hahn, *Automatic loop calculations with FeynArts, FormCalc and LoopTools*, *Nucl. Phys. Proc. Suppl.* **89** (2000) 231 [[hep-ph/0005029](#)] [[INSPIRE](#)].
- [90] T. Hahn, *New features in FormCalc 4*, *Nucl. Phys. Proc. Suppl.* **135** (2004) 333 [[hep-ph/0406288](#)] [[INSPIRE](#)].
- [91] T. Hahn, *New developments in FormCalc 4.1*, *eConf C* **050318** (2005) 0604 [[hep-ph/0506201](#)] [[INSPIRE](#)].
- [92] B. Chokoufe Nejad, T. Hahn, J.-N. Lang and E. Mirabella, *FormCalc 8: better algebra and vectorization*, *J. Phys. Conf. Ser.* **523** (2014) 012050 [[arXiv:1310.0274](#)] [[INSPIRE](#)].
- [93] A. Abada, D. Das and C. Weiland, *Enhanced Higgs mediated lepton flavour violating processes in the supersymmetric inverse seesaw model*, *JHEP* **03** (2012) 100 [[arXiv:1111.5836](#)] [[INSPIRE](#)].

- [94] R.N. Mohapatra, *Mechanism for understanding small neutrino mass in superstring theories*, *Phys. Rev. Lett.* **56** (1986) 561 [[INSPIRE](#)].
- [95] M. Malinsky, T. Ohlsson, Z.-Z. Xing and H. Zhang, *Non-unitary neutrino mixing and CP-violation in the minimal inverse seesaw model*, *Phys. Lett. B* **679** (2009) 242 [[arXiv:0905.2889](#)] [[INSPIRE](#)].
- [96] M.B. Gavela, T. Hambye, D. Hernandez and P. Hernández, *Minimal flavour seesaw models*, *JHEP* **09** (2009) 038 [[arXiv:0906.1461](#)] [[INSPIRE](#)].
- [97] A. Abada and M. Lucente, *Looking for the minimal inverse seesaw realisation*, *Nucl. Phys. B* **885** (2014) 651 [[arXiv:1401.1507](#)] [[INSPIRE](#)].
- [98] M. Hirsch, T. Kernreiter, J.C. Romao and A. Villanova del Moral, *Minimal supersymmetric inverse seesaw: neutrino masses, lepton flavour violation and LHC phenomenology*, *JHEP* **01** (2010) 103 [[arXiv:0910.2435](#)] [[INSPIRE](#)].
- [99] G. 't Hooft, *Naturalness, chiral symmetry, and spontaneous chiral symmetry breaking*, *NATO Sci. Ser. B* **59** (1980) 135 [[INSPIRE](#)].
- [100] M.C. Gonzalez-Garcia and J.W.F. Valle, *Fast decaying neutrinos and observable flavor violation in a new class of Majoron models*, *Phys. Lett. B* **216** (1989) 360 [[INSPIRE](#)].
- [101] CMS collaboration, *Search for new physics in the multijet and missing transverse momentum final state in proton-proton collisions at $\sqrt{s} = 8$ TeV*, *JHEP* **06** (2014) 055 [[arXiv:1402.4770](#)] [[INSPIRE](#)].
- [102] ATLAS collaboration, *Search for squarks and gluinos with the ATLAS detector in final states with jets and missing transverse momentum using $\sqrt{s} = 8$ TeV proton-proton collision data*, *JHEP* **09** (2014) 176 [[arXiv:1405.7875](#)] [[INSPIRE](#)].
- [103] E. Arganda, M.J. Herrero and A.M. Teixeira, *μ -e conversion in nuclei within the CMSSM seesaw: universality versus non-universality*, *JHEP* **10** (2007) 104 [[arXiv:0707.2955](#)] [[INSPIRE](#)].
- [104] J. Hisano, T. Moroi, K. Tobe and M. Yamaguchi, *Lepton flavor violation via right-handed neutrino Yukawa couplings in supersymmetric standard model*, *Phys. Rev. D* **53** (1996) 2442 [[hep-ph/9510309](#)] [[INSPIRE](#)].
- [105] E. Arganda, private communication.
- [106] Y. Kuno and Y. Okada, *Muon decay and physics beyond the standard model*, *Rev. Mod. Phys.* **73** (2001) 151 [[hep-ph/9909265](#)] [[INSPIRE](#)].
- [107] J.D. Vergados, *The neutrino mass and family, lepton and baryon nonconservation in gauge theories*, *Phys. Rept.* **133** (1986) 1 [[INSPIRE](#)].
- [108] J. Bernabeu, E. Nardi and D. Tommasini, *μ -e conversion in nuclei and Z' physics*, *Nucl. Phys. B* **409** (1993) 69 [[hep-ph/9306251](#)] [[INSPIRE](#)].
- [109] A. Faessler, T.S. Kosmas, S. Kovalenko and J.D. Vergados, *Constraints on R-parity violating supersymmetry from muon electron nuclear conversion*, [hep-ph/9904335](#) [[INSPIRE](#)].
- [110] R. Kitano, M. Koike and Y. Okada, *Detailed calculation of lepton flavor violating muon electron conversion rate for various nuclei*, *Phys. Rev. D* **66** (2002) 096002 [Erratum *ibid. D* **76** (2007) 059902] [[hep-ph/0203110](#)] [[INSPIRE](#)].

- [111] A. Crivellin, M. Hoferichter and M. Procura, *Improved predictions for $\mu \rightarrow e$ conversion in nuclei and Higgs-induced lepton flavor violation*, *Phys. Rev. D* **89** (2014) 093024 [[arXiv:1404.7134](#)] [[INSPIRE](#)].
- [112] H.C. Chiang, E. Oset, T.S. Kosmas, A. Faessler and J.D. Vergados, *Coherent and incoherent (μ^-, e^-) conversion in nuclei*, *Nucl. Phys. A* **559** (1993) 526 [[INSPIRE](#)].
- [113] T.S. Kosmas, S. Kovalenko and I. Schmidt, *Nuclear μ - e^- conversion in strange quark sea*, *Phys. Lett. B* **511** (2001) 203 [[hep-ph/0102101](#)] [[INSPIRE](#)].
- [114] A. Brignole, G. Degrassi, P. Slavich and F. Zwirner, *On the $O(\alpha_t^2)$ two loop corrections to the neutral Higgs boson masses in the MSSM*, *Nucl. Phys. B* **631** (2002) 195 [[hep-ph/0112177](#)] [[INSPIRE](#)].
- [115] G. Degrassi, P. Slavich and F. Zwirner, *On the neutral Higgs boson masses in the MSSM for arbitrary stop mixing*, *Nucl. Phys. B* **611** (2001) 403 [[hep-ph/0105096](#)] [[INSPIRE](#)].
- [116] A. Brignole, G. Degrassi, P. Slavich and F. Zwirner, *On the two loop sbottom corrections to the neutral Higgs boson masses in the MSSM*, *Nucl. Phys. B* **643** (2002) 79 [[hep-ph/0206101](#)] [[INSPIRE](#)].
- [117] A. Dedes and P. Slavich, *Two loop corrections to radiative electroweak symmetry breaking in the MSSM*, *Nucl. Phys. B* **657** (2003) 333 [[hep-ph/0212132](#)] [[INSPIRE](#)].
- [118] A. Dedes, G. Degrassi and P. Slavich, *On the two loop Yukawa corrections to the MSSM Higgs boson masses at large $\tan\beta$* , *Nucl. Phys. B* **672** (2003) 144 [[hep-ph/0305127](#)] [[INSPIRE](#)].
- [119] B.C. Allanach, A. Djouadi, J.L. Kneur, W. Porod and P. Slavich, *Precise determination of the neutral Higgs boson masses in the MSSM*, *JHEP* **09** (2004) 044 [[hep-ph/0406166](#)] [[INSPIRE](#)].
- [120] J.A. Casas and A. Ibarra, *Oscillating neutrinos and $\mu \rightarrow e, \gamma$* , *Nucl. Phys. B* **618** (2001) 171 [[hep-ph/0103065](#)] [[INSPIRE](#)].
- [121] L. Basso et al., *Proposal for generalised supersymmetry Les Houches accord for see-saw models and PDG numbering scheme*, *Comput. Phys. Commun.* **184** (2013) 698 [[arXiv:1206.4563](#)] [[INSPIRE](#)].
- [122] A. Abada, D. Das, A.M. Teixeira, A. Vicente and C. Weiland, *Tree-level lepton universality violation in the presence of sterile neutrinos: impact for R_K and R_π* , *JHEP* **02** (2013) 048 [[arXiv:1211.3052](#)] [[INSPIRE](#)].
- [123] D.V. Forero, M. Tortola and J.W.F. Valle, *Neutrino oscillations refitted*, [arXiv:1405.7540](#) [[INSPIRE](#)].
- [124] K.S. Babu and C. Kolda, *Higgs mediated $\tau \rightarrow 3\mu$ in the supersymmetric seesaw model*, *Phys. Rev. Lett.* **89** (2002) 241802 [[hep-ph/0206310](#)] [[INSPIRE](#)].
- [125] LHCb collaboration, *First evidence for the decay $B_s^0 \rightarrow \mu^+ \mu^-$* , *Phys. Rev. Lett.* **110** (2013) 021801 [[arXiv:1211.2674](#)] [[INSPIRE](#)].
- [126] S.P. Martin and M.T. Vaughn, *Two loop renormalization group equations for soft supersymmetry breaking couplings*, *Phys. Rev. D* **50** (1994) 2282 [Erratum *ibid. D* **78** (2008) 039903] [[hep-ph/9311340](#)] [[INSPIRE](#)].