

RECEIVED: June 28, 2020

REVISED: September 11, 2020

ACCEPTED: October 12, 2020

PUBLISHED: November 26, 2020

Six-point conformal blocks in the snowflake channel

Jean-François Fortin,^a Wen-Jie Ma^a and Witold Skiba^b

^a*Département de Physique, de Génie Physique et d'Optique, Université Laval,
1045 avenue de la Médecine, Québec, QC G1V 0A6, Canada*

^b*Department of Physics, Yale University,
217 Prospect Street, New Haven, CT 06520, U.S.A.*

E-mail: jean-francois.fortin@phy.ulaval.ca, wenjie.ma.1@ulaval.ca,
witold.skiba@yale.edu

ABSTRACT: We compute d -dimensional scalar six-point conformal blocks in the two possible topologies allowed by the operator product expansion. Our computation is a simple application of the embedding space operator product expansion formalism developed recently. Scalar six-point conformal blocks in the comb channel have been determined not long ago, and we present here the first explicit computation of the scalar six-point conformal blocks in the remaining inequivalent topology. For obvious reason, we dub the other topology the snowflake channel. The scalar conformal blocks, with scalar external and exchange operators, are presented as a power series expansion in the conformal cross-ratios, where the coefficients of the power series are given as a double sum of the hypergeometric type. In the comb channel, the double sum is expressible as a product of two ${}_3F_2$ -hypergeometric functions. In the snowflake channel, the double sum is expressible as a Kampé de Fériet function where both sums are intertwined and cannot be factorized. We check our results by verifying their consistency under symmetries and by taking several limits reducing to known results, mostly to scalar five-point conformal blocks in arbitrary spacetime dimensions.

KEYWORDS: Conformal Field Theory, Conformal and W Symmetry, Field Theories in Higher Dimensions

ARXIV EPRINT: [2004.02824](https://arxiv.org/abs/2004.02824)

Contents

1	Introduction	1
2	Scalar six-point conformal blocks	3
2.1	M -point correlation functions from the OPE	4
2.2	Scalar M -point correlation functions	5
2.3	Scalar M -point correlation functions in the comb channel	6
2.4	Scalar six-point correlation functions in the snowflake channel	7
3	Sanity checks	10
3.1	Symmetry properties	10
3.2	OPE limit	14
3.3	Limit of unit operator	16
4	Discussion and conclusion	16
A	Scalar five-point conformal blocks and the OPE	22
A.1	Proof of the equivalence	22
B	Snowflake and the OPE	27
B.1	Proof of the snowflake	27
B.2	An alternative form	32
C	Symmetry properties	36
C.1	Rotations of the triangle	36
C.2	Reflections of the triangle	38
C.3	Permutations of the dendrites	45

1 Introduction

The study of higher-point conformal blocks in conformal field theory (CFT) is a complicated subject without many explicit results. In a CFT, correlation functions, which are the natural observables of the theory, are given in terms of the CFT data and the conformal blocks. The CFT data, which consist of the spectrum of quasi-primary operators as well as the operator product expansion (OPE) coefficients, completely determine all correlation functions with up to three points. For higher-point correlation functions, the appearance of conformal cross-ratios, which are invariant under conformal transformations, leads to conformal blocks. The conformal blocks are functions of the conformal cross-ratios which are in principle fully constrained by conformal invariance.

Although conformal blocks are fixed by conformal invariance, they are notoriously difficult to compute in all generality. Several techniques have been developed over the years for the computation of four-point conformal blocks, which are the simplest blocks. For example, various methods use Casimir equations [1–3], the shadow formalism [4–6], the weight-shifting formalism [7, 8], integrability [9–13], AdS/CFT [14–19], and the OPE [20–35]. Another important reason why four-point conformal blocks have been studied extensively is the conformal bootstrap [36, 37], a way of constraining the CFT data solely from consistency of correlation functions under associativity. Indeed, it is known that four-point conformal blocks are sufficient to implement the full conformal bootstrap.

Conformal blocks with more than four points have not been studied in great detail as of now. Until very recently, the only results were for scalar M -point blocks in one and two spacetime dimensions as well as scalar five-point blocks in any spacetime dimensions [38–42]. Last year, the scalar M -point conformal blocks in the so-called comb channel were presented in [43, 44]. They showed that the scalar M -point conformal blocks in the comb channel can be expressed as a power series expansion in the conformal cross-ratios with the coefficients containing a product of $M - 4$ ${}_3F_2$ -hypergeometric functions. Although the techniques employed in the two references were different, AdS/CFT versus OPE, the two results have very similar forms even though the basis of conformal cross-ratios were distinct.

An interesting feature of higher-point correlation functions is that there exist several inequivalent topologies. Indeed, starting at six points, the use of OPE among pairs of operators in different sequences of operator pairings leads to different topologies, see figure 3. The comb channel [39] is only one of the many topologies that are possible for higher-point correlation functions.

One possible advantage of the OPE formalism used in [44] is that it is not limited to the comb channel. Indeed, it can be applied to any pair of operators in the M -point correlation function in any channel of interest. The net effect of the OPE is “adding” an operator to an $(M - 1)$ -point correlation function in a specific place of choice. This allows generation of possible multiple topologies with M points from one specific diagram with $M - 1$ points. After all, the existence of the many topologies is a consequence of the OPE, as was directly observed above. More importantly, the choice in the pairs of quasi-primary operators does not lead to any new computational complications.

With the OPE formalism [27, 30, 31], the OPE differential operator is applied on the known $(M - 1)$ -point correlation functions to generate M -point correlation functions. Since the action of the OPE differential operator on arbitrary products of conformal cross-ratios has been determined [30, 31], it is intuitively straightforward to proceed with the computation. It can however be technically laborious to express the final results in the most convenient way possible. Indeed, starting at five points, it is necessary to re-express the conformal cross-ratios of the original correlation function in terms of the conformal cross-ratios appropriate for the OPE differential operator, leading to several superfluous sums. Re-summations must be performed to recover a simple result for the final correlation function. For the computation of the scalar M -point conformal blocks, we found that, with an appropriate basis of conformal cross-ratios, all re-summations were easy

${}_2F_1$ -hypergeometric function re-summations [44]. For other channels, the necessary computations within the OPE formalism are exactly the same at the technical level, hence it should be straightforward to compute higher-point correlation functions in all topologies using the OPE formalism.

In this paper, we begin the investigation of higher-point correlation functions in all topologies by computing the scalar six-point conformal blocks in the remaining channel, see the bottom part of figure 3, which for obvious reason, we call the snowflake channel.¹ We show that the superfluous sums can be taken care of with simple ${}_2F_1$ -hypergeometric function re-summations (as for the comb channel) and some identities for ${}_3F_2$ -hypergeometric functions (absent in the comb channel).

This paper is organized as follows: section 2 presents a summary of the embedding space OPE formalism. The OPE is reviewed and the action of the scalar OPE differential operator is discussed. A general form for the contribution to the correlation functions of external scalars with scalar exchanges is presented, as well as the associated scalar conformal blocks. Moreover, the recurrence relation taking $(M-1)$ -point correlation functions to M -point correlation functions is introduced, and, after re-considering scalar M -point conformal blocks in the comb channel, the scalar six-point conformal block in the snowflake channel is given. Section 3 provides several consistency checks of the snowflake result. First, interesting symmetry properties of the scalar conformal blocks in the snowflake channel are proven. Then, the OPE limit and the limit of unit operator are used to ascertain that the snowflake result passes those tests. Finally, we conclude in section 4 while several appendices present most of the more technical proofs. Appendix A contains derivations of the scalar five-point conformal blocks of [43] directly from the OPE, showing that the results of [43] and [44] are equivalent for five points. Appendix B demonstrates how the snowflake result is obtained. An equivalent result is also presented and the proof of their equivalence is shown. Appendix C proves the identities that the scalar six-point conformal blocks in the snowflake channel verify under the symmetry group of the snowflake.

2 Scalar six-point conformal blocks

With the knowledge of the OPE and its explicit action on any function of conformal cross-ratios, one can in principle compute any correlation function starting from the known two-point functions or, for that matter, the only non-trivial one-point function. With this technique, starting at five points and above, it is necessary to re-express the initial correlation function in terms of the conformal cross-ratios appropriate for the OPE differential operator, and then rewrite the solution in terms of the most convenient conformal cross-ratios by re-summing as many superfluous sums as possible. This method is quite powerful, allowing the computation of conformal blocks in any channel. Although straightforward, it is however not always clear *a priori* what is the best choice of conformal cross-ratios that will lead to the simplest final answer. In this section, we quickly review the OPE and then sketch the derivation and state the result for the scalar six-point conformal blocks in both channels. Concrete proofs are left for the appendices.

¹In [42, 43], the snowflake channel is called the OPE channel.

2.1 M -point correlation functions from the OPE

The embedding space OPE introduced in [30, 31] states that the product of two quasi-primary operators can be expressed as

$$\begin{aligned}
 \mathcal{O}_i(\eta_1)\mathcal{O}_j(\eta_2) &= \sum_k \sum_{a=1}^{N_{ijk}} {}_a c_{ij}^k {}_a \mathcal{D}_{ij}^k(\eta_1, \eta_2) \mathcal{O}_k(\eta_2), \\
 {}_a \mathcal{D}_{ij}^k(\eta_1, \eta_2) &= \frac{1}{(\eta_1 \cdot \eta_2)^{p_{ijk}}} (\mathcal{T}_{12}^{N_i} \Gamma) (\mathcal{T}_{21}^{N_j} \Gamma) \cdot {}_a v_{ij}^{12k} \cdot \mathcal{D}_{12}^{(d, h_{ijk} - n_a/2, n_a)} (\mathcal{T}_{12 N_k} \Gamma)^*, \\
 p_{ijk} &= \frac{1}{2}(\tau_i + \tau_j - \tau_k), \quad h_{ijk} = -\frac{1}{2}(\chi_i - \chi_j + \chi_k), \\
 \tau_{\mathcal{O}} &= \Delta_{\mathcal{O}} - S_{\mathcal{O}}, \quad \chi_{\mathcal{O}} = \Delta_{\mathcal{O}} - \xi_{\mathcal{O}}, \quad \xi_{\mathcal{O}} = S_{\mathcal{O}} - \lfloor S_{\mathcal{O}} \rfloor,
 \end{aligned} \tag{2.1}$$

where the OPE differential operator and the remaining quantities (half-projectors, tensor structures, etc.) are introduced and detailed in [31]. Since all the calculations involve only scalar operators, the OPE can be simplified significantly as the tensor structures and half-projectors are trivial in that case. Considering an OPE of two scalar operators and neglecting any operators with spin on the right-hand side leads to

$$\begin{aligned}
 \mathcal{O}_i(\eta_1)\mathcal{O}_j(\eta_2) &= c_{ij}^k \frac{1}{(\eta_1 \cdot \eta_2)^{p_{ijk}}} \mathcal{D}_{12}^{(d, h_{ijk}, 0)} \mathcal{O}_k(\eta_2) + \dots, \\
 p_{ijk} &= \frac{1}{2}(\Delta_i + \Delta_j - \Delta_k), \quad h_{ijk} = -\frac{1}{2}(\Delta_i - \Delta_j + \Delta_k),
 \end{aligned} \tag{2.2}$$

where Δ_i is the scaling dimension of \mathcal{O}_i and c_{ij}^k is the OPE coefficient. The coordinates η_i are the embedding space coordinates in $d+2$ dimensions that are constrained to the light cone $\eta_i \cdot \eta_i = 0$ and are projectively identified $\eta_i \sim \lambda \eta_i$ for $\lambda > 0$.

For scalar operators another simplification occurs in the differential operator that has a particularly simple form when the Lorentz indices are absent, indicated by 0 in the third argument

$$\mathcal{D}_{ij}^{(d, h, 0)} = \left[\mathcal{D}_{ij}^2 \right]^h \equiv \left[(\eta_i \cdot \eta_j) \partial_j^2 - (d+2) \eta_j \cdot \partial_j \eta_i \cdot \partial_j \right]^h,$$

where $\partial_{jA} = \frac{\partial}{\partial \eta_j^A}$. It is more practical to rescale the scalar differential operator, \mathcal{D}_{ij}^2 , and introduce an operator that is homogeneous of degree 0 with respect to all the coordinates

$$\bar{\mathcal{D}}_{ij;kl;m}^2 = \frac{\eta_{ij} \eta_{kl}}{\eta_{ik} \eta_{il}} \mathcal{D}_{ij}^2,$$

with $\eta_{ij} \equiv \eta_i \cdot \eta_j$ for brevity. The action of $\bar{\mathcal{D}}_{ij;kl;m}^2$ on the conformal cross-ratios

$$\begin{aligned}
 x_m &= \frac{\eta_{ij} \eta_{kl} \eta_{im}}{\eta_{ik} \eta_{il} \eta_{jm}}, \\
 y_a &= 1 - \frac{\eta_{im} \eta_{ja}}{\eta_{ia} \eta_{jm}}, \quad 1 \leq a \leq M, \quad a \neq i, j, m, \\
 z_{ab} &= \frac{\eta_{ik} \eta_{il} \eta_{ab}}{\eta_{kl} \eta_{ia} \eta_{ib}}, \quad 1 \leq a < b \leq M, \quad a, b \neq i, j,
 \end{aligned} \tag{2.3}$$

has been obtained explicitly²

$$\begin{aligned}
 & \bar{\mathcal{D}}_{ij;kl;m}^{2h} x_m^{\bar{q}} \prod_{\substack{1 \leq a \leq M \\ a \neq i,j,m}} (1 - y_a)^{-q_a} \\
 &= x_m^{\bar{q}+h} \sum_{\{n_a, n_{am}, n_{ab}\} \geq 0} \frac{(-h)_{\bar{n}_m + \bar{n}} (q_m)_{\bar{n}_m} (\bar{q} + h)_{\bar{n} - \bar{n}}}{(\bar{q})_{\bar{n} + \bar{n}_m} (\bar{q} + 1 - d/2)_{\bar{n}_m + \bar{n}}} \\
 & \times \prod_{\substack{1 \leq a \leq M \\ a \neq i,j,m}} \frac{(q_a)_{n_a}}{n_{am}! (n_a - n_{am} - \bar{n}_a)!} y_a^{n_a} \left(\frac{x_m z_{am}}{y_a} \right)^{n_{am}} \prod_{\substack{1 \leq a < b \leq M \\ a, b \neq i,j,m}} \frac{1}{n_{ab}!} \left(\frac{x_m z_{ab}}{y_a y_b} \right)^{n_{ab}},
 \end{aligned} \tag{2.4}$$

where we have defined $\bar{q} = \sum_{\substack{1 \leq a \leq M \\ a \neq i,j}} q_a$ as well as

$$\begin{aligned}
 \bar{n} &= \sum_{\substack{1 \leq a \leq M \\ a \neq i,j,m}} n_a, & \bar{n}_m &= \sum_{\substack{1 \leq a \leq M \\ a \neq i,j,m}} n_{am}, \\
 \bar{n}_a &= \sum_{\substack{1 \leq b \leq M \\ b \neq i,j,m,a}} n_{ab}, & \bar{n} &= \sum_{\substack{1 \leq a < b \leq M \\ a, b \neq i,j,m}} n_{ab}.
 \end{aligned}$$

The OPE (2.1) was used to determine two-, three-, and four-points in [32–34]. Very specific rules, somewhat reminiscent of Feynman rules, were also developed in [35] to compute all necessary ingredients to implement the full conformal bootstrap at the level of four-point correlation functions.

2.2 Scalar M -point correlation functions

In general, it is possible to write the contribution to scalar M -point correlation functions, i.e. with scalar external and exchanged quasi-primary operators, from a specific channel as

$$I_{M(\Delta_{k_1}, \dots, \Delta_{k_{M-3}})}^{(\Delta_{i_2}, \dots, \Delta_{i_M}, \Delta_{i_1})} \Big|_{\text{channel}} = L_{M|\text{channel}}^{(\Delta_{i_2}, \dots, \Delta_{i_M}, \Delta_{i_1})} \left[\prod_{1 \leq a \leq M-3} (u_a^M)^{\frac{\Delta_{k_a}}{2}} \right] G_{M|\text{channel}}^{(d, h; \mathbf{p})}(\mathbf{u}^M, \mathbf{v}^M). \tag{2.5}$$

The scalar M -point conformal block is given by

$$\begin{aligned}
 G_{M|\text{channel}}^{(d, h; \mathbf{p})}(\mathbf{u}^M, \mathbf{v}^M) &= \sum_{\{m_a, m_{ab}\} \geq 0} C_{M|\text{channel}}^{(d, h; \mathbf{p})}(\mathbf{m}, \mathbf{m}) F_{M|\text{channel}}^{(d, h; \mathbf{p})}(\mathbf{m}, \mathbf{m}) \\
 & \times \prod_{1 \leq a \leq M-3} \frac{(u_a^M)^{m_a}}{m_a!} \prod_{1 \leq a < b \leq M-3} \frac{(1 - v_{ab}^M)^{m_{ab}}}{m_{ab}!},
 \end{aligned} \tag{2.6}$$

where the cross-ratios u_a and v_{ab} are defined below in (2.8).

The complete M -point correlation functions are sums of the different I_M , including exchanges of operators in non-trivial representations which are not discussed here.

In (2.5) the conformal dimensions of the external scalar quasi-primary operators are Δ_{i_a} while the conformal dimensions of the exchanged scalar quasi-primary operators are Δ_{k_a} . Moreover, the legs L_M are products of embedding space coordinates necessary to satisfy covariance under scale transformations while the conformal cross-ratios are denoted by the

²We note that the OPE differential operator used here is a simple rescaling of the one defined in [31].

vector \mathbf{u}^M of u_a^M and the matrix \mathbf{v}^M of v_{ab}^M . The scalar M -point conformal blocks (2.6) are written as sums over powers of conformal cross-ratios, with extra sums denoted by the function F_M .³

Finally, in the scalar M -point conformal blocks (2.6) the vectors \mathbf{h} and \mathbf{p} are generated by the action of the OPE (2.1) on the pairs of quasi-primary operators relevant to the channel of interest. This statement translates into

$$I_{M(\Delta_{k_1}, \dots, \Delta_{k_{M-3}})}^{(\Delta_{i_2}, \dots, \Delta_{i_M}, \Delta_{i_1})} \Big|_{\text{channel}} = \frac{1}{\eta_{1M}^{p_M}} \left(\frac{\eta_{kM} \eta_{lM}}{\eta_{1M} \eta_{kl}} \right)^{h_{M-1}} \bar{\mathcal{D}}_{M1;kl;m}^{2h_{M-1}} I_{M-1(\Delta_{k_1}, \dots, \Delta_{k_{M-4}})}^{(\Delta_{i_2}, \dots, \Delta_{i_{M-1}}, \Delta_{k_{M-3}})} \Big|_{\text{channel}}, \quad (2.7)$$

for some convenient choice of k , l and m with the appropriate channel on the r.h.s. to generate the desired channel on the l.h.s. For future convenience, we also define $\bar{p}_a = \sum_{b=2}^a p_b$ and $\bar{h}_a = \sum_{b=2}^a h_b$. The above equation is the essence of the OPE approach to computing correlation functions. Two operators in an M -point function are replaced by one operator appearing on the right-hand side of the OPE. This reduces the M -point function to an $(M-1)$ -point function. Reading the equation in the other direction, the differential operator present in the OPE generates the expression for the M -point function when it acts on a previously computed $(M-1)$ -point function.

In this context, the OPE (2.1) was also put to work in the computation of scalar M -point correlation functions in the comb channel in [44], to which we now turn.

2.3 Scalar M -point correlation functions in the comb channel

Acting repetitively with the OPE as in (2.7), we found in [44] the following scalar M -point conformal blocks (2.5) and (2.6) in the comb channel,

$$L_{M|\text{comb}}^{(\Delta_{i_2}, \dots, \Delta_{i_M}, \Delta_{i_1})} = \left(\frac{\eta_{34}}{\eta_{23} \eta_{24}} \right)^{\frac{\Delta_{i_2}}{2}} \left[\prod_{1 \leq a \leq M-2} \left(\frac{\eta_{a+1, a+3}}{\eta_{a+1, a+2} \eta_{a+2, a+3}} \right)^{\frac{\Delta_{i_{a+2}}}{2}} \right] \left(\frac{\eta_{M-1, M}}{\eta_{1, M-1} \eta_{1M}} \right)^{\frac{\Delta_{i_1}}{2}},$$

$$u_a^M = \frac{\eta_{1+a, 2+a} \eta_{3+a, 4+a}}{\eta_{1+a, 3+a} \eta_{2+a, 4+a}}, \quad v_{ab}^M = \frac{\eta_{2-a+b, 4+b}}{\eta_{2+b, 4+b}} \prod_{1 \leq c \leq a} \frac{\eta_{3+b-c, 4+b-c}}{\eta_{2+b-c, 4+b-c}}, \quad (2.8)$$

with

$$C_{M|\text{comb}}^{(d, \mathbf{h}; \mathbf{p})} = \frac{(p_3)_{m_1 + \text{tr}_0 \mathbf{m}} (p_2 + h_2)_{m_1 + \text{tr}_1 \mathbf{m}}}{(p_3)_{m_1 + \text{tr}_1 \mathbf{m}}} \left[\prod_{1 \leq a \leq M-3} \frac{(\bar{p}_{a+2} + \bar{h}_{a+2})_{m_a + m_{a+1} + \bar{m}_a + \bar{\bar{m}}_a}}{(\bar{p}_{a+2} + \bar{h}_{a+1})_{2m_a + \bar{m}_{a-1} + \bar{m}_a + \bar{\bar{m}}_a}} \right. \\ \left. \times (p_{a+2} - m_{a-1})_{m_a + \text{tr}_a \mathbf{m}} \frac{(-h_{a+2})_{m_a} (-h_{a+2} + m_a - m_{a+1})_{\bar{m}_{a-1}}}{(\bar{p}_{a+2} + \bar{h}_{a+1} + 1 - d/2)_{m_a}} \right],$$

$$\text{tr}_a \mathbf{m} = \sum_b m_{b, a+b}, \quad \bar{m}_a = \sum_{b \leq a} m_{ba}, \quad \bar{\bar{m}}_a = \sum_{b > a} (\bar{m}_b - \text{tr}_b \mathbf{m}), \quad (2.9)$$

³In general, F_M could be a function of both the vector $\mathbf{m} = (m_a)$ and the matrix $\mathbf{m} = (m_{ab})$. Thus the separation between C_M and F_M is somewhat arbitrary in (2.6). Here, F_6 in the snowflake channel is only a function of the vector \mathbf{m} , as is the case for F_M in the comb channel. That statement seems to generalize to all F_M , thus we conjecture that F_M can always be chosen such that it is a function of the vector \mathbf{m} only. Moreover, we construct $F_{6|\text{snowflake}}$ such that it has some interesting symmetry properties.

and

$$F_{M|\text{comb}}^{(d,h;p)}(\mathbf{m}) = \prod_{1 \leq a \leq M-4} {}_3F_2 \left[\begin{matrix} -m_a, -m_{a+1}, -\bar{p}_{a+2} - \bar{h}_{a+1} + d/2 - m_a \\ p_{a+3} - m_a, h_{a+2} + 1 - m_a \end{matrix}; 1 \right], \quad (2.10)$$

and finally

$$\begin{aligned} 2h_2 &= \Delta_{k_1} - \Delta_{i_2} - \Delta_{i_3}, & 2h_a &= \Delta_{k_{a-1}} - \Delta_{k_{a-2}} - \Delta_{i_{a+1}}, \\ p_2 &= \Delta_{i_3}, & 2p_3 &= \Delta_{i_2} + \Delta_{k_1} - \Delta_{i_3}, & 2p_a &= \Delta_{i_a} + \Delta_{k_{a-2}} - \Delta_{k_{a-3}}, \end{aligned} \quad (2.11)$$

where $k_{M-2} \equiv i_1$.

The computations were straightforward yet somewhat tedious. The choice of conformal cross-ratios was based on the OPE limit, a limit we use again later to check the validity of the scalar six-point conformal blocks in the snowflake channel discussed in the next subsection.

2.4 Scalar six-point correlation functions in the snowflake channel

As already mentioned, the OPE (2.1) or (2.2) can be applied on any pair of quasi-primary operators in the M -point correlation function, allowing the M -point correlation function to be expressed in terms of the derivative operator acting on the $(M-1)$ -point function in the corresponding channel. Hence, with the knowledge of the only five-point correlation function (that is in the comb channel), appropriately choosing the pair of quasi-primary operators on which the OPE acts leads to six-point correlation function in all channels. The desired channel follows from the choice of the OPE pair in the six-point function.

To reach the scalar six-point conformal blocks in the snowflake channel, we start from the four-point correlation functions and use the OPE appropriately. For $M=4$, there is only one channel with the topology of the comb, as shown in figure 1. One possible form for the contribution to scalar four-point correlation functions (2.5) and for scalar four-point conformal blocks (2.6) leads to [1, 2, 25, 26]

$$\begin{aligned} L_{4|\text{comb}}^{(\Delta_{i_2}, \dots, \Delta_{i_4}, \Delta_{i_1})} &= \left(\frac{\eta_{13}}{\eta_{12}\eta_{23}} \right)^{\frac{\Delta_{i_2}}{2}} \left(\frac{\eta_{12}}{\eta_{13}\eta_{23}} \right)^{\frac{\Delta_{i_3}}{2}} \left(\frac{\eta_{13}}{\eta_{14}\eta_{34}} \right)^{\frac{\Delta_{i_4}}{2}} \left(\frac{\eta_{34}}{\eta_{13}\eta_{14}} \right)^{\frac{\Delta_{i_1}}{2}}, \\ u_1^4 &= \frac{\eta_{14}\eta_{23}}{\eta_{12}\eta_{34}}, & v_{11}^4 &= \frac{\eta_{13}\eta_{24}}{\eta_{12}\eta_{34}}, \\ C_{4|\text{comb}}^{(d,h;p)} &= \frac{(-h_2)_{m_1+m_{11}}(p_2+h_3)_{m_1}(-h_3)_{m_1+m_{11}}(p_2+h_2)_{m_1}}{(p_2)_{2m_1+m_{11}}(p_2+1-d/2)_{m_1}}, \\ F_{4|\text{comb}}^{(d,h;p)} &= 1, \end{aligned} \quad (2.12)$$

$$\begin{aligned} 2h_2 &= \Delta_{i_3} - \Delta_{i_2} - \Delta_{k_1}, & 2h_3 &= \Delta_{i_1} - \Delta_{i_4} - \Delta_{k_1}, \\ p_2 &= \Delta_{k_1}, & 2p_3 &= \Delta_{i_2} + \Delta_{i_3} - \Delta_{k_1}, & 2p_4 &= \Delta_{i_4} + \Delta_{i_1} - \Delta_{k_1}. \end{aligned}$$

It is clear from (2.7) and figure 1 that there is only one channel for five-point conformal blocks since all external quasi-primary operators in four-point conformal blocks are topologically equivalent. This channel has the topology of the comb and is shown in figure 2.

$$I_{4|\text{comb}}^{(\Delta_{i_2}, \Delta_{i_3}, \Delta_{i_4}, \Delta_{i_1})} \Big|_{\Delta_{k_1}} = \mathcal{O}_{i_2} \begin{array}{c} \mathcal{O}_{i_3} \quad \mathcal{O}_{i_4} \\ | \quad | \\ \hline \mathcal{O}_{k_1} \end{array} \mathcal{O}_{i_1}$$

Figure 1. Scalar four-point conformal blocks.

$$I_{5|\text{comb}}^{(\Delta_{i_3}, \Delta_{i_4}, \Delta_{i_2}, \Delta_{i_5}, \Delta_{i_1})} \Big|_{\Delta_{k_1}, \Delta_{k_2}} = \mathcal{O}_{i_3} \begin{array}{c} \mathcal{O}_{i_4} \quad \mathcal{O}_{i_2} \quad \mathcal{O}_{i_5} \\ | \quad | \quad | \\ \hline \mathcal{O}_{k_1} \quad \mathcal{O}_{k_2} \end{array} \mathcal{O}_{i_1}$$

Figure 2. Scalar five-point conformal blocks.

It can be obtained from (2.7) with $k = 3$, $l = 4$ and $m = 4$, using the four-point correlation functions in the comb channel (2.12) where we first shifted $\mathcal{O}_{i_a}(\eta_a) \rightarrow \mathcal{O}_{i_{a-1}}(\eta_{a-1})$ with $\mathcal{O}_{i_0}(\eta_0) \equiv \mathcal{O}_{i_4}(\eta_4)$. The scalar five-point conformal blocks can be expressed as in (2.5) and (2.6) with the help of

$$\begin{aligned} L_{5|\text{comb}}^{(\Delta_{i_3}, \Delta_{i_4}, \Delta_{i_2}, \Delta_{i_5}, \Delta_{i_1})} &= \left(\frac{\eta_{14}}{\eta_{12}\eta_{24}} \right)^{\frac{\Delta_{i_2}}{2}} \left(\frac{\eta_{24}}{\eta_{23}\eta_{34}} \right)^{\frac{\Delta_{i_3}}{2}} \left(\frac{\eta_{23}}{\eta_{24}\eta_{34}} \right)^{\frac{\Delta_{i_4}}{2}} \left(\frac{\eta_{14}}{\eta_{15}\eta_{45}} \right)^{\frac{\Delta_{i_5}}{2}} \left(\frac{\eta_{45}}{\eta_{14}\eta_{15}} \right)^{\frac{\Delta_{i_1}}{2}}, \\ u_1^5 &= \frac{\eta_{12}\eta_{34}}{\eta_{14}\eta_{23}}, \quad u_2^5 = \frac{\eta_{15}\eta_{24}}{\eta_{12}\eta_{45}}, \quad v_{11}^5 = \frac{\eta_{13}\eta_{24}}{\eta_{14}\eta_{23}}, \quad v_{12}^5 = \frac{\eta_{14}\eta_{25}}{\eta_{12}\eta_{45}}, \quad v_{22}^5 = \frac{\eta_{24}\eta_{35}}{\eta_{23}\eta_{45}}, \\ C_{5|\text{comb}}^{(d, \mathbf{h}; \mathbf{p})} &= (-h_2)_{m_1+m_2+m_{11}+m_{22}} \frac{(-h_3)_{m_1+m_{11}+m_{22}} (p_2+h_2)_{m_1} (p_2+h_3)_{m_1}}{(p_2)_{2m_1+m_{11}+m_{22}} (p_2+1-d/2)_{m_1}} \\ &\quad \times \frac{(-h_4)_{m_2+m_{12}+m_{22}} (p_3-h_2+h_4)_{m_2+m_{11}} (p_3-m_1)_{m_2+m_{12}}}{(p_3-h_2)_{2m_2+m_{11}+m_{12}+m_{22}} (p_3-h_2+1-d/2)_{m_2}}, \\ F_{5|\text{comb}}^{(d, \mathbf{h}; \mathbf{p})} &= {}_3F_2 \left[\begin{array}{c} -m_1, -m_2, -p_2+d/2-m_1 \\ 1-p_2-h_2-m_1, p_3-m_1 \end{array} ; 1 \right], \\ 2h_2 &= \Delta_{i_2} - \Delta_{k_2} - \Delta_{k_1}, \quad 2h_3 = \Delta_{i_4} - \Delta_{i_3} - \Delta_{k_1}, \quad 2h_4 = \Delta_{i_1} - \Delta_{i_5} - \Delta_{k_2}, \\ p_2 &= \Delta_{k_1}, \quad 2p_3 = \Delta_{k_2} + \Delta_{i_2} - \Delta_{k_1}, \quad 2p_4 = \Delta_{i_3} + \Delta_{i_4} - \Delta_{k_1}, \quad 2p_5 = \Delta_{i_5} + \Delta_{i_1} - \Delta_{k_2}. \end{aligned} \tag{2.13}$$

We note here that the result (2.13) for the scalar five-point conformal blocks is equivalent to but different than the results (2.8), (2.9), (2.10), and (2.11) obtained in [44].⁴

For $M = 6$, there is now two possible topologies that can be obtained from five-point correlation functions. Indeed, from figure 2, it is clear that the external quasi-primary operator \mathcal{O}_{i_2} is different. Transforming the external quasi-primary operators \mathcal{O}_{i_1} , \mathcal{O}_{i_3} , \mathcal{O}_{i_4} , or \mathcal{O}_{i_5} into exchanged quasi-primary operators by appending two new external quasi-primary operators leads to the scalar six-point correlation function in the comb channel (see

⁴It is also different than the results of [39] and [43].

previous subsection). Doing the same with the quasi-primary operator \mathcal{O}_{i_2} gives instead the scalar six-point correlation in the snowflake channel. The difference can be seen in figure 3. For the scalar six-point correlation functions in the snowflake channel, we start from the scalar five-point conformal blocks (2.13) and shift the quasi-primary operators such that $\mathcal{O}_{i_a}(\eta_a) \rightarrow \mathcal{O}_{i_{a-1}}(\eta_{a-1})$ with the understanding that $\mathcal{O}_{i_0}(\eta_0) \equiv \mathcal{O}_{i_5}(\eta_5)$, and then use (2.7) with $k = 4$, $l = 5$ and $m = 5$, to get

$$\begin{aligned}
 L_{6|\text{snowflake}}^{(\Delta_{i_2}, \dots, \Delta_{i_6}, \Delta_{i_1})} &= \left(\frac{\eta_{13}}{\eta_{12}\eta_{23}} \right)^{\frac{\Delta_{i_2}}{2}} \left(\frac{\eta_{12}}{\eta_{13}\eta_{23}} \right)^{\frac{\Delta_{i_3}}{2}} \left(\frac{\eta_{35}}{\eta_{34}\eta_{45}} \right)^{\frac{\Delta_{i_4}}{2}} \\
 &\quad \times \left(\frac{\eta_{34}}{\eta_{35}\eta_{45}} \right)^{\frac{\Delta_{i_5}}{2}} \left(\frac{\eta_{15}}{\eta_{16}\eta_{56}} \right)^{\frac{\Delta_{i_6}}{2}} \left(\frac{\eta_{56}}{\eta_{15}\eta_{16}} \right)^{\frac{\Delta_{i_1}}{2}}, \\
 u_1^6 &= \frac{\eta_{15}\eta_{23}}{\eta_{12}\eta_{35}}, & u_2^6 &= \frac{\eta_{13}\eta_{45}}{\eta_{15}\eta_{34}}, & u_3^6 &= \frac{\eta_{16}\eta_{35}}{\eta_{13}\eta_{56}}, \\
 v_{11}^6 &= \frac{\eta_{13}\eta_{25}}{\eta_{12}\eta_{35}}, & v_{12}^6 &= \frac{\eta_{14}\eta_{35}}{\eta_{15}\eta_{34}}, & v_{22}^6 &= \frac{\eta_{13}\eta_{24}}{\eta_{12}\eta_{34}}, \\
 v_{13}^6 &= \frac{\eta_{15}\eta_{26}}{\eta_{12}\eta_{56}}, & v_{23}^6 &= \frac{\eta_{15}\eta_{36}}{\eta_{13}\eta_{56}}, & v_{33}^6 &= \frac{\eta_{35}\eta_{46}}{\eta_{34}\eta_{56}},
 \end{aligned} \tag{2.14}$$

with

$$\begin{aligned}
 C_{6|\text{snowflake}}^{(d, \mathbf{h}; \mathbf{p})} &= \frac{(p_2 + h_3)_{m_1+m_{23}} (p_3)_{-m_1+m_2+m_3+m_{12}+m_{33}} (-h_3)_{m_1+m_{11}+m_{22}+m_{13}}}{(p_2)_{2m_1+m_{11}+m_{13}+m_{22}+m_{23}} (p_2 + 1 - d/2)_{m_1}} \\
 &\quad \times \frac{(p_3 - h_2 + h_4)_{m_2+m_{11}} (p_2 + h_2)_{m_1-m_2+m_3+m_{13}+m_{23}} (-h_4)_{m_2+m_{12}+m_{22}+m_{33}}}{(p_3 - h_2)_{2m_2+m_{11}+m_{12}+m_{22}+m_{33}} (p_3 - h_2 + 1 - d/2)_{m_2}} \\
 &\quad \times \frac{(\bar{p}_3 + h_2 + h_5)_{m_3+m_{12}} (-h_2)_{m_1+m_2-m_3+m_{11}+m_{22}} (-h_5)_{m_3+m_{13}+m_{23}+m_{33}}}{(\bar{p}_3 + h_2)_{2m_3+m_{12}+m_{13}+m_{23}+m_{33}} (\bar{p}_3 + h_2 + 1 - d/2)_{m_3}},
 \end{aligned} \tag{2.15}$$

as well as

$$\begin{aligned}
 F_{6|\text{snowflake}}^{(d, \mathbf{h}; \mathbf{p})}(\mathbf{m}) &= \frac{(-p_3 + h_2 + d/2 - m_2)_{m_2}}{(p_3)_{-m_1} (-h_2 + m_1)_{-m_3}} \sum_{t_1, t_2 \geq 0} \frac{(-m_1)_{t_2} (-m_2)_{t_1} (-m_3)_{t_2}}{(p_3 - h_2 + 1 - d/2)_{t_1}} \\
 &\quad \times \frac{(p_3)_{t_1} (\bar{p}_3 - d/2)_{t_1} (-p_2 + d/2 - m_1)_{t_2} (1 + h_2 - m_1 + m_3)_{-t_1}}{(1 + h_2 - m_1)_{-t_1+t_2} (p_3 - m_1)_{t_1+t_2} t_1! t_2!} \\
 &= \frac{(-p_3 + h_2 + d/2 - m_2)_{m_2}}{(p_3)_{-m_1} (-h_2)_{-m_3}} \\
 &\quad \times F_{2,1,0}^{1,3,2} \left[\begin{matrix} \bar{p}_3 - d/2; -m_2, -h_2, p_3; -m_1, -m_3 \\ -h_2 - m_3, p_3 - m_1; p_3 - h_2 + 1 - d/2; - \end{matrix} \middle| 1, 1 \right],
 \end{aligned} \tag{2.16}$$

and finally

$$\begin{aligned}
 2h_2 &= \Delta_{k_3} - \Delta_{k_2} - \Delta_{k_1}, & 2h_3 &= \Delta_{i_3} - \Delta_{i_2} - \Delta_{k_1}, \\
 2h_4 &= \Delta_{i_5} - \Delta_{i_4} - \Delta_{k_2}, & 2h_5 &= \Delta_{i_1} - \Delta_{i_6} - \Delta_{k_3}, \\
 p_2 &= \Delta_{k_1}, & 2p_3 &= \Delta_{k_2} + \Delta_{k_3} - \Delta_{k_1}, & 2p_4 &= \Delta_{i_2} + \Delta_{i_3} - \Delta_{k_1}, \\
 2p_5 &= \Delta_{i_4} + \Delta_{i_5} - \Delta_{k_2}, & 2p_6 &= \Delta_{i_6} + \Delta_{i_1} - \Delta_{k_3}.
 \end{aligned} \tag{2.17}$$

$$\begin{aligned}
I_{6(\Delta_{k_1}, \Delta_{k_2}, \Delta_{k_3})}^{(\Delta_{i_2}, \dots, \Delta_{i_6}, \Delta_{i_1})} \Big|_{\text{comb}} &= \mathcal{O}_{i_2} \text{ --- } \begin{array}{c} \mathcal{O}_{i_3} \quad \mathcal{O}_{i_4} \quad \mathcal{O}_{i_5} \quad \mathcal{O}_{i_6} \\ | \quad | \quad | \quad | \\ \mathcal{O}_{k_1} \quad \mathcal{O}_{k_2} \quad \mathcal{O}_{k_3} \end{array} \text{ --- } \mathcal{O}_{i_1} \\
I_{6(\Delta_{k_1}, \Delta_{k_2}, \Delta_{k_3})}^{(\Delta_{i_2}, \dots, \Delta_{i_6}, \Delta_{i_1})} \Big|_{\text{snowflake}} &= \begin{array}{c} \mathcal{O}_{i_4} \quad \mathcal{O}_{i_5} \\ \diagdown \quad \diagup \\ \mathcal{O}_{k_2} \\ \diagup \quad \diagdown \\ \mathcal{O}_{i_3} \quad \mathcal{O}_{i_6} \\ \diagdown \quad \diagup \\ \mathcal{O}_{k_1} \quad \mathcal{O}_{k_3} \\ \diagdown \quad \diagup \\ \mathcal{O}_{i_2} \quad \mathcal{O}_{i_1} \end{array}
\end{aligned}$$

Figure 3. Scalar six-point conformal blocks in the comb (top) and snowflake (bottom) channels.

As mentioned previously, the function F_6 (2.16) in the snowflake channel is also a double sum of the hypergeometric type, same as in the comb channel. However, the snowflake double sum does not factorize into two hypergeometric functions, unlike the comb sum, compare (2.10) with $M = 6$. It can however be written as a Kampé de Fériet function $F_{2,1,0}^{1,3,2}$ as shown in appendix B (another expression in terms of a different Kampé de Fériet function $F_{1,1,1}^{2,1,1}$ is also given there). See (B.4) and [45, 46] for its definition.

In the following section, we study the snowflake results (2.14), (2.15), (2.16), and (2.17). We explicitly check the symmetry properties of G_6 and verify that it behaves properly under the OPE limit and the limit of unit operator. The proof of the snowflake results as well as an alternative form for the snowflake are shown in appendix B

3 Sanity checks

The scalar six-point conformal blocks obtained in the previous section must satisfy several properties. This section investigates the identities of G_6 (2.6) in the snowflake channel from the symmetries of the associated snowflake diagram figure 3. Then, the OPE limit and the limit of unit operator are taken to verify that the scalar six-point correlation functions reduce to the appropriate scalar five-point correlation functions.

3.1 Symmetry properties

The scalar M -point conformal blocks must verify several identities where the conformal cross-ratios and the vectors \mathbf{h} and \mathbf{p} are transformed. These identities are generated from the symmetries of the scalar M -point conformal blocks in the associated topology. The symmetries of the scalar M -point conformal blocks in the comb channel are relatively trivial. They correspond to $(\mathbb{Z}_2)^2 \rtimes \mathbb{Z}_2$, the semi-direct product of the direct product of




Figure 4. Symmetries of the scalar M -point conformal blocks in the comb channel. The figure shows the two generators, with reflections on the left and dendrite permutations on the right.




Figure 5. Symmetries of the scalar six-point conformal blocks in the snowflake channel. The figure shows rotations by $2\pi/3$ (left), reflections (middle), and dendrite permutations (right).

two cyclic groups \mathbb{Z}_2 of order two (for OPE, or dendrite, permutations depicted on the right of figure 4) and the cyclic group \mathbb{Z}_2 of order two (for reflection shown in the left part of figure 4). Here we present the identities of the scalar six-point conformal blocks in the snowflake channel. The proofs are left to appendix C.

The snowflake diagram figure 3 is invariant under the symmetry group generated by the three transformations shown in figure 5. Since the scalar six-point correlation functions in the snowflake channel I_6 are the same under symmetry transformations generated by the rotations, reflections and permutations described in figure 5, there are 47 identities that the scalar six-point conformal blocks G_6 should satisfy.

Although we dubbed it the snowflake channel, at first glance the symmetry group generated by rotations and reflections is only the dihedral group of order six, D_3 , which is the symmetry group of the triangle, not the hexagon expected for snowflakes. Including the OPE permutations, i.e. the permutations of the dendrites (or arms) of the snowflake, the full symmetry group of the snowflake diagram is however given by $(\mathbb{Z}_2)^3 \rtimes D_3$ where each cyclic group of order two corresponds to dendrite permutations. Since the order of this symmetry group is $|(\mathbb{Z}_2)^3 \rtimes D_3| = 48$, the snowflake diagram has a larger symmetry group than the hexagon, contrary to expectations.

Before proceeding, we verify that the transformations depicted in figure 5 indeed generate $(\mathbb{Z}_2)^3 \rtimes D_3$, the semi-direct product of $(\mathbb{Z}_2)^3$ (for dendrite permutations) and the dihedral group of order six. By defining the action of the symmetry generators on the external quasi-primary operators as R for the rotation (left diagram in figure 5), S for the reflection (center), and P for the permutation (right), it is easy to see that the dihedral part of symmetry group of the snowflake diagram has for presentation

$$\langle r, s | r^3 = s^2 = (rs)^2 = 1 \rangle, \tag{3.1}$$

with $r = R^{-1}$ and $s = S$. The presentation (3.1) corresponds to D_3 , with r and s representing rotations by $2\pi/3$ and reflections with respect to one of the three different axes, respectively. The $(\mathbb{Z}_2)^3 = \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$ part of the symmetry group is generated by $p_1 = P$, $p_2 = R^{-1}PR$, and $p_3 = RPR^{-1}$ with $p_i^2 = 1$ for all i . It is trivial to check that the p_i 's commute and that they correspond to dendrite permutations. To exclude the direct nature of the product, it suffices to observe that the generators p_i do not commute with the generators r and s . Having excluded $(\mathbb{Z}_2)^3 \times D_3$, it is easy to verify that the snowflake diagram has the symmetry group $(\mathbb{Z}_2)^3 \rtimes D_3$ of order 48.⁵

For rotations, we choose the generator where $\mathcal{O}_{i_a}(\eta_a) \rightarrow \mathcal{O}_{i_{a+2}}(\eta_{a+2})$ and $\Delta_{k_a} \rightarrow \Delta_{k_{a+1}}$ with $\mathcal{O}_{i_7}(\eta_7) \equiv \mathcal{O}_{i_1}(\eta_1)$, $\mathcal{O}_{i_8}(\eta_8) \equiv \mathcal{O}_{i_2}(\eta_2)$ as well as $\Delta_{k_4} \equiv \Delta_{k_1}$. Under this transformation, the legs and conformal cross-ratios (2.14) transform as

$$L_6 \prod_{1 \leq a \leq 3} (u_a^6)^{\frac{\Delta_{k_a}}{2}} \rightarrow L_6 \prod_{1 \leq a \leq 3} (u_a^6)^{\frac{\Delta_{k_a}}{2}},$$

$$\begin{aligned} u_1^6 &\rightarrow u_2^6, & u_2^6 &\rightarrow u_3^6, & u_3^6 &\rightarrow u_1^6, \\ v_{11}^6 &\rightarrow v_{12}^6, & v_{12}^6 &\rightarrow v_{23}^6, & v_{13}^6 &\rightarrow v_{22}^6, \\ v_{22}^6 &\rightarrow v_{33}^6, & v_{23}^6 &\rightarrow v_{11}^6, & v_{33}^6 &\rightarrow v_{13}^6, \end{aligned}$$

which imply the following identity,

$$\begin{aligned} G_{6|\text{snowflake}}^{(d, h_2, h_3, h_4, h_5; p_2, p_3, p_4, p_5, p_6)}(u_1^6, u_2^6, u_3^6; v_{11}^6, v_{12}^6, v_{13}^6, v_{22}^6, v_{23}^6, v_{33}^6) \\ = G_{6|\text{snowflake}}^{(d, -p_3, h_4, h_5, h_3; p_3 - h_2, p_2 + h_2, p_5, p_6, p_4)}(u_2^6, u_3^6, u_1^6; v_{12}^6, v_{23}^6, v_{22}^6, v_{33}^6, v_{11}^6, v_{13}^6). \end{aligned} \tag{3.2}$$

Using the decomposition (2.6) with (2.15) and (2.16), it is easy to see that C_6 does not change under this rotation generator, resulting in a non-trivial identity for F_6 [see (C.1)]. This identity can be translated into the language of Kampé de Fériet functions as discussed in the conclusion.

For reflections, we start with the generator acting as

$$\mathcal{O}_{i_2}(\eta_2) \leftrightarrow \mathcal{O}_{i_4}(\eta_4), \quad \mathcal{O}_{i_3}(\eta_3) \leftrightarrow \mathcal{O}_{i_5}(\eta_5), \quad \Delta_{k_1} \leftrightarrow \Delta_{k_2}.$$

For this reflection generator, the legs and conformal cross-ratios (2.14) transform as

$$L_6 \prod_{1 \leq a \leq 3} (u_a^6)^{\frac{\Delta_{k_a}}{2}} \rightarrow (v_{11}^6)^{h_3} (v_{12}^6)^{h_4} (v_{23}^6)^{h_5} L_6 \prod_{1 \leq a \leq 3} (u_a^6)^{\frac{\Delta_{k_a}}{2}},$$

$$\begin{aligned} u_1^6 &\rightarrow \frac{u_2^6}{v_{12}^6}, & u_2^6 &\rightarrow \frac{u_1^6}{v_{11}^6}, & u_3^6 &\rightarrow \frac{u_3^6}{v_{23}^6}, \\ v_{11}^6 &\rightarrow \frac{1}{v_{12}^6}, & v_{12}^6 &\rightarrow \frac{1}{v_{11}^6}, & v_{13}^6 &\rightarrow \frac{v_{33}^6}{v_{12}^6 v_{23}^6}, \\ v_{22}^6 &\rightarrow \frac{v_{22}^6}{v_{11}^6 v_{12}^6}, & v_{23}^6 &\rightarrow \frac{1}{v_{23}^6}, & v_{33}^6 &\rightarrow \frac{v_{13}^6}{v_{11}^6 v_{23}^6}, \end{aligned}$$

⁵The homomorphism from $D_3 \simeq S_3$ to the automorphism group of $(\mathbb{Z}_2)^3 = \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$, $\text{Aut}[(\mathbb{Z}_2)^3] = S_3$, associated to the semi-direct nature of the product is simply given by S_3 permutations of the three \mathbb{Z}_2 factors.

and that observation translates into the identity

$$\begin{aligned}
 & G_{6|\text{snowflake}}^{(d,h_2,h_3,h_4,h_5;p_2,p_3,p_4,p_5,p_6)}(u_1^6, u_2^6, u_3^6; v_{11}^6, v_{12}^6, v_{13}^6, v_{22}^6, v_{23}^6, v_{33}^6) \\
 &= (v_{11}^6)^{h_3} (v_{12}^6)^{h_4} (v_{23}^6)^{h_5} \\
 &\quad \times G_{6|\text{snowflake}}^{(d,h_2,h_4,h_3,h_5;p_3-h_2,p_2+h_2,p_5,p_4,p_6)} \left(\frac{u_2^6}{v_{12}^6}, \frac{u_1^6}{v_{11}^6}, \frac{u_3^6}{v_{23}^6}; \frac{1}{v_{12}^6}, \frac{1}{v_{11}^6}, \frac{v_{33}^6}{v_{12}^6 v_{23}^6}, \frac{v_{22}^6}{v_{11}^6 v_{12}^6}, \frac{1}{v_{23}^6}, \frac{v_{13}^6}{v_{11}^6 v_{23}^6} \right).
 \end{aligned} \tag{3.3}$$

Again, from the decomposition (2.6) with (2.15) and (2.16), we remark that F_6 in the form (C.2) does not change under this reflection generator, implying an identity for C_6 .

Finally, the generator $\mathcal{O}_{i_2}(\eta_2) \leftrightarrow \mathcal{O}_{i_3}(\eta_3)$ for dendrite permutations lead to

$$\begin{aligned}
 L_6 \prod_{1 \leq a \leq 3} (u_a^6)^{\frac{\Delta_{k_a}}{2}} &\rightarrow (v_{11}^6)^{h_2-h_4} (v_{22}^6)^{h_4} L_6 \prod_{1 \leq a \leq 3} (u_a^6)^{\frac{\Delta_{k_a}}{2}}, \\
 u_1^6 &\rightarrow \frac{u_1^6}{v_{11}^6}, & u_2^6 &\rightarrow \frac{u_2^6}{v_{22}^6}, & u_3^6 &\rightarrow u_3^6 v_{11}^6, \\
 v_{11}^6 &\rightarrow \frac{1}{v_{11}^6}, & v_{12}^6 &\rightarrow \frac{v_{11}^6 v_{12}^6}{v_{22}^6}, & v_{13}^6 &\rightarrow v_{23}^6, \\
 v_{22}^6 &\rightarrow \frac{1}{v_{22}^6}, & v_{23}^6 &\rightarrow v_{13}^6, & v_{33}^6 &\rightarrow \frac{v_{11}^6 v_{33}^6}{v_{22}^6},
 \end{aligned}$$

for the legs and conformal cross-ratios (2.14). Thus, the corresponding identity is

$$\begin{aligned}
 & G_{6|\text{snowflake}}^{(d,h_2,h_3,h_4,h_5;p_2,p_3,p_4,p_5,p_6)}(u_1^6, u_2^6, u_3^6; v_{11}^6, v_{12}^6, v_{13}^6, v_{22}^6, v_{23}^6, v_{33}^6) \\
 &= (v_{11}^6)^{h_2-h_4} (v_{22}^6)^{h_4} \\
 &\quad \times G_{6|\text{snowflake}}^{(d,h_2,-p_2-h_3,h_4,h_5;p_2,p_3,p_4,p_5,p_6)} \left(\frac{u_1^6}{v_{11}^6}, \frac{u_2^6}{v_{22}^6}, u_3^6 v_{11}^6; \frac{1}{v_{11}^6}, \frac{v_{11}^6 v_{12}^6}{v_{22}^6}, v_{23}^6, \frac{1}{v_{22}^6}, v_{13}^6, \frac{v_{11}^6 v_{33}^6}{v_{22}^6} \right).
 \end{aligned} \tag{3.4}$$

Once again, the decomposition (2.6) with (2.15) and (2.16) shows that F_6 (2.16) is invariant under this generator for dendrite permutations, resulting in a second identity for C_6 .

Therefore, the three symmetry transformations of figure 5 generate the symmetry group $(\mathbb{Z}_2)^3 \times D_3$ of order 48 (see appendix C). Each generator has an associated identity for the scalar six-point conformal blocks (2.6) in the snowflake channel, with (3.2), (3.3), and (3.4) being the identities for rotations, reflections, and dendrite permutations, respectively. From these three identities, it is straightforward to generate the remaining 45 identities of the snowflake symmetry group by composition. The proofs that our explicit solution (2.15) and (2.16) satisfies these symmetry transformations can be found in appendix C.

We now turn to the OPE limit and the limit of unit operator. Since we have already demonstrated that the scalar six-point conformal blocks in the snowflake channel obey several identities originating from the symmetry group of the snowflake diagram, it is only necessary to check the two limits once, all the other cases are equivalent by symmetry.

3.2 OPE limit

The OPE limit is defined as having two embedding space coordinates coincide. The two embedding space coordinates must correspond to an OPE in the associated topology. In this limit, the original M -point correlation function reduces to the proper $(M - 1)$ -point correlation function with a pre-factor originating from the OPE (2.1), as dictated by (2.7) for the scalar case.

For scalar six-point correlation functions in the snowflake channel depicted in figure 3, the possible OPE limits are $\eta_2 \rightarrow \eta_3$, $\eta_4 \rightarrow \eta_5$, and $\eta_6 \rightarrow \eta_1$. However, since I_6 is invariant under rotations as discussed above, it is only necessary to assess the behavior of I_6 in one OPE limit. Here, we check that in the limit $\eta_2 \rightarrow \eta_3$, we have

$$I_{6(\Delta_{k_1}, \Delta_{k_2}, \Delta_{k_3})}^{(\Delta_{i_2}, \Delta_{i_3}, \Delta_{i_4}, \Delta_{i_5}, \Delta_{i_6}, \Delta_{i_1})} \Big|_{\text{snowflake}} \rightarrow \frac{1}{\eta_{23}^{\frac{1}{2}(\Delta_2 + \Delta_3 - \Delta_{k_1})}} I_{5(\Delta_{k_2}, \Delta_{k_3})}^{(\Delta_{i_5}, \Delta_{i_4}, \Delta_{i_3}, \Delta_{i_6}, \Delta_{i_1})} \Big|_{\text{comb}}. \quad (3.5)$$

For this proof, we start from the alternative form of the scalar six-point correlation functions in the snowflake channel given in appendix B, which must lead in the OPE limit (3.5) to the scalar five-point correlation functions in the comb channel of [43] discussed in appendix A. This choice is of no consequence since we prove that these are equal to the results of section 2 in appendices B and A, respectively [see (2.14), (2.15), (2.16), and (2.17) for scalar six-point correlation functions in the snowflake channel and (2.13) for scalar five-point correlation functions in the comb channel].

In the OPE limit (3.5), we have

$$L_6^* \prod_{1 \leq a \leq 3} (u_a^{*6})^{\frac{\Delta_{k_a}}{2}} \rightarrow \frac{1}{(\eta_{23})^{\frac{1}{2}(\Delta_2 + \Delta_3 - \Delta_{k_1})}} (v_{34}^P)^{h_5} L_5^P \prod_{1 \leq a \leq 2} (u_a^P)^{\frac{\Delta_{k_a}}{2}},$$

$$u_1^{*6} \rightarrow 0, \quad u_2^{*6} \rightarrow u_1^P, \quad u_3^{*6} \rightarrow \frac{u_2^P}{v_{34}^P},$$

$$v_{1a}^{*6} \rightarrow 1 \quad (1 \leq a \leq 3),$$

$$v_{22}^{*6} \rightarrow v_{23}^P, \quad v_{23}^{*6} \rightarrow \frac{v_{24}^P}{v_{34}^P}, \quad v_{33}^{*6} \rightarrow \frac{1}{v_{34}^P},$$

where the conformal dimensions on the r.h.s. are the ones relevant for the five-point correlation functions, i.e.

$$p_3 - h_2 + h_4 \rightarrow p_3, \quad p_3 \rightarrow \bar{p}_3 + \bar{h}_3, \quad p_2 + h_2 \rightarrow p_4, \quad h_5 \rightarrow h_4, \quad h_2 \rightarrow h_3,$$

$$\bar{p}_3 + \bar{h}_2 + h_5 \rightarrow \bar{p}_4 + \bar{h}_4, \quad \bar{p}_3 + h_2 \rightarrow \bar{p}_4 + \bar{h}_3, \quad h_4 \rightarrow -p_2 - h_2, \quad p_3 - h_2 \rightarrow \bar{p}_3 + h_2.$$

Thus, the OPE limit (3.5) corresponds to the identity

$$(v_{34}^P)^{h_5} G_6^* = G_5^P,$$

with the appropriate changes for the vectors \mathbf{h} and \mathbf{p} .

Since

$$F_6^* = F_6 \rightarrow \frac{(p_2 + h_2 - m_2)_{m_2}}{(-h_2)_{-m_3}} {}_3F_2 \left[\begin{matrix} -m_2, -m_3, \bar{p}_3 - d/2 \\ p_2 + h_2 - m_2, -h_2 - m_3 \end{matrix}; 1 \right] = \frac{(p_2 + h_2 - m_2)_{m_2}}{(-h_2)_{-m_3}} F_5^P,$$

in the OPE limit (3.5), we have

$$\begin{aligned}
 (v_{34}^P)^{h_5} G_6^* &= \sum \frac{(p_3)_{m_2+m_3+m_{23}+m_{33}} (-h_2 - m_3)_{m_2+m_{22}} (-h_5)_{m_3+m_{23}+m_{33}} (\bar{p}_3 + h_2 + h_5)_{m_3}}{(\bar{p}_3 + h_2)_{2m_3+m_{23}+m_{33}} (\bar{p}_3 + h_2 + 1 - d/2)_{m_3}} \\
 &\times \frac{(p_2 + h_2 - m_2)_{m_3} (-h_4)_{m_2+m_{22}+m_{23}} (p_3 - h_2 + h_4)_{m_2+m_{33}}}{(p_3 - h_2)_{2m_2+m_{22}+m_{23}+m_{33}} (p_3 - h_2 + 1 - d/2)_{m_2}} \binom{m_{23}}{k_{23}} \binom{m_{33}}{k_{33}} \\
 &\times \binom{k_{23}}{m'_{24}} \binom{h_5 - m_3 - k_{23} - k_{33}}{m'_{34}} (-1)^{k_{23}+k_{33}+m'_{24}+m'_{34}} \\
 &\times \frac{(u_1^P)^{m_2}}{m_2!} \frac{(u_2^P)^{m_3}}{m_3!} \frac{(1 - v_{23}^P)^{m_{22}}}{m_{22}!} \frac{(1 - v_{24}^P)^{m'_{24}}}{m_{23}!} \frac{(1 - v_{34}^P)^{m'_{34}}}{m_{33}!} F_5^P,
 \end{aligned}$$

after expanding in the proper conformal cross-ratios u_a^P and $(1 - v_{ab}^P)$ of [43]. Here the vectors \mathbf{h} and \mathbf{p} are still the original ones. Thus to complete the proof, we need to evaluate the extra sums for which we did not explicitly write the indices of summation (to make the notation less cluttered) with the help of (A.1), and express the vectors \mathbf{h} and \mathbf{p} in terms of the five-point ones.

First, we sum over k_{22} and then k_{23} after changing the variable by $k_{23} \rightarrow k_{23} + m'_{24}$, which lead to

$$\begin{aligned}
 (v_{34}^P)^{h_5} G_6^* &= \sum \frac{(p_3)_{m_2+m_3+m_{23}+m_{33}} (-h_2 - m_3)_{m_2+m_{22}} (-h_5)_{m_3+m'_{24}+m'_{34}} (\bar{p}_3 + h_2 + h_5)_{m_3}}{(\bar{p}_3 + h_2)_{2m_3+m_{23}+m_{33}} (\bar{p}_3 + h_2 + 1 - d/2)_{m_3}} \\
 &\times \frac{(p_2 + h_2 - m_2)_{m_3} (-h_4)_{m_2+m_{22}+m_{23}} (p_3 - h_2 + h_4)_{m_2+m_{33}}}{(p_3 - h_2)_{2m_2+m_{22}+m_{23}+m_{33}} (p_3 - h_2 + 1 - d/2)_{m_2}} \frac{(-m'_{34})_{m_{23}+m_{33}-m'_{24}}}{m_{33}!(m_{23}-m'_{24})!} \\
 &\times \frac{(u_1^P)^{m_2}}{m_2!} \frac{(u_2^P)^{m_3}}{m_3!} \frac{(1 - v_{23}^P)^{m_{22}}}{m_{22}!} \frac{(1 - v_{24}^P)^{m'_{24}}}{m'_{24}!} \frac{(1 - v_{34}^P)^{m'_{34}}}{m'_{34}!} F_5^P.
 \end{aligned}$$

We then shift m_{23} by $m_{23} \rightarrow m_{23} + m'_{24}$, and then redefine $m_{33} = m - m_{23}$. With these changes, we can compute the sums over m_{23} and then m , which give

$$\begin{aligned}
 (v_{34}^P)^{h_5} G_6^* &= \sum \frac{(p_3)_{m_2+m_3+m'_{24}} (-h_2)_{m_2-m_3+m_{22}} (-h_5)_{m_3+m'_{24}+m'_{34}} (\bar{p}_3 + h_2 + h_5)_{m_3}}{(-h_2)_{-m_3} (\bar{p}_3 + h_2)_{2m_3+m'_{24}+m'_{34}} (\bar{p}_3 + h_2 + 1 - d/2)_{m_3}} \\
 &\times \frac{(p_2 + h_2)_{-m_2+m_3+m'_{34}} (-h_4)_{m_2+m_{22}+m'_{24}} (p_3 - h_2 + h_4)_{m_2}}{(p_2 + h_2)_{-m_2} (p_3 - h_2)_{2m_2+m_{22}+m'_{24}} (p_3 - h_2 + 1 - d/2)_{m_2}} \\
 &\times \frac{(u_1^P)^{m_2}}{m_2!} \frac{(u_2^P)^{m_3}}{m_3!} \frac{(1 - v_{23}^P)^{m_{22}}}{m_{22}!} \frac{(1 - v_{24}^P)^{m'_{24}}}{m'_{24}!} \frac{(1 - v_{34}^P)^{m'_{34}}}{m'_{34}!} F_5^P.
 \end{aligned}$$

Finally, changing the vectors \mathbf{h} and \mathbf{p} by their five-point counterparts and renaming $m_2 \rightarrow m_1$, $m_3 \rightarrow m_1$, $m_{22} \rightarrow m_{23}$, $m'_{24} \rightarrow m_{24}$, and $m'_{34} \rightarrow m_{34}$, we have

$$F_5^P = {}_3F_2 \left[\begin{matrix} -m_2, -m_3, \bar{p}_3 - d/2 \\ p_2 + h_2 - m_2, -h_2 - m_3 \end{matrix} ; 1 \right] \rightarrow {}_3F_2 \left[\begin{matrix} -m_1, -m_2, \bar{p}_4 + h_2 - d/2 \\ p_4 - m_1, -h_3 - m_2 \end{matrix} ; 1 \right],$$

as in (A.4) (hence its name), which indeed proves that $(v_{34}^P)^{h_5} G_6^* = G_5^P$, as expected.

3.3 Limit of unit operator

The limit of unit operator is defined by setting one external operator to the identity operator. In this limit, a M -point correlation function directly becomes the corresponding $(M - 1)$ -point correlation function.

For scalar six-point correlation functions in the snowflake channel, the symmetry properties of I_6 liberate us to verify the limit of unit operator for just one quasi-primary operator. We choose this quasi-primary operator to be $\mathcal{O}_{i_6}(\eta_6) \rightarrow \mathbb{1}$, for which $\Delta_{i_6} = 0$, $\Delta_{k_3} = \Delta_{i_1}$, and

$$I_6^{(\Delta_{i_2}, \Delta_{i_3}, \Delta_{i_4}, \Delta_{i_5}, \Delta_{i_6}, \Delta_{i_1})} \Big|_{\text{snowflake}} \rightarrow I_5^{(\Delta_{i_2}, \Delta_{i_3}, \Delta_{i_1}, \Delta_{i_4}, \Delta_{i_5})} \Big|_{\text{comb}}. \quad (3.6)$$

Since $h_5 = p_6 = 0$ and the remaining vector elements of \mathbf{h} and \mathbf{p} (2.17) translate directly into their five-point counterparts (2.13) in the limit of unit operator (3.6), the sums over m_3 , m_{13} , m_{23} , and m_{33} in (2.15) are trivial due to the Pochhammer symbol $(-h_5)_{m_3+m_{13}+m_{23}+m_{33}}$, forcing $m_3 = m_{13} = m_{23} = m_{33} = 0$. Therefore, the conformal cross-ratios u_3^6 , v_{13}^6 , v_{23}^6 , and v_{33}^6 disappear in the limit (3.6).

The remaining conformal cross-ratios (2.14) relate to the conformal cross-ratios (2.13) as in

$$u_a^6 \rightarrow u_a^5 \quad (1 \leq a \leq 2), \quad v_{ab}^6 \rightarrow v_{ab}^5 \quad (1 \leq a \leq b \leq 2),$$

and thus

$$L_6 \prod_{1 \leq a \leq 3} (u_a^6)^{\frac{\Delta_{k_a}}{2}} \rightarrow L_5 \prod_{1 \leq a \leq 2} (u_a^5)^{\frac{\Delta_{k_a}}{2}}.$$

These observations imply that $G_6 \rightarrow G_5$ in the limit of unit operator (3.6), which is straightforward to verify since $m_3 = m_{13} = m_{23} = m_{33} = 0$.

4 Discussion and conclusion

The field of research on d -dimensional higher-point correlation functions in CFT is a relatively uncharted territory. Although completely determined by conformal invariance, higher-point conformal blocks are notoriously difficult to compute in all generality. Moreover, there exists several inequivalent topologies for higher-point correlation functions, each of them having their associated set of identities originating from their topology. In this paper, we introduced all scalar six-point conformal blocks by computing the remaining topology, the scalar six-point conformal blocks in the snowflake channel.

Our results — presented in (2.14), (2.15), (2.16), and (2.17) — are obtained with the help of the embedding space OPE formalism developed in [30, 31]. They show that the embedding space OPE formalism is very powerful, leading to explicit results for any conformal higher-point correlation function of interest, after straightforward (yet somewhat tedious) re-summations of the hypergeometric type.

From the symmetry group of the snowflake diagram, we showed that scalar six-point conformal blocks in the snowflake channel have symmetry groups of order 48, larger than the hexagon symmetry group, contrary to expectations. We then showed that the result

verifies both the OPE limit (in which two embedding space coordinates coincide) and the limit of unit operator (where one external operator is set to the identity). Our snowflake result thus passes several non-trivial consistency checks, lending further credence to the embedding space OPE formalism.

Contrary to the scalar six-point conformal blocks in the comb channel, for which there are two extra sums that factorize into the product of two ${}_3F_2$ -hypergeometric functions, the scalar six-point conformal blocks in the snowflake channel have two extra sums that do not factorize. They can be written as a Kampé de Fériet function, and the snowflake invariance under rotations implies (C.1), that translates into the identities (for m_1 , m_2 , and m_3 non-negative integers and a_1 , a_2 , b_1 , b_2 , d_1 , and g_1 arbitrary)

$$\begin{aligned}
 & F_{2,1,0}^{1,3,2} \left[\begin{matrix} a_1; -m_2, b_1 + m_3, b_2 + m_1; -m_1, -m_3 \\ b_1, b_2; d_1; - \end{matrix} \middle| 1, 1 \right] \\
 &= \frac{(d_1 - a_1)_{m_2} (b_1 - a_1)_{m_3}}{(d_1)_{m_2} (b_1)_{m_3}} F_{2,1,0}^{1,3,2} \left[\begin{matrix} a_1; b_2 + m_1, 1 + a_1 - d_1, -m_3; -m_1, -m_2 \\ 1 + a_1 - d_1 - m_2, b_2; 1 + a_1 - b_1 - m_3; - \end{matrix} \middle| 1, 1 \right], \\
 & F_{1,1,1}^{2,1,1} \left[\begin{matrix} a_1, a_2; -m_2; -m_3 \\ a_2 - m_1; d_1; g_1 \end{matrix} \middle| 1, 1 \right] \\
 &= \frac{(a_2)_{-m_1} (d_1 - a_1)_{m_2}}{(a_2 - a_1)_{-m_1} (d_1)_{m_2}} F_{1,1,1}^{2,1,1} \left[\begin{matrix} a_1, 1 + a_1 - d_1; -m_1; -m_3 \\ 1 + a_1 - d_1 - m_2, 1 + a_1 - a_2; g_1 \end{matrix} \middle| 1, 1 \right],
 \end{aligned}$$

with

$$F_{2,1,0}^{1,3,2} \left[\begin{matrix} a_1; -m_2, 1+a_1-g_1, a_2; -m_1, -m_3 \\ 1+a_1-g_1-m_3, a_2-m_1; d_1; - \end{matrix} \middle| 1, 1 \right] = \frac{(g_1)_{m_3}}{(g_1-a_1)_{m_3}} F_{1,1,1}^{2,1,1} \left[\begin{matrix} a_1, a_2; -m_2; -m_3 \\ a_2-m_1; d_1; g_1 \end{matrix} \middle| 1, 1 \right],$$

which are obtained from repeated use of well-known ${}_3F_2$ -hypergeometric identities.⁶

It is obvious that the embedding space OPE formalism can be used to investigate higher-point correlation functions, including their symmetry groups. For example, starting from the scalar three-point correlation function, which has symmetry group D_3 , we conjecture that doubling the number of all external legs N times with the help of the OPE gives scalar (3×2^N) -point correlation functions with the symmetry groups $\left((\mathbb{Z}_2)^{3 \times 2^{N-1}} \times (\dots \times ((\mathbb{Z}_2)^3 \times D_3) \dots) \right)$.⁷ From the successive action of the OPE limit or the limit of unit operator, these maximally-symmetric topologies should lead to all topologies for smaller scalar higher-point correlation functions (starting from sufficiently large N). In general, starting from a specific M -point topology with symmetry group $H_{M|\text{channel}}$, this procedure of doubling the number of all external legs N times with the help of the OPE should generate $(M \times 2^N)$ -point topologies with symmetry groups $\left((\mathbb{Z}_2)^{M \times 2^{N-1}} \times (\dots \times ((\mathbb{Z}_2)^M \times H_{M|\text{channel}}) \dots) \right)$.

It is less clear what happens to diagrams with fewer symmetries. For example, scalar seven-point correlation functions with the topology resulting from scalar six-point correlation functions in the snowflake channel (called the extended snowflake channel below)

⁶Kampé de Fériet functions have not been studied as extensively as standard hypergeometric functions.

⁷Although this is only a conjecture, the order of the symmetry groups should be correct.

should have symmetry group of order sixteen only, smaller than for the snowflake diagram but larger than the scalar seven-point correlation functions in the comb channel. In fact, their symmetry group should be $\mathbb{Z}_2 \times ((\mathbb{Z}_2)^2 \rtimes \mathbb{Z}_2)$, the direct product of the symmetry group for the extra dendrite permutations and the symmetry group of the six-point comb diagram from which it can also be built. The symmetries are relatively easy to enumerate for a specific topology, but it is not clear how to write general expressions for the orders of the symmetry groups. Nevertheless, studying the cosets $S_M/H_{M|\text{channel}}$, with S_M the symmetric group of M elements, leads to interesting consequences.

For example, two- and three-point correlation functions have symmetry groups \mathbb{Z}_2 and D_3 , respectively, while four- and five-point correlation functions have symmetry groups $(\mathbb{Z}_2)^2 \rtimes \mathbb{Z}_2$ (the topologies are unique for fewer than six points). Since $\mathbb{Z}_2 \simeq S_2$ and $D_3 \simeq S_3$, the symmetry groups of two- and three-point conformal blocks correspond to the symmetry groups of the full two- and three-point correlation functions, and their cosets are trivial with only one element. This observation implies that there is no non-trivial information that can be derived from bootstrap considerations starting with two- and three-point correlation functions, as expected. This is not the case for $(M > 3)$ -point correlation functions. Indeed, for four-point correlation functions, the associated coset is $S_4/[(\mathbb{Z}_2)^2 \rtimes \mathbb{Z}_2]$ which has cardinality three. There are thus three elements, corresponding to the well-known s -, t -, and u -channels. For five-point correlation functions, the coset is $S_5/[(\mathbb{Z}_2)^2 \rtimes \mathbb{Z}_2]$ with cardinality fifteen, hence there should be fifteen different ways of expressing five-point correlation functions. For six-point correlation functions, the number of different expressions should depend on the channels. For the comb channel, there should be $|S_6/[(\mathbb{Z}_2)^2 \rtimes \mathbb{Z}_2]| = 90$ different expressions while for the snowflake channel there should only be $|S_6/[(\mathbb{Z}_2)^3 \rtimes D_6]| = 15$, for a total of 105. In the case of the seven-point conformal bootstrap, the comb channel and the extended snowflake channel should have coset cardinalities $|S_7/[(\mathbb{Z}_2)^2 \rtimes \mathbb{Z}_2]| = 630$ and $|S_7/[\mathbb{Z}_2 \times ((\mathbb{Z}_2)^2 \rtimes \mathbb{Z}_2)]| = 315$, respectively. Hence, there should be a total of 945 different ways of expressing seven-point correlation functions.

These numbers can also be understood from the counting of topologies. Each M -point topology is represented by an unrooted binary tree with $2M - 2$ nodes, where M nodes represent the external quasi-primary operators (vertices of degree one, which are the leaves of the tree) and $M - 2$ nodes correspond to the OPEs (vertices of degree three). It is transparent that unrooted binary trees with $2M - 2$ nodes have a total of $2M - 3$ edges, with M edges connecting the OPE nodes to the leaves and $M - 3$ edges representing the exchanged quasi-primary operators. The number of M -point topologies is thus given by the number of unrooted binary trees with M unlabeled leaves, denoted here by $T_0(M)$. Unfortunately, there is no simple closed-form formula for $T_0(M)$, but the first few integers in the sequence are $(1, 1, 1, 1, 2, 2, 4, 6, 11, \dots)$, starting at $M = 2$.⁸ On the contrary, the number of different ways of expressing the same full M -point correlation functions is given by the number of unrooted binary trees with M labeled leaves, denoted here by $T(M)$, which is given by $T(M) = (2M - 5)!!$. Comparing the symmetry groups and the number

⁸See *The On-line Encyclopedia of Integer Sequences* at <https://oeis.org/A000672> and <https://oeis.org/A129860> for more details.

of topologies, we thus conclude that we should have the following identity for M -point correlation functions,

$$\sum_{1 \leq i \leq T_0(M)} \frac{1}{|H_{M|\text{channel } i}|} = \frac{T(M)}{|S_M|} = \frac{(2M - 5)!!}{M!}.$$

This identity is in agreement with the partial results obtained above with M up to seven and we verified it for $M = 8, 9$ as well. It can be used to explore the orders of the symmetry groups of the different topologies. Moreover, the previous discussion implies that the number of ways of writing the same M -point correlation function is $T(M)$, although redundancies lead to a number of independent bootstrap equations which is much smaller, to which we now turn.

It is well known that the conformal bootstrap program can benefit from the knowledge of higher-point correlation functions. Indeed, it has been argued that bootstrapping higher-point correlation functions with external quasi-primary operators in the trivial representation is equivalent to the usual full conformal bootstrap of four-point correlation functions.⁹ We conjecture that the study of the symmetry groups of the different topologies of M -point conformal blocks also leads to interesting insights on the conformal bootstrap. Once again, let us denote the symmetry group of the scalar M -point conformal blocks with some specific topology as $H_{M|\text{channel}}$ and the symmetry group of the full M -point correlation functions, including the contributions of the non-trivial representations, as S_M (the symmetric group of M elements). As discussed above, the former depends on the topology of the particular channel under consideration while the latter corresponds to all the possible ways of re-arranging the M external quasi-primary operators in the full correlation function. We observe that the analysis of the symmetry groups $H_{M|\text{channel}}$ gives an intuitive picture of the M -point conformal bootstrap.

As already mentioned, there are $T(M) = (2M - 5)!!$ ways of writing the same full M -point correlation function. One can therefore write a total of $(2M - 5)!! - 1$ *a priori* independent equations for the M -point conformal bootstrap, up to redundancies implied by symmetries of different channels. In fact, it can be shown that most of the equations are redundant and one can choose specific equations to minimize the overall number of required conformal bootstrap equations. The smallest number of independent bootstrap equations N_B is equal to the greater of 1 and $T_0(M) - 1$. Namely, with a unique topology ($M = 4, 5$) or with two topologies ($M = 6, 7$) it is sufficient to have only one bootstrap equation, while with more than two topologies, the smallest number of bootstrap equations is equal to the number of topologies minus one.

This counting statement is easy to prove using the symmetries of different topologies. Let us start with the well-known case of four points. The only required bootstrap equation is the equality of the s - and t -channels illustrated in figure 6. The two diagrams identified by the bootstrap equality have the same symmetries $H_4 = (\mathbb{Z}_2)^2 \rtimes \mathbb{Z}_2$ since they belong to the same topology, but these symmetries are embedded differently in the full permutation group that acts on different operators in the two channels. In the s -channel, the non-trivial

⁹This statement must be modified accordingly when there are fermions in the theory.




Figure 6. Four-point bootstrap equation that is equality between the s - and t -channels.




Figure 7. Bootstrap equations for even M (top) and odd M (bottom) that generate the full permutation group S_M . For better readability, we denoted the external operators just by their subscripts, for example 1 instead of \mathcal{O}_{i_1} .

elements of the three \mathbb{Z}_2 's in $(\mathbb{Z}_2)^2 \rtimes \mathbb{Z}_2$ are σ_{12} , σ_{34} , and $\sigma_{13}\sigma_{24}$, where σ_{ab} denotes an exchange of external operators \mathcal{O}_{i_a} and \mathcal{O}_{i_b} . The full H_4 symmetry is obtained by multiplying these elements. By comparison, the non-trivial elements of the t -channel symmetry group are σ_{14} , σ_{23} , and $\sigma_{12}\sigma_{34}$. Equating the s - and t -channels allows one to use the symmetries of both diagrams to get all other bootstrap equations related by symmetry. After all, satisfying the equation means that all equations related by either set of symmetries of the diagram on the right-hand side or the left-hand side are also satisfied. Picking the following symmetries σ_{12} , σ_{23} , and σ_{34} from both channels, one generates the full permutation group S_4 since these choices comprise the generators of S_4 . Once we have the full permutation group, it is clear that one of these permutations includes the u -channel and there is no independent equation equating the u -channel to either the s - or t -channels. In other words, once the bootstrap equations between the s - and the t -channels are satisfied for all quasi-primary operators, then the bootstrap equations for all three channels are satisfied, therefore the u -channel is redundant.

Analogous use of symmetries allows us to prove the statement on the minimal number of independent bootstrap equations for any M . The illustration for even $M \geq 6$ and odd $M \geq 9$ is shown in figure 7. In both the even and odd cases we start with a single bootstrap equation that identifies two channels of different topologies. By the assignment of the external operators, the diagrams in the bootstrap equation contain $\sigma_{12}, \sigma_{23}, \dots, \sigma_{M-1, M}$ (as well as other symmetries that are unimportant for the argument) that generate the full permutation group S_M . Thus, any other diagram with different operator assignment of either of the two channels can be obtained by symmetries and automatically satisfies the bootstrap equations. This includes equalities between diagrams of the two topologies




Figure 8. Bootstrap equations for five points (top) and seven points (bottom) that generate the full permutation groups S_5 and S_7 , respectively. For better readability, we denoted the external operators just by their subscripts, for example 1 instead of \mathcal{O}_{i_1} .

chosen in figure 7 and, by the transitive property, also all equalities between diagrams of the same topology — either the one on the left or on the right of the equation. Diagrams in topologies that are not present in figure 7 are independent and each additional topology must be included in the set of bootstrap equations. However, since the original equation in figure 7 guarantees equality of all permutations, any operator assignment for the additional topologies will do. All other assignments will be equivalent by the action of S_M . Hence the smallest number of independent bootstrap equations is $T_0(M) - 1$.

One needs to consider $M = 5$, where there is only one topology, and $M = 7$ with two topologies, separately. This is because the choice of topologies for odd M in the bottom drawing in figure 7 requires topologies that are not present for $M = 5$ or identical for $M = 7$. Suitable equations for these cases are illustrated in figure 8. It is straightforward to check that the symmetries of these diagrams generate the full permutation groups S_5 and S_7 , accordingly. For $M = 5$ the \mathbb{Z}_2 's of the dendrite permutations obviously generate S_5 . For $M = 7$ the dendrite permutations generate S_6 acting on the set $\{1, 2, \dots, 6\}$. Additionally, the left-right reflection symmetry of the $M = 7$ comb diagram is realized by $\sigma_{67}\sigma_{34}\sigma_{25}$. This element can be combined with permutations contained in S_6 to yield σ_{67} , therefore generating the full permutation group S_7 . This completes the argument that one can satisfy all possible M -point bootstrap equations by suitably choosing only $N_B = \max\{1, T_0(M) - 1\}$ independent equations.

Coming back to our main result, it is unclear for now what type of generalizations occurs for the extra sums, encoded here in our function F_M . In the comb channel, we argued in [44] that the $M - 4$ extra sums appearing in scalar M -point correlation functions were necessary for the limit of unit operator to make sense. For $M = 6$, this argument cannot be used for the snowflake diagram. Nevertheless, (2.16) shows that in the snowflake channel, the number of extra sums is the same, although these extra sums do not factorize as in the comb channel. Are there three extra sums that do not factorize for the non-comb topology of the scalar seven-point correlation functions or does the number of sums depend on the topology?

An interesting avenue of research is to initiate the computation of higher-point correlation functions with spins, either for external quasi-primary operators or internal quasi-

primary operators, or both. Since the embedding space OPE formalism developed in [30, 31] treats all irreducible representations of the Lorentz group on the same footing, such computations should be feasible.

Finally, higher-point correlation functions can also be of use in the AdS/CFT correspondence. Indeed, scalar six-point conformal blocks in the snowflake channel correspond to some geodesic Witten diagrams (see for example [42]) and their knowledge might elucidate some of the kinematics in AdS.

Acknowledgments

The authors would like to thank Valentina Prilepina for useful discussions. The work of JFF is supported by NSERC and FRQNT. The work of WJM is supported by the China Scholarship Council and in part by NSERC and FRQNT.

A Scalar five-point conformal blocks and the OPE

The scalar higher-point correlation functions in the comb channel were obtained very recently in [43, 44]. In this appendix, we show how to obtain the scalar five-point conformal blocks of [43] from the OPE approach of [44]. The proof is a straightforward application of the re-summation formula

$${}_2F_1 \left[\begin{matrix} -n, b \\ c \end{matrix}; 1 \right] = \frac{(c-b)_n}{(c)_n}, \quad (\text{A.1})$$

for n a non-negative integer. Other useful identities used in the proof are the binomial identity

$$(1-v)^{a+b} = \sum_{i \geq 0} (-1)^i \binom{a+b}{i} v^i = \sum_{i, j \geq 0} (-1)^{i+j} \binom{a}{i} \binom{b}{j} v^{i+j}, \quad (\text{A.2})$$

and

$${}_3F_2 \left[\begin{matrix} -n, b, c \\ d, e \end{matrix}; 1 \right] = \frac{(d-b)_n}{(d)_n} {}_3F_2 \left[\begin{matrix} -n, b, e-c \\ b-d-n+1, e \end{matrix}; 1 \right], \quad (\text{A.3})$$

$${}_3F_2 \left[\begin{matrix} -n, b, c \\ d, 1+b+c-d-n \end{matrix}; 1 \right] = \frac{(d-b)_n (d-c)_n}{(d)_n (d-b-c)_n}.$$

The scalar five-point conformal blocks (2.13), which is our starting point to compute the scalar six-point conformal blocks in the snowflake channel, can be obtained similarly.

A.1 Proof of the equivalence

The comb channel of [43] is depicted in figure 9. To get figure 9, we need to shift the quasi-primary operators in [43] such that $\mathcal{O}_{i_a}(\eta_a) \rightarrow \mathcal{O}_{i_{a+1}}(\eta_{a+1})$ with $\mathcal{O}_{i_6}(\eta_6) \equiv \mathcal{O}_{i_1}(\eta_1)$. Using (2.7) with $k = 2$, $l = 3$ and $m = 2$ on the scalar four-point correlation functions




Figure 9. Scalar five-point conformal blocks of [43].

of [44], we obtain, after expanding with the binomial identity (A.2),

$$I_5 = L_5 \left(\frac{\eta_{23}\eta_{45}}{\eta_{24}\eta_{35}} \right)^{\frac{\Delta_{k_1}}{2}} \left(\frac{\eta_{34}\eta_{15}}{\eta_{14}\eta_{35}} \right)^{\frac{\Delta_{k_2}}{2}} \left(\frac{\eta_{14}\eta_{25}}{\eta_{12}\eta_{45}} \right)^{\bar{p}_4 + \bar{h}_4} \\ \times \sum C_4 K_5 \binom{n_{11}}{s_2} \frac{(-1)^{s_2}}{n_1! n_{11}!} \left(\frac{\eta_{23}\eta_{45}}{\eta_{24}\eta_{35}} \right)^{n_1} \left(\frac{\eta_{25}\eta_{34}}{\eta_{35}\eta_{24}} \right)^{s_2},$$

where the legs L_5 and the vectors \mathbf{h} and \mathbf{p} are defined in [44]. For notational simplicity, we omit the indices of summation n_1 , n_{11} , and s_2 on the sum. Here C_4 and K_5 are given by the analog of (2.12) found in [44] and (2.4), respectively, i.e.

$$C_4 = \frac{(-h_3)_{n_1} (p_2 + h_2)_{n_1} (\bar{p}_3 + \bar{h}_3)_{n_1 + n_{11}} (p_3)_{n_1 + n_{11}}}{(\bar{p}_3 + h_2)_{2n_1 + n_{11}} (\bar{p}_3 + h_2 + 1 - d/2)_{n_1}}, \\ K_5 = \sum_{\{r_a, r_{2a}, r_{ab}\} \geq 0} \frac{(-h_4)_{\bar{r}_2 + \bar{r}} (-s_2)_{\bar{r}_2} (\bar{p}_4 + \bar{h}_4)_{\bar{r} - \bar{r}} (p_4 - n_1)_{r_4} (\bar{p}_3 + \bar{h}_3 + n_1 + s_2)_{r_3}}{(\bar{p}_4 + \bar{h}_3)_{\bar{r} + \bar{r}_2} (\bar{p}_4 + \bar{h}_3 + 1 - d/2)_{\bar{r}_2 + \bar{r}} (r_4 - r_{24} - \bar{r}_4)! (r_3 - r_{23} - \bar{r}_3)!} \\ \times (x_2^5)^{\bar{r}_2 + \bar{r}} (y_3^5)^{r_3 - r_{23} - \bar{r}_3} (y_4^5)^{r_4 - r_{24} - \bar{r}_4} \frac{(z_{24}^5)^{r_{24}} (z_{34}^5)^{r_{34}}}{(r_{23})! (r_{24})! (r_{34})!}.$$

The legs and the conformal cross-ratios defined in [43] are given by

$$L_5^P = \left(\frac{\eta_{13}}{\eta_{12}\eta_{23}} \right)^{\frac{\Delta_2}{2}} \left(\frac{\eta_{12}}{\eta_{13}\eta_{23}} \right)^{\frac{\Delta_3}{2}} \left(\frac{\eta_{12}}{\eta_{14}\eta_{24}} \right)^{\frac{\Delta_4}{2}} \left(\frac{\eta_{12}}{\eta_{15}\eta_{25}} \right)^{\frac{\Delta_5}{2}} \left(\frac{\eta_{25}}{\eta_{12}\eta_{15}} \right)^{\frac{\Delta_1}{2}} \\ = L_5 \left(\frac{\eta_{14}\eta_{25}}{\eta_{12}\eta_{45}} \right)^{\frac{\Delta_1 - \Delta_5}{2}} \left(\frac{\eta_{12}\eta_{34}}{\eta_{13}\eta_{24}} \right)^{\frac{\Delta_3 - \Delta_2}{2}} \left(\frac{\eta_{12}\eta_{34}\eta_{45}}{\eta_{14}\eta_{24}\eta_{35}} \right)^{\frac{\Delta_4}{2}},$$

and

$$u_1^P = \frac{\eta_{14}\eta_{23}}{\eta_{13}\eta_{24}}, \quad u_2^P = \frac{\eta_{15}\eta_{24}}{\eta_{14}\eta_{25}}, \quad v_{23}^P = \frac{\eta_{12}\eta_{34}}{\eta_{13}\eta_{24}}, \quad v_{24}^P = \frac{\eta_{12}\eta_{35}}{\eta_{13}\eta_{25}}, \quad v_{34}^P = \frac{\eta_{12}\eta_{45}}{\eta_{14}\eta_{25}}.$$

Hence I_5 becomes

$$I_5 = L_5^P (v_{23}^P)^{p_3 + h_3} (v_{24}^P)^{-\bar{p}_3 - \bar{h}_3} (v_{34}^P)^{-p_4} (u_1^P)^{\frac{\Delta_{k_1}}{2}} (u_2^P)^{\frac{\Delta_{k_2}}{2}} \\ \times \sum C_4 K_5 \binom{n_{11}}{s_2} \frac{(-1)^{s_2}}{n_1! n_{11}!} \left(u_1^P \frac{v_{34}^P}{v_{24}^P} \right)^{n_1} \left(\frac{v_{23}^P}{v_{24}^P} \right)^{s_2}.$$

With the help of the following identities [see (2.3)]

$$x_2^5 = \frac{u_1^P u_2^P}{v_{24}^P}, \quad 1 - y_3^5 = \frac{1}{v_{24}^P}, \quad 1 - y_4^5 = \frac{1}{v_{34}^P}, \quad z_{24}^5 = \frac{v_{24}^P}{u_1^P v_{34}^P}, \quad z_{34}^5 = \frac{v_{23}^P}{u_1^P v_{34}^P},$$

we can express K_5 in terms of the conformal cross-ratios of [43], leading to

$$\begin{aligned}
 I_5 &= L_5^P (u_1^P)^{\frac{\Delta_{k_1}}{2}} (u_2^P)^{\frac{\Delta_{k_2}}{2}} \sum C_4 \binom{n_{11}}{s_2} \frac{(-1)^{s_2}}{n_1! n_{11}!} \left(1 - \frac{1}{v_{24}^P}\right)^{\sigma_3} \left(1 - \frac{1}{v_{34}^P}\right)^{\sigma_4} \\
 &\times (u_1^P)^{n_1+r_{23}} (u_2^P)^{m_2} (v_{23}^P)^{p_3+h_3+m_2+r_{34}} (v_{24}^P)^{-\bar{p}_3-\bar{h}_3-n_1-s_2-r_{23}-r_{34}} (v_{34}^P)^{n_1-p_4-m_2+r_{23}} \\
 &\times \frac{(-h_4)_{m_2} (-s_2)_{m_2-r_{34}} (\bar{p}_4 + \bar{h}_4)_{m_2+\sigma_3+\sigma_4}}{(\bar{p}_4 + \bar{h}_3)_{2m_2+\sigma_3+\sigma_4} (\bar{p}_4 + \bar{h}_3 + 1 - d/2)_{m_2}} \\
 &\times \frac{(p_4 - n_1)_{\sigma_4+m_2-r_{23}} (\bar{p}_3 + \bar{h}_3 + n_1 + s_2)_{\sigma_3+r_{23}+r_{34}}}{\sigma_3! \sigma_4! r_{23}! r_{24}! r_{34}!}.
 \end{aligned}$$

Thus, by combining the powers of the conformal cross-ratios, G_5^P (2.6) is given by

$$\begin{aligned}
 G_5^P &= \sum \binom{n_{11}}{s_2} \binom{\sigma_3}{l_3} \binom{\sigma_4}{l_4} \binom{p_3+h_3+s_2+r_{34}}{m_{23}} \\
 &\times \binom{-l_3-\bar{p}_3-\bar{h}_3-n_1-s_2-r_{23}-r_{34}}{m_{24}} \binom{n_1-l_4-p_4-m_2+r_{23}}{m_{34}} \\
 &\times \frac{(-h_4)_{m_2} (-s_2)_{m_2-r_{34}} (\bar{p}_4 + \bar{h}_4)_{m_2+\sigma_3+\sigma_4}}{(\bar{p}_4 + \bar{h}_3)_{2m_2+\sigma_3+\sigma_4} (\bar{p}_4 + \bar{h}_3 + 1 - d/2)_{m_2}} \\
 &\times \frac{(p_4 - n_1)_{\sigma_4+m_2-r_{23}} (\bar{p}_3 + \bar{h}_3 + n_1 + s_2)_{\sigma_3+r_{23}+r_{34}}}{\sigma_3! \sigma_4! r_{23}! r_{24}! r_{34}!} \\
 &\times \frac{(-1)^{s_2+l_3+l_4+m_{23}+m_{24}+m_{34}} C_4}{n_1! n_{11}!} (u_1^P)^{n_1+r_{23}} (u_2^P)^{m_2} (1-v_{23}^P)^{m_{23}} (1-v_{24}^P)^{m_{24}} (1-v_{34}^P)^{m_{34}},
 \end{aligned}$$

where all the superfluous sums must be appropriately taken care of to reach the result of [43].

Since the terms involving l_4 are

$$\begin{aligned}
 (-1)^{l_4} \binom{\sigma_4}{l_4} &= \frac{(-\sigma_4)_{l_4}}{l_4!}, \\
 (-1)^{m_{34}} \binom{n_1-l_4-p_4-m_2+r_{23}}{m_{34}} &= \frac{(p_4+m_2-n_1-r_{23}+m_{34})_{l_4} (p_4+m_2-n_1-r_{23})_{m_{34}}}{(p_4+m_2-n_1-r_{23})_{l_4} m_{34}!},
 \end{aligned}$$

they can be re-summed with the help of the identity (A.1), and we find that G_5^P becomes

$$\begin{aligned}
 G_5^P &= \sum \binom{n_{11}}{s_2} \binom{\sigma_3}{l_3} \binom{p_3+h_3+s_2+r_{34}}{m_{23}} \binom{-l_3-\bar{p}_3-\bar{h}_3-n_1-s_2-m_2+r_{34}}{m_{24}} (-m_{34})_{\sigma_4} \\
 &\times \frac{(-h_4)_{m_2} (-s_2)_{m_2-r_{34}} (\bar{p}_4 + \bar{h}_4)_{m_2+\sigma_3+\sigma_4}}{(\bar{p}_4 + \bar{h}_3)_{2m_2+\sigma_3+\sigma_4} (\bar{p}_4 + \bar{h}_3 + 1 - d/2)_{m_2}} \\
 &\times \frac{(p_4 - n_1)_{m_{34}+m_2-r_{23}} (\bar{p}_3 + \bar{h}_3 + n_1 + s_2)_{\sigma_3+r_{23}+r_{34}}}{\sigma_3! \sigma_4! r_{23}! r_{24}! r_{34}!} \\
 &\times \frac{(-1)^{s_2+l_3+m_{23}+m_{24}} C_4 (u_1^P)^{n_1+r_{23}} (u_2^P)^{m_2} (1-v_{23}^P)^{m_{23}} (1-v_{24}^P)^{m_{24}} (1-v_{34}^P)^{m_{34}}}{n_1! n_{11}! m_{34}!}.
 \end{aligned}$$

Similar steps can be performed to re-sum over l_3 , σ_3 , and finally σ_4 . The result is given by

$$\begin{aligned}
 G_5^P &= \sum C_4 \binom{n_{11}}{s_2} \frac{(-1)^{s_2+m_{23}}}{n_1! n_{11}!} \binom{p_3+h_3+s_2+r_{34}}{m_{23}} \\
 &\times \frac{(-h_4)_{m_2+m_{24}+m_{34}} (-s_2)_{m_2-r_{34}} (\bar{p}_4+\bar{h}_4)_{m_2}}{(\bar{p}_4+\bar{h}_3)_{2m_2+m_{24}+m_{34}} (\bar{p}_4+\bar{h}_3+1-d/2)_{m_2}} \\
 &\times \frac{(p_4-n_1)_{m_{34}+m_2-r_{23}} (\bar{p}_3+\bar{h}_3+n_1+s_2)_{m_{24}+r_{23}+r_{34}}}{r_{23}! (m_2-r_{23}-r_{34})! r_{34}!} \\
 &\times (u_1^P)^{n_1+r_{23}} (u_2^P)^{m_2} (1-v_{23}^P)^{m_{23}} \frac{(1-v_{24}^P)^{m_{24}}}{m_{24}!} \frac{(1-v_{34}^P)^{m_{34}}}{m_{34}!}.
 \end{aligned}$$

To evaluate the sum over r_{34} , we first change the variable as $s_2 \rightarrow s_2 + m_2 - r_{34}$. Then, the sum over r_{34} can also be simplified using the hypergeometric type re-summation (A.1), which implies that G_5 can be expressed as

$$\begin{aligned}
 G_5^P &= \sum C_4 \frac{(-1)^{s_2+m_{23}}}{n_1!} \binom{p_3+h_3+s_2+m_2}{m_{23}} \frac{(\bar{p}_3+\bar{h}_3+n_1+n_{11})_{m_2-r_{23}}}{s_2! (n_{11}-s_2-r_{23})! (m_2-r_{23})!} \\
 &\times \frac{(-h_4)_{m_2+m_{24}+m_{34}} (\bar{p}_4+\bar{h}_4)_{m_2} (p_4-n_1)_{m_{34}+m_2-r_{23}} (\bar{p}_3+\bar{h}_3+n_1+s_2+m_2)_{m_{24}+r_{23}}}{(\bar{p}_4+\bar{h}_3)_{2m_2+m_{24}+m_{34}} (\bar{p}_4+\bar{h}_3+1-d/2)_{m_2} r_{23}!} \\
 &\times (u_1^P)^{n_1+r_{23}} (u_2^P)^{m_2} (1-v_{23}^P)^{m_{23}} \frac{(1-v_{24}^P)^{m_{24}}}{m_{24}!} \frac{(1-v_{34}^P)^{m_{34}}}{m_{34}!}.
 \end{aligned}$$

Using the last identity of (A.2) for the sum over m_{23} leads to

$$\begin{aligned}
 G_5^P &= \sum C_4 \frac{(-1)^{s_2+k_{23}+l}}{n_1!} \binom{p_3+h_3+m_2}{k_{23}} \binom{s_2}{l} \frac{(\bar{p}_3+\bar{h}_3+n_1+n_{11})_{m_2-r_{23}}}{s_2! (n_{11}-s_2-r_{23})! (m_2-r_{23})!} \\
 &\times \frac{(-h_4)_{m_2+m_{24}+m_{34}} (\bar{p}_4+\bar{h}_4)_{m_2} (p_4-n_1)_{m_{34}+m_2-r_{23}} (\bar{p}_3+\bar{h}_3+n_1+s_2+m_2)_{m_{24}+r_{23}}}{(\bar{p}_4+\bar{h}_3)_{2m_2+m_{24}+m_{34}} (\bar{p}_4+\bar{h}_3+1-d/2)_{m_2} r_{23}!} \\
 &\times (u_1^P)^{n_1+r_{23}} (u_2^P)^{m_2} (1-v_{23}^P)^{k_{23}+l} \frac{(1-v_{24}^P)^{m_{24}}}{m_{24}!} \frac{(1-v_{34}^P)^{m_{34}}}{m_{34}!}.
 \end{aligned}$$

We can now make a redefinition of the variable such that $s_2 \rightarrow s_2 + l$ and re-sum over s_2 using (A.1) again. Expressing C_4 in terms of Pochhammer symbols, the scalar five-point conformal blocks become

$$\begin{aligned}
 G_5^P &= \sum \frac{(-1)^{k_{23}}}{n_1!} \binom{p_3+h_3+m_2}{k_{23}} (u_1^P)^{n_1+r_{23}} (u_2^P)^{m_2} (1-v_{23}^P)^{k_{23}+l} \frac{(1-v_{24}^P)^{m_{24}}}{m_{24}!} \frac{(1-v_{34}^P)^{m_{34}}}{m_{34}!} \\
 &\times \frac{(-h_4)_{m_2+m_{24}+m_{34}} (\bar{p}_4+\bar{h}_4)_{m_2}}{(\bar{p}_4+\bar{h}_3)_{2m_2+m_{24}+m_{34}} (\bar{p}_4+\bar{h}_3+1-d/2)_{m_2}} \frac{(p_4-n_1)_{m_{34}+m_2-r_{23}}}{r_{23}!} \\
 &\times \frac{(-h_3)_{n_1} (p_2+h_2)_{n_1} (\bar{p}_3+\bar{h}_3)_{m_2+m_{24}+n_1+r_{23}+l}}{(\bar{p}_3+h_2)_{2n_1+n_{11}} (\bar{p}_3+h_2+1-d/2)_{n_1}} \frac{(p_3)_{n_1+n_{11}}}{l! (n_{11}-r_{23}-l)!} \frac{(-m_{24}-r_{23})_{n_{11}-r_{23}-l}}{(m_2-r_{23})!}.
 \end{aligned}$$

Changing variables again as in $n_{11} \rightarrow n_{11} + r_{23} + l$, we can first evaluate the summation over n_{11} using (A.1), and then the summation over l with the help of the second identity

in (A.3) after changing $k_{23} \rightarrow k_{23} - l$, and finally replace $n_1 = m_1 - r_{23}$, leading to

$$\begin{aligned}
 G_5^P &= \sum \frac{1}{(m_1 - r_{23})!} (u_1^P)^{m_1} (u_2^P)^{m_2} \frac{(1 - v_{23}^P)^{m_{23}}}{m_{23}!} \frac{(1 - v_{24}^P)^{m_{24}}}{m_{24}!} \frac{(1 - v_{34}^P)^{m_{34}}}{m_{34}!} \\
 &\times \frac{(-h_4)_{m_2 + m_{24} + m_{34}} (\bar{p}_4 + \bar{h}_4)_{m_2}}{(\bar{p}_4 + \bar{h}_3)_{2m_2 + m_{24} + m_{34}} (\bar{p}_4 + \bar{h}_3 + 1 - d/2)_{m_2}} \frac{(p_4 - m_1 + r_{23})_{m_2 + m_{34} - r_{23}}}{r_{23}!} \\
 &\times \frac{(-h_3 + m_1 - m_2)_{m_{23}} (-h_3)_{m_1 - r_{23}} (p_2 + h_2)_{m_1 + m_{24}} (\bar{p}_3 + \bar{h}_3)_{m_1 + m_2 + m_{24}} (p_3)_{m_1}}{(\bar{p}_3 + h_2)_{2m_1 + m_{23} + m_{24}} (\bar{p}_3 + h_2 + 1 - d/2)_{m_1 - r_{23}}} \frac{1}{(m_2 - r_{23})!}.
 \end{aligned}$$

At this point, we are left with only one extra sum (over r_{23}), as expected. We now use the following relations,

$$\begin{aligned}
 (p_4 - m_1 + r_{23})_{m_2 + m_{34} - r_{23}} &= \frac{(-1)^{m_1} (p_3)_{m_2 - m_1 + m_{34}} (1 - p_4)_{m_1}}{(p_4 - m_1)_{r_{23}}}, \\
 \frac{(-h_3)_{m_1 - r_{23}}}{(\bar{p}_3 + h_2 + 1 - d/2)_{m_1 - r_{23}}} &= \frac{(-h_3)_{m_1} (-\bar{p}_3 - h_2 + d/2 - m_1)_{r_{23}}}{(\bar{p}_3 + h_2 + 1 - d/2)_{m_1} (1 + h_3 - m_1)_{r_{23}}},
 \end{aligned}$$

to re-sum the summation over r_{23} into a ${}_3F_2$ -hypergeometric function, given by

$${}_3F_2 \left[\begin{matrix} -m_1, -m_2, -\bar{p}_3 - h_2 + d/2 - m_1 \\ p_4 - m_1, 1 + h_3 - m_1 \end{matrix}; 1 \right].$$

With the help of the identity (A.3), this can be rewritten as

$$\frac{(1 + h_3)_{m_2}}{(1 + h_3 - m_1)_{m_2}} {}_3F_2 \left[\begin{matrix} -m_1, -m_2, \bar{p}_4 + h_2 - d/2 \\ p_4 - m_1, -h_3 - m_2 \end{matrix}; 1 \right],$$

which translates into the result of [43], i.e.

$$\begin{aligned}
 G_5^P &= \sum_{\{m_a, m_{ab} \geq 0\}} \frac{(p_2 + h_2)_{m_1 + m_{23} + m_{24}} (p_3)_{m_1} (-h_3)_{m_1 - m_2 + m_{23}} (p_4)_{m_2 - m_1 + m_{34}}}{(p_4)_{-m_1} (\bar{p}_3 + h_2)_{2m_1 + m_{23} + m_{24}} (\bar{p}_3 + h_2 + 1 - d/2)_{m_1}} \\
 &\times \frac{(\bar{p}_3 + \bar{h}_3)_{m_1 + m_2 + m_{24}} (-h_4)_{m_2 + m_{24} + m_{34}} (\bar{p}_4 + \bar{h}_4)_{m_2}}{(-h_3)_{-m_2} (\bar{p}_4 + \bar{h}_3)_{2m_2 + m_{24} + m_{34}} (\bar{p}_4 + \bar{h}_3 + 1 - d/2)_{m_2}} \\
 &\times \frac{(u_1^P)^{m_1} (u_2^P)^{m_2} (1 - v_{23}^P)^{m_{23}} (1 - v_{24}^P)^{m_{24}} (1 - v_{34}^P)^{m_{34}}}{m_1! m_2! m_{23}! m_{24}! m_{34}!} \\
 &\times {}_3F_2 \left[\begin{matrix} -m_1, -m_2, \bar{p}_4 + h_2 - d/2 \\ p_4 - m_1, -h_3 - m_2 \end{matrix}; 1 \right].
 \end{aligned} \tag{A.4}$$

This computation shows that the results of [43] and [44] are equivalent. Moreover, once a choice of conformal cross-ratios has been made, the OPE approach does lead to the correct result, after several re-summations.

The steps highlighted here can be repeated to obtain the scalar five-point conformal blocks in the comb channel discussed in (2.13). The proof of their equivalence at the level of (A.4) and (2.13) follows the one for the scalar six-point conformal blocks in the snowflake channel shown in appendix B.

B Snowflake and the OPE

In this appendix we expound the proof of the scalar six-point conformal blocks in the snowflake channel. We also present an alternative form for the scalar six-point conformal blocks in the snowflake channel, and prove that it is equivalent to the one introduced in the main text.

B.1 Proof of the snowflake

Starting from the scalar five-point correlation functions (2.13), shifted such that $\mathcal{O}_{i_a}(\eta_a) \rightarrow \mathcal{O}_{i_{a-1}}(\eta_{a-1})$ with $\mathcal{O}_{i_0}(\eta_0) \equiv \mathcal{O}_{i_5}(\eta_5)$, the legs and the conformal cross-ratios transform into

$$L_{5|\text{comb}}^{(\Delta_{i_2}, \dots, \Delta_{i_5}, \Delta_{k_3})} = \left(\frac{\eta_{13}}{\eta_{12}\eta_{23}} \right)^{\frac{\Delta_{i_2}}{2}} \left(\frac{\eta_{12}}{\eta_{13}\eta_{23}} \right)^{\frac{\Delta_{i_3}}{2}} \left(\frac{\eta_{35}}{\eta_{34}\eta_{45}} \right)^{\frac{\Delta_{i_4}}{2}} \left(\frac{\eta_{34}}{\eta_{35}\eta_{45}} \right)^{\frac{\Delta_{i_5}}{2}} \left(\frac{\eta_{35}}{\eta_{13}\eta_{15}} \right)^{\frac{\Delta_{k_3}}{2}},$$

$$u_1^5 = \frac{\eta_{15}\eta_{23}}{\eta_{12}\eta_{35}}, \quad u_2^5 = \frac{\eta_{13}\eta_{45}}{\eta_{15}\eta_{34}}, \quad v_{11}^5 = \frac{\eta_{13}\eta_{25}}{\eta_{12}\eta_{35}}, \quad v_{12}^5 = \frac{\eta_{14}\eta_{35}}{\eta_{15}\eta_{34}}, \quad v_{22}^5 = \frac{\eta_{13}\eta_{24}}{\eta_{12}\eta_{34}}.$$
(B.1)

Here, we already substituted $\mathcal{O}_{i_1}(\eta_1) \rightarrow \mathcal{O}_{k_3}(\eta_1)$ as needed for the recurrence relation (2.7). Moreover, the vectors \mathbf{h} and \mathbf{p} get transformed into (2.17), which also include the new elements appearing in the recurrence relation (2.7), while the functional forms of C_5 and F_5 (2.13) remain the same [they are the same functions but of the new vectors \mathbf{h} and \mathbf{p} (2.17), hence G_5 is the same function but of the new conformal cross-ratios and vectors].

Using the recurrence relation (2.7) with $k = 4$, $l = 5$, and $m = 5$, it is clear that the resulting scalar six-point correlation functions are in the snowflake channel. To proceed, we must act with the OPE differential operator as in (2.4) on the conformal cross-ratios [see (2.3)]

$$x_5^6 = \frac{\eta_{16}\eta_{45}}{\eta_{15}\eta_{46}}, \quad 1 - y_2^6 = \frac{\eta_{12}\eta_{56}}{\eta_{15}\eta_{26}}, \quad 1 - y_3^6 = \frac{\eta_{13}\eta_{56}}{\eta_{15}\eta_{36}}, \quad 1 - y_4^6 = \frac{\eta_{14}\eta_{56}}{\eta_{15}\eta_{46}},$$

$$z_{23}^6 = \frac{\eta_{23}\eta_{46}\eta_{56}}{\eta_{26}\eta_{36}\eta_{45}}, \quad z_{24}^6 = \frac{\eta_{24}\eta_{56}}{\eta_{26}\eta_{45}}, \quad z_{25}^6 = \frac{\eta_{25}\eta_{46}}{\eta_{26}\eta_{45}}, \quad z_{34}^6 = \frac{\eta_{34}\eta_{56}}{\eta_{36}\eta_{45}}, \quad z_{35}^6 = \frac{\eta_{35}\eta_{46}}{\eta_{36}\eta_{45}}.$$
(B.2)

This can be done easily by re-expressing (B.1) in terms of (B.2) as

$$u_1^5 = \frac{1 - y_5^6 z_{23}^6}{1 - y_2^6 z_{35}^6}, \quad u_2^5 = \frac{1 - y_3^6}{1 - y_5^6 z_{34}^6},$$

$$v_{11}^5 = \frac{1 - y_3^6 z_{25}^6}{1 - y_2^6 z_{35}^6}, \quad v_{12}^5 = \frac{1 - y_4^6 z_{35}^6}{1 - y_5^6 z_{34}^6}, \quad v_{22}^5 = \frac{1 - y_3^6 z_{24}^6}{1 - y_2^6 z_{34}^6}.$$

Defining a new set of conformal cross-ratios for the snowflake as in (2.14), we have

$$x_5^6 = \frac{u_2^6 v_3^6}{v_{33}^6}, \quad 1 - y_2^6 = \frac{1}{v_{13}^6}, \quad 1 - y_3^6 = \frac{1}{v_{23}^6}, \quad 1 - y_4^6 = \frac{v_{12}^6}{v_{33}^6},$$

$$z_{23}^6 = \frac{u_1^6 v_{33}^6}{u_2^6 v_{13}^6 v_{23}^6}, \quad z_{24}^6 = \frac{v_{22}^6}{u_2^6 v_{13}^6}, \quad z_{25}^6 = \frac{v_{11}^6 v_{33}^6}{u_2^6 v_{13}^6}, \quad z_{34}^6 = \frac{1}{u_2^6 v_{23}^6}, \quad z_{35}^6 = \frac{v_{33}^6}{u_2^6 v_{23}^6},$$

such that the action of the OPE differential operator on I_5 (2.13) gives

$$\begin{aligned}
 G_6 = & \sum \binom{n_{11}}{s_{11}} \binom{n_{12}}{s_{12}} \binom{n_{22}}{s_{22}} \frac{(-h_5)_{m_3} (p_3 - n_1 + n_2 + s_{12})_{\bar{r}_5} (\bar{p}_3 + h_2 + h_5)_{m_3 + \sigma_2 + \sigma_3 + \sigma_4}}{(\bar{p}_3 + h_2)_{2m_3 + \sigma_2 + \sigma_3 + \sigma_4} (\bar{p}_3 + h_2 + 1 - d/2)_{m_3}} \\
 & \times (-1)^{s_{11} + s_{12} + s_{22} + l_2 + l_3 + l_4 + m_{11} + m_{12} + m_{22} + m_{13} + m_{23} + m_{33}} \binom{s_{12} - r_{45} - \bar{r}_4 - l_4}{m_{33}} \binom{l_4}{m_{12}} \\
 & \times \binom{r_{25} + s_{11}}{m_{11}} \binom{r_{24} + s_{22}}{m_{22}} \binom{\sigma_2}{l_2} \binom{\sigma_3}{l_3} \binom{\sigma_4}{l_4} \binom{h_3 - n_1 - s_{11} - s_{22} - r_{25} - \bar{r}_2 - l_2}{m_{13}} \\
 & \times \binom{-p_2 - \bar{h}_3 + n_2 + s_{11} + s_{22} - r_{35} - \bar{r}_3 - l_3}{m_{23}} \\
 & \times \frac{(-h_3 + n_1 + s_{11} + s_{22})_{\sigma_2 + r_{25} + \bar{r}_2} (p_2 + \bar{h}_3 - n_2 - s_{11} - s_{22})_{\sigma_3 + r_{35} + \bar{r}_3} (-s_{12})_{\sigma_4 + r_{45} + \bar{r}_4}}{\sigma_2! \sigma_3! \sigma_4!} \\
 & \times \frac{(u_1^6)^{n_1 + r_{23}} (u_2^6)^{n_2 + r_{45}} (u_3^6)^{m_3} \prod_{1 \leq a < b \leq 3} (1 - v_{ab}^6)^{m_{ab}}}{r_{23}! r_{24}! r_{25}! r_{34}! r_{35}! r_{45}! n_1! n_{11}! n_{12}! n_{22}!} C_5 F_5,
 \end{aligned} \tag{B.3}$$

with the proper legs (2.14) and $m_3 = \bar{r}_5 + \bar{r}$. Here again, we omit the indices of summation on the sum for notational simplicity. Equation (B.3) corresponds to the scalar six-point conformal blocks in the snowflake channel, and we now aim to re-sum as many superfluous sums as possible to get to our final results (2.15) and (2.16).

The sum over l_2 is straightforward since the terms containing l_2 are

$$(-1)^{l_2} \binom{\sigma_2}{l_2} = \frac{(-\sigma_2)_{l_2}}{l_2!},$$

and

$$\begin{aligned}
 & \binom{h_3 - n_1 - s_{11} - s_{22} - r_{25} - \bar{r}_2 - l_2}{m_{13}} \\
 & = (-1)^{m_{13}} \frac{(-h_3 + n_1 + s_{11} + s_{22} + r_{25} + \bar{r}_2)_{m_{13}} (-h_3 + n_1 + s_{11} + s_{22} + r_{25} + \bar{r}_2 + m_{13})_{l_2}}{m_{13}! (-h_3 + n_1 + s_{11} + s_{22} + r_{25} + \bar{r}_2)_{l_2}}.
 \end{aligned}$$

Thus, the sum over l_2 can be performed using (A.1). Similarly, we can evaluate the sums over l_3 , l_4 , σ_2 , σ_3 , and σ_4 , in that order,¹⁰ using (A.1). This results in

$$\begin{aligned}
 G_6 = & \sum \binom{n_{11}}{s_{11}} \binom{n_{12}}{s_{12}} \binom{n_{22}}{s_{22}} \frac{(-h_5)_{m_3 + m_{13} + m_{23} + m_{33}} (p_3 - n_1 + n_2 + s_{12})_{\bar{r}_5} (\bar{p}_3 + h_2 + h_5)_{m_3 + m_{12}}}{(\bar{p}_3 + h_2)_{2m_3 + m_{12} + m_{13} + m_{23} + m_{33}} (\bar{p}_3 + h_2 + 1 - d/2)_{m_3}} \\
 & \times (-1)^{s_{11} + s_{12} + s_{22} + m_{11} + m_{22}} \binom{r_{25} + s_{11}}{m_{11}} \binom{r_{24} + s_{22}}{m_{22}} (-s_{12})_{m_{12} + m_{33} + r_{45} + \bar{r}_4} \\
 & \times \frac{(-h_3 + n_1 + s_{11} + s_{22})_{m_{13} + r_{25} + \bar{r}_2} (p_2 + \bar{h}_3 - n_2 - s_{11} - s_{22})_{m_{23} + r_{35} + \bar{r}_3}}{r_{23}! r_{24}! r_{25}! r_{34}! r_{35}! r_{45}! n_1! n_{11}! n_{12}! n_{22}!} \\
 & \times \frac{(u_1^6)^{n_1 + r_{23}} (u_2^6)^{n_2 + r_{45}} (u_3^6)^{m_3} \prod_{1 \leq a < b \leq 3} (1 - v_{ab}^6)^{m_{ab}}}{m_{12}! m_{13}! m_{23}! m_{33}!} C_5 F_5.
 \end{aligned}$$

¹⁰To evaluate the sums over l_4 and σ_4 , we must first change the variables such that $l_4 \rightarrow l_4 + m_{12}$ and $\sigma_4 \rightarrow \sigma_4 + m_{12}$, respectively.

Since $r_{35} = m_3 - r_{25} - r_{34} - r_{45} - r_{23} - r_{24}$, changing variables by $s_{12} \rightarrow s_{12} + m_{12} + m_{33} + r_{45} + r_{24} + r_{34}$ leads to

$$\begin{aligned}
 G_6 &= \sum \binom{n_{11}}{s_{11}} \binom{n_{22}}{s_{22}} \frac{(-h_5)_{m_3+m_{13}+m_{23}+m_{33}} (\bar{p}_3+h_2+h_5)_{m_3+m_{12}}}{(\bar{p}_3+h_2)_{2m_3+m_{12}+m_{13}+m_{23}+m_{33}} (\bar{p}_3+h_2+1-d/2)_{m_3}} \\
 &\times (-1)^{s_{11}+s_{12}+s_{22}+m_{11}+m_{22}} \binom{r_{25}+s_{11}}{m_{11}} \binom{r_{24}+s_{22}}{m_{22}} \\
 &\times \frac{(p_3-n_1+n_2+s_{12}+m_{12}+m_{33}+r_{45}+r_{24}+r_{34})_{m_3-r_{23}-r_{24}-r_{34}}}{(n_{12}-s_{12}-m_{12}-m_{33}-r_{45}-r_{24}-r_{34})! (m_3-r_{25}-r_{34}-r_{45}-r_{23}-r_{24})!} \\
 &\times \frac{(-h_3+n_1+s_{11}+s_{22})_{m_{13}+r_{25}+r_{23}+r_{24}} (p_2+\bar{h}_3-n_2-s_{11}-s_{22})_{m_3+m_{23}-r_{24}-r_{25}-r_{45}}}{s_{12}! r_{23}! r_{24}! r_{25}! r_{34}! r_{45}! n_1! n_{11}! n_{22}!} \\
 &\times \frac{(u_1^6)^{n_1+r_{23}} (u_2^6)^{n_2+r_{45}} (u_3^6)^{m_3} \prod_{1 \leq a \leq b \leq 3} (1-v_{ab}^6)^{m_{ab}}}{m_{12}! m_{13}! m_{23}! m_{33}!} C_5 F_5,
 \end{aligned}$$

and we can now sum over r_{34} , giving [after using (A.2)]

$$\begin{aligned}
 G_6 &= \sum \binom{n_{11}}{s_{11}} \binom{n_{22}}{s_{22}} \frac{(-h_5)_{m_3+m_{13}+m_{23}+m_{33}} (\bar{p}_3+h_2+h_5)_{m_3+m_{12}}}{(\bar{p}_3+h_2)_{2m_3+m_{12}+m_{13}+m_{23}+m_{33}} (\bar{p}_3+h_2+1-d/2)_{m_3}} \\
 &\times (-1)^{s_{11}+s_{12}+s_{22}+m_{11}+m_{22}} \binom{r_{25}}{k_{11}} \binom{s_{11}}{m_{11}-k_{11}} \binom{r_{24}}{k_{22}} \binom{s_{22}}{m_{22}-k_{22}} \\
 &\times \frac{(p_3-n_1+n_2+n_{12})_{m_3-r_{25}-r_{45}-r_{23}-r_{24}} (p_3-n_1+n_2+m_3+s_{12}+m_{12}+m_{33}-r_{23}-r_{25})_{r_{25}+r_{45}}}{(n_{12}-s_{12}-m_{12}-m_{33}-r_{45}-r_{24})! (m_3-r_{25}-r_{45}-r_{23}-r_{24})!} \\
 &\times \frac{(-h_3+n_1+s_{11}+s_{22})_{m_{13}+r_{25}+r_{23}+r_{24}} (p_2+\bar{h}_3-n_2-s_{11}-s_{22})_{m_3+m_{23}-r_{24}-r_{25}-r_{45}}}{s_{12}! r_{23}! r_{24}! r_{25}! r_{45}! n_1! n_{11}! n_{22}!} \\
 &\times \frac{(u_1^6)^{n_1+r_{23}} (u_2^6)^{n_2+r_{45}} (u_3^6)^{m_3} \prod_{1 \leq a \leq b \leq 3} (1-v_{ab}^6)^{m_{ab}}}{m_{12}! m_{13}! m_{23}! m_{33}!} C_5 F_5.
 \end{aligned}$$

We then change the variables by $r_{24} \rightarrow r_{24} + k_{22}$, and $r_{25} \rightarrow r_{25} + k_{11}$ and finally define $r_{24} = r - r_{25}$. With these changes, we can compute the sums over r_{25} , r , s_{12} , and n_{12} , with

$$\begin{aligned}
 G_6 &= \sum \binom{n_{11}}{s_{11}} \binom{n_{22}}{s_{22}} \frac{(-h_5)_{m_3+m_{13}+m_{23}+m_{33}} (\bar{p}_3+h_2+h_5)_{m_3+m_{12}}}{(\bar{p}_3+h_2)_{2m_3+m_{12}+m_{13}+m_{23}+m_{33}} (\bar{p}_3+h_2+1-d/2)_{m_3}} \\
 &\times (-1)^{s_{11}+s_{22}+m_{11}+m_{22}} \binom{s_{11}}{m_{11}-k_{11}} \binom{s_{22}}{m_{22}-k_{22}} \\
 &\times \frac{(p_2+h_2+n_1-n_2+m_{13}+m_{23}+2r_{23}+k_{11}+k_{22})_{m_3-r_{23}-r_{45}-k_{11}-k_{22}}}{(m_3-r_{45}-r_{23}-k_{11}-k_{22})!} \\
 &\times \frac{(-h_3+n_1+s_{11}+s_{22})_{m_{13}+r_{23}+k_{11}+k_{22}} (p_2+\bar{h}_3-n_2-s_{11}-s_{22})_{m_{23}+r_{23}}}{k_{11}! k_{22}! r_{23}! r_{45}! n_1! n_{11}! n_{22}!} \\
 &\times \frac{(-h_2)_{n_1+n_2+n_{11}+n_{22}} (-h_3)_{n_1+n_{11}+n_{22}} (p_2+h_2)_{n_1} (p_2+h_3)_{n_1}}{(p_2)_{2n_1+n_{11}+n_{22}} (p_2+1-d/2)_{n_1}}
 \end{aligned}$$

$$\begin{aligned} &\times \frac{(p_3 - n_1)_{m_2+m_3+m_{12}+m_{33}-r_{23}} (p_3 - h_2 + h_4)_{m_2+n_{11}+k_{11}} (-h_4)_{m_2+m_{12}+m_{33}+n_{22}+k_{22}}}{(p_3 - h_2)_{2m_2+m_{12}+m_{33}+n_{11}+n_{22}+k_{11}+k_{22}} (p_3 - h_2 + 1 - d/2)_{n_2}} \\ &\times \frac{(u_1^6)^{n_1+r_{23}} (u_2^6)^{n_2+r_{45}} (u_3^6)^{m_3} \prod_{1 \leq a \leq b \leq 3} (1 - v_{ab}^6)^{m_{ab}}}{m_{12}! m_{13}! m_{23}! m_{33}!} F_5, \end{aligned}$$

where we made C_5 explicit.

After changing the variables by $s_{22} \rightarrow s_{22} + m_{22} - k_{22}$ and $s_{11} \rightarrow s_{11} + m_{11} - k_{11}$, and also defining $s_{22} = s - s_{11}$, we can evaluate the summation over s_{11} , leading to

$$\begin{aligned} G_6 &= \sum \frac{(-1)^{k_{11}+k_{22}} (-n_{11}-n_{22}+m_{11}+m_{22}-k_{11}-k_{22})_s (-h_5)_{m_3+m_{13}+m_{23}+m_{33}} (\bar{p}_3+h_2+h_5)_{m_3+m_{12}}}{(\bar{p}_3+h_2)_{2m_3+m_{12}+m_{13}+m_{23}+m_{33}} (\bar{p}_3+h_2+1-d/2)_{m_3}} \\ &\times \frac{(p_2+h_2+n_1-n_2+m_{13}+m_{23}+2r_{23}+k_{11}+k_{22})_{m_3-r_{23}-r_{45}-k_{11}-k_{22}}}{(m_3-r_{45}-r_{23}-k_{11}-k_{22})!} \\ &\times \frac{(-h_3+n_1+s+m_{11}+m_{22}-k_{11}-k_{22})_{m_{13}+r_{23}+k_{11}+k_{22}}}{s!(n_{11}-m_{11}+k_{11})!(n_{22}-m_{22}+k_{22})!(m_{11}-k_{11})!(m_{22}-k_{22})!k_{11}!k_{22}!r_{23}!r_{45}!n_1!} \\ &\times (p_2+\bar{h}_3-n_2-s-m_{11}-m_{22}+k_{11}+k_{22})_{m_{23}+r_{23}} \\ &\times \frac{(-h_2)_{n_1+n_2+n_{11}+n_{22}} (-h_3)_{n_1+n_{11}+n_{22}} (p_2+h_2)_{n_1} (p_2+h_3)_{n_1}}{(p_2)_{2n_1+n_{11}+n_{22}} (p_2+1-d/2)_{n_1}} \\ &\times \frac{(p_3-n_1)_{m_2+m_3+m_{12}+m_{33}-r_{23}} (p_3-h_2+h_4)_{m_2+n_{11}+k_{11}} (-h_4)_{m_2+m_{12}+m_{33}+n_{22}+k_{22}}}{(p_3-h_2)_{2m_2+m_{12}+m_{33}+n_{11}+n_{22}+k_{11}+k_{22}} (p_3-h_2+1-d/2)_{n_2}} \\ &\times \frac{(u_1^6)^{n_1+r_{23}} (u_2^6)^{n_2+r_{45}} (u_3^6)^{m_3} \prod_{1 \leq a \leq b \leq 3} (1 - v_{ab}^6)^{m_{ab}}}{m_{12}! m_{13}! m_{23}! m_{33}!} F_5. \end{aligned}$$

Using the following identity,

$$\begin{aligned} &(p_2 + \bar{h}_3 - n_2 - s - m_{11} - m_{22} + k_{11} + k_{22})_{m_{23}+r_{23}} \\ &= \sum_j \binom{m_{23} + r_{23}}{j} (p_2 + \bar{h}_3 - n_2 - m_{11} - m_{22} + k_{11} + k_{22})_j (-s)_{m_{23}+r_{23}-j}, \end{aligned}$$

and changing the index of summation s by $s \rightarrow s + m_{23} + r_{23} - j$, the sum over s can be performed, leading to

$$\begin{aligned} G_6 &= \sum \frac{(-1)^{k_{11}+k_{22}+m_{23}+r_{23}-j} (-h_5)_{m_3+m_{13}+m_{23}+m_{33}} (\bar{p}_3+h_2+h_5)_{m_3+m_{12}}}{(\bar{p}_3+h_2)_{2m_3+m_{12}+m_{13}+m_{23}+m_{33}} (\bar{p}_3+h_2+1-d/2)_{m_3}} \\ &\times \frac{(p_2+h_2+n_1-n_2+m_{13}+m_{23}+2r_{23}+k_{11}+k_{22})_{m_3-r_{23}-r_{45}-k_{11}-k_{22}}}{(m_3-r_{45}-r_{23}-k_{11}-k_{22})!} \binom{m_{23}+r_{23}}{j} \\ &\times \frac{(-h_3)_{m_1+m_{11}+m_{22}+m_{13}+m_{23}+r_{23}-j} (p_2+\bar{h}_3-n_2-m_{11}-m_{22}+k_{11}+k_{22})_j}{(n_{11}-m_{11}+k_{11})!(n_{22}-m_{22}+k_{22})!(m_{11}-k_{11})!(m_{22}-k_{22})!k_{11}!k_{22}!r_{23}!r_{45}!n_1!} \\ &\times \frac{(-n_{11}-n_{22}-k_{11}-k_{22}+m_{11}+m_{22})_{m_{23}+r_{23}-j} (-h_2)_{n_1+n_2+n_{11}+n_{22}} (p_2+h_2)_{n_1} (p_2+h_3)_{n_1}}{(p_2)_{2n_1+n_{11}+n_{22}} (p_2+1-d/2)_{n_1}} \\ &\times \frac{(p_3-n_1)_{m_2+m_3+m_{12}+m_{33}-r_{23}} (p_3-h_2+h_4)_{m_2+n_{11}+k_{11}} (-h_4)_{m_2+m_{12}+m_{33}+n_{22}+k_{22}}}{(p_3-h_2)_{2m_2+m_{12}+m_{33}+n_{11}+n_{22}+k_{11}+k_{22}} (p_3-h_2+1-d/2)_{n_2}} \end{aligned}$$

$$\begin{aligned} & \times (-m_{13} - r_{23} - k_{11} - k_{22})_{n_{11} + n_{22} + k_{11} + k_{22} - m_{11} - m_{22} - m_{23} - r_{23} + j} \\ & \times \frac{(u_1^6)^{n_1 + r_{23}} (u_2^6)^{n_2 + r_{45}} (u_3^6)^{m_3} \prod_{1 \leq a \leq b \leq 3} (1 - v_{ab}^6)^{m_{ab}}}{m_{12}! m_{13}! m_{23}! m_{33}!} F_5. \end{aligned}$$

With $n_{22} = n - n_{11}$, it is now possible to sum over n_{11} , n , j , k_{11} , and k_{22} , giving

$$\begin{aligned} G_6 = & \sum \frac{(-h_5)_{m_3 + m_{13} + m_{23} + m_{33}} (\bar{p}_3 + h_2 + h_5)_{m_3 + m_{12}}}{(\bar{p}_3 + h_2)_{2m_3 + m_{12} + m_{13} + m_{23} + m_{33}} (\bar{p}_3 + h_2 + 1 - d/2)_{m_3}} \\ & \times \frac{(p_2 + h_2 + m_1 - t_1 - m_2 + t_2)_{m_3 + m_{13} + m_{23} + t_1 - t_2}}{(m_3 - t_2 - t_1)!} \frac{(-h_3)_{m_1 + m_{11} + m_{22} + m_{13}}}{t_1! t_2! (m_1 - t_1)! (m_2 - t_2)!} \\ & \times \frac{(-h_2 + m_1 + m_2 - m_3)_{m_{11} + m_{22}} (-h_2)_{m_1 - t_1 + m_2 - t_2} (p_2 + h_2)_{m_1 - t_1} (p_2 + h_3)_{m_1 + m_{23}}}{(p_2)_{2m_1 + m_{11} + m_{22} + m_{13} + m_{23}} (p_2 + 1 - d/2)_{m_1 - t_1}} \\ & \times \frac{(p_3 - m_1 + t_1)_{m_2 + m_3 + m_{12} + m_{33} - t_1} (p_3 - h_2 + h_4)_{m_2 + m_{11}} (-h_4)_{m_2 + m_{12} + m_{22} + m_{33}}}{(p_3 - h_2)_{2m_2 + m_{11} + m_{12} + m_{22} + m_{33}} (p_3 - h_2 + 1 - d/2)_{m_2 - t_2}} \\ & \times \frac{(u_1^6)^{m_1} (u_2^6)^{m_2} (u_3^6)^{m_3} \prod_{1 \leq a \leq b \leq 3} (1 - v_{ab}^6)^{m_{ab}}}{m_{11}! m_{12}! m_{22}! m_{13}! m_{23}! m_{33}!} F_5, \end{aligned}$$

after redefining

$$r_{23} = t_1, \quad r_{45} = t_2, \quad n_1 = m_1 - t_1, \quad n_2 = m_2 - t_2.$$

We are thus left with three extra sums, two sums over t_1 and t_2 , respectively, and one sum from F_5 , which is given by

$$F_5 = {}_3F_2 \left[\begin{matrix} -n_1, -n_2, -p_2 + d/2 - n_1 \\ 1 - p_2 - h_2 - n_1, p_3 - n_1 \end{matrix} ; 1 \right] = {}_3F_2 \left[\begin{matrix} -m_1 + t_1, -m_2 + t_2, -p_2 + d/2 - m_1 + t_1 \\ 1 - p_2 - h_2 - m_1 + t_1, p_3 - m_1 + t_1 \end{matrix} ; 1 \right].$$

Using (A.3) twice such that

$$\begin{aligned} & {}_3F_2 \left[\begin{matrix} -m_1 + t_1, -m_2 + t_2, -p_2 + d/2 - m_1 + t_1 \\ 1 - p_2 - h_2 - m_1 + t_1, p_3 - m_1 + t_1 \end{matrix} ; 1 \right] \\ & = \frac{(p_2 + h_2 - m_2 + t_2)_{m_1 - t_1}}{(p_2 + h_2)_{m_1 - t_1}} {}_3F_2 \left[\begin{matrix} -m_1 + t_1, -m_2 + t_2, \bar{p}_3 - d/2 \\ p_2 + h_2 - m_2 + t_2, p_3 - m_1 + t_1 \end{matrix} ; 1 \right] \\ & = \frac{(p_2 + h_2 - m_2 + t_2)_{m_1 - t_1} (-p_3 + h_2 + d/2 - m_2 + t_2)_{m_2 - t_2}}{(p_2 + h_2)_{m_1 - t_1} (p_2 + h_2 - m_2 + t_2)_{m_2 - t_2}} \\ & \quad \times {}_3F_2 \left[\begin{matrix} p_3, -m_2 + t_2, \bar{p}_3 - d/2 \\ p_3 - h_2 - d/2 + 1, p_3 - m_1 + t_1 \end{matrix} ; 1 \right], \end{aligned}$$

and re-summing over t_2 using (A.1) gives (2.15) and (2.16) (with $t_1 \rightarrow t_2$ and the index of summation from the ${}_3F_2$ -hypergeometric function chosen to be t_1). This completes the proof of the scalar six-point conformal blocks in the snowflake channel.

To express F_6 in terms of Kampé de Fériet functions [45, 46], which are defined as

$$F_{q,s,v}^{p,r,u} \left[\begin{matrix} \mathbf{a}; \mathbf{c}; \mathbf{f} \\ \mathbf{b}; \mathbf{d}; \mathbf{g} \end{matrix} \middle| x, y \right] = \sum_{m,n \geq 0} \frac{(\mathbf{a})_{m+n} (\mathbf{c})_m (\mathbf{f})_n x^m y^n}{(\mathbf{b})_{m+n} (\mathbf{d})_m (\mathbf{g})_n m! n!}, \quad (\text{B.4})$$

where

$$\begin{aligned} (\mathbf{a})_{m+n} &= (a_1)_{m+n} \cdots (a_p)_{m+n}, & (\mathbf{b})_{m+n} &= (b_1)_{m+n} \cdots (b_q)_{m+n}, \\ (\mathbf{c})_m &= (c_1)_m \cdots (c_r)_m, & (\mathbf{d})_m &= (d_1)_m \cdots (d_s)_m, \\ (\mathbf{f})_n &= (f_1)_n \cdots (f_u)_n, & (\mathbf{g})_n &= (g_1)_n \cdots (g_v)_n, \end{aligned}$$

we first rewrite (2.16) as

$$\begin{aligned} F_6 &= \frac{(-p_3+h_2+d/2-m_2)_{m_2}}{(p_3)_{-m_1}(-h_2+m_1)_{-m_3}} \sum_{t_1 \geq 0} \frac{(-m_2)_{t_1} (p_3)_{t_1} (\bar{p}_3-d/2)_{t_1} (-h_2+m_1)_{t_1}}{(-h_2+m_1-m_3)_{t_1} (p_3-m_1)_{t_1} (p_3-h_2+1-d/2)_{t_1} t_1!} \\ &\times {}_3F_2 \left[\begin{matrix} -m_1, -m_3, -p_2+d/2-m_1 \\ 1+h_2-m_1-t_1, p_3-m_1+t_1 \end{matrix}; 1 \right], \end{aligned}$$

by re-summing over t_2 . Using (A.3) on the ${}_3F_2$ -hypergeometric function, expanding the resulting ${}_3F_2$ -hypergeometric function, and combining Pochhammer symbols, we get to

$$F_6 = \frac{(-p_3+h_2+d/2-m_2)_{m_2}}{(p_3)_{-m_1}(-h_2)_{-m_3}} F_{2,1,0}^{1,3,2} \left[\begin{matrix} \bar{p}_3-d/2; -m_2, -h_2, p_3; -m_1, -m_3 \\ -h_2-m_3, p_3-m_1; p_3-h_2+1-d/2; - \end{matrix} \middle| 1, 1 \right],$$

as stated in (2.16).

Finally, by using (A.3) again on the ${}_3F_2$ -hypergeometric function from the summation over t_2 , it is possible to re-express F_6 in terms of a more symmetric Kampé de Fériet function as

$$\begin{aligned} F_6 &= \frac{1}{(p_3)_{-m_1}(-p_3+h_2+d/2)_{-m_2}(-\bar{p}_3-h_2+d/2)_{-m_3}} \\ &\times F_{1,1,1}^{2,1,1} \left[\begin{matrix} \bar{p}_3-d/2, p_3; -m_2; -m_3 \\ p_3-m_1; p_3-h_2+1-d/2; \bar{p}_3+h_2+1-d/2 \end{matrix} \middle| 1, 1 \right]. \end{aligned}$$

This form is more symmetric since it clearly shows that F_6 is made out of two intertwined ${}_3F_2$ -hypergeometric functions, which is reminiscent of the comb channel result.

B.2 An alternative form

Instead of starting from the scalar five-point correlation functions (2.13), it is also possible to repeat the steps above starting from (A.4). With the legs and the conformal cross-ratios

$$\begin{aligned} L_6^* &= \left(\frac{\eta_{13}}{\eta_{12}\eta_{23}} \right)^{\frac{\Delta_{i_2}}{2}} \left(\frac{\eta_{12}}{\eta_{13}\eta_{23}} \right)^{\frac{\Delta_{i_3}}{2}} \left(\frac{\eta_{15}}{\eta_{14}\eta_{45}} \right)^{\frac{\Delta_{i_4}}{2}} \left(\frac{\eta_{14}}{\eta_{15}\eta_{45}} \right)^{\frac{\Delta_{i_5}}{2}} \left(\frac{\eta_{12}}{\eta_{16}\eta_{26}} \right)^{\frac{\Delta_{i_6}}{2}} \left(\frac{\eta_{26}}{\eta_{12}\eta_{16}} \right)^{\frac{\Delta_{i_1}}{2}}, \\ u_1^{*6} &= \frac{\eta_{23}\eta_{15}\eta_{34}}{\eta_{13}\eta_{24}\eta_{35}}, & u_2^{*6} &= \frac{\eta_{12}\eta_{34}\eta_{45}}{\eta_{14}\eta_{35}\eta_{24}}, & u_3^{*6} &= \frac{\eta_{16}\eta_{24}\eta_{35}}{\eta_{26}\eta_{15}\eta_{34}}, \\ v_{11}^{*6} &= \frac{\eta_{12}\eta_{34}}{\eta_{13}\eta_{24}}, & v_{12}^{*6} &= \frac{\eta_{34}\eta_{25}}{\eta_{24}\eta_{35}}, & v_{13}^{*6} &= \frac{\eta_{12}\eta_{36}}{\eta_{13}\eta_{26}}, \\ v_{22}^{*6} &= \frac{\eta_{15}\eta_{34}}{\eta_{14}\eta_{35}}, & v_{23}^{*6} &= \frac{\eta_{12}\eta_{46}}{\eta_{14}\eta_{26}}, & v_{33}^{*6} &= \frac{\eta_{12}\eta_{56}}{\eta_{15}\eta_{26}}, \end{aligned} \tag{B.5}$$

the resulting scalar six-point conformal blocks in the snowflake channel are

$$\begin{aligned}
 C_6^* &= \frac{(p_3-h_2+h_4)_{m_2+m_{12}+m_{33}} (-h_3)_{m_1+m_{12}} (p_3)_{-m_1+m_2+m_3+m_{23}+m_{33}} (p_2+h_3)_{m_1+m_{11}+m_{13}}}{(-h_2)_{m_1+m_2-m_3+m_{12}} (p_2)_{2m_1+m_{11}+m_{12}+m_{13}} (p_2+1-d/2)_{m_1}} \\
 &\times \frac{(p_3-h_2+h_4)_{m_2+m_{12}+m_{33}} (p_2+h_2)_{m_1-m_2+m_3+m_{13}} (-h_4)_{m_2+m_{22}+m_{23}}}{(p_3-h_2)_{2m_2+m_{12}+m_{22}+m_{23}+m_{33}} (p_3-h_2+1-d/2)_{m_2}} \\
 &\times \frac{(\bar{p}_3+h_2+h_5)_{m_3} (-h_2)_{m_1+m_2-m_3+m_{12}+m_{22}} (-h_5)_{m_3+m_{13}+m_{23}+m_{33}}}{(\bar{p}_3+h_2)_{2m_3+m_{13}+m_{23}+m_{33}} (\bar{p}_3+h_2+1-d/2)_{m_3}},
 \end{aligned} \tag{B.6}$$

with F_6^* given by (2.16).

We note here that although the alternative forms for F_M are the same ($F_6^* = F_6$), the alternative forms for C_M (B.6) and (2.15) are not the same function ($C_6^* \neq C_6$). However, the two results must be equivalent as shown below.

First, it is easy to see that the conformal cross-ratios (B.5) and (2.14) are related such that

$$\begin{aligned}
 u_1^{*6} &= \frac{u_1^6}{v_{22}^6}, & u_2^{*6} &= \frac{u_2^6}{v_{12}^6 v_{22}^6}, & u_3^{*6} &= \frac{u_3^6 v_{22}^6}{v_{13}^6}, \\
 v_{11}^{*6} &= \frac{1}{v_{22}^6}, & v_{12}^{*6} &= \frac{v_{11}^6}{v_{22}^6}, & v_{13}^{*6} &= \frac{v_{23}^6}{v_{13}^6}, \\
 v_{22}^{*6} &= \frac{1}{v_{12}^6}, & v_{23}^{*6} &= \frac{v_{33}^6}{v_{12}^6 v_{13}^6}, & v_{33}^{*6} &= \frac{1}{v_{13}^6}.
 \end{aligned}$$

Hence, since the scalar six-point correlation functions I_6 must be the same, we get the identity

$$G_6 = (v_{12}^6)^{h_4} (v_{13}^6)^{h_5} (v_{22}^6)^{h_2} G_6^* \equiv G_{v_6}^*, \tag{B.7}$$

for the scalar six-point conformal blocks in the snowflake channel (2.6). To prove (B.7), we re-express $G_{v_6}^*$ in terms of the conformal cross-ratios for G_6 , expand in the conformal cross-ratios of the latter, and evaluate the superfluous sums.

Using the fact that $F_6^* = F_6$, we first obtain

$$\begin{aligned}
 G_{v_6}^* &= \sum \frac{(-h_2)_{m_1+m_2-m_3+m_{12}+m_{22}} (-h_5)_{m_3+m_{13}+m_{23}+m_{33}} (\bar{p}_3+h_2+h_5)_{m_3}}{(\bar{p}_3+h_2)_{2m_3+m_{13}+m_{23}+m_{33}} (\bar{p}_3+h_2+1-d/2)_{m_3}} \\
 &\times \frac{(p_2+h_3)_{m_1+m_{11}+m_{13}} (p_2+h_2)_{m_1-m_2+m_3+m_{13}} (p_3)_{m_2-m_1+m_3+m_{23}+m_{33}}}{(p_2)_{2m_1+m_{11}+m_{12}+m_{13}} (p_2+1-d/2)_{m_1}} \\
 &\times \frac{(-h_2)_{m_1+m_2-m_3+m_{11}+m_{12}} (p_3-h_2+h_4)_{m_2+m_{12}+m_{33}}}{(p_3-h_2)_{2m_2+m_{12}+m_{22}+m_{23}+m_{33}} (p_3-h_2+1-d/2)_{m_2}} \\
 &\times \frac{(-h_4)_{m_2+m_{22}+m_{23}} (-h_3)_{m_1+m_{12}}}{(-h_2)_{m_1+m_2-m_3+m_{12}}} \left[\prod_{1 \leq a \leq 3} \frac{(u_a^6)^{m_a}}{m_a!} \right] (v_{12}^6)^{h_4-m_2} (v_{13}^6)^{h_5-m_3} (v_{22}^6)^{h_2+m_3-m_1-m_2} \\
 &\times \frac{\left(1-\frac{1}{v_{22}^6}\right)^{m_{11}} \left(1-\frac{v_{11}^6}{v_{22}^6}\right)^{m_{12}} \left(1-\frac{v_{23}^6}{v_{13}^6}\right)^{m_{13}} \left(1-\frac{1}{v_{12}^6}\right)^{m_{22}} \left(1-\frac{v_{33}^6}{v_{12}^6 v_{13}^6}\right)^{m_{23}} \left(1-\frac{1}{v_{13}^6}\right)^{m_{33}}}{m_{11}! m_{12}! m_{13}! m_{22}! m_{23}! m_{33}!} F_6,
 \end{aligned}$$

in terms of the conformal cross-ratios (2.14). Expanding in terms of u_a^6 and $1 - v_{ab}^6$, G_{v6}^* becomes

$$\begin{aligned}
 G_{v6}^* &= \sum \frac{(-h_2)_{m_1+m_2-m_3+m_{12}+m_{22}}(-h_5)_{m_3+m_{13}+m_{23}+m_{33}}(\bar{p}_3+h_2+h_5)_{m_3}}{(\bar{p}_3+h_2)_{2m_3+m_{13}+m_{23}+m_{33}}(\bar{p}_3+h_2+1-d/2)_{m_3}} \\
 &\times \frac{(p_2+h_3)_{m_1+m_{11}+m_{13}}(p_2+h_2)_{m_1-m_2+m_3+m_{13}}(p_3)_{m_2-m_1+m_3+m_{23}+m_{33}}}{(p_2)_{2m_1+m_{11}+m_{12}+m_{13}}(p_2+1-d/2)_{m_1}} \\
 &\times \frac{(-h_2)_{m_1+m_2-m_3+m_{11}+m_{12}}(p_3-h_2+h_4)_{m_2+m_{12}+m_{33}}}{(p_3-h_2)_{2m_2+m_{12}+m_{22}+m_{23}+m_{33}}(p_3-h_2+1-d/2)_{m_2}} \\
 &\times \frac{(-h_4)_{m_2+m_{22}+m_{23}}(-h_3)_{m_1+m_{12}}}{(-h_2)_{m_1+m_2-m_3+m_{12}}} \prod_{1 \leq a \leq b \leq 3} \binom{m_{ab}}{k_{ab}} \binom{k_{12}}{m'_{11}} \binom{h_4-m_2-k_{22}-k_{23}}{m'_{12}} \\
 &\times \binom{h_5-m_3-k_{13}-k_{23}-k_{33}}{m'_{13}} \binom{h_2+m_3-m_1-m_2-k_{11}-k_{12}}{m'_{22}} \binom{k_{13}}{m'_{23}} \binom{k_{23}}{m'_{33}} \\
 &\times (-1)^{\sum_{1 \leq a \leq b \leq 3} (k_{ab}+m'_{ab})} \prod_{1 \leq a \leq 3} \frac{(u_a^6)^{m_a}}{m_a!} \prod_{1 \leq a \leq b \leq 3} \frac{(1-v_{ab}^6)^{m'_{ab}}}{m_{ab}!} F_6.
 \end{aligned}$$

We only need to evaluate the extra sums now to recover C_6 (2.15) and thus prove (B.7).

We start by observing that the terms containing k_{11} are given by

$$(-1)^{k_{11}} \binom{m_{11}}{k_{11}} = \frac{(-m_{11})_{k_{11}}}{k_{11}!},$$

and

$$\begin{aligned}
 &\binom{h_2+m_3-m_1-m_2-k_{11}-k_{12}}{m'_{22}} \\
 &= (-1)^{m'_{22}} \frac{(-h_2+m_1+m_2-m_3+k_{12}+m'_{22})_{k_{11}}(-h_2+m_1+m_2-m_3+k_{12})_{m'_{22}}}{(-h_2+m_1+m_2-m_3+k_{12})_{k_{11}} m'_{22}!}.
 \end{aligned}$$

Thus the summation over k_{11} can be done with the help of (A.1). The same can be said for the summations over all of the k_{ab} , except for k_{23} .¹¹ After these steps, G_{v6}^* becomes

$$\begin{aligned}
 G_{v6}^* &= \sum \frac{(-h_2)_{m_1+m_2-m_3+m_{12}+m_{22}}(-h_5)_{m_3+m_{13}+m_{23}+m_{33}}(\bar{p}_3+h_2+h_5)_{m_3}}{(\bar{p}_3+h_2)_{2m_3+m_{13}+m_{23}+m_{33}}(\bar{p}_3+h_2+1-d/2)_{m_3}} \\
 &\times \frac{(p_2+h_3)_{m_1+m_{11}+m_{13}}(p_2+h_2)_{m_1-m_2+m_3+m_{13}}(p_3)_{m_2-m_1+m_3+m_{23}+m_{33}}}{(p_2)_{2m_1+m_{11}+m_{12}+m_{13}}(p_2+1-d/2)_{m_1}} \\
 &\times \frac{(-h_2)_{m_1+m_2-m_3+m'_{11}+m'_{22}}(p_3-h_2+h_4)_{m_2+m_{12}+m_{33}}}{(p_3-h_2)_{2m_2+m_{12}+m_{22}+m_{23}+m_{33}}(p_3-h_2+1-d/2)_{m_2}} \\
 &\times \frac{(-h_4)_{m_2+m_{22}+m_{23}}(-h_3)_{m_1+m_{12}}}{(-h_2)_{m_1+m_2-m_3+m_{12}}} \binom{m_{23}}{k_{23}} \binom{k_{23}}{m'_{33}}
 \end{aligned}$$

¹¹To evaluate the sums over k_{12} and k_{13} , we must first change the variables by $k_{12} \rightarrow k_{12} + m'_{11}$ and $k_{13} \rightarrow k_{13} + m'_{23}$, respectively.

$$\begin{aligned}
 & \times \frac{m_{12}!(-m'_{22})_{m_{11}+m_{12}-m'_{11}}(-h_5+m_3+m_{33}+m_{13}+k_{23})_{m'_{13}+m'_{23}-m_{13}-m_{33}}}{m'_{11}!m'_{12}!m'_{13}m'_{22}!m'_{23}!} \\
 & \times \frac{m_{13}!(-m'_{12})_{m_{22}}(-h_4+m_2+m_{22}+k_{23})_{m'_{12}-m_{22}}(-m'_{13})_{m_{13}+m_{33}-m'_{23}}}{(m_{12}-m'_{11})!(m_{13}-m'_{23})!} \\
 & \times (-1)^{k_{23}+m'_{33}} \prod_{1 \leq a \leq 3} \frac{(u_a^6)^{m_a}}{m_a!} \prod_{1 \leq a \leq b \leq 3} \frac{(1-v_{ab}^6)^{m'_{ab}}}{m_{ab}!} F_6.
 \end{aligned}$$

We can now proceed with the summation over m_{11} , getting to

$$\begin{aligned}
 G_{v6}^* &= \sum \frac{(-h_2)_{m_1+m_2-m_3+m_{12}+m_{22}}(-h_5)_{m_3+m_{13}+m_{23}+m_{33}}(\bar{p}_3+h_2+h_5)_{m_3}}{(\bar{p}_3+h_2)_{2m_3+m_{13}+m_{23}+m_{33}}(\bar{p}_3+h_2+1-d/2)_{m_3}} \\
 & \times \frac{(p_2+h_3)_{m_1+m_{13}}(p_2+h_2)_{m_1-m_2+m_3+m_{13}}(p_3)_{m_2-m_1+m_3+m_{23}+m_{33}}}{(p_2)_{2m_1+m'_{11}+m'_{22}+m_{13}}(p_2+1-d/2)_{m_1}} \\
 & \times \frac{(-h_2)_{m_1+m_2-m_3+m'_{11}+m'_{22}}(p_3-h_2+h_4)_{m_2+m_{12}+m_{33}}}{(p_3-h_2)_{2m_2+m_{12}+m_{22}+m_{23}+m_{33}}(p_3-h_2+1-d/2)_{m_2}} \\
 & \times \frac{(-h_4)_{m_2+m_{22}+m_{23}}(-h_3)_{m_1+m'_{11}+m'_{22}}}{(-h_2)_{m_1+m_2-m_3+m_{12}}} \binom{m_{23}}{k_{23}} \binom{k_{23}}{m'_{33}} \\
 & \times \frac{m_{11}!m_{12}!(-m'_{22})_{m_{12}-m'_{11}}(-h_5+m_3+m_{33}+m_{13}+k_{23})_{m'_{13}+m'_{23}-m_{13}-m_{33}}}{m'_{11}!m'_{12}!m'_{13}m'_{22}!m'_{23}!} \\
 & \times \frac{m_{13}!(-m'_{12})_{m_{22}}(-h_4+m_2+m_{22}+k_{23})_{m'_{12}-m_{22}}(-m'_{13})_{m_{13}+m_{33}-m'_{23}}}{(m_{12}-m'_{11})!(m_{13}-m'_{23})!} \\
 & \times (-1)^{k_{23}+m'_{33}} \prod_{1 \leq a \leq 3} \frac{(u_a^6)^{m_a}}{m_a!} \prod_{1 \leq a \leq b \leq 3} \frac{(1-v_{ab}^6)^{m'_{ab}}}{m_{ab}!} F_6.
 \end{aligned}$$

To evaluate the summation over k_{23} , we first change the variable by $k_{23} \rightarrow k_{23} + m'_{33}$. Then, using the following identity

$$\begin{aligned}
 & (-h_4+m_2+m_{22}+m'_{33}+k_{23})_{m'_{12}-m_{22}} \\
 &= \sum_j \binom{m'_{12}-m_{22}}{j} (-h_4+m_2+m_{22}+m_{23})_{m'_{12}-m_{22}-j} (-m_{23}+m'_{33}+k_{23})_j,
 \end{aligned}$$

we can re-sum over k_{23} , leading to

$$\begin{aligned}
 G_{v6}^* &= \sum \frac{(-h_2)_{m_1+m_2-m_3+m_{12}+m_{22}}(\bar{p}_3+h_2+h_5)_{m_3}}{(\bar{p}_3+h_2)_{2m_3+m_{13}+m_{23}+m_{33}}(\bar{p}_3+h_2+1-d/2)_{m_3}} \\
 & \times \frac{(p_2+h_3)_{m_1+m_{13}}(p_2+h_2)_{m_1-m_2+m_3+m_{13}}(p_3)_{m_2-m_1+m_3+m_{23}+m_{33}}}{(p_2)_{2m_1+m'_{11}+m'_{22}+m_{13}}(p_2+1-d/2)_{m_1}} \\
 & \times \frac{(-h_2)_{m_1+m_2-m_3+m'_{11}+m'_{22}}(p_3-h_2+h_4)_{m_2+m_{12}+m_{33}}}{(p_3-h_2)_{2m_2+m_{12}+m_{22}+m_{23}+m_{33}}(p_3-h_2+1-d/2)_{m_2}} \\
 & \times \frac{(-h_4)_{m_2+m'_{12}+m_{23}-j}(-h_3)_{m_1+m'_{11}+m'_{22}}}{(-h_5+m_3+m_{13}+m_{23}+m_{33})_{-j}(-h_2)_{m_1+m_2-m_3+m_{12}}}
 \end{aligned}$$

$$\begin{aligned}
 & \times \frac{m_{11}!m_{12}!(-m'_{22})_{m_{12}-m'_{11}}(-h_5)_{m_3+m'_{13}+m'_{23}+m'_{33}}}{m'_{11}!m'_{12}!m'_{13}m'_{22}!m'_{23}!m'_{33}!} \\
 & \times \frac{m_{13}!m_{23}!(-m'_{12})_{m_{22}+j}(-m'_{13}-m'_{23}+m_{13}+m_{33})_{m_{23}-m'_{33}-j}(-m'_{13})_{m_{13}+m_{33}-m'_{23}}}{j!(m_{12}-m'_{11})!(m_{23}-m'_{33}-j)!(m_{13}-m'_{23})!} \\
 & \times \prod_{1 \leq a \leq 3} \frac{(u_a^6)^{m_a}}{m_a!} \prod_{1 \leq a \leq b \leq 3} \frac{(1-v_{ab}^6)^{m'_{ab}}}{m'_{ab}!} F_6.
 \end{aligned}$$

At this point, we sum over m_{22} and m_{12} after we change the variable such that $m_{12} \rightarrow m_{12} + m'_{11}$. This gives

$$\begin{aligned}
 G_{v6}^* &= \sum \frac{(\bar{p}_3 + h_2 + h_5)_{m_3}}{(\bar{p}_3 + h_2)_{2m_3+m_{13}+m_{23}+m_{33}}(\bar{p}_3 + h_2 + 1 - d/2)_{m_3}} \\
 & \times \frac{(p_2 + h_3)_{m_1+m_{13}}(p_2 + h_2)_{m_1-m_2+m_3+m_{13}}(p_3)_{m_2-m_1+m_3+m'_{12}+m_{23}+m_{33}-j}}{(p_2)_{2m_1+m'_{11}+m'_{22}+m_{13}}(p_2 + 1 - d/2)_{m_1}} \\
 & \times \frac{(-h_2)_{m_1+m_2-m_3+m'_{11}+m'_{22}}(p_3 - h_2 + h_4)_{m_2+m'_{11}+m_{33}}}{(p_3 - h_2)_{2m_2+m'_{11}+m'_{12}+m'_{22}+m_{23}+m_{33}-j}(p_3 - h_2 + 1 - d/2)_{m_2}} \\
 & \times \frac{(-h_5)_{m_3+m'_{13}+m'_{23}+m'_{33}}(-h_4)_{m_2+m'_{12}+m'_{22}+m_{23}-j}(-h_3)_{m_1+m'_{11}+m'_{22}}}{m_{33}!(-h_5 + m_3 + m_{13} + m_{23} + m_{33})_{-j}} \\
 & \times \frac{(-m'_{12})_j(-m'_{13}-m'_{23}+m_{13}+m_{33})_{m_{23}-m'_{33}-j}(-m'_{13})_{m_{13}+m_{33}-m'_{23}}}{j!(m_{23}-m'_{33}-j)!(m_{13}-m'_{23})!} \\
 & \times \prod_{1 \leq a \leq 3} \frac{(u_a^6)^{m_a}}{m_a!} \prod_{1 \leq a \leq b \leq 3} \frac{(1-v_{ab}^6)^{m'_{ab}}}{m'_{ab}!} F_6.
 \end{aligned}$$

Changing the variable by $m_{23} \rightarrow m_{23} + m'_{33} + j$ and then redefining $m_{23} = m - m_{33}$, we can finally perform the re-summations over m_{33} , j , m , and m_{13} (where we first make the substitution $m_{13} \rightarrow m_{13} + m'_{23}$). Redefining m'_{ab} by m_{ab} , we are thus left with $G_{v6}^* = G_6$, completing the proof.

C Symmetry properties

This appendix presents the proofs of the symmetry properties of the scalar six-point conformal blocks in the snowflake channel, which are generated by rotations, reflections and dendrite permutations as in (3.2), (3.3), and (3.4), respectively. Collectively, these symmetries generate the $(\mathbb{Z}_2)^3 \times D_3$ group discussed in section 3.1.

C.1 Rotations of the triangle

To prove invariance under the rotation generator R (3.2), it is only necessary to check that (2.16) verifies

$$F_{6|\text{snowflake}}^{(d,h_2,h_3,h_4,h_5;p_2,p_3,p_4,p_5,p_6)}(m_1, m_2, m_3) = F_{6|\text{snowflake}}^{(d,-p_3,h_4,h_5,h_3;p_3-h_2,p_2+h_2,p_5,p_6,p_4)}(m_2, m_3, m_1). \quad (\text{C.1})$$

To simplify the notation, we denote (C.1) as $F_6 = F_{6R}$.

First, we rewrite (2.16) as

$$F_6 = \frac{(-p_3 + h_2 + d/2 - m_2)_{m_2}}{(p_3)_{-m_1}(-h_2 + m_1 + m_2)_{-m_3}} \sum_{t_1, t_2 \geq 0} \frac{(-m_1)_{t_2}(-m_2)_{t_1}(-m_3)_{t_2}}{(p_3 - h_2 - d/2 + 1)_{t_1}} \\ \times \frac{(p_3)_{t_1}(\bar{p}_3 - d/2)_{t_1}(-p_2 + d/2 - m_1)_{t_2}(1 + h_2 - m_1 + m_3 - m_2)_{m_2 - t_1}}{(1 + h_2 - m_1 - m_2)_{m_2 - t_1 + t_2}(p_3 - m_1)_{t_1 + t_2} t_1! t_2!},$$

and then use the identity

$$\frac{(1 + h_2 - m_1 + m_3 - m_2)_{m_2 - t_1}}{(1 + h_2 - m_1 - m_2)_{m_2 - t_1 + t_2}} = \frac{(1 + h_2 - m_1 + m_3 - m_2)_{m_2 - t_1}}{(1 + h_2 - m_1 - m_2 + t_2)_{m_2 - t_1}(1 + h_2 - m_1 - m_2)_{t_2}} \\ = \sum_{t_3 \geq 0} \frac{(-m_3 + t_2)_{t_3}(-m_2 + t_1)_{t_3}}{(1 + h_2 - m_1 - m_2 + t_2)_{t_3}(1 + h_2 - m_1 - m_2)_{t_2} t_3!},$$

to express F_6 as

$$F_6 = \frac{(-p_3 + h_2 + d/2 - m_2)_{m_2}}{(p_3)_{-m_1}(-h_2 + m_1 + m_2)_{-m_3}} \sum_{t_1, t_2, t_3 \geq 0} \frac{(-m_1)_{t_2}(-m_2)_{t_1 + t_3}(-m_3)_{t_2 + t_3}}{(p_3 - h_2 - d/2 + 1)_{t_1}} \\ \times \frac{(p_3)_{t_1}(\bar{p}_3 - d/2)_{t_1}(-p_2 + d/2 - m_1)_{t_2}}{(1 + h_2 - m_1 - m_2)_{t_2 + t_3}(p_3 - m_1)_{t_1 + t_2} t_1! t_2! t_3!}.$$

We now modify the sum over t_1 with the help of (A.3),

$${}_3F_2 \left[\begin{matrix} -m_2 + t_3, \bar{p}_3 - d/2, p_3 \\ p_3 - h_2 - d/2 + 1, p_3 - m_1 + t_2 \end{matrix} ; 1 \right] = \frac{(p_2 + h_2 - m_2 + t_3)_{m_2 - t_3}}{(-p_3 + h_2 + d/2 - m_2 + t_3)_{m_2 - t_3}} \\ \times {}_3F_2 \left[\begin{matrix} -m_2 + t_3, \bar{p}_3 - d/2, -m_1 + t_2 \\ p_2 + h_2 - m_2 + t_3, p_3 - m_1 + t_2 \end{matrix} ; 1 \right],$$

to get

$$F_6 = \sum_{t_2, t_3 \geq 0} \frac{(-m_1)_{t_2}(-m_2)_{t_3}(-m_3)_{t_2 + t_3}}{(p_3)_{-m_1}(p_2 + h_2)_{-m_2}(-h_2 + m_1 + m_2)_{-m_3} t_2! t_3!} \\ \times \frac{(-p_2 + d/2 - m_1)_{t_2}(-p_3 + h_2 + d/2 - m_2)_{t_3}}{(p_3 - m_1)_{t_2}(p_2 + h_2 - m_2)_{t_3}(1 + h_2 - m_1 - m_2)_{t_2 + t_3}} \\ \times {}_3F_2 \left[\begin{matrix} -m_1 + t_2, \bar{p}_3 - d/2, -m_2 + t_3 \\ p_3 - m_1 + t_2, p_2 + h_2 - m_2 + t_3 \end{matrix} ; 1 \right],$$

or

$$F_6 = \sum_{t_1, t_2, t_3 \geq 0} \frac{(-m_1)_{t_1 + t_2}(-m_2)_{t_1 + t_3}(-m_3)_{t_2 + t_3}}{(p_3)_{-m_1}(p_2 + h_2)_{-m_2}(-h_2 + m_1 + m_2)_{-m_3} t_1! t_2! t_3!} \\ \times \frac{(\bar{p}_3 - d/2)_{t_1}(-p_2 + d/2 - m_1)_{t_2}(-p_3 + h_2 + d/2 - m_2)_{t_3}}{(p_3 - m_1)_{t_1 + t_2}(p_2 + h_2 - m_2)_{t_1 + t_3}(1 + h_2 - m_1 - m_2)_{t_2 + t_3}}. \quad (C.2)$$

This is the simplest form to check the invariance of F_6 under the reflection generator (3.3).

We can use (A.3) again to obtain

$$F_6 = \sum_{t_2, t_3 \geq 0} \frac{(-m_1)_{t_2}(-m_2)_{t_3}(-m_3)_{t_2 + t_3}}{(-p_2 + d/2)_{-m_1}(p_2 + h_2)_{-m_2}(-h_2 + m_1 + m_2)_{-m_3} t_2! t_3!} \\ \times \frac{(-p_3 + h_2 + d/2 - m_2)_{t_3}}{(p_2 + h_2 - m_2)_{t_3}(1 + h_2 - m_1 - m_2)_{t_2 + t_3}} {}_3F_2 \left[\begin{matrix} -m_1 + t_2, \bar{p}_3 - d/2, p_2 + h_2 \\ p_2 + 1 - d/2, p_2 + h_2 - m_2 + t_3 \end{matrix} ; 1 \right]. \quad (C.3)$$

It is now possible to expand the ${}_3F_2$ -hypergeometric function as a summation over t_1 and evaluate the sum over t_2 with the help of (A.1), leading to

$$F_6 = \sum_{t_1, t_3 \geq 0} \frac{(-m_1)_{t_1} (-m_2)_{t_3} (-m_3)_{t_3}}{(-p_2 + d/2)_{-m_1} (p_2 + h_2)_{-m_2} (-h_2 + m_1 + m_2)_{-m_3} t_1! t_3!} \\ \times \frac{(1 + h_2 - m_1 - m_2 + m_3)_{m_1 - t_1} (\bar{p}_3 - d/2)_{t_1} (p_2 + h_2)_{t_1} (-p_3 + h_2 + d/2 - m_2)_{t_3}}{(p_2 + 1 - d/2)_{t_1} (p_2 + h_2 - m_2)_{t_1 + t_3} (1 + h_2 - m_1 - m_2)_{m_1 - t_1 + t_3}}.$$

Expressing the sum over t_3 as a ${}_3F_2$ -hypergeometric function, we get

$$F_6 = \sum_{t_1 \geq 0} \frac{(-h_2 + m_2 - m_3 + t_1)_{m_3}}{(-p_2 + d/2)_{-m_1} (p_2 + h_2)_{-m_2}} \frac{(-m_1)_{t_1} (\bar{p}_3 - d/2)_{t_1} (p_2 + h_2)_{t_1}}{(p_2 + 1 - d/2)_{t_1} (p_2 + h_2 - m_2)_{t_1} t_1!} \\ \times {}_3F_2 \left[\begin{matrix} -m_2, -m_3, -p_3 + h_2 + d/2 - m_2 \\ 1 + h_2 - m_2 - t_1, p_2 + h_2 - m_2 + t_1 \end{matrix} ; 1 \right].$$

Using (A.3) once more leads to

$$F_6 = \sum_{t_1 \geq 0} \frac{(-h_2 - m_3 + t_1)_{m_3}}{(-p_2 + d/2)_{-m_1} (p_2 + h_2)_{-m_2}} \frac{(-m_1)_{t_1} (\bar{p}_3 - d/2)_{t_1} (p_2 + h_2)_{t_1}}{(p_2 + 1 - d/2)_{t_1} (p_2 + h_2 - m_2)_{t_1} t_1!} \\ \times {}_3F_2 \left[\begin{matrix} -m_2, -m_3, \bar{p}_3 - d/2 + t_1 \\ -h_2 - m_3 + t_1, p_2 + h_2 - m_2 + t_1 \end{matrix} ; 1 \right] \\ = \sum_{t_1 \geq 0} \frac{(-h_2 - m_3 + t_1)_{m_3}}{(-p_2 + d/2)_{-m_1} (p_2 + h_2)_{-m_2}} \frac{(-m_1)_{t_1} (\bar{p}_3 - d/2)_{t_1} (p_2 + h_2)_{t_1}}{(p_2 + 1 - d/2)_{t_1} (p_2 + h_2 - m_2)_{t_1} t_1!} \\ \times {}_3F_2 \left[\begin{matrix} -m_2, -m_3, \bar{p}_3 - d/2 + t_1 \\ p_2 + h_2 - m_2 + t_1, -h_2 - m_3 + t_1 \end{matrix} ; 1 \right],$$

where in the last equality we simply changed the order of the two bottom parameters of the ${}_3F_2$ -hypergeometric function. With this change, we can finally re-use (A.3) to get

$$F_6 = \sum_{t_1 \geq 0} \frac{(p_2 + h_2 - m_2 + m_3 + t_1)_{m_2} (-h_2 - m_3 + t_1)_{m_3}}{(-p_2 + d/2)_{-m_1} (p_2 + h_2)_{-m_2}} \frac{(-m_1)_{t_1} (\bar{p}_3 - d/2)_{t_1} (p_2 + h_2)_{t_1}}{(p_2 + 1 - d/2)_{t_1} (p_2 + h_2 - m_2)_{m_2 + t_1} t_1!} \\ \times {}_3F_2 \left[\begin{matrix} -m_2, -m_3, -\bar{p}_3 - h_2 + d/2 - m_3 \\ 1 - p_2 - h_2 - m_3 - t_1, -h_2 - m_3 + t_1 \end{matrix} ; 1 \right],$$

which is nothing else than $F_6 = F_{6R}$, completing the proof of (C.1) and (3.2).

C.2 Reflections of the triangle

Invariance under the reflection generator S implies the identity (3.3), which we rewrite as $G_6 = G_{6S}$ to simplify the notation. Before proceeding, we observe that, under the reflection generator, $F_6 \rightarrow F_6$ from the definition (C.2). Expressing G_{6S} in terms of the

original conformal cross-ratios and expanding, we get

$$\begin{aligned}
 G_{6S} = & \sum \frac{(p_3 - h_2 + h_4)_{m'_2+m_{23}} (p_2 + h_3)_{m'_1+m_{11}} (\bar{p}_3 + h_2 + h_5)_{m'_3+m_{12}}}{(\bar{p}_3 + h_2)_{2m'_3+m_{12}+m_{13}+m_{23}+m_{33}} (\bar{p}_3 + h_2 + 1 - d/2)_{m'_3}} \\
 & \times \frac{(p_2 + h_2)_{m'_1+m'_3-m'_2+m_{12}+m_{33}} (p_3)_{m'_2-m'_1+m'_3+m_{13}+m_{23}} (-h_2)_{m'_1+m'_2-m'_3+m_{11}+m_{22}}}{(p_3 - h_2)_{2m'_2+m_{11}+m_{22}+m_{13}+m_{23}} (p_3 - h_2 + 1 - d/2)_{m'_2}} \\
 & \times \frac{(-h_4)_{m'_2+m_{11}+m_{22}+m_{13}} (-h_3)_{m'_1+m_{12}+m_{22}+m_{33}} (-h_5)_{m'_3+m_{13}+m_{23}+m_{33}}}{(p_2)_{2m'_1+m_{11}+m_{12}+m_{22}+m_{33}} (p_2 + 1 - d/2)_{m'_1}} \\
 & \times \binom{m_{11}}{k_{11}} \binom{m_{12}}{k_{12}} \binom{m_{13}}{k_{13}} \binom{m_{22}}{k_{22}} \binom{m_{23}}{k_{23}} \binom{m_{33}}{k_{33}} \binom{h_3 - m'_1 - k_{12} - k_{22} - k_{33}}{m'_{11}} \\
 & \times \binom{h_4 - m'_2 - k_{11} - k_{13} - k_{22}}{m'_{12}} \binom{k_{33}}{m'_{13}} \binom{k_{22}}{m'_{22}} \binom{h_5 - m'_3 - k_{13} - k_{23} - k_{33}}{m'_{23}} \binom{k_{13}}{m'_{33}} \\
 & \times (-1)^{\sum_{1 \leq a \leq b \leq 3} (k_{ab} + m'_{ab})} F_6 \prod_{1 \leq a \leq 3} \frac{(u_a^6)^{m'_a}}{m'_a!} \prod_{1 \leq a \leq b \leq 3} \frac{(1 - v_{ab}^6)^{m'_{ab}}}{m_{ab}!},
 \end{aligned}$$

where we must re-sum all the extra sums.

We start by evaluating the summations over k_{11} , k_{12} , and k_{23} using (A.1), getting to

$$\begin{aligned}
 G_{6S} = & \sum \frac{(p_3 - h_2 + h_4)_{m'_2+m_{23}} (p_2 + h_3)_{m'_1+m_{11}} (\bar{p}_3 + h_2 + h_5)_{m'_3+m_{12}}}{(\bar{p}_3 + h_2)_{2m'_3+m_{12}+m_{13}+m_{23}+m_{33}} (\bar{p}_3 + h_2 + 1 - d/2)_{m'_3}} \\
 & \times \frac{(p_2 + h_2)_{m'_1+m'_3-m'_2+m_{12}+m_{33}} (p_3)_{m'_2-m'_1+m'_3+m_{13}+m_{23}} (-h_2)_{m'_1+m'_2-m'_3+m_{11}+m_{22}}}{(p_3 - h_2)_{2m'_2+m_{11}+m_{22}+m_{13}+m_{23}} (p_3 - h_2 + 1 - d/2)_{m'_2}} \\
 & \times \frac{(-h_4)_{m'_2+m_{11}+m_{22}+m_{13}} (-h_3)_{m'_1+m_{12}+m_{22}+m_{33}} (-h_5)_{m'_3+m_{13}+m_{23}+m_{33}}}{(p_2)_{2m'_1+m_{11}+m_{12}+m_{22}+m_{33}} (p_2 + 1 - d/2)_{m'_1}} \\
 & \times \binom{m_{13}}{k_{13}} \binom{m_{22}}{k_{22}} \binom{m_{33}}{k_{33}} \frac{(-m'_{11})_{m_{12}} (-h_3 + m'_1 + m_{12} + k_{22} + k_{33})_{m'_{11}-m_{12}}}{m'_{11}!} \\
 & \times \frac{(-m'_{12})_{m_{11}} (-h_4 + m'_2 + m_{11} + k_{13} + k_{22})_{m'_{12}-m_{11}} \binom{k_{33}}{m'_{13}} \binom{k_{22}}{m'_{22}} \binom{k_{13}}{m'_{33}}}{m'_{12}!} \\
 & \times \frac{(-m'_{23})_{m_{23}} (-h_5 + m'_3 + m_{23} + k_{13} + k_{33})_{m'_{23}-m_{23}} (-1)^{k_{13}+k_{22}+k_{33}+m'_{13}+m'_{22}+m'_{33}}}{m'_{23}!} \\
 & \times F_6 \prod_{1 \leq a \leq 3} \frac{(u_a^6)^{m'_a}}{m'_a!} \prod_{1 \leq a \leq b \leq 3} \frac{(1 - v_{ab}^6)^{m'_{ab}}}{m_{ab}!}.
 \end{aligned}$$

We then change the variables by $k_{13} \rightarrow k_{13} + m'_{33}$, $k_{22} \rightarrow k_{22} + m'_{22}$, and $k_{33} \rightarrow k_{33} + m'_{33}$, and use the identity

$$\begin{aligned}
 & (-h_4 + m'_2 + m'_{22} + m'_{33} + m_{11} + k_{13} + k_{22})_{m'_{12}-m_{11}} \\
 & = \sum_{j_1} \binom{m'_{12}-m_{11}}{j_1} (-h_4 + m'_2 + m'_{22} + m_{11} + m_{13} + k_{22})_{m'_{12}-m_{11}-j_1} (-m_{13} + m'_{33} + k_{13})_{j_1},
 \end{aligned}$$

to re-sum over k_{13} , leading to

$$\begin{aligned}
 G_{6S} = & \sum \frac{(p_3 - h_2 + h_4)_{m'_2 + m_{23}} (p_2 + h_3)_{m'_1 + m_{11}} (\bar{p}_3 + h_2 + h_5)_{m'_3 + m_{12}}}{(\bar{p}_3 + h_2)_{2m'_3 + m_{12} + m_{13} + m_{23} + m_{33}} (\bar{p}_3 + h_2 + 1 - d/2)_{m'_3}} \\
 & \times \frac{(p_2 + h_2)_{m'_1 + m'_3 - m'_2 + m_{12} + m_{33}} (p_3)_{m'_2 - m'_1 + m'_3 + m_{13} + m_{23}} (-h_2)_{m'_1 + m'_2 - m'_3 + m_{11} + m_{22}}}{(p_3 - h_2)_{2m'_2 + m_{11} + m_{22} + m_{13} + m_{23}} (p_3 - h_2 + 1 - d/2)_{m'_2}} \\
 & \times \frac{(-h_4)_{m'_2 + m_{11} + m_{22} + m_{13}} (-h_3)_{m'_1 + m_{12} + m_{22} + m_{33}} (-h_5)_{m'_3 + m_{13} + m_{23} + m_{33}}}{(p_2)_{2m'_1 + m_{11} + m_{12} + m_{22} + m_{33}} (p_2 + 1 - d/2)_{m'_1}} \\
 & \times \frac{(-m'_{11})_{m_{12}} (-h_3 + m'_1 + m'_{13} + m'_{22} + m_{12} + k_{22} + k_{33})_{m'_{11} - m_{12}}}{m_{11}! (m_{13} - m'_{33} - j_1)!} \\
 & \times \frac{(-m'_{12})_{m_{11} + j_1} (-h_4 + m'_2 + m'_{22} + m_{11} + m_{13} + k_{22})_{m'_{12} - m_{11} - j_1}}{k_{33}! m_{12}! (m_{22} - m'_{22} - k_{22})!} \\
 & \times \frac{(-m'_{23})_{m_{13} + m_{23} - m'_{33} - j_1} (-h_5 + m'_3 + m'_{13} + m_{13} + m_{23} + k_{33} - j_1)_{m'_{23} + m'_{33} - m_{13} - m_{23} + j_1}}{j_1! k_{22}! m_{23}! (m_{33} - m'_{13} - k_{33})!} \\
 & \times (-1)^{k_{22} + k_{33}} F_6 \prod_{1 \leq a \leq 3} \frac{(u_a^6)^{m'_a}}{m'_a!} \prod_{1 \leq a \leq b \leq 3} \frac{(1 - v_{ab}^6)^{m'_{ab}}}{m'_{ab}!}.
 \end{aligned}$$

Similarly, using the identity

$$\begin{aligned}
 & (-h_3 + m'_1 + m'_{13} + m'_{22} + m_{12} + k_{22} + k_{33})_{m'_{11} - m_{12}} \\
 & = \sum_{j_2} \binom{m'_{11} - m_{12}}{j_2} (-h_3 + m'_1 + m'_{22} + m_{12} + m_{33} + k_{22})_{m'_{11} - m_{12} - j_2} (-m_{33} + m'_{13} + k_{33})_{j_2},
 \end{aligned}$$

we can compute the sum over k_{33} and we get

$$\begin{aligned}
 G_{6S} = & \sum \frac{(p_3 - h_2 + h_4)_{m'_2 + m_{23}} (p_2 + h_3)_{m'_1 + m_{11}} (\bar{p}_3 + h_2 + h_5)_{m'_3 + m_{12}}}{(\bar{p}_3 + h_2)_{2m'_3 + m_{12} + m_{13} + m_{23} + m_{33}} (\bar{p}_3 + h_2 + 1 - d/2)_{m'_3}} \\
 & \times \frac{(p_2 + h_2)_{m'_1 + m'_3 - m'_2 + m_{12} + m_{33}} (p_3)_{m'_2 - m'_1 + m'_3 + m_{13} + m_{23}} (-h_2)_{m'_1 + m'_2 - m'_3 + m_{11} + m_{22}}}{(p_3 - h_2)_{2m'_2 + m_{11} + m_{22} + m_{13} + m_{23}} (p_3 - h_2 + 1 - d/2)_{m'_2}} \\
 & \times \frac{(-h_4)_{m'_2 + m_{11} + m_{22} + m_{13}} (-h_3)_{m'_1 + m_{12} + m_{22} + m_{33}} (-h_5)_{m'_3 + m'_{13} + m'_{23} + m'_{33}}}{(p_2)_{2m'_1 + m_{11} + m_{12} + m_{22} + m_{33}} (p_2 + 1 - d/2)_{m'_1}} \\
 & \times \frac{(-m'_{11})_{m_{12} + j_2} (-h_3 + m'_1 + m'_{22} + m_{12} + m_{33} + k_{22})_{m'_{11} - m_{12} - j_2}}{m_{11}! (m_{13} - m'_{33} - j_1)!} \\
 & \times \frac{(-m'_{12})_{m_{11} + j_1} (-h_4 + m'_2 + m'_{22} + m_{11} + m_{13} + k_{22})_{m'_{12} - m_{11} - j_1}}{m_{12}! (m_{22} - m'_{22} - k_{22})!} \\
 & \times \frac{(-m'_{23})_{m_{13} + m_{23} + m_{33} - m'_{13} - m'_{33} - j_1 - j_2}}{(-h_5 + m'_3 + m_{13} + m_{23} + m_{33})_{-j_1 - j_2} j_1! j_2! k_{22}! m_{23}! (m_{33} - m'_{13} - j_2)!} \\
 & \times (-1)^{k_{22}} F_6 \prod_{1 \leq a \leq 3} \frac{(u_a^6)^{m'_a}}{m'_a!} \prod_{1 \leq a \leq b \leq 3} \frac{(1 - v_{ab}^6)^{m'_{ab}}}{m'_{ab}!}.
 \end{aligned}$$

Once again, we introduce

$$\begin{aligned}
 & (-h_3 + m'_1 + m'_{22} + m_{12} + m_{33} + k_{22})_{m'_{11} - m_{12} - j_2} \\
 &= \sum_{j_3} \binom{m'_{11} - m_{12} - j_2}{j_3} (-h_3 + m'_1 + m_{12} + m_{22} + m_{33})_{m'_{11} - m_{12} - j_2 - j_3} (-m_{22} + m'_{22} + k_{22})_{j_3},
 \end{aligned}$$

and sum over k_{22} to obtain

$$\begin{aligned}
 G_{6S} &= \sum \frac{(p_3 - h_2 + h_4)_{m'_2 + m_{23}} (p_2 + h_3)_{m'_1 + m_{11}} (\bar{p}_3 + h_2 + h_5)_{m'_3 + m_{12}}}{(\bar{p}_3 + h_2)_{2m'_3 + m_{12} + m_{13} + m_{23} + m_{33}} (\bar{p}_3 + h_2 + 1 - d/2)_{m'_3}} \\
 &\times \frac{(p_2 + h_2)_{m'_1 + m'_3 - m'_2 + m_{12} + m_{33}} (p_3)_{m'_2 - m'_1 + m'_3 + m_{13} + m_{23}} (-h_2)_{m'_1 + m'_2 - m'_3 + m_{11} + m_{22}}}{(p_3 - h_2)_{2m'_2 + m_{11} + m_{22} + m_{13} + m_{23}} (p_3 - h_2 + 1 - d/2)_{m'_2}} \\
 &\times \frac{(-h_4)_{m'_2 + m'_{12} + m'_{22} + m_{13} - j_1} (-h_3)_{m'_1 + m'_{11} + m_{22} + m_{33} - j_2 - j_3} (-h_5)_{m'_3 + m'_{13} + m'_{23} + m'_{33}}}{(p_2)_{2m'_1 + m_{11} + m_{12} + m_{22} + m_{33}} (p_2 + 1 - d/2)_{m'_1}} \\
 &\times \frac{(-m'_{12})_{m_{11} + m_{22} - m'_{22} + j_1 - j_3}}{(-h_4 + m'_2 + m_{22} + m_{11} + m_{13})_{-j_3} m_{12}! (m_{22} - m'_{22} - j_3)!} \\
 &\times \frac{(-m'_{23})_{m_{13} + m_{23} + m_{33} - m'_{13} - m'_{33} - j_1 - j_2}}{(-h_5 + m'_3 + m_{13} + m_{23} + m_{33})_{-j_1 - j_2} j_1! j_2! j_3! m_{23}! (m_{33} - m'_{13} - j_2)!} \\
 &\times \frac{(-m'_{11})_{m_{12} + j_2 + j_3}}{m_{11}! (m_{13} - m'_{33} - j_1)!} F_6 \prod_{1 \leq a \leq 3} \frac{(u_a^6)^{m'_a}}{m'_a!} \prod_{1 \leq a \leq b \leq 3} \frac{(1 - v_{ab}^6)^{m'_{ab}}}{m'_{ab}!}.
 \end{aligned}$$

At this point, we define $m_{22} = m - m_{11}$ and evaluate the summation over m_{11} , always using (A.1), to find

$$\begin{aligned}
 G_{6S} &= \sum \frac{(p_3 - h_2 + h_4)_{m'_2 + m_{23}} (p_2 + h_3)_{m'_1} (\bar{p}_3 + h_2 + h_5)_{m'_3 + m_{12}}}{(\bar{p}_3 + h_2)_{2m'_3 + m_{12} + m_{13} + m_{23} + m_{33}} (\bar{p}_3 + h_2 + 1 - d/2)_{m'_3}} \\
 &\times \frac{(p_2 + h_2)_{m'_1 + m'_3 - m'_2 + m_{12} + m_{33}} (p_3)_{m'_2 - m'_1 + m'_3 + m_{13} + m_{23}} (-h_2)_{m'_1 + m'_2 - m'_3 + m}}{(p_3 - h_2)_{2m'_2 + m + m_{13} + m_{23}} (p_3 - h_2 + 1 - d/2)_{m'_2}} \\
 &\times \frac{(-h_4)_{m'_2 + m'_{12} + m'_{22} + m_{13} - j_1} (-h_3)_{m'_1 + m'_{11} + m'_{22} + m_{33} - j_2} (-h_5)_{m'_3 + m'_{13} + m'_{23} + m'_{33}}}{(p_2)_{2m'_1 + m + m_{12} + m_{33}} (p_2 + 1 - d/2)_{m'_1}} \\
 &\times \frac{(p_2 + 2m'_1 + m'_{11} + m'_{22} + m_{33} - j_2)_{m - m'_{22} - j_3} (-m'_{12})_{m - m'_{22} + j_1 - j_3}}{(-h_4 + m'_2 + m + m_{13})_{-j_3} m_{12}! (m - m'_{22} - j_3)!} \\
 &\times \frac{(-m'_{23})_{m_{13} + m_{23} + m_{33} - m'_{13} - m'_{33} - j_1 - j_2}}{(-h_5 + m'_3 + m_{13} + m_{23} + m_{33})_{-j_1 - j_2} m_{23}! (m_{33} - m'_{13} - j_2)!} \\
 &\times \frac{(-m'_{11})_{m_{12} + j_2 + j_3}}{(m_{13} - m'_{33} - j_1)!} F_6 \prod_{1 \leq a \leq 3} \frac{(u_a^6)^{m'_a}}{m'_a!} \prod_{1 \leq a \leq b \leq 3} \frac{(1 - v_{ab}^6)^{m'_{ab}}}{m'_{ab}!}.
 \end{aligned}$$

We now express the summation over j_3 as a ${}_3F_2$ -hypergeometric function, use (A.3), and re-expand the ${}_3F_2$ -hypergeometric function as a sum over j_3 to rewrite G_{6S} as

$$\begin{aligned}
 G_{6S} = & \sum \frac{(p_3-h_2+h_4)_{m'_2+m_{23}}(p_2+h_3)_{m'_1}(\bar{p}_3+h_2+h_5)_{m'_3+m_{12}}}{(\bar{p}_3+h_2)_{2m'_3+m_{12}+m_{13}+m_{23}+m_{33}}(\bar{p}_3+h_2+1-d/2)_{m'_3}} \\
 & \times \frac{(p_2+h_2)_{m'_1+m'_3-m'_2+m_{12}+m_{33}}(p_3)_{m'_2-m'_1+m'_3+m_{13}+m_{23}}(-h_2)_{m'_1+m'_2-m'_3+m}}{(p_3-h_2)_{2m'_2+m+m_{13}+m_{23}}(p_3-h_2+1-d/2)_{m'_2}} \\
 & \times \frac{(-h_4)_{m'_2+m'_{12}+m'_{22}+m_{13}-j_1+j_3}(-h_3)_{m'_1+m'_{11}+m'_{22}+m_{33}-j_2}(-h_5)_{m'_3+m'_{13}+m'_{23}+m'_{33}}}{(p_2)_{2m'_1+m'_{22}+m_{12}+m_{33}+j_3}(p_2+1-d/2)_{m'_1}} \\
 & \times \frac{(-m'_{12})_{m-m'_{22}+j_1-j_3}(-m'_{23})_{m_{13}+m_{23}+m_{33}-m'_{13}-m'_{33}-j_1-j_2}}{(-h_5+m'_3+m_{13}+m_{23}+m_{33})_{-j_1-j_2}j_1!j_2!j_3!m_{12}!(m-m'_{22}-j_3)!m_{23}!(m_{33}-m'_{13}-j_2)!} \\
 & \times \frac{(-m'_{11})_{m_{12}+j_2+j_3}}{(m_{13}-m'_{33}-j_1)!} F_6 \prod_{1 \leq a \leq 3} \frac{(u_a^6)^{m'_a}}{m'_a!} \prod_{1 \leq a \leq b \leq 3} \frac{(1-v_{ab}^6)^{m'_{ab}}}{m'_{ab}!}.
 \end{aligned}$$

After changing m by $m \rightarrow m+m'_{22}+j_3$, we evaluate the summation over m , leading to

$$\begin{aligned}
 G_{6S} = & \sum \frac{(p_3-h_2+h_4)_{m'_2+m_{23}}(p_2+h_3)_{m'_1}(\bar{p}_3+h_2+h_5)_{m'_3+m_{12}}}{(\bar{p}_3+h_2)_{2m'_3+m_{12}+m_{13}+m_{23}+m_{33}}(\bar{p}_3+h_2+1-d/2)_{m'_3}} \\
 & \times \frac{(p_2+h_2)_{m'_1+m'_3-m'_2+m_{12}+m_{33}}(p_3)_{m'_2-m'_1+m'_3+m'_{12}+m_{13}+m_{23}-j_1}(-h_2)_{m'_1+m'_2-m'_3+m'_{22}+j_3}}{(p_3-h_2)_{2m'_2+m'_{12}+m'_{22}+m_{13}+m_{23}-j_1+j_3}(p_3-h_2+1-d/2)_{m'_2}} \\
 & \times \frac{(-h_4)_{m'_2+m'_{12}+m'_{22}+m_{13}-j_1+j_3}(-h_3)_{m'_1+m'_{11}+m'_{22}+m_{33}-j_2}(-h_5)_{m'_3+m'_{13}+m'_{23}+m'_{33}}}{(p_2)_{2m'_1+m'_{22}+m_{12}+m_{33}+j_3}(p_2+1-d/2)_{m'_1}} \\
 & \times \frac{(-m'_{12})_{j_1}(-m'_{23})_{m_{13}+m_{23}+m_{33}-m'_{13}-m'_{33}-j_1-j_2}}{(-h_5+m'_3+m_{13}+m_{23}+m_{33})_{-j_1-j_2}j_1!j_2!j_3!m_{12}!m_{23}!(m_{33}-m'_{13}-j_2)!} \\
 & \times \frac{(-m'_{11})_{m_{12}+j_2+j_3}}{(m_{13}-m'_{33}-j_1)!} F_6 \prod_{1 \leq a \leq 3} \frac{(u_a^6)^{m'_a}}{m'_a!} \prod_{1 \leq a \leq b \leq 3} \frac{(1-v_{ab}^6)^{m'_{ab}}}{m'_{ab}!}.
 \end{aligned}$$

We then change the variable m_{13} by $m_{13} \rightarrow m_{13}+m'_{33}+j_1$ and compute the summation over j_1 . As a result, we get

$$\begin{aligned}
 G_{6S} = & \sum \frac{(p_3-h_2+h_4)_{m'_2+m_{23}}(p_2+h_3)_{m'_1}(\bar{p}_3+h_2+h_5)_{m'_3+m'_{12}+m_{12}}}{(\bar{p}_3+h_2)_{2m'_3+m'_{12}+m'_{33}+m_{12}+m_{13}+m_{23}+m_{33}}(\bar{p}_3+h_2+1-d/2)_{m'_3}} \\
 & \times \frac{(p_2+h_2)_{m'_1+m'_3-m'_2+m_{12}+m_{33}}(p_3)_{m'_2-m'_1+m'_3+m'_{12}+m'_{33}+m_{13}+m_{23}}(-h_2)_{m'_1+m'_2-m'_3+m'_{22}+j_3}}{(p_3-h_2)_{2m'_2+m'_{12}+m'_{22}+m'_{33}+m_{13}+m_{23}+j_3}(p_3-h_2+1-d/2)_{m'_2}} \\
 & \times \frac{(-h_4)_{m'_2+m'_{12}+m'_{22}+m'_{33}+m_{13}+j_3}(-h_3)_{m'_1+m'_{11}+m'_{22}+m_{33}-j_2}(-h_5)_{m'_3+m'_{13}+m'_{23}+m'_{33}}}{(p_2)_{2m'_1+m'_{22}+m_{12}+m_{33}+j_3}(p_2+1-d/2)_{m'_1}} \\
 & \times \frac{(-m'_{23})_{m_{13}+m_{23}+m_{33}-m'_{13}-j_2}}{(-h_5+m'_3+m'_{33}+m_{13}+m_{23}+m_{33})_{-j_2}j_2!j_3!m_{12}!m_{23}!(m_{33}-m'_{13}-j_2)!} \\
 & \times \frac{(-m'_{11})_{m_{12}+j_2+j_3}}{(m_{13})!} F_6 \prod_{1 \leq a \leq 3} \frac{(u_a^6)^{m'_a}}{m'_a!} \prod_{1 \leq a \leq b \leq 3} \frac{(1-v_{ab}^6)^{m'_{ab}}}{m'_{ab}!}.
 \end{aligned}$$

We now redefine $m_{23} = n - m_{13}$ and evaluate the summation over m_{13} with (A.1), which implies

$$\begin{aligned}
 G_{6S} = & \sum \frac{(p_3 - h_2 + h_4)_{m'_2} (p_2 + h_3)_{m'_1} (\bar{p}_3 + h_2 + h_5)_{m'_3 + m'_{12} + m_{12}}}{(\bar{p}_3 + h_2)_{2m'_3 + m'_{12} + m'_{33} + m_{12} + n + m_{33}} (\bar{p}_3 + h_2 + 1 - d/2)_{m'_3}} \\
 & \times \frac{(p_2 + h_2)_{m'_1 + m'_3 - m'_2 + m_{12} + m_{33}} (p_3)_{m'_2 - m'_1 + m'_3 + m'_{12} + m'_{33} + n} (-h_2)_{m'_1 + m'_2 - m'_3 + m'_{22} + j_3}}{(p_3 - h_2)_{2m'_2 + m'_{12} + m'_{22} + m'_{33} + j_3} (p_3 - h_2 + 1 - d/2)_{m'_2}} \\
 & \times \frac{(-h_4)_{m'_2 + m'_{12} + m'_{22} + m'_{33} + j_3} (-h_3)_{m'_1 + m'_{11} + m'_{22} + m_{33} - j_2} (-h_5)_{m'_3 + m'_{13} + m'_{23} + m'_{33}}}{(p_2)_{2m'_1 + m'_{22} + m_{12} + m_{33} + j_3} (p_2 + 1 - d/2)_{m'_1}} \\
 & \times \frac{(-m'_{23})_{n + m_{33} - m'_{13} - j_2}}{(-h_5 + m'_3 + m'_{33} + n + m_{33})_{-j_2} j_2! j_3! m_{12}! (m_{33} - m'_{13} - j_2)!} \\
 & \times \frac{(-m'_{11})_{m_{12} + j_2 + j_3}}{n!} F_6 \prod_{1 \leq a \leq 3} \frac{(u_a^6)^{m'_a}}{m'_a!} \prod_{1 \leq a \leq b \leq 3} \frac{(1 - v_{ab}^6)^{m'_{ab}}}{m'_{ab}!}.
 \end{aligned}$$

We can proceed with the summation over j_2 , which gives a ${}_3F_2$ -hypergeometric function, use (A.3) once more, and re-expand with the same index of summation to rewrite G_{6S} as

$$\begin{aligned}
 G_{6S} = & \sum \frac{(p_3 - h_2 + h_4)_{m'_2} (p_2 + h_3)_{m'_1} (\bar{p}_3 + h_2 + h_5)_{m'_3 + m'_{12} + m_{12}}}{(\bar{p}_3 + h_2)_{2m'_3 + m'_{12} + m'_{33} + m_{12} + n + m_{33}} (\bar{p}_3 + h_2 + 1 - d/2)_{m'_3}} \\
 & \times \frac{(p_2 + h_2)_{m'_1 + m'_3 - m'_2 + m_{12} + m_{33}} (p_3)_{m'_2 - m'_1 + m'_3 + m'_{12} + m'_{33} + n} (-h_2)_{m'_1 + m'_2 - m'_3 + m'_{22} + j_3}}{(p_3 - h_2)_{2m'_2 + m'_{12} + m'_{22} + m'_{33} + j_3} (p_3 - h_2 + 1 - d/2)_{m'_2}} \\
 & \times \frac{(-h_4)_{m'_2 + m'_{12} + m'_{22} + m'_{33} + j_3} (-h_3)_{m'_1 + m'_{11} + m'_{13} + m'_{22}} (-h_5)_{m'_3 + m'_{13} + m'_{23} + m'_{33} + j_2}}{(p_2)_{2m'_1 + m'_{22} + m_{12} + m_{33} + j_3} (p_2 + 1 - d/2)_{m'_1}} \\
 & \times \frac{(-h_3 + m'_1 + m'_{13} + m'_{22} + m_{12} + j_2 + j_3)_{m_{33} - m'_{13} - j_2} (-m'_{23})_{m_{33} - m'_{13} - j_2}}{j_2! j_3! m_{12}! (m_{33} - m'_{13} - j_2)!} \\
 & \times \frac{(-m'_{11})_{m_{12} + j_2 + j_3}}{n!} F_6 \prod_{1 \leq a \leq 3} \frac{(u_a^6)^{m'_a}}{m'_a!} \prod_{1 \leq a \leq b \leq 3} \frac{(1 - v_{ab}^6)^{m'_{ab}}}{m'_{ab}!}.
 \end{aligned}$$

This allows us to sum over n , leading to

$$\begin{aligned}
 G_{6S} = & \sum \frac{(p_3 - h_2 + h_4)_{m'_2} (p_2 + h_3)_{m'_1} (\bar{p}_3 + h_2 + h_5)_{m'_3 + m'_{12} + m_{12}}}{(\bar{p}_3 + h_2)_{2m'_3 + m'_{12} + m'_{13} + m'_{23} + m'_{33} + m_{12} + j_2} (\bar{p}_3 + h_2 + 1 - d/2)_{m'_3}} \\
 & \times \frac{(p_2 + h_2)_{m'_1 + m'_3 - m'_2 + m'_{13} + m'_{23} + m_{12} + j_2} (p_3)_{m'_2 - m'_1 + m'_3 + m'_{12} + m'_{33}} (-h_2)_{m'_1 + m'_2 - m'_3 + m'_{22} + j_3}}{(p_3 - h_2)_{2m'_2 + m'_{12} + m'_{22} + m'_{33} + j_3} (p_3 - h_2 + 1 - d/2)_{m'_2}} \\
 & \times \frac{(-h_4)_{m'_2 + m'_{12} + m'_{22} + m'_{33} + j_3} (-h_3)_{m'_1 + m'_{11} + m'_{13} + m'_{22}} (-h_5)_{m'_3 + m'_{13} + m'_{23} + m'_{33} + j_2}}{(p_2)_{2m'_1 + m'_{22} + m_{12} + m_{33} + j_3} (p_2 + 1 - d/2)_{m'_1}} \\
 & \times \frac{(-h_3 + m'_1 + m'_{13} + m'_{22} + m_{12} + j_2 + j_3)_{m_{33} - m'_{13} - j_2} (-m'_{23})_{m_{33} - m'_{13} - j_2}}{(m_{33} - m'_{13} - j_2)!} \\
 & \times \frac{(-m'_{11})_{m_{12} + j_2 + j_3}}{j_2! j_3! m_{12}!} F_6 \prod_{1 \leq a \leq 3} \frac{(u_a^6)^{m'_a}}{m'_a!} \prod_{1 \leq a \leq b \leq 3} \frac{(1 - v_{ab}^6)^{m'_{ab}}}{m'_{ab}!}.
 \end{aligned}$$

We then change m_{33} by $m_{33} \rightarrow m_{33} + m'_{13} + j_2$ and sum over m_{33} to get

$$\begin{aligned}
 G_{6S} = & \sum \frac{(p_3 - h_2 + h_4)_{m'_2} (p_2 + h_3)_{m'_1 + m'_{23}} (\bar{p}_3 + h_2 + h_5)_{m'_3 + m'_{12} + m_{12}}}{(\bar{p}_3 + h_2)_{2m'_3 + m'_{12} + m'_{13} + m'_{23} + m'_{33} + m_{12} + j_2} (\bar{p}_3 + h_2 + 1 - d/2)_{m'_3} \\
 & \times \frac{(p_2 + h_2)_{m'_1 + m'_3 - m'_2 + m'_{13} + m'_{23} + m_{12} + j_2} (p_3)_{m'_2 - m'_1 + m'_3 + m'_{12} + m'_{33}} (-h_2)_{m'_1 + m'_2 - m'_3 + m'_{22} + j_3}}{(p_3 - h_2)_{2m'_2 + m'_{12} + m'_{22} + m'_{33} + j_3} (p_3 - h_2 + 1 - d/2)_{m'_2}} \\
 & \times \frac{(-h_4)_{m'_2 + m'_{12} + m'_{22} + m'_{33} + j_3} (-h_3)_{m'_1 + m'_{11} + m'_{13} + m'_{22}} (-h_5)_{m'_3 + m'_{13} + m'_{23} + m'_{33} + j_2}}{(p_2)_{2m'_1 + m'_{13} + m'_{22} + m'_{23} + m_{12} + j_2 + j_3} (p_2 + 1 - d/2)_{m'_1}} \\
 & \times \frac{(-m'_{11})_{m_{12} + j_2 + j_3}}{j_2! j_3! m_{12}!} F_6 \prod_{1 \leq a \leq 3} \frac{(u_a^6)^{m'_a}}{m'_a!} \prod_{1 \leq a \leq b \leq 3} \frac{(1 - v_{ab}^6)^{m'_{ab}}}{m'_{ab}!}.
 \end{aligned}$$

We now sum over m_{12} , leading to a ${}_3F_2$ -hypergeometric function, and use (A.3) one last time to rewrite G_{6S} as

$$\begin{aligned}
 G_{6S} = & \sum \frac{(p_3 - h_2 + h_4)_{m'_2} (p_2 + h_3)_{m'_1 + m'_{23}} (\bar{p}_3 + h_2 + h_5)_{m'_3 + m'_{12}}}{(\bar{p}_3 + h_2)_{2m'_3 + m'_{12} + m'_{13} + m'_{23} + m'_{33} + m_{12} + j_2} (\bar{p}_3 + h_2 + 1 - d/2)_{m'_3} \\
 & \times \frac{(p_2 + h_2)_{m'_1 + m'_3 - m'_2 + m'_{13} + m'_{23} + m_{12} + j_2} (p_3)_{m'_2 - m'_1 + m'_3 + m'_{12} + m'_{33}} (-h_2)_{m'_1 + m'_2 - m'_3 + m'_{11} + m'_{22} - m_{12} - j_2}}{(p_3 - h_2)_{2m'_2 + m'_{12} + m'_{22} + m'_{33} + j_3} (p_3 - h_2 + 1 - d/2)_{m'_2}} \\
 & \times \frac{(-1)^{m_{12}} (-h_4)_{m'_2 + m'_{12} + m'_{22} + m'_{33} + j_3} (-h_3)_{m'_1 + m'_{11} + m'_{13} + m'_{22}} (-h_5)_{m'_3 + m'_{13} + m'_{23} + m'_{33} + m_{12} + j_2}}{(p_2)_{2m'_1 + m'_{11} + m'_{13} + m'_{22} + m'_{23}} (p_2 + 1 - d/2)_{m'_1}} \\
 & \times \frac{(-m'_{11})_{m_{12} + j_2 + j_3}}{j_2! j_3! m_{12}!} F_6 \prod_{1 \leq a \leq 3} \frac{(u_a^6)^{m'_a}}{m'_a!} \prod_{1 \leq a \leq b \leq 3} \frac{(1 - v_{ab}^6)^{m'_{ab}}}{m'_{ab}!}.
 \end{aligned}$$

This transformation allows us to sum over j_3 following (A.1), which leads to

$$\begin{aligned}
 G_{6S} = & \sum \frac{(p_3 - h_2 + h_4)_{m'_2 + m'_{11} - m_{12} - j_2} (p_2 + h_3)_{m'_1 + m'_{23}} (\bar{p}_3 + h_2 + h_5)_{m'_3 + m'_{12}}}{(\bar{p}_3 + h_2)_{2m'_3 + m'_{12} + m'_{13} + m'_{23} + m'_{33} + m_{12} + j_2} (\bar{p}_3 + h_2 + 1 - d/2)_{m'_3} \\
 & \times \frac{(p_2 + h_2)_{m'_1 + m'_3 - m'_2 + m'_{13} + m'_{23} + m_{12} + j_2} (p_3)_{m'_2 - m'_1 + m'_3 + m'_{12} + m'_{33}} (-h_2)_{m'_1 + m'_2 - m'_3 + m'_{11} + m'_{22} - m_{12} - j_2}}{(p_3 - h_2)_{2m'_2 + m'_{11} + m'_{12} + m'_{22} + m'_{33} - m_{12} - j_2} (p_3 - h_2 + 1 - d/2)_{m'_2}} \\
 & \times \frac{(-1)^{m_{12}} (-h_4)_{m'_2 + m'_{12} + m'_{22} + m'_{33}} (-h_3)_{m'_1 + m'_{11} + m'_{13} + m'_{22}} (-h_5)_{m'_3 + m'_{13} + m'_{23} + m'_{33} + m_{12} + j_2}}{(p_2)_{2m'_1 + m'_{11} + m'_{13} + m'_{22} + m'_{23}} (p_2 + 1 - d/2)_{m'_1}} \\
 & \times \frac{(-m'_{11})_{m_{12} + j_2}}{j_2! m_{12}!} F_6 \prod_{1 \leq a \leq 3} \frac{(u_a^6)^{m'_a}}{m'_a!} \prod_{1 \leq a \leq b \leq 3} \frac{(1 - v_{ab}^6)^{m'_{ab}}}{m'_{ab}!}.
 \end{aligned}$$

We are thus left with two extra sums (over m_{12} and j_2). However, they are both trivial. Indeed, by redefining $j_2 \rightarrow j_2 - m_{12}$ and using the binomial identity

$$\sum_{m_{12}} (-1)^{m_{12}} \frac{1}{m_{12}! (j_2 - m_{12})!} = \frac{1}{j_2!} (1 - 1)^{j_2},$$

we find that $j_2 = 0$ and thus

$$\begin{aligned}
 G_{6S} = & \sum \frac{(p_3 - h_2 + h_4)_{m'_2 + m'_{11}} (p_2 + h_3)_{m'_1 + m'_{23}} (\bar{p}_3 + h_2 + h_5)_{m'_3 + m'_{12}}}{(\bar{p}_3 + h_2)_{2m'_3 + m'_{12} + m'_{13} + m'_{23} + m'_{33}} (\bar{p}_3 + h_2 + 1 - d/2)_{m'_3} \\
 & \times \frac{(p_2 + h_2)_{m'_1 + m'_3 - m'_2 + m'_{13} + m'_{23}} (p_3)_{m'_2 - m'_1 + m'_3 + m'_{12} + m'_{33}} (-h_2)_{m'_1 + m'_2 - m'_3 + m'_{11} + m'_{22}}}{(p_3 - h_2)_{2m'_2 + m'_{11} + m'_{12} + m'_{22} + m'_{33}} (p_3 - h_2 + 1 - d/2)_{m'_2}}
 \end{aligned}$$

$$\begin{aligned} & \times \frac{(-h_4)_{m'_2+m'_{12}+m'_{22}+m'_{33}} (-h_3)_{m'_1+m'_{11}+m'_{13}+m'_{22}} (-h_5)_{m'_3+m'_{13}+m'_{23}+m'_{33}}}{(p_2)_{2m'_1+m'_{11}+m'_{13}+m'_{22}+m'_{23}} (p_2+1-d/2)_{m'_1}} \\ & \times F_6 \prod_{1 \leq a \leq 3} \frac{(u_a^6)^{m'_a}}{m'_a!} \prod_{1 \leq a < b \leq 3} \frac{(1-v_{ab}^6)^{m'_{ab}}}{m'_{ab}!}, \end{aligned}$$

which implies $G_{6S} = G_6$ and proves (3.3).

C.3 Permutations of the dendrites

Finally, we present the proof of the invariance of the scalar six-point correlation functions under dendrite permutation P (3.4). For this proof, we use the alternative form (B.6) for which $F_6^* = F_6$ (2.16), where $F_6^* \rightarrow F_6^*$ trivially under P . Equation (3.4) thus becomes

$$\begin{aligned} & G_{6|\text{snowflake}}^*(d, h_2, h_3, h_4, h_5; p_2, p_3, p_4, p_5, p_6) (u_1^*, u_2^*, u_3^*; v_{11}^*, v_{12}^*, v_{13}^*, v_{22}^*, v_{23}^*, v_{33}^*) \\ & = (v_{11}^*)^{h_2} (v_{12}^*)^{h_2} (v_{13}^*)^{h_5} \\ & \times G_{6|\text{snowflake}}^*(d, h_2, -p_2-h_3, h_4, h_5; p_2, p_3, p_4, p_5, p_6) \left(\frac{u_1^*}{v_{11}^* v_{12}^*}, \frac{u_2^*}{v_{11}^* v_{12}^*}, \frac{u_3^* v_{11}^* v_{12}^*}{v_{13}^*}, \frac{1}{v_{11}^*}, \frac{1}{v_{12}^*}, \frac{1}{v_{13}^*}, \frac{v_{22}^*}{v_{12}^*}, \frac{v_{23}^*}{v_{13}^*}, \frac{v_{33}^*}{v_{13}^*} \right). \end{aligned} \quad (\text{C.4})$$

For notational simplicity, we rewrite (C.4) as $G_6^* = G_{6P}^*$.

First, by expanding G_{6P}^* in terms of the initial conformal cross-ratios, we obtain

$$\begin{aligned} G_{6P}^* & = \sum \frac{(-h_2)_{m_1+m_2-m_3+m_{12}+m_{22}} (-h_5)_{m_3+m_{13}+m_{23}+m_{33}} (\bar{p}_3+h_2+h_5)_{m_3}}{(\bar{p}_3+h_2)_{2m_3+m_{13}+m_{23}+m_{33}} (\bar{p}_3+h_2+1-d/2)_{m_3}} \\ & \times \frac{(-h_3)_{m_1+m_{11}+m_{13}} (p_2+h_2)_{m_1-m_2+m_3+m_{13}} (p_3)_{m_2-m_1+m_3+m_{23}+m_{33}}}{(p_2)_{2m_1+m_{11}+m_{12}+m_{13}} (p_2+1-d/2)_{m_1}} \\ & \times \frac{(-h_2)_{m_1+m_2-m_3+m_{11}+m_{12}} (p_3-h_2+h_4)_{m_2+m_{12}+m_{33}} (-h_4)_{m_2+m_{22}+m_{23}} (p_2+h_3)_{m_1+m_{12}}}{(p_3-h_2)_{2m_2+m_{12}+m_{22}+m_{23}+m_{33}} (p_3-h_2+1-d/2)_{m_2} (-h_2)_{m_1+m_2-m_3+m_{12}}} \\ & \times \binom{m_{11}}{k_{11}} \binom{m_{12}}{k_{12}} \binom{m_{13}}{k_{13}} \binom{m_{22}}{k_{22}} \binom{m_{23}}{k_{23}} \binom{m_{33}}{k_{33}} \binom{h_2+m_3-m_1-m_2-k_{11}}{m'_{11}} \\ & \times \binom{h_2+m_3-m_1-m_2-k_{12}-k_{22}}{m'_{12}} \binom{h_5-m_3-k_{13}-k_{23}-k_{33}}{m'_{13}} \binom{k_{22}}{m'_{22}} \binom{k_{23}}{m'_{23}} \binom{k_{33}}{m'_{33}} \\ & \times (-1)^{\sum_{1 \leq a < b \leq 3} (k_{ab}+m'_{ab})} F_6^* \prod_{1 \leq a \leq 3} \frac{(u_a^*)^{m_a}}{m_a!} \prod_{1 \leq a < b \leq 3} \frac{(1-v_{ab}^*)^{m'_{ab}}}{m_{ab}!}, \end{aligned}$$

where all superfluous sums must be evaluated.

After evaluating the summations over all the k_{ab} (with change of variables $k_{ab} \rightarrow k_{ab}+m'_{ab}$ for $2 \leq a \leq b \leq 3$), we find

$$\begin{aligned} G_{6P}^* & = \sum \frac{(-h_2)_{m_1+m_2-m_3+m'_{12}+m'_{22}} (-h_5)_{m_3+m'_{13}+m'_{23}+m'_{33}} (\bar{p}_3+h_2+h_5)_{m_3}}{(\bar{p}_3+h_2)_{2m_3+m_{13}+m_{23}+m_{33}} (\bar{p}_3+h_2+1-d/2)_{m_3}} \\ & \times \frac{(-h_3)_{m_1+m_{11}+m_{13}} (p_2+h_2)_{m_1-m_2+m_3+m_{13}} (p_3)_{m_2-m_1+m_3+m_{23}+m_{33}}}{(p_2)_{2m_1+m_{11}+m_{12}+m_{13}} (p_2+1-d/2)_{m_1}} \\ & \times \frac{(-h_2)_{m_1+m_2-m_3+m_{11}+m_{12}} (p_3-h_2+h_4)_{m_2+m_{12}+m_{33}} (-h_4)_{m_2+m_{22}+m_{23}} (p_2+h_3)_{m_1+m_{12}}}{(p_3-h_2)_{2m_2+m_{12}+m_{22}+m_{23}+m_{33}} (p_3-h_2+1-d/2)_{m_2} (-h_2)_{m_1+m_2-m_3+m_{12}}} \end{aligned}$$

$$\begin{aligned} & \times \frac{(-m'_{11})_{m_{11}}(-h_2-m_3+m_1+m_2+m_{11})_{m'_{11}-m_{11}}(-m'_{12})_{m_{12}+m_{22}-m'_{22}}}{m_{11}!(m_{22}-m'_{22})!(m_{23}-m'_{23})!(m_{33}-m'_{33})!} \frac{(-m'_{12})_{m_{12}+m_{22}-m'_{22}}}{m_{12}!} \\ & \times \frac{(-m'_{13})_{m_{13}+m_{23}+m_{33}-m'_{23}-m'_{33}}}{m_{13}!} F_6^* \prod_{1 \leq a \leq 3} \frac{(u_a^{*6})^{m_a}}{m_a!} \prod_{1 \leq a \leq b \leq 3} \frac{(1-v_{ab}^{*6})^{m'_{ab}}}{m'_{ab}!}. \end{aligned}$$

Re-summing over m_{22} , after the change the variable $m_{22} \rightarrow m_{22}+m'_{22}$, we obtain

$$\begin{aligned} G_{6P}^* &= \sum \frac{(-h_2)_{m_1+m_2-m_3+m'_{12}+m'_{22}}(-h_5)_{m_3+m'_{13}+m'_{23}+m'_{33}}(\bar{p}_3+h_2+h_5)_{m_3}}{(\bar{p}_3+h_2)_{2m_3+m_{13}+m_{23}+m_{33}}(\bar{p}_3+h_2+1-d/2)_{m_3}} \\ & \times \frac{(-h_3)_{m_1+m_{11}+m_{13}}(p_2+h_2)_{m_1-m_2+m_3+m_{13}}(p_3)_{m_2-m_1+m_3+m_{23}+m_{33}}}{(p_2)_{2m_1+m_{11}+m_{12}+m_{13}}(p_2+1-d/2)_{m_1}} \\ & \times \frac{(-h_2)_{m_1+m_2-m_3+m_{11}+m_{12}}(p_3-h_2+h_4)_{m_2+m'_{12}+m_{33}}(-h_4)_{m_2+m'_{22}+m_{23}}(p_2+h_3)_{m_1+m_{12}}}{(p_3-h_2)_{2m_2+m'_{12}+m'_{22}+m_{23}+m_{33}}(p_3-h_2+1-d/2)_{m_2}} \frac{(-h_2)_{m_1+m_2-m_3+m_{12}}}{(-h_2)_{m_1+m_2-m_3+m_{12}}} \\ & \times \frac{(-m'_{11})_{m_{11}}(-h_2-m_3+m_1+m_2+m_{11})_{m'_{11}-m_{11}}(-m'_{12})_{m_{12}}}{m_{11}!(m_{23}-m'_{23})!(m_{33}-m'_{33})!} \frac{(-m'_{12})_{m_{12}}}{m_{12}!} \\ & \times \frac{(-m'_{13})_{m_{13}+m_{23}+m_{33}-m'_{23}-m'_{33}}}{m_{13}!} F_6^* \prod_{1 \leq a \leq 3} \frac{(u_a^{*6})^{m_a}}{m_a!} \prod_{1 \leq a \leq b \leq 3} \frac{(1-v_{ab}^{*6})^{m'_{ab}}}{m'_{ab}!}. \end{aligned}$$

Now, we redefine variables such that $m_{23} \rightarrow m_{23}+m'_{23}$ and $m_{33} \rightarrow m_{33}+m'_{33}$, and we define $m_{33} = m-m_{23}$ to evaluate the sums over m_{23} , m , and m_{13} , always with the help of (A.1),

$$\begin{aligned} G_{6P}^* &= \sum \frac{(-h_2)_{m_1+m_2-m_3+m'_{12}+m'_{22}}(-h_5)_{m_3+m'_{13}+m'_{23}+m'_{33}}(\bar{p}_3+h_2+h_5)_{m_3}}{(\bar{p}_3+h_2)_{2m_3+m'_{13}+m'_{23}+m'_{33}}(\bar{p}_3+h_2+1-d/2)_{m_3}} \\ & \times \frac{(-h_3)_{m_1+m_{11}}(p_2+h_2)_{m_1-m_2+m_3+m'_{13}}(p_3)_{m_2-m_1+m_3+m'_{23}+m'_{33}}}{(p_2)_{2m_1+m_{11}+m_{12}+m'_{13}}(p_2+1-d/2)_{m_1}} \\ & \times \frac{(-h_2)_{m_1+m_2-m_3+m_{11}+m_{12}}(p_3-h_2+h_4)_{m_2+m'_{12}+m'_{33}}(-h_4)_{m_2+m'_{22}+m'_{23}}(p_2+h_3)_{m_1+m_{12}+m'_{13}}}{(p_3-h_2)_{2m_2+m'_{12}+m'_{22}+m'_{23}+m'_{33}}(p_3-h_2+1-d/2)_{m_2}} \frac{(-h_2)_{m_1+m_2-m_3+m_{12}}}{(-h_2)_{m_1+m_2-m_3+m_{12}}} \\ & \times \frac{(-m'_{11})_{m_{11}}(-h_2-m_3+m_1+m_2+m_{11})_{m'_{11}-m_{11}}(-m'_{12})_{m_{12}}}{m_{11}!} \frac{(-m'_{12})_{m_{12}}}{m_{12}!} \\ & \times F_6^* \prod_{1 \leq a \leq 3} \frac{(u_a^{*6})^{m_a}}{m_a!} \prod_{1 \leq a \leq b \leq 3} \frac{(1-v_{ab}^{*6})^{m'_{ab}}}{m'_{ab}!}. \end{aligned}$$

The summation over m_{11} corresponds to a ${}_3F_2$ -hypergeometric function which can be transformed with the help of (A.3), leading to

$$\begin{aligned} & {}_3F_2 \left[\begin{matrix} -m'_{11}, -h_3+m_1, -h_2+m_1+m_2-m_3+m_{12} \\ p_2+2m_1+m_{12}+m'_{13}, -h_2+m_1+m_2-m_3 \end{matrix}; 1 \right] = \frac{(p_2+h_3+m_1+m_{12}+m'_{13})_{m'_{11}}}{(p_2+2m_1+m_{12}+m'_{13})_{m'_{11}}} \\ & \times {}_3F_2 \left[\begin{matrix} -m'_{11}, -h_3+m_1, -m_{12} \\ 1-p_2-h_3-m_1-m_{12}-m'_{11}-m'_{13}, -h_2+m_1+m_2-m_3 \end{matrix}; 1 \right]. \end{aligned}$$

Hence, we find that

$$\begin{aligned}
 G_{6P}^* &= \sum \frac{(-h_2)_{m_1+m_2-m_3+m'_{12}+m'_{22}}(-h_5)_{m_3+m'_{13}+m'_{23}+m'_{33}}(\bar{p}_3+h_2+h_5)_{m_3}}{(\bar{p}_3+h_2)_{2m_3+m'_{13}+m'_{23}+m'_{33}}(\bar{p}_3+h_2+1-d/2)_{m_3}} \\
 &\times \frac{(-h_3)_{m_1+m_{11}}(p_2+h_2)_{m_1-m_2+m_3+m'_{13}}(p_3)_{m_2-m_1+m_3+m'_{23}+m'_{33}}}{(p_2)_{2m_1+m'_{11}+m_{12}+m'_{13}}(p_2+1-d/2)_{m_1}} \\
 &\times \frac{(-1)^{m_{11}}(-m_{12})_{m_{11}}(-h_4)_{m_2+m'_{22}+m'_{23}}(p_2+h_3)_{m_1+m_{12}-m_{11}+m'_{11}+m'_{13}}(p_3-h_2+h_4)_{m_2+m'_{12}+m'_{33}}}{(p_3-h_2)_{2m_2+m'_{12}+m'_{22}+m'_{23}+m'_{33}}(p_3-h_2+1-d/2)_{m_2}} \\
 &\times \frac{(-m'_{11})_{m_{11}}(-h_2-m_3+m_1+m_2+m_{11})_{m'_{11}-m_{11}}(-m'_{12})_{m_{12}}}{m_{11}! m_{12}!} \\
 &\times F_6^* \prod_{1 \leq a \leq 3} \frac{(u_a^{*6})^{m_a}}{m_a!} \prod_{1 \leq a \leq b \leq 3} \frac{(1-v_{ab}^{*6})^{m'_{ab}}}{m'_{ab}!}.
 \end{aligned}$$

After evaluating the summation over m_{12} (with $m_{12} \rightarrow m_{12}+m_{11}$ first) with the help of (A.1), the result becomes

$$\begin{aligned}
 G_{6P}^* &= \sum \frac{(-h_2)_{m_1+m_2-m_3+m'_{12}+m'_{22}}(-h_5)_{m_3+m'_{13}+m'_{23}+m'_{33}}(\bar{p}_3+h_2+h_5)_{m_3}}{(\bar{p}_3+h_2)_{2m_3+m'_{13}+m'_{23}+m'_{33}}(\bar{p}_3+h_2+1-d/2)_{m_3}} \\
 &\times \frac{(-h_3)_{m_1+m'_{12}}(p_2+h_2)_{m_1-m_2+m_3+m'_{13}}(p_3)_{m_2-m_1+m_3+m'_{23}+m'_{33}}}{(p_2)_{2m_1+m'_{11}+m'_{12}+m'_{13}}(p_2+1-d/2)_{m_1}} \\
 &\times \frac{(-h_4)_{m_2+m'_{22}+m'_{23}}(p_2+h_3)_{m_1+m'_{11}+m'_{13}}(p_3-h_2+h_4)_{m_2+m'_{12}+m'_{33}}}{(p_3-h_2)_{2m_2+m'_{12}+m'_{22}+m'_{23}+m'_{33}}(p_3-h_2+1-d/2)_{m_2}} \\
 &\times \frac{(-m'_{12})_{m_{11}}(-m'_{11})_{m_{11}}(-h_2-m_3+m_1+m_2+m_{11})_{m'_{11}-m_{11}}}{m_{11}!} \\
 &\times F_6^* \prod_{1 \leq a \leq 3} \frac{(u_a^{*6})^{m_a}}{m_a!} \prod_{1 \leq a \leq b \leq 3} \frac{(1-v_{ab}^{*6})^{m'_{ab}}}{m'_{ab}!}.
 \end{aligned}$$

Finally, we evaluate the summation over m_{11} and the result is

$$\begin{aligned}
 G_{6P}^* &= \sum \frac{(-h_2)_{m_1+m_2-m_3+m'_{12}+m'_{22}}(-h_5)_{m_3+m'_{13}+m'_{23}+m'_{33}}(\bar{p}_3+h_2+h_5)_{m_3}}{(\bar{p}_3+h_2)_{2m_3+m'_{13}+m'_{23}+m'_{33}}(\bar{p}_3+h_2+1-d/2)_{m_3}} \\
 &\frac{(p_3-h_2+h_4)_{m_2+m'_{12}+m'_{33}}(-h_3)_{m_1+m'_{12}}(p_2+h_2)_{m_1-m_2+m_3+m'_{13}}(p_3)_{m_2-m_1+m_3+m'_{23}+m'_{33}}}{(p_2)_{2m_1+m'_{11}+m'_{12}+m'_{13}}(p_2+1-d/2)_{m_1}} \\
 &\frac{(-h_2)_{m_1+m_2-m_3+m'_{11}+m'_{12}}(-h_4)_{m_2+m'_{22}+m'_{23}}(p_2+h_3)_{m_1+m'_{11}+m'_{13}}}{(-h_2)_{m_1+m_2-m_3+m'_{12}}(p_3-h_2)_{2m_2+m'_{12}+m'_{22}+m'_{23}+m'_{33}}(p_3-h_2+1-d/2)_{m_2}} \\
 &\times F_6^* \prod_{1 \leq a \leq 3} \frac{(u_a^{*6})^{m_a}}{m_a!} \prod_{1 \leq a \leq b \leq 3} \frac{(1-v_{ab}^{*6})^{m'_{ab}}}{m'_{ab}!},
 \end{aligned}$$

proving (C.4) and thus (3.4).

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] F.A. Dolan and H. Osborn, *Conformal partial waves and the operator product expansion*, *Nucl. Phys. B* **678** (2004) 491 [[hep-th/0309180](#)] [[INSPIRE](#)].
- [2] F.A. Dolan and H. Osborn, *Conformal Partial Waves: Further Mathematical Results*, [arXiv:1108.6194](#) [[INSPIRE](#)].
- [3] P. Kravchuk, *Casimir recursion relations for general conformal blocks*, *JHEP* **02** (2018) 011 [[arXiv:1709.05347](#)] [[INSPIRE](#)].
- [4] S. Ferrara and G. Parisi, *Conformal covariant correlation functions*, *Nucl. Phys. B* **42** (1972) 281 [[INSPIRE](#)].
- [5] S. Ferrara, A.F. Grillo, G. Parisi and R. Gatto, *The shadow operator formalism for conformal algebra. Vacuum expectation values and operator products*, *Lett. Nuovo Cim.* **4** (1972) 115 [[INSPIRE](#)].
- [6] D. Simmons-Duffin, *Projectors, Shadows, and Conformal Blocks*, *JHEP* **04** (2014) 146 [[arXiv:1204.3894](#)] [[INSPIRE](#)].
- [7] D. Karateev, P. Kravchuk and D. Simmons-Duffin, *Weight Shifting Operators and Conformal Blocks*, *JHEP* **02** (2018) 081 [[arXiv:1706.07813](#)] [[INSPIRE](#)].
- [8] M.S. Costa and T. Hansen, *AdS Weight Shifting Operators*, *JHEP* **09** (2018) 040 [[arXiv:1805.01492](#)] [[INSPIRE](#)].
- [9] M. Isachenkov and V. Schomerus, *Superintegrability of d-dimensional Conformal Blocks*, *Phys. Rev. Lett.* **117** (2016) 071602 [[arXiv:1602.01858](#)] [[INSPIRE](#)].
- [10] V. Schomerus, E. Sobko and M. Isachenkov, *Harmony of Spinning Conformal Blocks*, *JHEP* **03** (2017) 085 [[arXiv:1612.02479](#)] [[INSPIRE](#)].
- [11] V. Schomerus and E. Sobko, *From Spinning Conformal Blocks to Matrix Calogero-Sutherland Models*, *JHEP* **04** (2018) 052 [[arXiv:1711.02022](#)] [[INSPIRE](#)].
- [12] M. Isachenkov and V. Schomerus, *Integrability of conformal blocks. Part I. Calogero-Sutherland scattering theory*, *JHEP* **07** (2018) 180 [[arXiv:1711.06609](#)] [[INSPIRE](#)].
- [13] I. Burić, V. Schomerus and M. Isachenkov, *Conformal Group Theory of Tensor Structures*, *JHEP* **10** (2020) 004 [[arXiv:1910.08099](#)] [[INSPIRE](#)].
- [14] E. Hijano, P. Kraus, E. Perlmutter and R. Snively, *Witten Diagrams Revisited: The AdS Geometry of Conformal Blocks*, *JHEP* **01** (2016) 146 [[arXiv:1508.00501](#)] [[INSPIRE](#)].
- [15] M. Nishida and K. Tamaoka, *Geodesic Witten diagrams with an external spinning field*, *Prog. Theor. Exp. Phys.* **2017** (2017) 053B06 [[arXiv:1609.04563](#)] [[INSPIRE](#)].
- [16] A. Castro, E. Lladrés and F. Rejon-Barrera, *Geodesic Diagrams, Gravitational Interactions and OPE Structures*, *JHEP* **06** (2017) 099 [[arXiv:1702.06128](#)] [[INSPIRE](#)].
- [17] E. Dyer, D.Z. Freedman and J. Sully, *Spinning Geodesic Witten Diagrams*, *JHEP* **11** (2017) 060 [[arXiv:1702.06139](#)] [[INSPIRE](#)].
- [18] H.-Y. Chen, E.-J. Kuo and H. Kyono, *Anatomy of Geodesic Witten Diagrams*, *JHEP* **05** (2017) 070 [[arXiv:1702.08818](#)] [[INSPIRE](#)].
- [19] C. Sleight and M. Taronna, *Spinning Witten Diagrams*, *JHEP* **06** (2017) 100 [[arXiv:1702.08619](#)] [[INSPIRE](#)].

- [20] S. Ferrara, A.F. Grillo and R. Gatto, *Manifestly conformal covariant operator-product expansion*, *Lett. Nuovo Cim.* **2** (1971) 1363 [INSPIRE].
- [21] S. Ferrara, R. Gatto and A.F. Grillo, *Conformal invariance on the light cone and canonical dimensions*, *Nucl. Phys. B* **34** (1971) 349 [INSPIRE].
- [22] S. Ferrara, A.F. Grillo and R. Gatto, *Manifestly conformal-covariant expansion on the light cone*, *Phys. Rev. D* **5** (1972) 3102 [INSPIRE].
- [23] S. Ferrara, P. Gatto and A.F. Grilla, *Conformal algebra in space-time and operator product expansion*, in *Springer Tracts in Modern Physics* **67**, Springer (1973), pp. 1–64 [INSPIRE].
- [24] S. Ferrara, A.F. Grillo, G. Parisi and R. Gatto, *Covariant expansion of the conformal four-point function*, *Nucl. Phys. B* **49** (1972) 77 [Erratum *ibid.* **53** (1973) 643] [INSPIRE].
- [25] S. Ferrara, A.F. Grillo, R. Gatto and G. Parisi, *Analyticity properties and asymptotic expansions of conformal covariant Green's functions*, *Nuovo Cim. A* **19** (1974) 667 [INSPIRE].
- [26] F.A. Dolan and H. Osborn, *Conformal four point functions and the operator product expansion*, *Nucl. Phys. B* **599** (2001) 459 [hep-th/0011040] [INSPIRE].
- [27] J.-F. Fortin and W. Skiba, *Conformal Bootstrap in Embedding Space*, *Phys. Rev. D* **93** (2016) 105047 [arXiv:1602.05794] [INSPIRE].
- [28] J.-F. Fortin and W. Skiba, *Conformal Differential Operator in Embedding Space and its Applications*, *JHEP* **07** (2019) 093 [arXiv:1612.08672] [INSPIRE].
- [29] V. Comeau, J.-F. Fortin and W. Skiba, *Further Results on a Function Relevant for Conformal Blocks*, arXiv:1902.08598 [INSPIRE].
- [30] J.-F. Fortin and W. Skiba, *A recipe for conformal blocks*, arXiv:1905.00036 [INSPIRE].
- [31] J.-F. Fortin and W. Skiba, *New methods for conformal correlation functions*, *JHEP* **06** (2020) 028 [arXiv:1905.00434] [INSPIRE].
- [32] J.-F. Fortin, V. Prilepina and W. Skiba, *Conformal two-point correlation functions from the operator product expansion*, *JHEP* **04** (2020) 114 [arXiv:1906.12349] [INSPIRE].
- [33] J.-F. Fortin, V. Prilepina and W. Skiba, *Conformal Three-Point Correlation Functions from the Operator Product Expansion*, arXiv:1907.08599 [INSPIRE].
- [34] J.-F. Fortin, V. Prilepina and W. Skiba, *Conformal Four-Point Correlation Functions from the Operator Product Expansion*, *JHEP* **08** (2020) 115 [arXiv:1907.10506] [INSPIRE].
- [35] J.-F. Fortin, W.-J. Ma, V. Prilepina and W. Skiba, *Efficient Rules for All Conformal Blocks*, arXiv:2002.09007 [INSPIRE].
- [36] S. Ferrara, A.F. Grillo and R. Gatto, *Tensor representations of conformal algebra and conformally covariant operator product expansion*, *Annals Phys.* **76** (1973) 161 [INSPIRE].
- [37] A.M. Polyakov, *Nonhamiltonian approach to conformal quantum field theory*, *Zh. Eksp. Teor. Fiz.* **66** (1974) 23 [*Sov. Phys. JETP* **39** (1974) 9] [INSPIRE].
- [38] K.B. Alkalaev and V.A. Belavin, *From global to heavy-light: 5-point conformal blocks*, *JHEP* **03** (2016) 184 [arXiv:1512.07627] [INSPIRE].
- [39] V. Rosenhaus, *Multipoint Conformal Blocks in the Comb Channel*, *JHEP* **02** (2019) 142 [arXiv:1810.03244] [INSPIRE].
- [40] V. Gonçalves, R. Pereira and X. Zhou, *20' Five-Point Function from $AdS_5 \times S^5$ Supergravity*, *JHEP* **10** (2019) 247 [arXiv:1906.05305] [INSPIRE].

- [41] S. Parikh, *Holographic dual of the five-point conformal block*, *JHEP* **05** (2019) 051 [[arXiv:1901.01267](#)] [[INSPIRE](#)].
- [42] C.B. Jepsen and S. Parikh, *Propagator identities, holographic conformal blocks, and higher-point AdS diagrams*, *JHEP* **10** (2019) 268 [[arXiv:1906.08405](#)] [[INSPIRE](#)].
- [43] S. Parikh, *A multipoint conformal block chain in d dimensions*, *JHEP* **05** (2020) 120 [[arXiv:1911.09190](#)] [[INSPIRE](#)].
- [44] J.-F. Fortin, W. Ma and W. Skiba, *Higher-Point Conformal Blocks in the Comb Channel*, *JHEP* **07** (2020) 213 [[arXiv:1911.11046](#)] [[INSPIRE](#)].
- [45] H. Exton, *Multiple hypergeometric functions and applications*, in *Mathematics & its Applications*, Halsted Press, Chichester U.K. (1976).
- [46] H.M. Srivastava and P.W. Karlsson, *Multiple Gaussian Hypergeometric Series*, in *Ellis Horwood Series in Mathematics and its Applications*, Halsted Press, Chichester U.K. (1985).