RECEIVED: September 1, 2021

REVISED: October 24, 2021

ACCEPTED: October 25, 2021

PUBLISHED: November 18, 2021

Hidden relations of central charges and OPEs in holographic CFT

Yue-Zhou Li,^{a,b} H. Lü^{a,c} and Liang Ma^a

^aCenter for Joint Quantum Studies and Department of Physics, School of Science,
Tianjin University,
Tianjin 300350, China

^bDepartment of Physics, McGill University,
3600 Rue University, Montréal, QC Canada

^cPeng Huanwu Center for Fundamental Theory,
Hefei, Anhui 230026, China

E-mail: liyuezhou@physics.mcgill.ca, mrhonglu@gmail.com,
liangma@tju.edu.cn

ABSTRACT: It is known that the (a, c) central charges in four-dimensional CFTs are linear combinations of the three independent OPE coefficients of the stress-tensor three-point function. In this paper, we adopt the holographic approach using AdS gravity as an effect field theory and consider higher-order corrections up to and including the cubic Riemann tensor invariants. We derive the holographic central charges and OPE coefficients and show that they are invariant under the metric field redefinition. We further discover a hidden relation among the OPE coefficients that two of them can be expressed in terms of the third using differential operators, which are the unit radial vector and the Laplacian of a four-dimensional hyperbolic space whose radial variable is an appropriate length parameter that is invariant under the field redefinition. Furthermore, we prove that the consequential relation $c = 1/3\ell_{\text{eff}}\partial a/\partial\ell_{\text{eff}}$ and its higher-dimensional generalization are valid for massless AdS gravity constructed from the most general Riemann tensor invariants.

KEYWORDS: AdS-CFT Correspondence, Gauge-gravity correspondence, Classical Theories of Gravity, Conformal Field Theory

ARXIV EPRINT: [2108.05384](https://arxiv.org/abs/2108.05384)

Contents

1	Introduction	1
2	Holographic central charges	3
2.1	Generalities of higher derivative gravity	3
2.2	General formula of holographic central charges	4
2.3	Proof of the central charge relation	6
3	Holographic CFT as AdS EFT	6
3.1	General arguments	6
3.2	Example: to the cubic order and OPE relations	7
3.3	Further comments: a -theorem and functional relations?	11
4	Conclusion	12
A	Relation of central charges in general dimensions	13
A.1	Holographic central charges	13
A.2	Complete the proof of the relation	15
A.3	Two explicit examples	15
A.3.1	A polynomial example	16
A.3.2	A fractional example	17
B	Field redefinitions and invariant central charges	17
B.1	Effective AdS	17
B.2	Field redefinitions	18
B.3	Relation of central charges	19
C	Stress-tensor three-point functions	21
C.1	Three-point structures	21
C.2	OPEs in $d = 4$	22

1 Introduction

In a curved space, a conformal field theory (CFT) suffers from an anomaly that the trace of the stress tensor acquires a nonvanishing expectation value [1, 2]. In $d = 4$, this conformal anomaly reads (we discard the Maxwell part)

$$\langle T_i^i \rangle = -\frac{a}{16\pi^2} E^{(4)} + \frac{c}{16\pi^2} I^{(4)}, \quad (1.1)$$

where $E^{(4)}$ and $I^{(4)}$ represent the Euler density and the Weyl tensor squared of the geometric background respectively. Central charges a and c are important characteristics of

the CFT, especially for the stress-tensor sector. Specifically, a -charge measures massless degrees of freedom and is the value of the a -function evaluated at the fixed point [3–5], and c -charge is proportional to the canonical normalization of stress-tensor two-point function [6], i.e., $C_T|_{d=4} = 40/\pi^4 c$, where

$$\langle T_{ij}(x)T_{kl}(y) \rangle = \frac{C_T \mathcal{I}_{ijkl}(x-y)}{(x-y)^{2d}}. \quad (1.2)$$

They encode the coefficient λ_{TTT} of the operator product expansion (OPE) of the stress-tensor [7], can further probe the conformal collider physics [8, 9] and the averaged null energy condition (ANEC) [10].

In $\mathcal{N} = 4$ super Yang-Mills (SYM) theory, the two central charges are identical, but they are not in general the same, e.g., $\mathcal{N} = 1, 2$ supersymmetric theories [8]. It is insightful to study the large N CFTs with sparse gap ($\Delta_{\text{gap}} \gg 1$), which are expected to have weakly-coupled local gravity duals in AdS [11]. The central charges are encoded in the pure gravity sector, based on the holographic dictionary [12–14]. The causality analysis at the Regge limit in AdS gravity imposes CEMZ bound [15]

$$\frac{|a-c|}{c} \lesssim \frac{1}{\Delta_{\text{gap}}^2}, \quad (1.3)$$

which was later obtained purely from the sparse CFTs [16–18] and extended to more general three-point functions (e.g. [19–21]).

The above known relations among the central charges and also the OPEs are algebraic in nature. By studying “massless” higher-curvature gravities that have only massless graviton in AdS, a linear differential relation was proposed and it survived many checks [22]. It is

$$c = \frac{1}{3} \ell_{\text{eff}} \frac{\partial a}{\partial \ell_{\text{eff}}}, \quad (1.4)$$

where ℓ_{eff} is the (effective) radius of the AdS vacuum. In this paper, we consider pure gravity of the most general Riemann tensor invariants and derive the explicit general holographic formulae for the (a, c) charges that enable us to prove this relation.

AdS gravity that corresponds to the large- N and sparse CFT can also be approached as an effective field theory (EFT) and higher-order curvature invariants arise order-by-order perturbatively. In this EFT approach, there are no massive modes within the appropriate cutoff and therefore it is necessarily to consider all possible such terms. However, we actually have less nontrivial higher-order terms since there is a redundancy that one can perform field redefinitions of the metric in terms of curvature tensors order-by-order without altering the physics. (This is referred to as the “equivalence theorem” [23].) It is thus useful to express the relation eq. (1.4) in terms of the length parameter $\ell_{\text{inv}}^{(0)}$ that is invariant under the field redefinition, namely

$$c = \frac{1}{3} \ell_{\text{inv}}^{(0)} \frac{\partial a}{\partial \ell_{\text{inv}}^{(0)}}. \quad (1.5)$$

We consider AdS gravity up to and including the cubic order of the Riemann tensor polynomials. At this order, we can derive the three OPE coefficients λ_{TTT} as well as the (a, c)

charges holographically. We verify that they are indeed invariant under the field redefinitions, and therefore so is eq. (1.5). In doing so, we confirm the known result that the central charges are linear combinations of the OPE coefficients. Furthermore, we discover further hidden differential relations among the OPE coefficients by differential operators that live in the four-dimensional hyperbolic space whose radial coordinate is $\ell_{\text{inv}}^{(0)}$.

The paper is organized as follows. In section 2, we consider pure gravity constructed from general Riemann tensor invariants. With the assumption that the theory admits an AdS vacuum, we derive the explicit formulae for (a, c) central charges and derive the differential relation (1.4) for massless gravities. In section 3, we treat the higher-order curvature invariants perturbatively and consider the most general polynomial invariants up to and including the cubic order. We derive the proper $\ell_{\text{inv}}^{(0)}$ that is invariant under the field redefinition and becomes ℓ_{eff} when the theory reduces to massless gravity. We prove the relation (1.5). We find further differential relations among the OPE coefficients of the three-point function of the stress tensor. We conclude the paper in section 4. In appendix A, we generalise our four-dimensional results to general dimensions. In appendix B, we present the field redefinitions in general dimensions and derive the invariant length parameters. In appendix C, we study the OPE coefficients of the three-point function of the stress tensor.

2 Holographic central charges

2.1 Generalities of higher derivative gravity

For our purpose, we consider pure gravity theories constructed from general Riemann tensor invariants in $(d + 1)$ spacetime dimensions:

$$S = \frac{1}{16\pi G_N^{(d+1)}} \int d^{d+1}x \sqrt{-g} \mathcal{L}(R_{\mu\nu\rho\sigma}, g^{\mu\nu}). \quad (2.1)$$

We assume that the theory admits an AdS vacuum of radius ℓ_{eff} . It is instructive to introduce tensors $P_{\mu\nu\rho\sigma}$ and $C_{\alpha\beta\gamma\eta}^{\mu\nu\rho\sigma}$ [24, 25]

$$P_{\mu\nu\rho\sigma} = \frac{\partial \mathcal{L}}{\partial R^{\mu\nu\rho\sigma}} \Big|_{g_{\alpha\beta}}, \quad C_{\alpha\beta\gamma\eta}^{\mu\nu\rho\sigma} = \frac{\partial P^{\mu\nu\rho\sigma}}{\partial R^{\alpha\beta\gamma\eta}}. \quad (2.2)$$

When evaluated on the vacuum, the tensor structures are rigidly fixed

$$\begin{aligned} \bar{P}^{\mu\nu\rho\sigma} &= 2\xi_0 \bar{g}^{\mu[\rho} \bar{g}^{\sigma]\nu}, \\ \bar{C}_{\alpha\beta\gamma\eta}^{\mu\nu\rho\sigma} &= \xi_1 \left(\delta_\alpha^{[\mu} \delta_\beta^{\nu]} \delta_\gamma^{[\rho} \delta_\eta^{\sigma]} + \delta_\alpha^{[\rho} \delta_\beta^{\sigma]} \delta_\gamma^{[\mu} \delta_\eta^{\nu]} \right) + \xi_2 \left((\bar{g}^{\mu\rho} \bar{g}^{\nu\sigma} - \bar{g}^{\mu\sigma} \bar{g}^{\nu\rho}) (\bar{g}_{\alpha\gamma} \bar{g}_{\beta\eta} - \bar{g}_{\alpha\eta} \bar{g}_{\beta\gamma}) \right) \\ &\quad + 4\xi_3 \left(\delta_\tau^{[\mu} \bar{g}^{\nu]\rho} \delta_\varepsilon^{[\sigma} \delta_{[\alpha}^{\tau} \bar{g}_{\beta]\gamma} \delta_{[\eta]}^{\varepsilon]} \right). \end{aligned} \quad (2.3)$$

In holography, as we will see shortly, the coefficients $\xi_{i=0\sim 3}$ contain all the information pertaining to the central charges and hence OPEs λ_{TTT} . Some useful identities were

proved [26] and we quote them here

$$\bar{\mathcal{L}}(\ell_{\text{eff}}) = -\frac{4d}{\ell_{\text{eff}}^2} \xi_0, \quad \bar{\mathcal{L}}'(\ell_{\text{eff}}) = \frac{4d(d+1)}{\ell_{\text{eff}}^3} \xi_0, \quad (2.4)$$

$$\mathfrak{h}(\ell_{\text{eff}}) \equiv \bar{\mathcal{L}}(\ell_{\text{eff}}) + \frac{\ell_{\text{eff}}}{d+1} \bar{\mathcal{L}}'(\ell_{\text{eff}}) = 0, \quad (2.5)$$

$$\mathfrak{h}'(\ell_{\text{eff}}) = \frac{4d(4\xi_1 + 4d\xi_3 + 4d(d+1)\xi_2 + (d-1)\xi_0\ell_{\text{eff}}^2)}{\ell_{\text{eff}}^5}. \quad (2.6)$$

Here, a prime denotes a derivative with respect to ℓ_{eff} , but with subtleties that should be clarified. To be precise, the Lagrangian in (2.1) should be written as $\mathcal{L} = \mathcal{L}(R_{\mu\nu\rho\sigma}, g^{\mu\nu}, \alpha)$ where α denotes all the coupling constants, including the bare cosmological constant Λ_0 . Off shell, these coupling constants are all independent of ℓ_{eff} , and they become related by the on-shell condition, namely the left equation in (2.4). The derivative is implemented off shell, but assuming that the metric is AdS of radius ℓ_{eff} , namely [24, 25]

$$\mathcal{L}'(\ell_{\text{eff}}) = \frac{\partial \bar{\mathcal{L}}}{\partial \bar{R}^{\mu\nu\rho\sigma}} \frac{\partial \bar{R}^{\mu\nu\rho\sigma}}{\partial \ell_{\text{eff}}}. \quad (2.7)$$

We consider in general theories with a bare cosmological constant and we solve the on-shell condition by express the Λ_0 in terms of ℓ_{eff} and other coupling constants. It is clear that the derivative in (2.7) does not explicitly involve in Λ_0 . The rule for higher derivatives follows straightforwardly.

Gravity of Riemann invariants is in general a 4'th-order derivative theory and the spectrum contains not only the usual massless graviton $h_{ab}^{(0)}$, but also a massive spin-2 $h_{ab}^{(M)}$ and a massive scalar h . The effective Newton constant $G_{\text{Neff}}^{(d+1)}$ and the masses of additional modes can be expressed in terms of $\xi_{i=0\sim 3}$ [24, 25]

$$\frac{1}{G_{\text{Neff}}^{(d+1)}} = \frac{2}{G_N^{(d+1)}} \left(\xi_0 + \frac{2}{\ell_{\text{eff}}^2} (d-2) \xi_1 \right), \quad (2.8)$$

$$m_s^2 = \frac{\xi_0 \ell_{\text{eff}}^2 (d-1) + 4(\xi_1 + d(d+1)\xi_2 + d\xi_3)}{(2\xi_1 + (d+1)\xi_3 + 4d\xi_2) \ell_{\text{eff}}^2}, \quad (2.9)$$

$$m_g^2 = -\frac{\xi_0 \ell_{\text{eff}}^2 + 2(d-2)\xi_1}{(2\xi_1 + \xi_3) \ell_{\text{eff}}^2}. \quad (2.10)$$

It is important to note that first equation above is the coefficient of the kinetic term of the graviton and hence it does not explicitly depend on the bare cosmological constant Λ_0 . In our proof of eq. (1.4), we need to decouple both massive modes by requiring

$$2\xi_1 + (d+1)\xi_3 + 4d\xi_2 = 0, \quad 2\xi_1 + \xi_3 = 0. \quad (2.11)$$

The resulting theory is phrased as massless gravity in [22] that include quasi-topological gravities [27–30].

2.2 General formula of holographic central charges

The conformal anomaly in CFT_4 can be reconstructed from the AdS bulk by considering FG expansion [31, 32]

$$ds^2 = \frac{\ell_{\text{eff}}^2}{4\rho^2} d\rho^2 + \frac{1}{\rho} g_{ij} dx^i dx^j, \quad g_{ij} = g_{0ij} + g_{1ij}\rho + g_{2ij}\rho^2 + \dots, \quad (2.12)$$

where $\rho \rightarrow 0$ is the AdS boundary. We introduce the ultraviolet (UV) cutoff parameter $\epsilon > 0$ and evaluate the boundary integrals at $\rho = \epsilon$. Correspondingly, the action can be expanded as

$$S = \frac{1}{16\pi G_N^{(5)}} \int_{\rho \rightarrow \epsilon} d^5x \sqrt{-\hat{g}_5} \mathcal{L} = \frac{1}{16\pi G_N^{(5)}} \int d^4x \sqrt{-g_0} \int_{\rho \rightarrow \epsilon} d\rho \left(\cdots + \frac{A}{\rho} + \cdots \right), \quad (2.13)$$

where the coefficient A of ρ^{-1} is interpreted as conformal anomaly, because ρ^{-1} gives rise to a logarithmic dependence of the UV cutoff

$$S_{\text{anomaly}} = - \int d^4x \sqrt{-g_0} \frac{1}{2} \langle T_i^i \rangle \log \epsilon, \quad (2.14)$$

which manifestly breaks the scaling invariance.

For convenience, we consider a reduced FG expansion: we explicitly construct an AdS with $S^2 \times S^2$ boundary [33]

$$ds^2 = \frac{\ell_{\text{eff}}^2}{4\rho^2} d\rho^2 + \frac{1}{\rho} \left(f_1(\rho) d\Omega_1^2 + f_2(\rho) d\Omega_2^2 \right), \quad (2.15)$$

and (f_1, f_2) can be closely solved as the truncated FG expansion

$$f_1 = f_{10} + f_{11}\rho + f_{12}\rho^2 + \cdots, \quad f_2 = f_{20} + f_{21}\rho + f_{22}\rho^2 + \cdots, \quad (2.16)$$

where

$$f_{11} = \frac{1}{6}(\gamma - 2)\ell_{\text{eff}}^2, \quad f_{21} = \frac{(1 - 2\gamma)\ell_{\text{eff}}^2}{6\gamma}. \quad (2.17)$$

The constants f_{10} and f_{20} simply represent the radii of the two spheres. We define the ratio $\gamma = f_{10}/f_{20}$, but take $f_{10} = f_{20} = 1$. The Euler density gives the topological number of $S^2 \times S^2$, whilst the Weyl-squared on the boundary depends on γ :

$$E^{(4)} = 8, \quad I^{(4)} = \frac{4}{3} \left(2 + \gamma + \frac{1}{\gamma} \right). \quad (2.18)$$

We can then readily find

$$A = \frac{1 + \gamma^2}{6\gamma} \ell_{\text{eff}}^3 \left(\xi_0 + \frac{4\xi_1}{\ell_{\text{eff}}^2} \right) - \frac{2}{3} \ell_{\text{eff}}^3 \left(\xi_0 - \frac{2\xi_1}{\ell_{\text{eff}}^2} \right), \quad (2.19)$$

from which we can read off the central charges

$$a = \frac{\pi \xi_0 \ell_{\text{eff}}^3}{4G_N^{(5)}}, \quad c = \frac{\pi \ell_{\text{eff}}^3}{4G_N^{(5)}} \left(\xi_0 + \frac{4\xi_1}{\ell_{\text{eff}}^2} \right) = \frac{\pi \ell_{\text{eff}}^3}{8G_{\text{Neff}}^{(5)}}. \quad (2.20)$$

We can then show generally $C_T|_{d=4} = 40/\pi^4 c$ in holography since C_T is necessarily proportional to $1/G_{\text{Neff}}^{(5)}$. (For explicit low-lying examples, see e.g., [34, 35].) In appendix A, we will generalize our results to arbitrary dimensions and prove generally that the a -charge above can be read off from the entanglement entropy of spherical entangling surfaces, demonstrated in some explicit low-lying examples [36–38].

2.3 Proof of the central charge relation

The identities (2.6) and (2.8) imply that the relation between $G_{\text{Neff}}^{(5)}$ and $\mathfrak{h}'(\ell_{\text{eff}})$:

$$\frac{1}{G_{\text{Neff}}^{(5)}} = \frac{\ell_{\text{eff}}^3}{24G_N^{(5)}} \left(\mathfrak{h}'(\ell_{\text{eff}}) + \frac{128}{\ell_{\text{eff}}^2} (\xi_1 - 10\xi_2 - 2\xi_3) \right). \quad (2.21)$$

The term $\mathfrak{h}'(\ell_{\text{eff}})$ can also be obtained by a derivative of eq. (2.5), followed by substituting eq. (2.4). We have

$$\begin{aligned} c &= \frac{\pi \ell_{\text{eff}}^3}{12G_N^{(5)}} \left(3\xi_0 + \ell_{\text{eff}} \frac{\partial \xi_0}{\partial \ell_{\text{eff}}} + \frac{8}{\ell_{\text{eff}}^2} (\xi_1 - 10\xi_2 - 2\xi_3) \right) \\ &= \frac{\pi}{12G_N^{(5)}} \ell_{\text{eff}} \frac{\partial (\xi_0 \ell_{\text{eff}}^3)}{\partial \ell_{\text{eff}}} + \frac{2\pi}{3G_N^{(5)}} (\xi_1 - 10\xi_2 - 2\xi_3) \\ &= \frac{1}{3} \ell_{\text{eff}} \frac{\partial a}{\partial \ell_{\text{eff}}} + \frac{2\pi}{3G_N^{(5)}} (\xi_1 - 10\xi_2 - 2\xi_3). \end{aligned} \quad (2.22)$$

The second term above vanishes after imposing the massless conditions (2.11). We therefore prove the universal relation (1.4) of holographic central charges for massless AdS gravities. It should be emphasized that since the massless conditions (2.11) in general involve ℓ_{eff} , we should take the ℓ_{eff} derivative of a before imposing the massless conditions such that the relevant coupling constants are independent of ℓ_{eff} .

In appendix A, we generalize the relation to arbitrary dimensions and the subtleties of the ℓ_{eff} derivative are further clarified with explicit examples.

3 Holographic CFT as AdS EFT

3.1 General arguments

In this section, we use the EFT approach to the weakly-coupled bulk gravity, where higher-order Riemann tensor polynomials are perturbative corrections to Einstein gravity with a bare negative cosmological constant

$$\Lambda_0 = -\frac{d(d-1)}{2\ell_0^2}. \quad (3.1)$$

For the theory to be valid below the energy scale $M \gg 1/\ell_0$, graviton is the only light particle and the massive states are all beyond M , i.e.,

$$\begin{aligned} 2\xi_1 + (d+1)\xi_3 + 4d\xi_2 &= \frac{\#_1}{M^2} + \frac{\#_2}{M^4\ell_0^2} + \cdots, \\ 2\xi_1 + \xi_3 &= \frac{\#'_1}{M^2} + \frac{\#'_2}{M^4\ell_0^2} + \cdots. \end{aligned} \quad (3.2)$$

Furthermore, we assume the following hierarchy [39]

$$\frac{1}{G_N^{(d+1)}} \gg M^{d-1} \gg \frac{1}{\ell_0^{d-1}}, \quad \ell_0 \sim \mathcal{O}(1). \quad (3.3)$$

We consider only the tree-level gravitational physics, of the leading order in $\mathcal{O}(G_N^{(d+1)})$, which is dual to the large C_T ($C_T \sim N^2$) limit of the boundary CFT. Schematically, we may write the effective Lagrangian as (we only consider parity-even gravity)

$$\mathcal{L}(R_{\mu\nu\rho\sigma}, g^{\mu\nu}) = R - 2\Lambda_0 + \sum_{n=2} \sum_i g_{n,i} [R_i^{(n)}], \quad (3.4)$$

where $g_{n,i}$'s are Wilson coefficients associated with higher derivative terms $[R_i^{(n)}]$ that denote i 'th curvature invariant operators with $2n$ derivatives. In flat space, dimensional analysis indicates the scaling behavior $g_{n,i} \sim 1/M^{2(n-1)}$; however, it becomes more complicated in AdS. The subtlety arises from the fact that $g_{(n+1),i}$ can also enter the coupling g_n under field redefinitions. Thus the most general power-counting for each $g_{n,i}$ involves all $1/(M^{2(n-1-k)}\ell_0^{2k})$ terms with $k = 0, 1, 2, \dots$.

In this framework, $\#$ and $\#'$ are dimensionless quantities that depend on Wilson coefficients of higher derivative terms; however, they are not invariant under the field redefinition, and neither does ℓ_0 . It is always possible to find a field redefinition that brings eq. (3.2) to eq. (2.11) [29]. As physical observables, central charges should be invariant under the field redefinitions. It is thus natural to expect the relation eq. (1.4) should have a general form for AdS EFTs, where ℓ_{eff} is replaced by a certain field-redefinition invariant quantity $\ell_{\text{inv}}^{(0)}$ that coincides with ℓ_{eff} for massless gravity. This leads to eq. (1.5).

3.2 Example: to the cubic order and OPE relations

For a concrete example, we truncate the AdS EFT to the cubic order and show the field redefinition invariance of central charges and the validity of the relation eq. (1.5). Cubic is also the minimum order that fully enumerates three-point structures of the stress tensor. (We shall return to this point later.) The complete set of quadratic and cubic operators is

$$\begin{aligned} [R^{(2)}] &= \left\{ R^2, R_{\mu\nu}R^{\mu\nu}, R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma} \right\}, \\ [R^{(3)}] &= \left\{ R^3, RR_{\mu\nu}R^{\mu\nu}, R_{\nu}^{\mu}R_{\rho}^{\nu}R_{\mu}^{\rho}, R_{\mu\nu\rho\sigma}R^{\mu\rho}R^{\nu\sigma}, RR_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}, R^{\mu\nu}R_{\mu\rho\sigma\eta}R_{\nu}^{\rho\sigma\eta}, \right. \\ &\quad \left. R^{\mu\nu}_{\rho\sigma}R^{\rho\sigma}_{\alpha\beta}R^{\alpha\beta}_{\mu\nu}, R^{\mu}_{\nu}R^{\rho}_{\sigma}R^{\nu}_{\alpha}R^{\sigma}_{\beta}R^{\alpha}_{\mu}R^{\beta}_{\rho}, R_{\mu\nu}\square R^{\mu\nu}, R\square R \right\}. \end{aligned} \quad (3.5)$$

The contributions to central charges from this set were obtained in [33] and we review them in appendix B. To the cubic order, the Wilson coefficients can be expressed in terms of three sets of dimensionless parameters ($q_i, \tilde{q}_i, c_i$):

$$\begin{aligned} g_{2,i} &= \frac{q_i}{M^2} + \frac{\tilde{q}_i}{M^4\ell_0^2}, & i &= 1, 2, 3, \\ g_{3,i} &= \frac{c_i}{M^4}, & i &= 1, 2, \dots, 10. \end{aligned} \quad (3.6)$$

We now show the rigidity of central charges under the general field redefinition:

$$g_{\mu\nu} \rightarrow d_0 g_{\mu\nu} + \sum_i d_{1i} [R_i^{(1)}]_{\mu\nu} + \sum_i d_{2i} [R_i^{(2)}]_{\mu\nu}, \quad (3.7)$$

where the d_n coefficients are given explicitly in appendix B, and

$$\begin{aligned} [R^{(1)}]_{\mu\nu} &= \{R_{\mu\nu}, Rg_{\mu\nu}\}, \\ [R^{(2)}]_{\mu\nu} &= \left\{ R_{\mu\rho\sigma\gamma} R_{\nu}{}^{\rho\sigma\gamma}, R_{\mu\rho\nu\sigma} R^{\rho\sigma}, R_{\mu\nu} R, R_{\mu\rho} R_{\nu}^{\rho}, R^2 g_{\mu\nu}, \right. \\ &\quad \left. R_{\rho\sigma} R^{\rho\sigma} g_{\mu\nu}, R_{\rho\sigma\gamma\eta} R^{\rho\sigma\gamma\eta} g_{\mu\nu}, \nabla_{\mu} \nabla_{\nu} R, \square R_{\mu\nu}, \square R g_{\mu\nu} \right\}. \end{aligned} \quad (3.8)$$

Note that $g_{\mu\nu}$ on the r.h.s. of eq. (3.7) shall be further scaled by d_0 such that the Einstein Hilbert term is invariant under the field redefinition. (There is no such scaling needed for the Minkowski vacuum.) The consequence is that we have to introduce $\tilde{q}_i$ to absorb the descendants of the higher-order c_i terms under the field redefinitions. Under this scheme, the bare ℓ_0 must also vary under the field redefinition. The explicit rule how Wilson coefficients vary is recorded in appendix B (see eq. (B.9)), and here we simply quote our findings. The coefficients q_3 , c_7 and c_8 are explicitly invariant, and there is one more invariant combination

$$\tilde{q}'_3 \equiv \tilde{q}_3 - 4(5c_5 + c_6) + \frac{8}{3}(5q_1 + q_2)q_3. \quad (3.9)$$

We can construct six invariant length parameters

$$\begin{aligned} \ell_{\text{inv}}^{(1)} &= \ell_0 - \frac{10(5q_1 + q_2)}{3M^2\ell_0} - \frac{10(5\tilde{q}_1 + \tilde{q}_2)}{3M^4\ell_0^3} + \frac{10(4(25c_1 + 5c_2 + c_3 + c_4) - 3(5q_1 + q_2)^2)}{3M^4\ell_0^3}, \\ \ell_{\text{inv}}^{(2)} &= \frac{q_3}{M^2\ell_0} + \frac{10(5q_1 + q_2)q_3}{3M^4\ell_0^3}, \quad \ell_{\text{inv}}^{(3)} = \frac{\tilde{q}'_3}{M^4\ell_0^3}, \\ \ell_{\text{inv}}^{(4)} &= \frac{q_3^2}{M^4\ell_0^3}, \quad \ell_{\text{inv}}^{(5)} = \frac{c_7}{M^4\ell_0^3}, \quad \ell_{\text{inv}}^{(6)} = \frac{c_8}{M^4\ell_0^3}. \end{aligned} \quad (3.10)$$

Any linear combinations of the above could be the invariant ‘‘AdS radius’’ ℓ_{inv} , as long as the leading term is ℓ_0 . Thus we find that the central charges are indeed invariant, but there is an ambiguity in the expression:

$$a = \frac{\ell_{\text{inv}}^3}{G_N^{(5)}} \left(\tilde{\#}_1 + \frac{\tilde{\#}_2}{M^2\ell_{\text{inv}}^2} + \frac{\tilde{\#}_3}{M^4\ell_{\text{inv}}^4} \right), \quad c = \frac{\ell_{\text{inv}}^3}{G_N^{(5)}} \left(\tilde{\#}'_1 + \frac{\tilde{\#}'_2}{M^2\ell_{\text{inv}}^2} + \frac{\tilde{\#}'_3}{M^4\ell_{\text{inv}}^4} \right), \quad (3.11)$$

where $\tilde{\#}_i$ and $\tilde{\#}'_i$ are invariant dimensionless coefficients, depending on the choice of a fiducial length parameter. Part of this ambiguity stems from the choice of the OPE basis for the stress tensor, which we will discuss presently. For now, we simply aim to find a particular ℓ_{inv} that validates (1.5). The logic is straightforward, we would like to make a field redefinition, bring gravity theory to massless gravity with standard hierarchy $g_{2,i} \sim 1/M^2$, $g_{3,i} \sim 1/M^4$ (i.e., $\tilde{q}_i = 0$). To satisfy the differential relation for massless gravity, ℓ_{inv} is fixed to be $\ell_{\text{eff}}|_{\text{massless}}$. (In this order-by-order approach, we impose the massless condition at each order that gives two ℓ_{eff} -independent linear relations of the coupling constants. The subtlety issue of $\partial/\partial_{\text{eff}}$ of the general Riemann tensor theory does not arise.) This procedure uniquely determines

$$\ell_{\text{inv}}^{(0)} = \left(1, \frac{7}{3}, \frac{1}{3}, -\frac{7}{6}, -\frac{26}{3}, \frac{11}{2} \right) \cdot \ell_{\text{inv}}^{(i)}, \quad (3.12)$$

Figure 1. Einstein, quadratic and cubic terms enumerate the stress-tensor three-point structures.

where the dot $\cdot$ denotes the internal product of two vectors. Adopting this $\ell_{\text{inv}}^{(0)}$ for the (a, c) expressions, we find that the invariant dimensionless coefficients are

$$\begin{aligned}\tilde{\#}_1 = \tilde{\#}'_1 = 1, \quad \tilde{\#}_2 = 3\tilde{\#}'_2 = -12q_3, \\ \tilde{\#}_3 = -3\tilde{\#}'_3 = 16q_3^2 - 6\tilde{q}'_3 + 36c_7 - 9c_8.\end{aligned}\tag{3.13}$$

The specific ratios $\tilde{\#}_i/\tilde{\#}'_i$ above lead to the differential relation eq. (1.5).

It is known in CFTs that central charges can be expressed in terms of λ_{TTT} , i.e., OPEs of $\langle TTT \rangle$ [7]. According to conformal symmetry, there are only three parity-even conformal invariant structures in $\langle TTT \rangle$. (In $d = 3$, there are only two structures.) The three-point function $\langle TTT \rangle$ is thus determined by the three tensor structures multiplied by the intrinsic OPE λ_{TTT} . The three structures are enumerated by Einstein, quadratic and cubic gravity vertices, as depicted in figure 1. All higher derivative terms either repeat these λ_{TTT} 's or encode higher point contact structures. For this reason, we may argue that the results we obtain about the OPE coefficients may be universal.

One can think of λ_{TTT} as a three-dimensional vector, and there is ambiguity in the choice of the OPE basis, which can explain part of the choices in defining ℓ_{inv} , since λ_{TTT} must be expressible in terms of these invariant lengths. However, there are only three OPEs, but we have five free parameters to define a seemingly valid ℓ_{inv} . In fact, one can immediately see that the coefficients of $\ell_{\text{inv}}^{(2)}$ and $\ell_{\text{inv}}^{(3)}$ must be equal in the three-point function from the bulk Witten diagram, because q_3 and $\tilde{q}_3$ combine to give $g_{2,3}$. However, the field redefinition appears to allow different coefficients. In other words, there exist ℓ_{inv} 's that cannot express any OPE by ℓ_{inv}^3 . More specifically, those ℓ_{inv} 's that can represent OPEs via ℓ_{inv}^3 have to include the combination

$$\# \left(\ell_{\text{inv}}^{(2)} + \ell_{\text{inv}}^{(3)} \right) - \frac{1}{18} \left(5 + 6\# + 18\#^2 \right) \ell_{\text{inv}}^{(4)},\tag{3.14}$$

where $\#$ is pure number. To resolve this issue, we note first that at the perturbative order considered, we have $\ell_{\text{inv}}^{(4)} = (\ell_{\text{inv}}^{(2)})^2/\ell_0$, and therefore, $\ell_{\text{inv}}^{(4)}$ should not be considered as an independent length parameter. Secondly, we may also take the view that the $\tilde{q}_i$ terms are introduced passively in order to absorb the descendants of the higher order terms from the field redefinition. We should therefore set the invariant quantity $\tilde{q}'_3$ to zero to avoid introducing artificial variables. This leads to the precise balancing between three independent length parameters and three OPE coefficients. In fact we can use the field redefinition from the beginning to remove all terms associated with the Ricci tensor and

Ricci scalar, in which case, we have only three nontrivial parameters $(q_3, 2c_7 + c_8, \ell_{\text{eff}})$. (In five bulk dimensions, the cubic Lovelock gravity is trivial for which we have $2c_7 + c_8 = 0$.) Our approach of using ℓ_{inv} yields the same result, but makes the procedure more covariant under the field redefinition.

To elaborate this further, we note that the three-point basis proposed in [7] is often used in the literature, e.g., [34, 35, 40–42], where OPEs are denoted as $\mathcal{A}, \mathcal{B}, \mathcal{C}$, and in $d = 4$ one has [7]

$$\begin{aligned} a &= \frac{\pi^6}{2880} (13\mathcal{A} - 2\mathcal{B} - 40\mathcal{C}) \equiv \tilde{\mathcal{A}}, \\ c &= \frac{\pi^6}{480} (9\mathcal{A} - \mathcal{B} - 10\mathcal{C}) \equiv \tilde{\mathcal{C}}. \end{aligned} \quad (3.15)$$

OPEs $\mathcal{A}, \mathcal{B}, \mathcal{C}$ can be explicitly calculated in our example by using ANEC operator [8], which are recorded in appendix C. These results establish holographically the relations in (3.15).

Another natural choice is the orthogonal basis [43, 44], where the orthogonality is defined with respect to three-point pairing [45], because the orthogonal basis in the flat-space limit precisely corresponds to R , $[R^{(2)}]$ and $[R^{(3)}]$ respectively. The details of these structures are lengthy and we leave them recorded in appendix C. In general dimensions $d > 3$, we find

$$\begin{aligned} \lambda_{TTT}^{(1)} &= \frac{-(\mathcal{A}(d^2 + d - 2)) + 4\mathcal{C}(d + 1) + 2\mathcal{B}}{d(d^2 - 4)}, \\ \lambda_{TTT}^{(2)} &= \frac{2(d - 1)(2\mathcal{A} + 4\mathcal{C} - \mathcal{B})}{(d - 2)d(d + 3)}, \\ \lambda_{TTT}^{(3)} &= \frac{\mathcal{A}(-d^2 + d + 4) + 4\mathcal{C}(d + 1) - 2d\mathcal{B}}{d(-d^2 + d + 2)}. \end{aligned} \quad (3.16)$$

In the orthogonal basis, we can immediately observe $\lambda_{TTT}^{(1)} \propto c$. We can thus rotate to a $\ell_{\text{inv}}^{\text{orth}}$

$$\ell_{\text{inv}}^{\text{orth}} = \left(1, 1, 1, -\frac{29}{18}, -\frac{38}{3}, \frac{13}{2}\right) \cdot \ell_{\text{inv}}^{(i)}, \quad (3.17)$$

for which

$$c = \frac{\pi(\ell_{\text{inv}}^{\text{orth}})^3}{8G_N^{(5)}} = -\frac{\pi^6}{20} \lambda_{TTT}^{(1)}. \quad (3.18)$$

The a -charge is a bit more complicated, with

$$\tilde{\#}_1 = 1, \quad \tilde{\#}_2 = -8q_3, \quad \tilde{\#}_3 = \frac{4}{3} (37c_7 - 9c_8 - 6\tilde{q}_3' + 16q_3^2), \quad (3.19)$$

which satisfies

$$a = -\frac{\pi^6}{2160} (54\lambda_{TTT}^{(1)} + 35\lambda_{TTT}^{(2)} + 15\lambda_{TTT}^{(3)}). \quad (3.20)$$

Regardless the basis, the linear relations among a , c and three λ_{TTT} 's show that we may treat three λ_{TTT} 's as independent, and (a, c) charges are derived quantities. The detail coefficients of the relation depend on the basis that we would like to choose.

Hidden relation emerges when we choose a specific invariant length $\ell_{\text{inv}}^{(0)}$. It leads to a linear differential relation between (a, c) . We can further establish a new differential relation between a -charge and the third linearly independent combination of the OPE coefficients¹

$$\tilde{\mathcal{B}} \equiv \frac{\pi^6}{2016} (9\mathcal{A} - 2\mathcal{B} - 2\mathcal{C}) = \square_{\ell_{\text{inv}}^{(0)}} a, \quad (3.21)$$

where

$$\nabla_\ell \equiv \frac{1}{3}\ell \frac{\partial}{\partial \ell}, \quad \square_\ell \equiv \nabla_\ell (\nabla_\ell - 1). \quad (3.22)$$

(See appendix C for detail derivation.) This relation, together with eq. (1.5) have a profound implication: even through stress-tensor OPEs are algebraically independent, they are related to each other via differential operators from the holographic point of view! Furthermore, we have a geometric implication of these differential relations. The differential operator $\square_\ell$ is a Laplacian of the four-dimensional hyperbolic space of radius $1/3$, with radial coordinate ℓ :

$$ds_4^2 = \frac{9(d\ell^2 + dx^2 + dy^2 + dz^2)}{\ell^2}, \quad (3.23)$$

which is Euclidean AdS_4 in planar coordinates. In particular, ∇_ℓ is the radial vector of unit length in the hyperbolic space. We expect that there is a systematic way to generate more differential relations among OPEs, and they might be organized under the hidden symmetry associated with the isometry group of the metric (3.23), and we leave the construction of this interesting systematics for future study.

3.3 Further comments: a -theorem and functional relations?

Holographic analysis indicates that there is differential relation between central charges. It can be established for general Riemann tensor gravity or for perturbative EFT. Perturbatively, the relation is established by differentiating with $\ell_{\text{inv}}^{(0)}$ that asymptotes ℓ_0 in the $M \rightarrow \infty$ limit. However CFTs *a priori* do not know the parameter ℓ_0 , it is therefore hard to make sense of the differential relation from the CFT perspective. Here we argue that an analogous differential relation can be established if we extend central charges to radius-dependent functions by deforming away from AdS, exactly following the same approach as addressing the holographic a -theorem [46]. In this approach, one simply replaces $\ell_{\text{inv}}^{(0)}$ by $A'(r)$, where $r \rightarrow \infty$ is the AdS boundary and $A(r)$ flows to $\ell_{\text{inv}}^{(0)}$ in pure AdS. We find the following differential relation is valid

$$\frac{da(r)}{dr} = 3c(r) \frac{A''(r)}{A'(r)}, \quad (3.24)$$

which can be deduced from eq. (3.24) by using the differential chain rule. As was proved in [47], at the conformal perturbation level to all loops, one has

$$\frac{da(\mu)}{d\mu} = G_{ij} \beta^i \beta^j, \quad (3.25)$$

¹We are grateful to Andrs Anabalon for pointing out that the $\tilde{\mathcal{B}}$ in our first version is not independent, but equal to $(c - a)/9$.

where $G_{ij} > 0$ and β^i 's are β -functions. Here μ is the energy scale that flows to UV and corresponds to r in the AdS bulk. This is exactly the a -theorem, and we conclude

$$G_{ij}\beta^i\beta^j = 3c(\mu)\frac{A''(\mu)}{A'(\mu)}, \quad (3.26)$$

which is consistent because of $A''(r)|_{\text{AdS}} = 0$. Since $c(r) > 0$ for any unitary theory, we thus find holographic a -theorem is validated provided $A''(r)/A'(r) > 0$. It is interesting to study the relation between $A''(r)/A'(r) > 0$ and the null energy condition in a generic AdS EFT in the future. More nontrivially, eq. (3.21) then implies a novel differential relation among the OPE $\tilde{\mathcal{B}}$ -function and the (a, c) functions, which are themselves OPE functions:

$$c'(\mu) - a'(\mu) = \frac{a'(\mu)}{c(\mu)}\tilde{\mathcal{B}}(\mu). \quad (3.27)$$

4 Conclusion

Two related topics were addressed in this paper. One is that we proved the differential relation (1.4) of [22] and its higher-dimensional generalization for massless gravities that are constructed from the most general Riemann tensor invariants. The second is to approach this relation perturbatively, order by order. This allowed us to find more differential relations among the OPE coefficients of the three-point function $\langle TTT \rangle$ of the stress tensor.

We considered AdS gravity extended with general Riemann tensor invariants up to and including the cubic order. We showed that the (a, c) central charges and three OPE coefficients $(\mathcal{A}, \mathcal{B}, \mathcal{C})$ were all invariant under field redefinitions of the metric. We reconfirmed the known fact that both central charges could be expressed as certain linear combinations of the OPE coefficients. By recombining these OPE coefficients to $(\tilde{\mathcal{A}}, \tilde{\mathcal{B}}, \tilde{\mathcal{C}})$, we find that two of them can be expressed as differentials of the third:

$$\tilde{\mathcal{C}} = \nabla_{\ell_{\text{inv}}^{(0)}}\tilde{\mathcal{A}}, \quad \tilde{\mathcal{B}} = \square_{\ell_{\text{inv}}^{(0)}}\tilde{\mathcal{A}}. \quad (4.1)$$

where ∇_{ℓ} and $\square_{\ell}$ are the unit radial vector and the Laplacian of the four-dimensional hyperbolic space (3.23) respectively. The central charge differential relation (1.5) is the consequence of the first relation above. However, there is a difference: the relations in (4.1) are perturbative results since OPE coefficients can only be defined perturbatively. The central-charge relation (1.5), on the other hand, can be promoted to (1.4) for general massless gravities whose higher-order couplings are not necessarily small. The hidden relations tantalizingly suggest that there may exist a deep organizing principle for the correlation functions of CFTs, associated with the isometry group of the metric (3.23), which can be viewed as Euclidean AdS₄ in planar coordinates.

However, the understanding of the differential relations from the CFT perspective remains illusive, since $\ell_{\text{inv}}^{(0)}$ does not have an immediate translation to any quantity in CFTs. We argue that we can use the analogous approach in the holographic a -theorem and obtain the corresponding differential relations of (a, c) functions but also those of the OPE coefficients. However, the subject requires further investigation.

Acknowledgments

We would like to thank Zhan-Feng Mai for useful conversation and initial collaboration on proof of relation of central charges, and Simon Caron-Huot, Alexander Maloney for discussions. The work is supported in part by the National Natural Science Foundation of China (NSFC) grants No. 11875200 and No. 11935009. Y.-Z.L. is also supported in part by the Fonds de Recherche du Québec - Nature et Technologies and by the Simons Collaboration on the Nonperturbative Bootstrap. H.L. and L.M. benefit also from NSFC grants No. 11947301 and No. 12047502.

Appendix

This appendix includes: generalization of the relation of central charges to general dimensions (section A), details of field redefinition of Wilson coefficients up to cubic order in general dimensions (section B), three-point structures of $\langle TTT \rangle$ in general dimensions and the calculation of corresponding OPEs from ANEC operator in $d = 4$ (section C).

A Relation of central charges in general dimensions

A.1 Holographic central charges

The differential relation of the central charges can be generalized to general dimensions $d \geq 3$. However, there is a subtlety since in higher dimensions, there are more central charges and they can be very complicated. Nevertheless, the central charge C_T and a can be universally defined. (The identification of C_T with a suitable c was only done holographically in [49] for $d = 6$.) It is also important to note even though there is no conformal anomaly in odd d , C_T is always well-defined

$$C_T = \frac{\pi^{-\frac{d}{2}-1} \Gamma(d+2)}{8(d-1) \Gamma\left(\frac{d}{2}\right)} \frac{\ell_{\text{eff}}^{d-1}}{G_{N\text{eff}}^{(d+1)}}. \quad (\text{A.1})$$

The substitute of a -charge in odd d is the entanglement entropy (EE) for spherical entangling surface S^{d-2} [38]. It is shown that EE for spherical entanglement surface can be mapped to thermal entropy of $R \times H^{d-1}$ via an appropriate conformal map. In holography, EE for half of S^{d-1} of $R \times S^{d-1}$ boundary of global AdS boils down to black hole entropy of hyperbolic topological black hole in $d+1$ -dimension [38]

$$ds_{d+1}^2 = -f dt^2 + \frac{dr^2}{f} + r^2 d\Omega_{d-1,k=-1}^2, \quad f = \frac{r^2}{\ell_{\text{eff}}^2} - 1, \quad (\text{A.2})$$

which is locally AdS. According to the Wald formalism [48], we have

$$S = -\frac{\omega_{k,d-1} \ell_{\text{eff}}^{d-1}}{8G_N^{(d+1)}} \left(P_{abcd} \epsilon^{ab} \epsilon^{cd} \right)_{r=\ell_{\text{eff}}}, \quad (\text{A.3})$$

where $\bar{P}_{abcd}\epsilon^{ab}\epsilon^{cd} = -4\xi_0$ and $\omega_{k,d-1}$ denotes the entangling surface and it is divergent in even d , . It is instructive to simply divide by $\omega_{k,d-1}$ and define the density

$$a^* = \frac{\Omega_{d-1}}{4\pi\omega_{k,d-1}} S = \frac{\pi^{\frac{d}{2}-1}\xi_0\ell_{\text{eff}}^{d-1}}{4\Gamma\left(\frac{d}{2}\right)G_N^{(d+1)}}, \quad (\text{A.4})$$

where Ω_{d-1} is the volume of $d-1$ -sphere. It is clear from the Wald formalism that a^* does not explicitly depend on the bare cosmological constant, which we solve for in terms of ℓ_{eff} and other coupling constants in the theory.

In even d , this reproduces a known fact that EE for spherical entangling surface is proportional to a -charge. To show this, we apply the reduced the FG trick in general even dimensions by restricting to S^{d-1} boundary topology.

$$ds^2 = \frac{\ell_{\text{eff}}^2}{4\rho^2} d\rho^2 + \frac{f(\rho)}{\rho} d\Omega_d^2, \quad (\text{A.5})$$

which kills all the Weyl invariants. We expand the action

$$S = \frac{1}{16\pi G_N^{(d+1)}} \int d^{d+1}x \sqrt{-\hat{g}_{d+1}} \mathcal{L} \quad (\text{A.6})$$

around the AdS boundary $\rho \rightarrow 0$

$$\mathcal{L} = \mathcal{L}_{\text{vac}} + \mathcal{L}_1|_{\rho \rightarrow 0} \rho + \mathcal{L}_2|_{\rho \rightarrow 0} \rho^2 + \cdots + \mathcal{L}_{\frac{d}{2}}|_{\rho \rightarrow 0} \rho^{\frac{d}{2}}, \quad (\text{A.7})$$

$$\sqrt{-\hat{g}_{d+1}} = \frac{\ell_{\text{eff}}}{2} \rho^{-\frac{d+2}{2}} \left(a_0 + a_1 \rho + a_2 \rho^2 + \cdots + a_{\frac{d}{2}} \rho^{\frac{d}{2}} \right) \sim \frac{\ell_{\text{eff}}}{2} \rho^{-\frac{d+2}{2}} f^{\frac{d}{2}} \quad (\text{A.8})$$

and then collect all the terms contributing to ρ^{-1} . The ansatz for f is

$$f = f_0 + f_1 \rho + f_2 \rho^2 + f_3 \rho^3 + \cdots + f_{\frac{d}{2}} \rho^{\frac{d}{2}}. \quad (\text{A.9})$$

Performing variation with respect to the coefficients f_i and then applying the variation principle, we find

$$f_1 = -\frac{\ell_{\text{eff}}^2}{2}, \quad f_2 = \frac{\ell_{\text{eff}}^4}{16f_0}, \quad f_i = 0, \quad 3 \leq i \leq \frac{d-2}{2}. \quad (\text{A.10})$$

Thus we have

$$\mathcal{L} = \mathcal{L}_{\text{vac}} + \mathcal{L}_{\frac{d}{2}} \rho^{\frac{d}{2}}, \quad \mathcal{L}_{\text{vac}} = -\frac{4d}{\ell_{\text{eff}}^2} \xi_0, \quad \mathcal{L}_{\frac{d}{2}} = \frac{2d^2}{\ell_{\text{eff}}^2} \frac{f_{\frac{d}{2}}}{f_0} \xi_0, \quad (\text{A.11})$$

$$a_{\frac{d}{2}} = -\frac{\ell_{\text{eff}}^d}{d} - \frac{(-4)^{\frac{d}{2}}}{2} \frac{\Gamma(\frac{d+2}{2})^2}{\Gamma(d+1)} f_{\frac{d}{2}} f_0^{\frac{d-2}{2}}, \quad a_0 = -\frac{(-4)^{\frac{d}{2}}}{d} \frac{\Gamma(\frac{d+2}{2})^2}{\Gamma(d+1)} f_0^{\frac{d}{2}}. \quad (\text{A.12})$$

We find precisely

$$a = a^*, \quad \text{for even } d. \quad (\text{A.13})$$

It is of interest to note that from this action procedure to calculate the a -charge, we need to know explicitly the on-shell condition $\mathcal{L}_{\text{vac}} = -\frac{4d}{\ell_{\text{eff}}^2} \xi_0$, while the equation (A.4) does not explicitly involve this relation.

A.2 Complete the proof of the relation

The proof is analogous to the $D = 5$ example presented in the main text. It follows from (2.6) and (2.8) that we have

$$\frac{1}{G_{\text{Neff}}^{(d+1)}} = \frac{\ell_{\text{eff}}^3}{2d(d-1)G_N^{(d+1)}} \left(\mathfrak{h}'(\ell_{\text{eff}}) + \frac{8d^2}{\ell_{\text{eff}}^5} \xi \right), \quad (\text{A.14})$$

where $\xi \equiv (d-3)\xi_1 - 2(d+1)\xi_2 - 2\xi_3$. We therefore have

$$C_T = \frac{\Omega_{d-1}\Gamma(d+2)}{16\pi^{d+1}(d-1)} \frac{\ell_{\text{eff}}^{d-1}}{G_{\text{Neff}}^{(d+1)}} = \frac{\Omega_{d-1}\Gamma(d+2)\ell_{\text{eff}}^{d+2}}{32\pi^{d+1}d(d-1)^2G_N^{(d+1)}} \left(\mathfrak{h}'(\ell_{\text{eff}}) + \frac{8d^2}{\ell_{\text{eff}}^5} \xi \right). \quad (\text{A.15})$$

Taking a derivative of eq. (2.5) gives:

$$\begin{aligned} C_T &= \frac{\Omega_{d-1}\Gamma(d+2)}{32\pi^{d+1}d(d-1)^2G_N^{(d+1)}} \left[\ell_{\text{eff}}^{d+2} \bar{\mathcal{L}}'(\ell_{\text{eff}}) - 8\xi_0 d \ell_{\text{eff}}^{d-1} + 4d \ell_{\text{eff}}^d \frac{\partial \xi_0}{\partial \ell_{\text{eff}}} + 8d^2 \ell_{\text{eff}}^{d-3} \xi \right] \\ &= \frac{\Gamma(d+2)}{\pi^d(d-1)^2} \left\{ \ell_{\text{eff}} \frac{\partial}{\partial \ell_{\text{eff}}} \left[\frac{\Omega_{d-1}\xi_0 \ell_{\text{eff}}^{d-1}}{8\pi G_N^{(d+1)}} \right] + \frac{\Omega_{d-1}d}{4\pi G_N^{(d+1)}} \ell_{\text{eff}}^{d-3} \xi \right\} \\ &= \frac{\Gamma(d+2)}{\pi^d(d-1)^2} \ell_{\text{eff}} \frac{\partial a^*}{\partial \ell_{\text{eff}}} + \frac{\Omega_{d-1}\Gamma(d+2)d}{4\pi^{d+1}(d-1)^2G_N^{(d+1)}} \xi. \end{aligned} \quad (\text{A.16})$$

After imposing the massless conditions that imply $\xi = 0$, we obtain the universal central charge relation

$$C_T = \frac{\Gamma(d+2)}{\pi^d(d-1)^2} \ell_{\text{eff}} \frac{\partial a^*}{\partial \ell_{\text{eff}}}. \quad (\text{A.17})$$

Again we should emphasize that the derivative with respect to ℓ_{eff} should be done before imposing the massless conditions such that all the coupling constants are independent of ℓ_{eff} . We shall illustrate this with two explicit examples in the next subsection.

This differential relation was first proposed in [22], and was independently noticed in [26], where a^* was replaced by a quantity from the free energy of sphere. As was explained in the main text, from the EFT approach of AdS gravity, we only have to replace ℓ_{eff} by a suitable $\ell_{\text{inv}}^{(0)}$ and the differential relation holds. We record $\ell_{\text{inv}}^{(0)}$ for the cubic example in appendix B.

A.3 Two explicit examples

Here, we address the subtleties of taking a derivative with respect to ℓ_{eff} using two explicit examples. In this paper we consider a general class of theories of the type

$$\mathcal{L} = R - 2\Lambda_0 + \tilde{\mathcal{L}}(R_{\mu\nu\rho\sigma}, g^{\mu\nu}, \alpha), \quad (\text{A.18})$$

where α denotes all the dimensionful coupling constants of the Riemann tensor invariants. We assume that the theory admits an AdS vacuum of radius ℓ_{eff} and the on-shell condition, i.e. the first equation in (2.4). This enables us to express Λ_0 as a function of ℓ_{eff} and α which is independent of ℓ_{eff} . Our derivative with respect to ℓ_{eff} is thus taken such that $\partial\alpha/\partial\ell_{\text{eff}} = 0$. (In fact, the derivatives of various quantities with respect to ℓ_{eff} do not involve Λ_0 explicitly either.) The massless conditions, on the other hand, relate coupling constants that could involve ℓ_{eff} and therefore the conditions should be imposed after take the derivatives.

A.3.1 A polynomial example

We first consider the Einstein-quadratic-cubic-AdS theory in five dimensions ($d = 4$). The theory was studied in detail in section 3 as an effective theory. Here we treat it as a classical theory where the coupling constants are not necessarily small. The three quadratic and eight cubic Riemann tensor invariants are summarized in (3.5). We use (q_1, q_2, q_3) , $(c_1, c_2, \dots, c_8)$ to denote the quadratic and cubic coupling constants respectively. We have

$$\Lambda_0 = -\frac{6}{\ell_{\text{eff}}^2} + \frac{4(10q_1 + 2q_2 + q_3)}{\ell_{\text{eff}}^4} + \frac{2(400c_1 + 80c_2 + 16c_3 + 16c_4 + 40c_5 + 8c_6 + 4c_7 + 3c_8)}{\ell_{\text{eff}}^6}, \quad (\text{A.19})$$

and (a, c) central charges

$$\begin{aligned} a &= \frac{\pi}{8G_N^{(5)}} \left(\ell_{\text{eff}}^3 - 4(10q_1 + 2q_2 + q_3)\ell_{\text{eff}} \right. \\ &\quad \left. + \frac{3(400c_1 + 80c_2 + 16c_3 + 16c_4 + 40c_5 + 8c_6 + 4c_7 + 3c_8)}{\ell_{\text{eff}}} \right), \\ c &= \frac{\pi}{8G_N^{(5)}} \left(\ell_{\text{eff}}^3 - 4(10q_1 + 2q_2 - q_3)\ell_{\text{eff}} \right. \\ &\quad \left. + \frac{1200c_1 + 240c_2 + 48c_3 + 48c_4 - 40c_5 - 8c_6 - 36c_7 + 21c_8}{\ell_{\text{eff}}} \right). \end{aligned} \quad (\text{A.20})$$

Taking a derivative of a with respect to ℓ_{eff} , while treating q_i and c_i couplings as independent, we have

$$\begin{aligned} c &= \frac{1}{3}\ell_{\text{eff}} \frac{\partial a}{\partial \ell_{\text{eff}}} + \frac{\pi}{3G_N^{(5)}\ell_{\text{eff}}} \xi, \\ \xi &= 3(200c_1 + 40c_2 + 8c_3 + 8c_4 - 4c_7 + 3c_8) - 2(5q_1 + q_2 - q_3)\ell_{\text{eff}}^2. \end{aligned} \quad (\text{A.21})$$

From the massless conditions

$$\begin{aligned} q_3 &= -q_1 - \frac{q_2}{2} + \frac{3}{2\ell_{\text{eff}}^2}(40c_1 + 12c_2 + 4c_3 + 3c_4 + 16c_5 + 4c_6 + 4c_7), \\ c_8 &= \frac{\ell_{\text{eff}}^2}{3}(4q_1 + q_2) - \frac{1}{3}(240c_1 + 52c_2 + 12c_3 + 11c_4 + 16c_5 + 4c_6), \end{aligned} \quad (\text{A.22})$$

we can easily check that $\xi = 0$, giving rise to (1.4). In the past literature [22, 33] the massless conditions are imposed at each order independently, such that one has two linear relations for q_i 's, as well as for c_i 's, which then do not involve ℓ_{eff} . One can therefore impose the massless conditions before taking the ℓ_{eff} derivatives in these cases. In the next, we present a nontrivial example where the massless conditions necessarily involve ℓ_{eff} .

A.3.2 A fractional example

Here we consider a more complicated fractional theory in five dimensions, with the Lagrangian

$$\mathcal{L} = R - 2\Lambda_0 + \frac{\alpha}{1 + \beta_1 R^2 + \beta_2 R_{\mu\nu} R^{\mu\nu} + \beta_3 R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma}}. \quad (\text{A.23})$$

The on-shell condition for the bare cosmological constant is

$$\Lambda_0 = -\frac{6}{\ell_{\text{eff}}^2} + \frac{\alpha \ell_{\text{eff}}^4 (720\beta_1 + 144\beta_2 + 72\beta_3 + \ell_{\text{eff}}^4)}{2(400\beta_1 + 80\beta_2 + 40\beta_3 + \ell_{\text{eff}}^4)^2}. \quad (\text{A.24})$$

Using the techniques outlined in the main text, we can easily obtain the (a, c) central charges:

$$\begin{aligned} a &= \frac{\pi}{8G_N^{(5)}} \left(\ell_{\text{eff}}^3 + \frac{4\alpha(10\beta_1 + 2\beta_2 + \beta_3)\ell_{\text{eff}}^9}{(\ell_{\text{eff}}^4 + 40(10\beta_1 + 2\beta_2 + \beta_3))^2} \right), \\ c &= \frac{\pi}{8G_N^{(5)}} \left(\ell_{\text{eff}}^3 + \frac{4\alpha(10\beta_1 + 2\beta_2 - \beta_3)\ell_{\text{eff}}^9}{(\ell_{\text{eff}}^4 + 40(10\beta_1 + 2\beta_2 + \beta_3))^2} \right). \end{aligned} \quad (\text{A.25})$$

Again these can be obtained without explicitly solving for Λ_0 . We therefore have

$$\begin{aligned} c &= \frac{1}{3}\ell_{\text{eff}} \frac{\partial a}{\partial \ell_{\text{eff}}} + \frac{2\pi\alpha\ell_{\text{eff}}^9}{3G_N^{(5)}(\ell_{\text{eff}}^4 + 40(10\beta_1 + 2\beta_2 + \beta_3))^3} \xi, \\ \xi &= (\beta_3 - 5\beta_1 - \beta_2)\ell_{\text{eff}}^4 + 120(50\beta_1^2 + 5(4\beta_2 + 3\beta_3)\beta_1 + 2\beta_2^2 + \beta_3^2 + 3\beta_2\beta_3). \end{aligned} \quad (\text{A.26})$$

In the above, the derivative with respect to ℓ_{eff} was taken by treating (α, β_i) parameters as being independent of ℓ_{eff} . The massless conditions are

$$\beta_2 = -4\beta_3, \quad \beta_1 = \frac{\ell_{\text{eff}}^4 + 1560\beta_3 + \sqrt{\ell_{\text{eff}}^8 - 1680\beta_3\ell_{\text{eff}}^4 + 14400\beta_3^2}}{2400}. \quad (\text{A.27})$$

We find that ξ vanishes identically under these conditions, giving rise to (1.5). In this example, we see that the massless condition will necessarily make the couplings depending on ℓ_{eff} and therefore, the derivative must be taken before the massless condition.

B Field redefinitions and invariant central charges

Here, we consider the perturbative approach to AdS gravity in general dimensions. We consider Einstein gravity with a negative bare cosmological constant, extended with Riemann tensor invariants (3.4), up to and including six derivatives total.

B.1 Effective AdS

First, we present the general holographic central charges of effective AdS of radius ℓ_{eff} . The equations of motion relate the bare cosmological constant Λ_0 and ℓ_{eff}

$$\begin{aligned} \Lambda_0 &= -\frac{d(d-1)}{2\ell_{\text{eff}}^2} + \frac{d(d-3)}{2\ell_{\text{eff}}^4} (d(d+1)g_{2,1} + dg_{2,2} + 2g_{2,3}) - \frac{d(d-5)}{2\ell_{\text{eff}}^6} \left(d^2(d+1)^2 g_{3,1} \right. \\ &\quad \left. + d^2(d+1)g_{3,2} + d^2g_{3,3} + d^2g_{3,4} + 2d(d+1)g_{3,5} + 2dg_{3,6} + 4g_{3,7} + (d-1)g_{3,8} \right). \end{aligned} \quad (\text{B.1})$$

a -charge and C_T charge are explicitly given as follows [33]

$$a = \frac{\Omega_{d-1}}{16\pi G_N^{(d+1)}} \left(\ell_{\text{eff}}^{d-1} - 2(d(d+1)g_{2,1} + dg_{2,2} + 2g_{2,3})\ell_{\text{eff}}^{d-3} + 3\left(d^2(d+1)^2g_{3,1} + d^2(d+1)g_{3,2} + d^2g_{3,3} + d^2g_{3,4} + 2d(d+1)g_{3,5} + 2dg_{3,6} + 4g_{3,7} + (d-1)g_{3,8}\right)\ell_{\text{eff}}^{d-5} \right), \quad (\text{B.2})$$

$$C_T = \frac{\Omega_{d-1}\Gamma(d+2)}{16\pi^{d+1}(d-1)G_N^{(d+1)}} \left(\ell_{\text{eff}}^{d-1} - 2(d(d+1)g_{2,1} + dg_{2,2} - 2(d-3)g_{2,3})\ell_{\text{eff}}^{d-3} + \left(3d^2(d+1)^2g_{3,1} + 3d^2(d+1)g_{3,2} + 3d^2g_{3,3} + 3d^2g_{3,4} - 2d(d+1)(2d-7)g_{3,5} - 2d(2d-7)g_{3,6} - 12(2d-5)g_{3,7} + 3(3d-5)g_{3,8}\right)\ell_{\text{eff}}^{d-5} \right). \quad (\text{B.3})$$

In $d = 4$, we have $C_T = 40/\pi^4 c$.

B.2 Field redefinitions

The general field redefinition that is relevant to the cubic-order of curvature tensor is given by (3.7) with

$$d_0 = 1 + \frac{\alpha_0}{M^2\ell_0^2} + \frac{\beta_0}{M^4\ell_0^4}, \quad d_{1i} = \frac{\alpha_i}{M^2} + \frac{\tilde{\alpha}_i}{M^4\ell_0^2}, \quad d_{2i} = \frac{\beta_i}{M^4}. \quad (\text{B.4})$$

Note that the scaling coefficient d_0 of the metric is to ensure that the Einstein Hilbert term is invariant under the field redefinition. This requires that

$$\alpha_0 = -d(\alpha_1 + (d+1)\alpha_2), \quad \beta_0 = -d(\tilde{\alpha}_1 + (d+1)\tilde{\alpha}_2) - \frac{1}{4}d^2(d+1)(\alpha_1 + (d+1)\alpha_2)^2, \quad (\text{B.5})$$

The bare cosmological constant is not invariant under the field redefinition; it is shifted by

$$\Lambda_0 \rightarrow \tilde{\Lambda} = \Lambda_0 + \frac{\Lambda_q}{M^2\ell_0^2} + \frac{\Lambda_c}{M^4\ell_0^4}, \quad (\text{B.6})$$

with

$$\Lambda_q = -\frac{d^2(d^2-1)(\alpha_1 + (d+1)\alpha_2)}{4\ell_0^2}, \quad \Lambda_c = -\frac{d^2(d^2-1)(\tilde{\alpha}_1 + (d+1)\tilde{\alpha}_2)}{4\ell_0^2} - \frac{d^3(d^2-1)(d+2)(\alpha_1 + (d+1)\alpha_2)^2}{8\ell_0^2}. \quad (\text{B.7})$$

Thus we have

$$\ell_0 \rightarrow \ell_0 + \frac{d(d+1)}{4M^2\ell_0}(\alpha_1 + (d+1)\alpha_2) - \frac{d^2(d-3)(d+1)}{32M^4\ell_0^3}(\alpha_1 + (d+1)\alpha_2)^2 + \frac{d(d+1)}{4M^4\ell_0^3}(\tilde{\alpha}_1 + (d+1)\tilde{\alpha}_2). \quad (\text{B.8})$$

Note that the coefficients $g_{n,i}$ are given by (3.6). The rule of the field redefinition for each coupling coefficients $g_{n,i}$ is explicitly presented below

$$\begin{aligned}
 q_1 &\rightarrow q_1 + \frac{1}{2}\alpha_1 + \frac{d-1}{2}\alpha_2, & q_2 &\rightarrow q_2 - \alpha_1, \\
 \tilde{q}_1 &\rightarrow \tilde{q}_1 + \frac{1}{2}\tilde{\alpha}_1 + \frac{d-1}{2}\tilde{\alpha}_2 + \frac{d(d-1)}{2}\beta_3 + \frac{d(d-1)(d+1)}{2}\beta_5 - \frac{d(d-3)}{2}\alpha_1q_1 \\
 &\quad - \frac{d(d-5)}{8}\alpha_1^2 - \frac{d(d-3)(d+1)}{2}\alpha_2q_1 - \frac{d(d^2-4d-1)}{4}\alpha_1\alpha_2 \\
 &\quad - \frac{d(d-1)(d-5)(d+1)}{8}\alpha_2^2, \\
 \tilde{q}_2 &\rightarrow \tilde{q}_2 - \tilde{\alpha}_1 + \frac{d(d-1)}{2}\beta_2 + \frac{d(d-1)}{2}\beta_4 + \frac{d(d-1)(d+1)}{2}\beta_6 \\
 &\quad - \frac{d(d-3)}{2}\alpha_1q_2 + \frac{d(d-5)}{4}\alpha_1^2 - \frac{d(d-3)(d+1)}{2}\alpha_2q_2 + \frac{d(d-3)(d+1)}{2}\alpha_1\alpha_2 \\
 \tilde{q}_3 &\rightarrow \tilde{q}_3 + \frac{d(d-1)}{2}\beta_1 + \frac{d(d-1)(d+1)}{2}\beta_7 - \frac{d(d-3)}{2}(\alpha_1 + (d+1)\alpha_2)q_3, \\
 c_1 &\rightarrow c_1 + \frac{1}{2}\beta_3 + \frac{d-1}{2}\beta_5 + \frac{1}{2}\alpha_1q_1 + \frac{1}{8}\alpha_1^2 + \frac{d-3}{2}\alpha_2q_1 + \frac{d-3}{4}\alpha_1\alpha_2 + \frac{(d-1)(d-3)}{8}\alpha_2^2, \\
 c_2 &\rightarrow c_2 + \frac{1}{2}\beta_2 - \beta_3 + \frac{1}{2}\beta_4 + \frac{d-1}{2}\beta_6 - \frac{1}{2}\alpha_1(4q_1 - q_2) - \frac{3}{4}\alpha_1^2 + \frac{d-3}{4}\alpha_2q_2 - \frac{d-3}{2}\alpha_1\alpha_2, \\
 c_3 &\rightarrow c_3 - \beta_4 + 4\alpha_1q_3 + \alpha_1^2, & c_4 &\rightarrow c_4 - \beta_2 - 2\alpha_1(q_2 + 2q_3), \\
 c_5 &\rightarrow c_5 + \frac{1}{2}\beta_1 + \frac{d-1}{2}\beta_7 + \frac{1}{2}(\alpha_1 + (d-3)\alpha_2)q_3, & c_6 &\rightarrow c_6 - \beta_1 - 2\alpha_1q_3, \\
 c_9 &\rightarrow c_9 - \beta_9 - \alpha_1(q_2 + 4q_3), \\
 c_{10} &\rightarrow c_{10} + \frac{1}{2}\beta_9 + \frac{d-1}{2}\beta_{10} - (\alpha_1 + 2d\alpha_2)q_1 - \frac{d+1}{2}\alpha_2q_2 + \alpha_1q_3 - 2\alpha_2q_3. \tag{B.9}
 \end{aligned}$$

It is not hard to use this rule to bring any cubic gravity to, e.g., quasi-topological gravity and Einsteinian cubic gravity, analogous to flat-space [29].

Invariant quantities include q_3, c_7, c_8 , together with

$$\tilde{q}'_3 = \tilde{q}_3 - d((d+1)c_5 + c_6) + \frac{2d(d-3)}{d-1}((d+1)q_1 + q_2)q_3. \tag{B.10}$$

The $d = 4$ case was present in the main text.

B.3 Relation of central charges

In the previous subsection, we obtain the transformation rules for each coupling coefficients, including the bare cosmological constant, under the general field redefinition. We are interested in showing that the physical quantities such as central charges are indeed invariant under the field redefinitions. It is clear that the coupling coefficients q_3, c_7, c_8 are manifestly invariant. For general extended gravity up to and including the cubic orders,

we find a total of six field-redefinition invariant lengths $\ell_{\text{inv}}^{(i)}$

$$\begin{aligned}\ell_{\text{inv}}^{(1)} &= \ell_0 - \frac{d(d+1)}{2(d-1)} \frac{((d+1)q_1 + q_2)}{M^2 \ell_0} - \frac{d(d+1)}{2(d-1)} \frac{((d+1)\tilde{q}_1 + \tilde{q}_2)}{M^4 \ell_0^3} \\ &\quad + \frac{d^2(d+1)}{2(d-1)} \frac{(d+1)^2 c_1 + (d+1)c_2 + c_3 + c_4 - \frac{(5d-11)}{4(d-1)}((d+1)q_1 + q_2)^2}{M^4 \ell_0^3}, \\ \ell_{\text{inv}}^{(2)} &= \frac{q_3}{M^2 \ell_0} + \frac{d(d+1)}{2(d-1)} \frac{((d+1)q_1 + q_2)q_3}{M^4 \ell_0^3}, \quad \ell_{\text{inv}}^{(3)} = \frac{\tilde{q}_3'}{M^4 \ell_0^3}, \\ \ell_{\text{inv}}^{(4)} &= \frac{q_3^2}{M^4 \ell_0^3}, \quad \ell_{\text{inv}}^{(5)} = \frac{c_7}{M^4 \ell_0^3}, \quad \ell_{\text{inv}}^{(6)} = \frac{c_8}{M^4 \ell_0^3}.\end{aligned}\tag{B.11}$$

From the perturbative point of view, the length scale ℓ_{inv} can be chosen to be $\ell_{\text{inv}}^{(1)}$ plus a linear combination of $\ell_{\text{inv}}^{(i)}$, ($i = 2, 3, 4, 5, 6$), with any numerical coefficients. Regardless the choice of ℓ_{inv} , we find that the central charges can always be expressed as

$$\begin{aligned}a &= \frac{\Omega_{d-1}}{16\pi G_N^{(d+1)}} \left(\ell_{\text{inv}}^{d-1} \tilde{\#}_1 + \ell_{\text{inv}}^{d-3} \frac{\tilde{\#}_2}{M^2} + \ell_{\text{inv}}^{d-5} \frac{\tilde{\#}_3}{M^4} \right), \\ C_T &= \frac{\Omega_{d-1} \Gamma(d+2)}{16\pi^{d+1} (d-1) G_N^{(d+1)}} \left(\ell_{\text{inv}}^{d-1} \tilde{\#}'_1 + \ell_{\text{inv}}^{d-3} \frac{\tilde{\#}'_2}{M^2} + \ell_{\text{inv}}^{d-5} \frac{\tilde{\#}'_3}{M^4} \right),\end{aligned}\tag{B.12}$$

where the coefficients $(\tilde{\#}_i, \tilde{\#}'_i)$ are invariant under the field redefinition, thereby proving that the central charges are invariant.

There exists a particular choice of ℓ_{inv} , which we name $\ell_{\text{inv}}^{(0)}$. For this choice, $\ell_{\text{inv}}^{(0)}$ reduces to ℓ_{eff} when the field redefinition brings us to massless gravity. It is given by

$$\begin{aligned}\ell_{\text{inv}}^{(0)} &= (y_1, y_2, y_3, y_4, y_5, y_6) \cdot \ell_{\text{inv}}^{(i)} \\ y_1 &= 1 \quad y_2 = \frac{(d-3)(2d-1)}{d-1}, \quad y_3 = \frac{d^2 - 4d + 1}{d-1}, \\ y_4 &= -\frac{(d-3)^2(2d-5)(2d-1)}{2(d-1)^2}, \quad y_5 = -\frac{2(3d^2 - 10d + 5)}{d-1}, \quad y_6 = \frac{1}{2}(4d-5).\end{aligned}\tag{B.13}$$

In terms of $\ell_{\text{inv}}^{(0)}$, we find

$$\begin{aligned}\tilde{\#}_1 &= \tilde{\#}'_1 = 1, \quad \tilde{\#}_2 = \frac{d-1}{d-3} \tilde{\#}'_2 = -2(d-1)(d-2)q_3, \\ \tilde{\#}_3 &= \frac{d-1}{d-5} \tilde{\#}'_3 = \frac{d-2}{2} \left(12(d-1)c_7 - 3(d-1)c_8 - 2(d-1)\tilde{q}_3' + 4d(d-3)^2 q_3^2 \right),\end{aligned}\tag{B.14}$$

which manifestly satisfies the relation

$$C_T = \frac{\Gamma(d+2)}{\pi^d (d-1)^2} \ell_{\text{inv}}^{(0)} \frac{\partial a^*}{\partial \ell_{\text{inv}}^{(0)}}.\tag{B.15}$$

Thus the hidden relation between C_T and a^* manifests itself when the invariant length $\ell_{\text{inv}}^{(0)}$ is chosen. As we saw in the main text, more hidden relations exist between the central charge and OPE coefficients under this choice of invariant length.

C Stress-tensor three-point functions

C.1 Three-point structures

Three-point function of stress-tensor $\langle TTT \rangle$ was first obtained in [7]. The expression is quite cumbersome, we contract the three-point function with graviton polarizations (which are null $\epsilon^2 = 0$), where the indices can be recovered by using Toda operator [50]. We further fix the conformal frame $(0, x, \infty)$ to simplify the structures

$$\begin{aligned}
 \langle TTT \rangle = & \frac{1}{2} \mathcal{A} x^{-d-4} \left(\epsilon_2 \cdot \epsilon_3 \left(2x^2 \epsilon_1 \cdot \epsilon_3 \left(x^2 \epsilon_1 \cdot \epsilon_2 - 2x \cdot \epsilon_1 x \cdot \epsilon_2 \right) - (d^2 - 4) (x \cdot \epsilon_1)^2 x \cdot \epsilon_2 x \cdot \epsilon_3 \right) \right. \\
 & + \left. (d^2 - 4) x \cdot \epsilon_1 x \cdot \epsilon_2 x \cdot \epsilon_3 (\epsilon_1 \cdot \epsilon_2 x \cdot \epsilon_3 - \epsilon_1 \cdot \epsilon_3 x \cdot \epsilon_2) \right) \\
 & + \frac{1}{4} \mathcal{B} x^{-d-6} \left(2x^2 \epsilon_2 \cdot \epsilon_3 x \cdot \epsilon_1 \left(x \cdot \epsilon_3 \left(-(d-2) x \cdot \epsilon_1 x \cdot \epsilon_2 - 2x^2 \epsilon_1 \cdot \epsilon_2 \right) + 2x^2 \epsilon_1 \cdot \epsilon_3 x \cdot \epsilon_2 \right) \right. \\
 & + x \cdot \epsilon_2 x \cdot \epsilon_3 \left(x \cdot \epsilon_1 x \cdot \epsilon_3 \left(2(d+2) x^2 \epsilon_1 \cdot \epsilon_2 + (d-2)(d+4) x \cdot \epsilon_1 x \cdot \epsilon_2 \right) \right. \\
 & - \left. \left. 2x^2 \epsilon_1 \cdot \epsilon_3 \left((d-2) x \cdot \epsilon_1 x \cdot \epsilon_2 + 2x^2 \epsilon_1 \cdot \epsilon_2 \right) \right) \right) \\
 & + \frac{1}{2} \mathcal{C} x^{-d-6} \left(4x^2 x \cdot \epsilon_1 x \cdot \epsilon_2 x \cdot \epsilon_3 (d \epsilon_1 \cdot \epsilon_3 x \cdot \epsilon_2 - (d+2) \epsilon_1 \cdot \epsilon_2 x \cdot \epsilon_3) \right. \\
 & + (x \cdot \epsilon_1)^2 \left(4dx^2 \epsilon_2 \cdot \epsilon_3 x \cdot \epsilon_2 x \cdot \epsilon_3 - (d-2)(d+4) (x \cdot \epsilon_2)^2 (x \cdot \epsilon_3)^2 + 2x^4 (\epsilon_2 \cdot \epsilon_3)^2 \right) \\
 & + \left. 2x^4 \left((\epsilon_1 \cdot \epsilon_2)^2 (x \cdot \epsilon_3)^2 + (\epsilon_1 \cdot \epsilon_3)^2 (x \cdot \epsilon_2)^2 \right) \right).
 \end{aligned} \tag{C.1}$$

In general, there are three independent structures with parameters $\mathcal{A}, \mathcal{B}$ and $\mathcal{C}$, which are interpreted as OPE coefficients of the associated OPE basis

$$\langle TTT \rangle = (\mathcal{A}, \mathcal{B}, \mathcal{C}) \cdot \langle TTT \rangle^{(i)}. \tag{C.2}$$

We can arbitrarily rotate the OPE basis, i.e., three-point structures $\langle TTT \rangle^{(i)}$, and read off the corresponding OPEs. One choice is the orthogonal basis [43, 44]

$$\langle TTT \rangle = \sum_{i=1}^3 \lambda_{TTT}^{(i)} \langle TTT \rangle_{\text{orth}}^{(i)}, \tag{C.3}$$

where the orthogonality is defined with respect to three-point pairing [45]. These are

$$\begin{aligned}
 \langle TTT \rangle_{\text{orth}}^{(1)} &= \\
 & \frac{d(d^2-4)}{16(d+1)(d+3)} x^{-d-6} \left(4x^2 x \cdot \epsilon_1 ((d+4) \epsilon_1 \cdot \epsilon_3 x \cdot \epsilon_2 - (d+2) \epsilon_1 \cdot \epsilon_2 x \cdot \epsilon_3) \right. \\
 & \times \left(dx \cdot \epsilon_2 x \cdot \epsilon_3 + 2x^2 \epsilon_2 \cdot \epsilon_3 \right) + 2x^4 \left(\epsilon_1 \cdot \epsilon_3 \left((d+4) \epsilon_1 \cdot \epsilon_3 (x \cdot \epsilon_2)^2 - 4x^2 \epsilon_1 \cdot \epsilon_2 \epsilon_2 \cdot \epsilon_3 \right) \right. \\
 & \left. + (d+4) (\epsilon_1 \cdot \epsilon_2)^2 (x \cdot \epsilon_3)^2 - 4(d+2) \epsilon_1 \cdot \epsilon_3 \epsilon_1 \cdot \epsilon_2 x \cdot \epsilon_2 x \cdot \epsilon_3 \right) \\
 & \left. + (d+4) (x \cdot \epsilon_1)^2 \left(4dx^2 \epsilon_2 \cdot \epsilon_3 x \cdot \epsilon_2 x \cdot \epsilon_3 + (d-2) d (x \cdot \epsilon_2)^2 (x \cdot \epsilon_3)^2 + 2x^4 (\epsilon_2 \cdot \epsilon_3)^2 \right) \right), \\
 \langle TTT \rangle_{\text{orth}}^{(2)} &= \\
 & - \frac{(d-2)d}{16(d-1)^2} x^{-d-6} \left(-2x^4 \left(\epsilon_1 \cdot \epsilon_3 \left((d^2+d-4) \epsilon_1 \cdot \epsilon_3 (x \cdot \epsilon_2)^2 \right. \right. \right. \\
 & \left. \left. + 4(d+1) x^2 \epsilon_1 \cdot \epsilon_2 \epsilon_2 \cdot \epsilon_3 \right) + (d^2+d-4) (\epsilon_1 \cdot \epsilon_2)^2 (x \cdot \epsilon_3)^2 - 4(d+1) \epsilon_1 \cdot \epsilon_3 \epsilon_1 \cdot \epsilon_2 x \cdot \epsilon_2 x \cdot \epsilon_3 \right) \\
 & \left. + (x \cdot \epsilon_1)^2 \left(-2(d^2+d-4) x^4 (\epsilon_2 \cdot \epsilon_3)^2 \right. \right. \\
 & \left. \left. + 4(d-3)(d+2) x^2 \epsilon_2 \cdot \epsilon_3 x \cdot \epsilon_2 x \cdot \epsilon_3 + (d-3)(d-2)(d+2)(d+4) (x \cdot \epsilon_2)^2 (x \cdot \epsilon_3)^2 \right) \right. \\
 & \left. + 4x^2 x \cdot \epsilon_1 (\epsilon_1 \cdot \epsilon_2 x \cdot \epsilon_3 + \epsilon_1 \cdot \epsilon_3 x \cdot \epsilon_2) \left(2(d+1) x^2 \epsilon_2 \cdot \epsilon_3 + (d-3)(d+2) x \cdot \epsilon_2 x \cdot \epsilon_3 \right) \right), \\
 \langle TTT \rangle_{\text{orth}}^{(3)} &= \\
 & \frac{(d-2)d}{16(d-1)} x^{-d-6} \left(4x^2 x \cdot \epsilon_1 (d \epsilon_1 \cdot \epsilon_2 x \cdot \epsilon_3 - (d-2) \epsilon_1 \cdot \epsilon_3 x \cdot \epsilon_2) \right. \\
 & \times \left((d+2) x \cdot \epsilon_2 x \cdot \epsilon_3 - 2x^2 \epsilon_2 \cdot \epsilon_3 \right) + 2x^4 \left(\epsilon_1 \cdot \epsilon_3 \left((d-2) \epsilon_1 \cdot \epsilon_3 (x \cdot \epsilon_2)^2 + 4x^2 \epsilon_1 \cdot \epsilon_2 \epsilon_2 \cdot \epsilon_3 \right) \right. \\
 & \left. + (d-2) (\epsilon_1 \cdot \epsilon_2)^2 (x \cdot \epsilon_3)^2 - 4d \epsilon_1 \cdot \epsilon_3 \epsilon_1 \cdot \epsilon_2 x \cdot \epsilon_2 x \cdot \epsilon_3 \right) + (d-2) (x \cdot \epsilon_1)^2 \left(\right. \\
 & \left. - 4(d+2) x^2 \epsilon_2 \cdot \epsilon_3 x \cdot \epsilon_2 x \cdot \epsilon_3 + (d+2)(d+4) (x \cdot \epsilon_2)^2 (x \cdot \epsilon_3)^2 + 2x^4 (\epsilon_2 \cdot \epsilon_3)^2 \right) \Big). \quad (\text{C.4})
 \end{aligned}$$

In the flat-space limit of $d = 3$, this orthogonal basis reduces to three-point vertices R , $[R^{(2)}]$ (which is vanishing in $d = 3$) and $[R^{(3)}]$. It is easy to show that $\langle TTT \rangle_{\text{orth}}^{(2)} \equiv 0$ in $d = 3$ by parameterizing $\epsilon = (e_1, e_2, i\sqrt{e_1^2 + e_2^2})$. It is then straightforward to find eq. (3.16).

C.2 OPEs in $d = 4$

Directly evaluating stress-tensor three-point function from Witten diagram is highly challenging because of unmanageable bulk tensor structures. Nevertheless, one can instead consider ANEC operator

$$\mathcal{E}(x^+, \vec{x}^\perp) = \int dx^- T_{--}(x^+, x^-, \vec{x}^\perp), \quad (\text{C.5})$$

where $x^\pm$ are lightcone directions. ANEC operator and more general light-ray operators [51] are especially useful to constrain AdS gravity, e.g., bound a/c and suggest EFT prescription [8, 9, 52], and can even sharply provide superconvergence sum rule [53]. To encode stress-tensor three-point function, we can evaluate the expectation value ANEC under states excited by the stress tensors [8]

$$\frac{\langle T\mathcal{E}T \rangle|_{x^+=0}}{\langle TT \rangle} \sim 1 + t_2 \left(\frac{(\epsilon_1 \cdot n)(\epsilon_2 \cdot n)}{\epsilon_1 \cdot \epsilon_2} - \frac{1}{d-1} \right) + t_4 \left(\frac{(\epsilon_1 \cdot n)^2(\epsilon_2 \cdot n)^2}{(\epsilon_1 \cdot \epsilon_2)^2} - \frac{2}{d^2-1} \right), \quad (\text{C.6})$$

where $\vec{x}^\perp = \vec{n}/(1+n^{d-1})$. The holographic side is much simplified, since the ANEC operator excites a shock wave in the bulk, namely

$$ds^2 = \frac{\ell_{\text{eff}}^2}{\tilde{r}^2} d\tilde{r}^2 + \tilde{r}^2 \left(dx_i dx^i + \delta(x^+) W(\rho, x) (dx^+)^2 \right), \quad (\text{C.7})$$

where

$$W(\rho, x) \sim \tilde{r}^{-4} \left(|x_1 - x'_1|^2 + |x_2 - x'_2|^2 + \ell_{\text{eff}}^2/\tilde{r}^2 \right)^{-3}, \quad (\text{C.8})$$

and we have made the following coordinate transformation from the Poincare AdS

$$x^+ \rightarrow -\frac{1}{x^+}, \quad x^- \rightarrow x^- - \frac{|\vec{x}^\perp|^2}{x^+} - \frac{z^2}{x^+}, \quad \vec{x}^\perp \rightarrow \frac{\vec{x}^\perp}{x^+}, \quad \tilde{r} = \frac{\ell_{\text{eff}} x^+}{z}. \quad (\text{C.9})$$

Consequently, we only have to find the contribution that the shock wave couples to two gravitons. After calculation, we find

$$t_2 = -\frac{24}{C_T} \left(40g_{3,5} + 8g_{3,6} + 108g_{3,7} + 45g_{3,8} - 2\ell_{\text{eff}}^2 g_{2,3} \right), \quad t_4 = \frac{2160}{C_T} (2g_{3,7} + g_{3,8}). \quad (\text{C.10})$$

(The generalize the result was obtained in [42].) We can readily verify that they are field redefinition invariant. Use the identities [8]

$$t_2 = \frac{15(5\mathcal{A} + 4\mathcal{B} - 12\mathcal{C})}{9\mathcal{A} - \mathcal{B} - 10\mathcal{C}}, \quad t_4 = -\frac{15(17\mathcal{A} + 32\mathcal{B} - 80\mathcal{C})}{4(9\mathcal{A} - \mathcal{B} - 10\mathcal{C})}, \quad C_T = \frac{\pi^2}{12}(9\mathcal{A} - \mathcal{B} - 10\mathcal{C}), \quad (\text{C.11})$$

we find, in terms of $\ell_{\text{inv}}^{(0)}$, that

$$\begin{aligned} \mathcal{A} &= -\frac{64 \left(\ell_{\text{inv}}^{(0)} \right)^3}{9\pi^5 G_N^{(5)}} \left(1 - \frac{24q_3}{M^2 \left(\ell_{\text{inv}}^{(0)} \right)^2} + \frac{6}{7M^4 \left(\ell_{\text{inv}}^{(0)} \right)^4} (32(2c_7 + c_8) - 21\hat{q}_3) \right), \\ \mathcal{B} &= -\frac{196 \left(\ell_{\text{inv}}^{(0)} \right)^3}{9\pi^5 G_N^{(5)}} \left(1 - \frac{636q_3}{49M^2 \left(\ell_{\text{inv}}^{(0)} \right)^2} + \frac{6}{343M^4 \left(\ell_{\text{inv}}^{(0)} \right)^4} (2918(2c_7 + c_8) - 399\hat{q}_3) \right), \\ \mathcal{C} &= \frac{92 \left(\ell_{\text{inv}}^{(0)} \right)^3}{9\pi^5 G_N^{(5)}} \left(1 - \frac{336q_3}{23M^2 \left(\ell_{\text{inv}}^{(0)} \right)^2} + \frac{6}{161M^4 \left(\ell_{\text{inv}}^{(0)} \right)^4} (142(2c_7 + c_8) - 231\hat{q}_3) \right). \end{aligned} \quad (\text{C.12})$$

where

$$\hat{q}_3 = \tilde{q}_3 + \frac{8}{3}(5q_1 + q_2 - q_3)q_3 - 20c_5 - 4c_6 - \frac{30}{7}c_7 + \frac{33}{14}c_8, \quad (\text{C.13})$$

is an invariant combination under the field redefinition. Before proceeding further, we would like to recall that $\ell_{\text{inv}}^{(0)}$ is the invariant expression associated with the effective AdS radius of massless gravities, satisfying the massless conditions $q_1 = q_3, q_2 = -4q_3$ and $\tilde{q}_3 = 0$, (the Gauss-Bonnet combination), together with [33]

$$\begin{aligned} c_1 &= \frac{1}{200}(16c_3 + 6c_4 + 160c_5 + 40c_6 + 52c_7 - 9c_8), \\ c_2 &= \frac{1}{20}(-12c_3 - 7c_4 - 80c_5 - 20c_6 - 24c_7 + 3c_8). \end{aligned} \quad (\text{C.14})$$

There are six-parameter family of cubic massless gravities, but the (a, c) charges depend only on (c_5, c_6, c_7, c_8) [33]. The same is true for the OPE coefficients $(\mathcal{A}, \mathcal{B}, \mathcal{C})$. In fact they are grouped into two combinations $(2c_7 + c_8)$ and $\hat{q}_3$. From the perturbative perspective, the field redefinition can be used to reduced the family of theories further down to one cubic theory, e.g. with a non-vanishing c_8 , since the cubic Lovelock combination with $2c_7 + c_8 = 0$ vanishes identically in $D = 5$ bulk dimensions. We can thus consider a fiducial $\ell_{\text{inv}}^{(0)}$ associated with quasi-topological gravity ($c_5 = 3/8, c_6 = -9/7, c_7 = 0, c_8 = 1$), for which $\hat{q}_3$ vanishes. Since the central charges can be expressed as linear combinations of the OPE coefficients. It is instructive to define the following linearly independent combinations

$$\begin{aligned} \tilde{\mathcal{A}} &\equiv \frac{\pi^6}{2880}(13\mathcal{A} - 2\mathcal{B} - 40\mathcal{C}) = a, \\ \tilde{\mathcal{B}} &\equiv \frac{\pi^6}{2016}(9\mathcal{A} - 2\mathcal{B} - 2\mathcal{C}), \\ \tilde{\mathcal{C}} &\equiv \frac{\pi^6}{480}(9\mathcal{A} - \mathcal{B} - 10\mathcal{C}) = c. \end{aligned} \quad (\text{C.15})$$

There are hidden linear differential relations among these OPE coefficients, given by eq. (4.1). The above choice of the linear combinations for $\mathcal{A}$ and $\mathcal{C}$ is unique, but it is not for the $\tilde{\mathcal{B}}$. If we instead choose $\hat{q}_3 = z(2c_7 + c_8)$, we can define

$$\tilde{\mathcal{B}} = \frac{\pi^6}{362880}((1620 - 119z)\mathcal{A} - 8(45 + 28z)\mathcal{B} - 40(9 - 14z)\mathcal{C}), \quad (\text{C.16})$$

which also satisfies (4.1). Since $(\tilde{\mathcal{A}}, \tilde{\mathcal{B}}, \tilde{\mathcal{C}})$ are linearly independent, different choices of z specifies different basis for the OPE coefficients, but the differential relations (4.1) is unchanged. Thus the (a, c) relation (1.4) is valid for all massless gravity even in the non-perturbative sense where the coupling constants are not necessarily small, the statements in (4.1) is valid only perturbatively where the OPE coefficients can be defined.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] M.J. Duff, *Observations on Conformal Anomalies*, *Nucl. Phys. B* **125** (1977) 334 [[INSPIRE](#)].
- [2] M.J. Duff, *Twenty years of the Weyl anomaly*, *Class. Quant. Grav.* **11** (1994) 1387 [[hep-th/9308075](#)] [[INSPIRE](#)].

- [3] A.B. Zamolodchikov, *Irreversibility of the Flux of the Renormalization Group in a 2D Field Theory*, *JETP Lett.* **43** (1986) 730 [[INSPIRE](#)].
- [4] J.L. Cardy, *Is There a c Theorem in Four-Dimensions?*, *Phys. Lett. B* **215** (1988) 749 [[INSPIRE](#)].
- [5] Z. Komargodski and A. Schwimmer, *On Renormalization Group Flows in Four Dimensions*, *JHEP* **12** (2011) 099 [[arXiv:1107.3987](#)] [[INSPIRE](#)].
- [6] H. Osborn and A.C. Petkou, *Implications of conformal invariance in field theories for general dimensions*, *Annals Phys.* **231** (1994) 311 [[hep-th/9307010](#)] [[INSPIRE](#)].
- [7] J. Erdmenger and H. Osborn, *Conserved currents and the energy momentum tensor in conformally invariant theories for general dimensions*, *Nucl. Phys. B* **483** (1997) 431 [[hep-th/9605009](#)] [[INSPIRE](#)].
- [8] D.M. Hofman and J. Maldacena, *Conformal collider physics: Energy and charge correlations*, *JHEP* **05** (2008) 012 [[arXiv:0803.1467](#)] [[INSPIRE](#)].
- [9] D.M. Hofman, D. Li, D. Meltzer, D. Poland and F. Rejon-Barrera, *A Proof of the Conformal Collider Bounds*, *JHEP* **06** (2016) 111 [[arXiv:1603.03771](#)] [[INSPIRE](#)].
- [10] T. Hartman, S. Kundu and A. Tajdini, *Averaged Null Energy Condition from Causality*, *JHEP* **07** (2017) 066 [[arXiv:1610.05308](#)] [[INSPIRE](#)].
- [11] I. Heemskerk, J. Penedones, J. Polchinski and J. Sully, *Holography from Conformal Field Theory*, *JHEP* **10** (2009) 079 [[arXiv:0907.0151](#)] [[INSPIRE](#)].
- [12] J.M. Maldacena, *The Large N limit of superconformal field theories and supergravity*, *Adv. Theor. Math. Phys.* **2** (1998) 231 [[hep-th/9711200](#)] [[INSPIRE](#)].
- [13] S.S. Gubser, I.R. Klebanov and A.M. Polyakov, *Gauge theory correlators from noncritical string theory*, *Phys. Lett. B* **428** (1998) 105 [[hep-th/9802109](#)] [[INSPIRE](#)].
- [14] E. Witten, *Anti-de Sitter space and holography*, *Adv. Theor. Math. Phys.* **2** (1998) 253 [[hep-th/9802150](#)] [[INSPIRE](#)].
- [15] X.O. Camanho, J.D. Edelstein, J. Maldacena and A. Zhiboedov, *Causality Constraints on Corrections to the Graviton Three-Point Coupling*, *JHEP* **02** (2016) 020 [[arXiv:1407.5597](#)] [[INSPIRE](#)].
- [16] N. Afkhami-Jeddi, T. Hartman, S. Kundu and A. Tajdini, *Einstein gravity 3-point functions from conformal field theory*, *JHEP* **12** (2017) 049 [[arXiv:1610.09378](#)] [[INSPIRE](#)].
- [17] M.S. Costa, T. Hansen and J. Penedones, *Bounds for OPE coefficients on the Regge trajectory*, *JHEP* **10** (2017) 197 [[arXiv:1707.07689](#)] [[INSPIRE](#)].
- [18] N. Afkhami-Jeddi, T. Hartman, S. Kundu and A. Tajdini, *Shockwaves from the Operator Product Expansion*, *JHEP* **03** (2019) 201 [[arXiv:1709.03597](#)] [[INSPIRE](#)].
- [19] D. Meltzer and E. Perlmutter, *Beyond $a = c$: gravitational couplings to matter and the stress tensor OPE*, *JHEP* **07** (2018) 157 [[arXiv:1712.04861](#)] [[INSPIRE](#)].
- [20] N. Afkhami-Jeddi, S. Kundu and A. Tajdini, *A Conformal Collider for Holographic CFTs*, *JHEP* **10** (2018) 156 [[arXiv:1805.07393](#)] [[INSPIRE](#)].
- [21] N. Afkhami-Jeddi, S. Kundu and A. Tajdini, *A Bound on Massive Higher Spin Particles*, *JHEP* **04** (2019) 056 [[arXiv:1811.01952](#)] [[INSPIRE](#)].

- [22] Y.-Z. Li, H. Lü and Z.-F. Mai, *Universal Structure of Covariant Holographic Two-Point Functions In Massless Higher-Order Gravities*, *JHEP* **10** (2018) 063 [[arXiv:1808.00494](#)] [[INSPIRE](#)].
- [23] J.S.R. Chisholm, *Change of variables in quantum field theories*, *Nucl. Phys.* **26** (1961) 469 [[INSPIRE](#)].
- [24] P. Bueno and P.A. Cano, *Einsteinian cubic gravity*, *Phys. Rev. D* **94** (2016) 104005 [[arXiv:1607.06463](#)] [[INSPIRE](#)].
- [25] P. Bueno, P.A. Cano, V.S. Min and M.R. Visser, *Aspects of general higher-order gravities*, *Phys. Rev. D* **95** (2017) 044010 [[arXiv:1610.08519](#)] [[INSPIRE](#)].
- [26] P. Bueno, P.A. Cano, R.A. Hennigar and R.B. Mann, *Universality of Squashed-Sphere Partition Functions*, *Phys. Rev. Lett.* **122** (2019) 071602 [[arXiv:1808.02052](#)] [[INSPIRE](#)].
- [27] J. Oliva and S. Ray, *A new cubic theory of gravity in five dimensions: Black hole, Birkhoff's theorem and C-function*, *Class. Quant. Grav.* **27** (2010) 225002 [[arXiv:1003.4773](#)] [[INSPIRE](#)].
- [28] R.C. Myers and B. Robinson, *Black Holes in Quasi-topological Gravity*, *JHEP* **08** (2010) 067 [[arXiv:1003.5357](#)] [[INSPIRE](#)].
- [29] P. Bueno, P.A. Cano, J. Moreno and A. Murcia, *All higher-curvature gravities as Generalized quasi-topological gravities*, *JHEP* **11** (2019) 062 [[arXiv:1906.00987](#)] [[INSPIRE](#)].
- [30] P. Bueno, P.A. Cano and R.A. Hennigar, *(Generalized) quasi-topological gravities at all orders*, *Class. Quant. Grav.* **37** (2020) 015002 [[arXiv:1909.07983](#)] [[INSPIRE](#)].
- [31] M. Henningson and K. Skenderis, *The Holographic Weyl anomaly*, *JHEP* **07** (1998) 023 [[hep-th/9806087](#)] [[INSPIRE](#)].
- [32] S. Nojiri and S.D. Odintsov, *On the conformal anomaly from higher derivative gravity in AdS/CFT correspondence*, *Int. J. Mod. Phys. A* **15** (2000) 413 [[hep-th/9903033](#)] [[INSPIRE](#)].
- [33] Y.-Z. Li, H. Lü and J.-B. Wu, *Causality and a-theorem Constraints on Ricci Polynomial and Riemann Cubic Gravities*, *Phys. Rev. D* **97** (2018) 024023 [[arXiv:1711.03650](#)] [[INSPIRE](#)].
- [34] A. Buchel, J. Escobedo, R.C. Myers, M.F. Paulos, A. Sinha and M. Smolkin, *Holographic GB gravity in arbitrary dimensions*, *JHEP* **03** (2010) 111 [[arXiv:0911.4257](#)] [[INSPIRE](#)].
- [35] R.C. Myers, M.F. Paulos and A. Sinha, *Holographic studies of quasi-topological gravity*, *JHEP* **08** (2010) 035 [[arXiv:1004.2055](#)] [[INSPIRE](#)].
- [36] R.C. Myers and A. Sinha, *Seeing a c-theorem with holography*, *Phys. Rev. D* **82** (2010) 046006 [[arXiv:1006.1263](#)] [[INSPIRE](#)].
- [37] R.C. Myers and A. Sinha, *Holographic c-theorems in arbitrary dimensions*, *JHEP* **01** (2011) 125 [[arXiv:1011.5819](#)] [[INSPIRE](#)].
- [38] H. Casini, M. Huerta and R.C. Myers, *Towards a derivation of holographic entanglement entropy*, *JHEP* **05** (2011) 036 [[arXiv:1102.0440](#)] [[INSPIRE](#)].
- [39] S. Caron-Huot, D. Mazac, L. Rastelli and D. Simmons-Duffin, *AdS Bulk Locality from Sharp CFT Bounds*, [[arXiv:2106.10274](#)] [[INSPIRE](#)].
- [40] X.O. Camanho, J.D. Edelstein and J.M. Sánchez De Santos, *Lovelock theory and the AdS/CFT correspondence*, *Gen. Rel. Grav.* **46** (2014) 1637 [[arXiv:1309.6483](#)] [[INSPIRE](#)].

- [41] P. Bueno, P.A. Cano and A. Ruipérez, *Holographic studies of Einsteinian cubic gravity*, *JHEP* **03** (2018) 150 [[arXiv:1802.00018](#)] [[INSPIRE](#)].
- [42] Y.-Z. Li, *Holographic Studies of The Generic Massless Cubic Gravities*, *Phys. Rev. D* **99** (2019) 066014 [[arXiv:1901.03349](#)] [[INSPIRE](#)].
- [43] S. Caron-Huot and Y.-Z. Li, *Helicity basis for three-dimensional conformal field theory*, [arXiv:2102.08160](#) [[INSPIRE](#)].
- [44] Y.-Z. Li, *Notes on flat-space limit of AdS/CFT*, *JHEP* **09** (2021) 027 [[arXiv:2106.04606](#)] [[INSPIRE](#)].
- [45] D. Karateev, P. Kravchuk and D. Simmons-Duffin, *Harmonic Analysis and Mean Field Theory*, *JHEP* **10** (2019) 217 [[arXiv:1809.05111](#)] [[INSPIRE](#)].
- [46] D.Z. Freedman, S.S. Gubser, K. Pilch and N.P. Warner, *Renormalization group flows from holography supersymmetry and a c theorem*, *Adv. Theor. Math. Phys.* **3** (1999) 363 [[hep-th/9904017](#)] [[INSPIRE](#)].
- [47] H. Osborn, *Derivation of a Four-dimensional c Theorem*, *Phys. Lett. B* **222** (1989) 97 [[INSPIRE](#)].
- [48] R.M. Wald, *Black hole entropy is the Noether charge*, *Phys. Rev. D* **48** (1993) R3427 [[gr-qc/9307038](#)] [[INSPIRE](#)].
- [49] H. Lü and R. Wen, *Holographic (a, c)-charges and Their Universal Relation in d = 6 from Massless Higher-order Gravities*, *Phys. Rev. D* **99** (2019) 126003 [[arXiv:1901.11037](#)] [[INSPIRE](#)].
- [50] M.S. Costa, J. Penedones, D. Poland and S. Rychkov, *Spinning Conformal Correlators*, *JHEP* **11** (2011) 071 [[arXiv:1107.3554](#)] [[INSPIRE](#)].
- [51] M. Kologlu, P. Kravchuk, D. Simmons-Duffin and A. Zhiboedov, *The light-ray OPE and conformal colliders*, *JHEP* **01** (2021) 128 [[arXiv:1905.01311](#)] [[INSPIRE](#)].
- [52] A. Belin, D.M. Hofman and G. Mathys, *Einstein gravity from ANEC correlators*, *JHEP* **08** (2019) 032 [[arXiv:1904.05892](#)] [[INSPIRE](#)].
- [53] M. Kologlu, P. Kravchuk, D. Simmons-Duffin and A. Zhiboedov, *Shocks, Superconvergence, and a Stringy Equivalence Principle*, *JHEP* **11** (2020) 096 [[arXiv:1904.05905](#)] [[INSPIRE](#)].