

RECEIVED: November 18, 2016

REVISED: December 7, 2016

ACCEPTED: December 9, 2016

PUBLISHED: December 21, 2016

## On CFT and quantum chaos

---

**Gustavo J. Turiaci<sup>a</sup> and Herman Verlinde<sup>a,b</sup>**

<sup>a</sup>*Physics Department, Princeton University,  
Princeton NJ 08544, U.S.A.*

<sup>b</sup>*Princeton Center for Theoretical Science, Princeton University,  
Princeton NJ 08544, U.S.A.*

*E-mail:* [turiaci@princeton.edu](mailto:turiaci@princeton.edu), [verlinde@princeton.edu](mailto:verlinde@princeton.edu)

**ABSTRACT:** We make three observations that help clarify the relation between CFT and quantum chaos. We show that any 1+1-D system in which conformal symmetry is non-linearly realized exhibits two main characteristics of chaos: maximal Lyapunov behavior and a spectrum of Ruelle resonances. We use this insight to identify a lattice model for quantum chaos, built from parafermionic spin variables with an equation of motion given by a Y-system. Finally we point to a relation between the spectrum of Ruelle resonances of a CFT and the analytic properties of OPE coefficients between light and heavy operators. In our model, this spectrum agrees with the quasi-normal modes of the BTZ black hole.

**KEYWORDS:** AdS-CFT Correspondence, Conformal Field Theory, Spontaneous Symmetry Breaking

**ARXIV EPRINT:** [1603.03020](https://arxiv.org/abs/1603.03020)

---

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Lyapunov from Goldstone</b>	<b>5</b>
<b>3</b>	<b>A chaotic lattice model</b>	<b>8</b>
<b>4</b>	<b>Ruelle resonances</b>	<b>11</b>
4.1	Ruelle resonances in CFT	11
4.2	Resonances and OPE coefficients	13
4.3	Ruelle from Liouville	15
<b>5</b>	<b>Conclusions</b>	<b>16</b>
<b>A</b>	<b>DOZZ three point function</b>	<b>17</b>


---

## 1 Introduction

There has been considerable recent interest in the manifestation of many body quantum chaos in strongly coupled conformal field theory [1–4]. Characteristics of chaotic systems, such as Lyapunov behavior, scrambling and Ruelle resonances, can be effectively isolated by studying out-of-time ordered correlation functions [5–15]. AdS/CFT duality relates these characteristics to evident properties of wave perturbations near black hole horizons, such as exponential redshifts, gravitational shockwaves and quasi-normal modes.

Many body quantum chaos is interesting in its own right, but usually hard to quantify. Identifying simple models or general mechanisms that exhibit aspects of quantum chaos is therefore a worthwhile goal. In this note we make three interrelated observations that may help 1) identify a new class of toy models in the form of a simple lattice model built out of parafermionic spin variables 2) clarify the relationship between maximal quantum chaos and the non-linear realization of conformal symmetry at finite temperature, 3) relate the spectrum of Ruelle resonances to analytic properties of OPE coefficients in the CFT. We now briefly describe each of the three components of our story.

**1) A discrete model of many body quantum chaos.** Useful many body systems that may exhibit chaos are quantum spin chains and matrix models. Another interesting example is the SYK model, which is solvable at strong coupling, maximally chaotic, and exhibits emergent conformal symmetry at low energies [5–8]. Our model of interest combines ingredients and properties of both examples, with the added feature that its Lyapunov behavior can be exhibited via weakly coupled effective field theory. The model



**Figure 1.** The discrete model is defined on a rhombic lattice. We indicated the center  $(\sigma, \tau)$  of the diamond  $(\sigma \pm 1, \tau \pm 1)$ . The equation of motion (1.3) expresses the variable at the top of the diamond in terms of the other three.

described below is a minor specialization of the class of integrable lattice models introduced by Faddeev, Kashaev and Volkov [16–20].

The model is assembled from a collection of  $\mathbb{Z}_N$  parafermionic operators  $f_n$ , labeled by an integer  $1 \leq n \leq L$  with  $L$  some large odd integer. We identify  $f_{L+1} \equiv f_1$ , so the integers  $n$  label points on a 1D periodic lattice. The  $f_n$  satisfy the algebra

$$f_{2n+1}f_{2n} = q^2 f_{2n}f_{2n+1}, \quad q = e^{i\pi/N}, \quad (1.1)$$

while  $[f_n, f_m] = 0$  for  $|m - n| \geq 2$ . This parafermion algebra can be realized on a finite dimensional Hilbert space  $\mathcal{H} = V_1 \otimes V_2 \otimes \dots \otimes V_L$  with  $V_n$  an  $N$ -dimensional vector space attached to the link between site  $n$  and  $n + 1$ , on which  $f_n$  and  $f_{n+1}$  act via appropriate clock and shift matrices. In the end, we imagine taking the continuum limit  $L \rightarrow \infty$ . The integer  $N$  is assumed to be large but finite.<sup>1</sup>

The time-evolution is discrete and specified as follows [16–18]. We relabel the variables  $f_n$  by means of two integers  $f_{\sigma, \tau}$  with  $\sigma + \tau = \text{even}$ , via  $f_{2r, 0} = f_{2r}$  and  $f_{2r+1, 1} = f_{2r+1}$ . The relabeled variables specify the initial condition of the model. The time evolution will generate a discrete, cylindrical 1+1-D space time formed by a rhombic lattice. The time evolution proceeds via a local propagation rule [16–18]. We can focus on a single diamonds shaped lattice cell

$$Y_N \equiv f_{\sigma, \tau+1}, \quad Y_S \equiv f_{\sigma, \tau-1}, \quad Y_W \equiv f_{\sigma-1, \tau}, \quad Y_E \equiv f_{\sigma+1, \tau}. \quad (1.2)$$

The evolution equation of the model reads

$$Y_N Y_S = \frac{Y_W Y_E}{(1 + Y_W)(1 + Y_E)} \quad (1.3)$$

Equation (1.3) is the simplest example of a Y-system. It specifies the variable  $Y_N$  at the top of the diamond shaped lattice cell in terms of the other three variables  $Y_E, Y_W$  and  $Y_S$ , see figure 1. The Y-system (1.3) defines an integrable lattice model, that can be recognized

<sup>1</sup>As we will see shortly,  $N$  will be proportional to the central charge of the low energy effective CFT.

as a discretized version of 2D hyperbolic geometry [16–18]. The exchange relation (1.1) amounts to a quantization of this hyperbolic geometry.<sup>2</sup>

The lattice model is a well defined quantum system, albeit one with a discrete time evolution. The model has been constructed [16–18] so that in the large  $L$  and IR limit, it describes a 2D continuum CFT with a non-linearly realized conformal symmetry with central charge  $c = 1 + 6(b + b^{-1})^2$  with  $b^2 = 1/N$ . As we will explain, this CFT exhibits maximal Lyapunov behavior, and an infinite set of Ruelle resonances match the quasi-normal frequencies of the BTZ black hole [21].

It may seem surprising that an integrable model can display properties characteristic of many body quantum chaos. To address this potential worry, one could choose to perturb the system away from integrability, e.g. by introducing frustration or by adding disorder. Since the features of quantum chaos will already become apparent in the unperturbed model, we will not go select among the list of such possible modifications<sup>3</sup> and instead focus on this idealized case, while ignoring the role of exact integrability. Indeed, we can note that there are other systems, such as  $\mathcal{N} = 4$  SYM theory at large  $N$ , that are believed to be both integrable and chaotic. We will return to this point in the concluding section.

**2) Lyapunov from Goldstone.** A central part of our reasoning consists of a new physical derivation of the Lyapunov behavior of an irrational CFT at finite temperature. The idea is as follows. 1+1-D CFTs are characterized by an infinite conformal symmetry group, given by reparametrizations of the lightcone coordinates  $u$  and  $v$

$$(u, v) \rightarrow (\xi(u), \eta(v)) \tag{1.4}$$

This conformal symmetry is broken by the conformal anomaly and by the presence of a finite energy density at finite temperature (and by the UV-cut-off). For a CFT with a dense asymptotic energy spectrum, it is then natural to expect that the conformal symmetry is non-linearly realized in terms of a light Goldstone mode.

This motivates us to consider the effective field theory of the relevant Goldstone excitation, described by the chiral field  $\xi(u)$  in (1.4) that parameterizes the conformal group. The effective Lagrangian is uniquely fixed by symmetries, and given by the geometric action of the Virasoro group [22]. In section 2, we will use this insight to derive the commutation relations of the Goldstone fields  $\xi(u)$  and  $\eta(v)$ . We will find that the thermal expectation value of the commutators squared

$$\langle [\xi(u), \xi(0)]^2 \rangle \sim e^{2\lambda u}, \quad \langle [\eta(v), \eta(0)]^2 \rangle \sim e^{2\lambda v}, \tag{1.5}$$

initially grow exponentially with the time separation, with a temperature dependent Lyapunov exponent  $\lambda = 2\pi/\beta$ . In fact, we will derive the somewhat more precise result that,

---

<sup>2</sup>In some way, one may view the model as a many body analogue of a hyperbolic billiard.

<sup>3</sup>One could add disorder e.g. by using the freedom of normalization of the  $f_n$  to set  $f_n^\dagger f_n = \kappa_n \mathbb{1}_{N \times N}$ , with  $\kappa_n$  random real numbers picked from a narrow probability distribution centered around  $\overline{\kappa_n} = \kappa$ . Alternatively, one could add frustration e.g. by including a next-to-neighbor interaction in the time step rule (1.2) and (1.3) via  $f_{\sigma, \tau+1} f_{\sigma, \tau-1} = (1 + \epsilon f_{\sigma+3, \tau}^{-1})(1 + \epsilon f_{\sigma-3, \tau}^{-1}) / (1 + \epsilon f_{\sigma+1, \tau}^{-1})(1 + \epsilon f_{\sigma-1, \tau}^{-1})$ .

inside a thermal expectation value, the commutator between two generic local operators takes the form<sup>4</sup>

$$[W(t_1), V(t_2)] \simeq \epsilon e^{\lambda t_{12}} \partial_{t_1} W(t_1) \partial_{t_2} V(t_2) \tag{1.6}$$

with  $\epsilon$  some constant proportional to  $1/c$ . This result, which holds for time-like separations in the intermediate range  $c \gg \lambda t_{12} \gg 1$ , matches with the bulk interpretation of the commutator as resulting from a near horizon gravitational shockwave interaction [1, 2, 23, 24].

**3) Ruelle resonances as poles in OPE coefficients.** A main characteristic of a chaotic system is that it thermalizes: out of time ordered correlation functions decay to zero at late times. The approach toward equilibrium is governed by Ruelle resonances [25]. They appear as poles in the Fourier transform of the thermal two-point function, or in systems that obey the ETH [26–30], the matrix element between two excited states with total energy  $M$

$$G(\omega) = \int dt \langle M | \mathcal{O}(t) \mathcal{O}(0) | M \rangle e^{i\omega t} \tag{1.7}$$

The Ruelle resonances of holographic 2D CFTs are well studied [21, 31]. As argued in [32], the matrix element reduces (for small  $t$ ) to the thermal 2-point function. Its Fourier transform  $G(\omega)$  has poles at resonant frequencies

$$\omega = -\frac{4\pi i}{\beta}(n + h), \tag{1.8}$$

that coincide with the quasi-normal modes of the BTZ black hole [21]. By factorizing the matrix element (1.7) in the intermediate channel, we can write

$$G(\omega) = \sum_{|i\rangle \in \mathcal{H}_{CFT}} \delta(M + \omega - E_i) |\langle M | \mathcal{O} | i \rangle|^2 \tag{1.9}$$

$$= \rho(M + \omega) |\langle M | \mathcal{O} | M + \omega \rangle|^2 \tag{1.10}$$

where we used that in the Cardy regime, we can replace the spectral density  $\rho(E) = \sum_{|i\rangle} \delta(E - E_i)$  by a continuous distribution, and label the CFT states by their energy. We learn that the Ruelle resonances dictate the analytic structure of the matrix element of a light operator  $\mathcal{O}$  between two highly excited states. This indicates that the resonances must show up as poles in the OPE coefficient of a light operator and two heavy operators. Or in AdS-dual terms, the quasi-normal modes should show up as poles in the absorption and emission amplitudes of wave perturbations by a BTZ black hole.

In section 4 we will show that the analytic continuation of the OPE coefficients of the continuum limit of our model indeed has poles located at the expected frequencies (1.8). This supports the statement that the continuum limit of the model is ergodic.

---

<sup>4</sup>Here for simplicity we only consider the time dependence of the correlator. In general, the left- and right-moving sectors each may have their own temperature and Lyapunov exponents  $\lambda_{l,r} = 2\pi/\beta_{l,r}$ .

## 2 Lyapunov from Goldstone

Consider an irrational 2D CFT with central charge  $c \gg 1$  with an asymptotic density of states given by the Cardy formula, and with a sparse low energy spectrum. We place the CFT on a circle, parameterized by a periodic coordinate  $x$  with period  $2\pi$ . We introduce light-cone coordinates  $(u, v) = (t - x, t + x)$ .

Consider a finite energy state with a constant expectation value for, say, the left-moving energy momentum tensor

$$\langle T(u) \rangle = L_0 \gg \frac{c}{12} \tag{2.1}$$

In this regime, we can associate to the state a finite inverse temperature  $\frac{\beta}{2\pi} = \sqrt{\frac{c}{24L_0}}$ .

Let us perform a general conformal transformation (1.4). We require that

$$\xi(u + 2\pi) = \xi(u) + 2\pi \tag{2.2}$$

The expectation value of the energy momentum tensor transforms non-trivially

$$\langle T(u) \rangle = L_0 \xi'^2(u) + \frac{c}{12} S_\xi(u) \tag{2.3}$$

with  $S_\xi$  the Schwarzian derivative

$$S_\xi(u) = \frac{1}{2} \left( \frac{\xi''(u)}{\xi'(u)} \right)^2 - \left( \frac{\xi''(u)}{\xi'(u)} \right)' \tag{2.4}$$

The spontaneous breaking of conformal symmetry is displayed via the  $\xi$ -dependence of this expectation value. Indeed, we can compare the relation (2.3) with the expression for the energy-momentum tensor of a fluid. The first term is analogous to the usual kinetic energy  $\frac{1}{2}\rho v^2$ , whereas the second term in (2.3) is the familiar vacuum contribution due to the conformal anomaly. It has a well-known physical explanation in terms of the Hawking-Unruh effect: the coordinate change from  $u$  to  $\xi(u)$  reshuffles the positive frequency (annihilation) and negative frequency (creation) modes, and thus alters the notion of the vacuum state.

Our physical assumption is that, for irrational CFTs at large  $c$  and in the Cardy regime, it becomes accurate to treat the coordinate transformation  $\xi(u)$  as a Goldstone field, in terms of which the conformal symmetry is non-linearly realized. Adopting this logic, we thus promote  $\xi(u)$  to an operator, that acts within the Hilbert subspace spanned by all states with energy density close to  $L_0$ , and their descendants. Within this subspace, we can remove the expectation value in (2.3) and elevate the equality in (2.3) to an *operator identity*

$$T(u) = L_0 \xi'^2(u) + \frac{c}{12} S_\xi(u). \tag{2.5}$$

As we will see shortly, the expression (2.5) for the energy-momentum tensor in terms of  $\xi(u)$  is familiar from the geometric quantization of  $\text{Diff}(S^1)$ , the group of (chiral) conformal transformations in 2D.

A cautious reader may view equation (2.5) simply as a (in)convenient parameterization of the energy momentum tensor  $T(u)$ . Our assumption, however, is that the symmetry parameter  $\xi(u)$  acts as a genuine local quantum field that creates and annihilates local

physical excitations. Given that  $\xi(u)$  is a scalar and  $T(u)$  is the generator of conformal transformations, we know that<sup>5</sup>

$$[T(u_1), \xi(u_2)] = \hbar \xi'(u_2) \delta(u_{12}) \quad \hbar \equiv \frac{6}{c} \quad (2.6)$$

$$[T(u_1), T(u_2)] = -\hbar(T(u_1) + T(u_2))\delta'(u_{12}) + \frac{\hbar}{2}\delta'''(u_{12}). \quad (2.7)$$

The emergence of a light Goldstone mode at finite temperature can be explained as a physical consequence of the fact that an irrational CFT in the Cardy regime has an extremely dense energy spectrum.

Equations (2.6) and (2.7) become semi-classical in the large  $c$  limit. From equation (2.1) we see that the field  $\xi(u)$  has expectation value

$$\langle \xi(u) \rangle = u \quad (2.8)$$

So semi-classically, we can think of the Goldstone field as:  $\xi(u) = u +$  small fluctuations.

We are now ready to state the main technical result of this section: *The three relations (2.5), (2.6) and (2.7) uniquely determine the commutation relation of the Goldstone field  $\xi(u)$ , and are sufficient to derive the Lyapunov growth of commutators.*

Working to leading order in  $1/c$ , one finds that [33–35]

$$[\xi(u_1), \xi(u_2)] = \frac{\epsilon(u_{12})}{L_0} + \frac{\sinh(\lambda\tau(u_1, u_2))}{L_0 \sinh \pi\lambda} \quad (2.9)$$

$$\tau(u_1, u_2) = \xi(u_1) - \xi(u_2) - \pi\epsilon(u_{12}), \quad \lambda = \sqrt{\frac{24L_0}{c}} \quad (2.10)$$

with  $\epsilon(x)$  the stair step function, defined via  $\epsilon'(x) = 2\delta(x)$  with  $\delta(x)$  the periodic delta-function:  $\epsilon(x) = 2n + 1$  for  $x \in (2\pi n, 2\pi(n + 1))$ . The same argument and derivation goes through for the right-movers. So we also have a right-moving Goldstone mode  $\eta(v) = v +$  small fluctuations, that satisfies the analogous commutation relation (2.9).<sup>6</sup> The left- and right-moving Goldstone fields commute  $[\xi(u), \eta(v)] = 0$ .

A detailed derivation of equation (2.9) and (2.10) can be found in [33–35]. Here we give a short summary. The constituent relation (2.5) between the energy-momentum tensor and the field  $\xi(u)$  can be decomposed as

$$T(u) = \varphi'^2(u) - 2\varphi''(u), \quad (2.11)$$

$$\varphi(u) = \frac{\lambda}{2} \xi(u) + \frac{1}{2} \log(\lambda\xi'(u)). \quad (2.12)$$

The commutation relations (2.6) and (2.7) then follow from the free field commutator

$$[\varphi(u_1), \varphi(u_2)] = \hbar \epsilon(u_{12}), \quad (2.13)$$

---

<sup>5</sup>Here we absorb a factor of  $\hbar \equiv 6/c$  in the definition of  $T(u)$ . This is a customary step, that exhibits the fact that the commutation relations (2.6) and (2.7) become semi-classical at large  $c$ .

<sup>6</sup>For simplicity we will assume that the left and right movers have the same temperature.

with  $\hbar = 6/c$ . So our task has been simplified: all we need to do is use relation (2.12) to solve for  $\xi(u)$  in terms of  $\varphi(u)$ , and use the chain rule to deduce the commutator of  $\xi(u_1)$  and  $\xi(u_2)$  from the free field commutator (2.13) of  $\varphi$ .

The free field  $\varphi(u)$  is periodic up to a shift

$$\varphi(u + 2\pi) = \varphi(u) + \pi\lambda. \tag{2.14}$$

Using this fact, equation (2.12) integrates to [34, 35]

$$\xi(u) = \frac{1}{\lambda} \log \left( \int_0^{2\pi} dy \frac{e^{2\varphi(u+y) - \lambda\pi}}{\sinh \pi\lambda} \right) \tag{2.15}$$

With this relation and equation (2.13) in hand, it is now a relatively straightforward calculation to derive the result (2.9) and (2.10).

Let us turn to the physical consequences of equations (2.9) and (2.10). We observe that  $\lambda$  is equal to the maximal Lyapunov exponent  $\lambda = 2\pi/\beta$ . We will assume that  $\lambda \gg 1$ , i.e. the thermal wave length is very short compared to the size of the spatial circle. The second term in the commutator (2.9), and its right-mover counter part, thus grows exponentially with the coordinate differences  $u_{12}$  and  $v_{12}$  over the range

$$\beta \ll |u_{12}| < 2\pi, \quad \beta \ll |v_{12}| < 2\pi. \tag{2.16}$$

We will restrict our attention to this coordinate range. In this regime, equation (2.9) implies that the commutator between two local functions  $\hat{f}(u_2) \equiv f(\xi(u_2))$  and  $\hat{g} \equiv g(\xi(u_1))$  of the Goldstone fields satisfy

$$[\hat{f}(u_1), \hat{g}(u_2)] \simeq e^{\lambda(|u_{12}| - 2\pi)} \hat{f}'(u_1) \hat{g}'(u_2). \tag{2.17}$$

Here we used equation (2.8) to replace  $\xi(u) \rightarrow u$  on the r.h.s. We would like to translate equation (2.17) into a statement about the commutator between local CFT operators.

Consider some local CFT operator  $\mathcal{O}(u, v)$ . Under the conformal transformation (1.4) it transforms as

$$\mathcal{O}(u, v) \rightarrow \xi'(u)^{h_L} \eta'(v)^{h_R} \mathcal{O}(\xi(u), \eta(v)) \tag{2.18}$$

Hence local operators are indeed non-trivial functions of the dynamical Goldstone fields.

It is logical to take this observation one step further, and, similarly as we did for the energy-momentum tensor, assume that local operators  $\mathcal{O}(u, v)$  can be represented as c-number valued functions of the operator valued fields  $\xi(u)$  and  $\eta(v)$  and their derivatives. The collection of these functions is determined by the spectrum and operator algebra of the CFT. Their form is constrained by the locality requirement that space-like separated operators commute. This condition is very restrictive: it prescribes that primary local operators are all of the form [34–38]

$$\mathcal{O}_h(u, v) = (f(u, v))^h, \tag{2.19}$$

$$f(u, v) = \frac{\lambda^2 \xi'(u) \eta'(v)}{4 \sinh^2 \left( \frac{\lambda}{2} (\xi(u) - \eta(v)) \right)}. \tag{2.20}$$



Equations (2.9) and (2.10) can then be used to compute the commutation relations between time-like separated operators, as follows.

The accepted test for Lyapunov growth of the commutator between two local operators  $W$  and  $V$  is to compute the expectation value

$$\langle W_\epsilon(u, v) [W(u, v), V(0, 0)] V_\epsilon(0, 0) \rangle \tag{2.21}$$

where the subscript  $\epsilon$  indicates a small displacement. This expectation value is equal to the difference between a time ordered and an out-of-time-ordered (OTO) correlation function. The OTO correlation function is obtained via analytic continuation of the time ordered correlation functions, where one circles, say, the coordinate  $u$  around the origin. This operation amounts to analytic continuation of the left-moving conformal blocks to the second Riemann sheet. Of course, we could also choose to do the analytic continuation using the coordinate  $v$ . This would have given the same final result.

The full-circle-monodromy  $\mathbf{M}$  of a conformal block is the square  $\mathbf{M} = \mathbf{R}^2$  of half-circle-monodromy known as the R-operation. The R-operator, acting on the left conformal blocks, re-orders the left-moving parts of the operators  $W$  and  $V$ . In the linearized regime, i.e. to leading order in  $1/c$ , we can write  $\mathbf{R} \simeq \mathbf{1} - \mathbf{r}$  with  $\mathbf{r}$  the perturbative operation that takes the commutator between the left-moving parts of  $W$  and  $V$ . The full-circle-monodromy is  $\mathbf{M} \simeq \mathbf{R}^2 = \mathbf{1} - 2\mathbf{r}$  and thus the full commutator inside (2.21) is equal to acting with  $(\mathbf{1} - \mathbf{M}) = 2\mathbf{r}$  on the two operators  $W$  and  $V$ . From equation (2.17) we then deduce that

$$[W(u_1, v_1), V(u_2, v_2)] \simeq 2e^{\lambda(u_{12}-u_0)} \partial_{u_1} W(u_1, v_1) \partial_{u_2} V(u_2, v_2) \tag{2.22}$$

This result, which holds for time like separation in the regime (2.16), displays the maximal Lyapunov behavior and the linearized gravitational effect of an early incoming perturbation (created by  $V$ ) on the arrival time of the outgoing signal (detected by  $W$ ).<sup>7</sup>

We end with a brief comment on the extension to higher orders. As indicated by the description of the monodromy moves, one expects that the commutator (2.22) exponentiates to a non-perturbative exchange relation. Fourier transforming the left-moving coordinate via  $W_\alpha(v) = \int du e^{i\alpha u} W(u, v)$ , this exchange algebra is expected to take the following form

$$W_\alpha(v_1) V_{\omega-\alpha}(v_2) = \sum_\beta M_\alpha^\beta V_\beta(v_2) W_{\omega-\beta}(v_1). \tag{2.23}$$

If we assume that the bulk interaction is dominated by gravity, then AdS/CFT makes a precise prediction for the monodromy matrix  $M_\alpha^\beta$  [3]. The prediction precisely matches with the monodromy matrix of Liouville CFT [3].

### 3 A chaotic lattice model

In this section, we will connect the FKV lattice model, defined by equations (1.1), (1.2) and (1.3), with the above effective CFT derivation of Lyapunov behavior.

---

<sup>7</sup>In equation (2.22) we assumed that the conformal dimensions of  $V$  and  $W$  are small, allowing us to neglect the Jacobian from (2.18). Including this contribution would not change the Lyapunov exponent.

The motivation for studying the lattice model is two-fold. First, the geometric theory of the Goldstone fields  $\xi(u)$  and  $\eta(v)$  is an effective theory, that only becomes accurate at finite temperature and long distance scales. Like all effective field theories, it does not define a fully consistent CFT by itself, nor does it have a unique UV completion. There are two ways in which one can try to embed an effective field theory into a self-consistent quantum system: a) look for an explicit UV completion, or b) introduce an explicit UV regulator. Approach b) is more practical.

A second motivation is that one can hope that the lattice model, by virtue of being more well defined, may allow for more explicit dynamical understanding of the underlying mechanism for chaos. Indeed, it turns out that the lattice Liouville model can be formulated in a way that preserves the geometric appeal of the continuum theory [16–18]

The Y-system (1.3) and the expression (2.19) of local operators in terms of the function (2.20) both have a direct connection with hyperbolic geometry. To see this, we first note that the 1+1-D metric defined by

$$ds^2 = f(u, v)dudv = \frac{\lambda^2 d\xi d\eta}{(2 \sinh(\frac{\lambda}{2}(\xi - \eta)))^2} \tag{3.1}$$

describes a hyperbolic space-time with constant negative curvature. The authors of [16–18] gave a beautiful discretized description of this 2D hyperbolic metric as follows.

We can write equation (3.1) as

$$f(u, v) = \frac{1}{\Delta^2} \frac{(e^{\lambda\xi(u+\Delta)} - e^{\lambda\xi(u-\Delta)})(e^{\lambda\eta(v+\Delta)} - e^{\lambda\eta(v-\Delta)})}{(e^{\lambda\xi(u+\Delta)} - e^{\lambda\eta(v+\Delta)})(e^{\lambda\xi(u-\Delta)} - e^{\lambda\eta(v-\Delta)})} \tag{3.2}$$

with  $\Delta$  an infinitesimal coordinate shift. Note that this expression for  $f(u, v)$  looks like a cross-ratio. So it is invariant under Möbius transformations. Now consider the values of  $f(u, v)$  in four nearby points, separated by null shifts  $\Delta$

$$\begin{aligned} f_{\sigma, \tau-1} &= f(u, v), & f_{\sigma+1, \tau} &= f(u + \Delta, v), \\ f_{\sigma-1, \tau} &= f(u, v + \Delta), & f_{\sigma, \tau+1} &= f(u + \Delta, v + \Delta). \end{aligned} \tag{3.3}$$

These four cross-ratios depend on six functions  $e^{\lambda\xi(u)}$ ,  $e^{\lambda\xi(u\pm\Delta)}$ ,  $e^{\lambda\eta(v)}$ , and  $e^{\lambda\eta(v\pm\Delta)}$ , but thanks to the Möbius invariance, only three of the six functions are independent. Therefore, the four cross-ratios (1.2) satisfy one relation [16–18]. Putting  $\Delta = 1$ , it reads

$$f_{\sigma, \tau+1} f_{\sigma, \tau-1} = \frac{f_{\sigma+1, \tau} f_{\sigma-1, \tau}}{(1 + f_{\sigma+1, \tau})(1 + f_{\sigma-1, \tau})}. \tag{3.4}$$

This confirms that the equation of motion of the FKV lattice model is a discretization of the hyperbolic metric (3.1). The parafermionic algebra

$$f_n f_{n\pm 1} = q^2 f_{n+1} f_n \tag{3.5}$$

defines a quantization of the space of discretized hyperbolic metrics.

Our new observation is that this lattice model can serve as a useful prototype of quantum chaos. The most direct way to substantiate this claim would be compute an out-of-time ordered four-point function of local operators

$$\langle f_{\sigma,\tau+t+1} [f_{\sigma,\tau+t-1}, f_{\sigma,\tau+1}] f_{\sigma,\tau-1} \rangle_{\beta} \quad (3.6)$$

at finite temperature, as a function of the time difference  $t$ . While this would in principle be doable, we will leave this task to future work. Instead we will cut the computation short, by banking on the results of [16–18, 20] that show that the above lattice model in the large  $L$  limit approaches continuum Liouville CFT. Together with the result of the previous section, this is sufficient to demonstrate that the continuum limit of the lattice model displays maximal Lyapunov behavior.

For completeness, let us display a few more elements of the dictionary. Working to leading order at large  $N$

$$\begin{aligned} e^{\varphi_n^+} e^{\varphi_m^+} &= e^{\varphi_m^+} e^{\varphi_n^+} q^{2\epsilon_{nm}} & e^{\varphi(u_2)} e^{\varphi(u_1)} &= e^{\varphi(u_1)} e^{\varphi(u_2)} e^{\hbar\epsilon(u_{12})} \\ e^{\varphi_{n+L}^+} &= e^{2\pi\lambda} e^{\varphi_n^+} & e^{\varphi(u+4\pi)} &= e^{2\pi\lambda} e^{\varphi(u)} \\ \frac{L}{2\pi} e^{\varphi_n^+} &= e^{\frac{\lambda}{2}\xi_n} - e^{\frac{\lambda}{2}\xi_{n-1}} & e^{\varphi(u)} &= \partial_u e^{\frac{\lambda}{2}\xi(u)} \end{aligned} \quad (3.7)$$

The right column lists the formulas (2.13), (2.14) and (2.12) that were used to derive the commutation relation (2.9) of the left-moving Goldstone variable  $\xi(u)$ . The left column is the lattice version of the same set of relations, with  $\epsilon_{nm}$  the discretized stair-step function. We can write a parallel set of formulas that represent the right-moving modes  $\varphi(v)$  and  $\eta(v)$  in terms of lattice variables  $\varphi_n^-$  and  $\eta_n$ .

Lattice variables  $\varphi_n^{\pm}$  that satisfy the exchange relation in (3.7) are obtained from the local operators  $f_n$  in two steps [16–18, 20]. First we define two mutually commuting sets of chiral operators  $w_n^{\pm}$  via

$$w_n^+ = q f_{2n+1} f_{2n+2}^{-1}, \quad w_n^- = q f_{2n+1} f_{2n}^{-1}. \quad (3.8)$$

These satisfy the algebra  $w_n^{\pm} w_m^{\pm} = q^{\pm 2\omega_{mn}} w_m^{\pm} w_n^{\pm}$ , with  $\omega_{mn} = \text{sgn}(m-n)\delta_{|m-n|,1}$ . The chiral variables  $\varphi_n^{\pm}$  are then defined as

$$\varphi_n^{\pm} = \sum_m \epsilon_{nm} \log w_m^{\pm}, \quad \varphi_{n+L}^{\pm} = \varphi_n^{\pm} + 2\pi\lambda, \quad 2\pi\lambda = \frac{1}{L} \sum_{n=1}^L \log w_n^{\pm}. \quad (3.9)$$

At the initial time  $\tau = 0$ , we can recover the single valued local parafermionic operators  $f_{2n}$  from the non-local chiral variables via

$$f_{2n} = e^{\varphi_n^-} e^{\varphi_n^+} \quad (3.10)$$

This is the lattice version of the relation  $e^{2\phi(u,v)} = e^{\varphi(u)+\varphi(v)}$  that expresses a non-chiral free field vertex operator into the product of the two chiral vertex operators. We note, however, that the time evolution (3.4) does not amount to free field propagation.

Among many other non-trivial results, [16–18] and [20] give an explicit construction of a unitary time evolution operator  $U$  that implements the time step (3.4)

$$f_{\sigma,\tau+1} = U^\dagger f_{\sigma,\tau-1} U. \tag{3.11}$$

This time evolution does not preserve the chiral factorization (3.10). However, it is shown that there exists a Bäcklund operator  $B$  that solves the time evolution via

$$f_{\sigma,\tau} = B^{-1} e^{\varphi_{\frac{1}{2}(\sigma-\tau)}^+} e^{\varphi_{\frac{1}{2}(\sigma+\tau)}^-} B \tag{3.12}$$

This Bäcklund operation is causal but highly non-local, and no explicit representation of  $B$  is known at present. Indeed, as exemplified by this equation, all non-trivial dynamics of the Liouville lattice model is encoded in the way in which the two chiral sectors get mixed and become entangled under the time evolution step (3.4). Our results are evidence that this mixing and entangling is happening in a maximally efficient way.

Our argument that the lattice model exhibits maximal Lyapunov growth is a copy of the effective CFT derivation presented in section 2. The three relations in the left column of equation (3.7) specify the commutation relations of the  $\xi_n$  variables, in the same way as the right column fixes the commutator algebra of  $\xi(u)$ . The commutator algebra is expected to approach the continuum result (2.9) in the large  $L$  limit. Our working assumption is that the exact solution (3.12) of the lattice model leads to an expression of the local operators  $f_{\sigma,\tau}$  in terms of the chiral modes  $\xi_n$  and  $\eta_n$  that mirrors formula (3.2). Via the same reasoning as in section 2, this expression can then be used to verify that the lattice model is local and to establish that the OTO four point function (3.6) grows exponentially with time.

## 4 Ruelle resonances

In this section we will expand on the topic of Ruelle resonances, which provide another signature of chaos and ergodicity. We will briefly review these concepts and then use the intuition for large  $c$  irrational conformal field theories to translate the knowledge about these resonances into concrete CFT data. We will introduce a notion of OPE coefficients (of light operators between heavy states) as analytic functions of energy. We will see that the presence of Ruelle resonances, in combination with the conformal bootstrap and AdS/CFT, impose stringent constraints on the form of these analytic OPE functions. We will then verify that the known OPE coefficients of the effective CFT of section 2 and the continuum limit of the lattice model of section 3 satisfy all these physical requirements.

### 4.1 Ruelle resonances in CFT

Ruelle resonances are poles in the Fourier transform of linear response functions that govern thermalization, the decay process towards thermal equilibrium after a quench. Consider a small perturbation produced by a local operator  $\mathcal{O}_b(x)$  to the Hamiltonian

$$\delta H = \int J(x) \mathcal{O}_b(x). \tag{4.1}$$

Here  $J(x)$  is an external source. Then one can study how this perturbation influences the time evolution of the expectation value of some other operator  $\langle \mathcal{O}_a(0) \rangle$ , which for convenience we place at  $x = 0$ . By expanding the evolution operator to linear order

$$\delta \langle \mathcal{O}_a(0) \rangle = \int dx' G_{ab}^{\text{ret}}(x') J(x'), \tag{4.2}$$

where  $G_{ab}^{\text{ret}}(x) = \theta(t) \langle [\mathcal{O}_a(x), \mathcal{O}_b(0)] \rangle$  (with  $t =$  time component of  $x$ ) is the retarded Green's function.  $G_{ab}^{\text{ret}}(x)$  may be expressed in terms of two point functions as

$$G_{ab}^{\text{ret}}(x) = \theta(t) (G_{ab}^+(x) - G_{ab}^-(x)), \tag{4.3}$$

with  $G_{ab}^+(x) = \langle \mathcal{O}_a(x) \mathcal{O}_b(0) \rangle$  the time ordered two point function and  $G_{ab}^-(x) = \langle \mathcal{O}_b(0) \mathcal{O}_a(x) \rangle$  the out-of-time-ordered two point function. Equation (4.2) is the basis of linear response theory, from which one can deduce transport properties such as the Kubo formula. Response functions are usually analyzed in the frequency domain. The Ruelle resonances appear as poles in the complex frequency plane. The imaginary part of the location of the poles determines the relaxation time. The leading behavior in  $\delta \langle \mathcal{O}_b \rangle(t)$  is governed the viscous hydrodynamical mode with the smallest imaginary part.

We are interested in studying this response function in a pure state microcanonical ensemble, defined by some highly excited CFT state  $|M\rangle$  with a large scale dimension  $M \gg \frac{c}{12}$ , so deep in the Cardy regime. The two-point functions of interest are given by the matrix elements of the two light operators  $\mathcal{O}_a$  and  $\mathcal{O}_b$  between two heavy states

$$\begin{aligned} G_{ab}^+(u, v) &= \langle M | \mathcal{O}_a(u, v) \mathcal{O}_b(0) | M \rangle \\ G_{ab}^-(u, v) &= \langle M | \mathcal{O}_b(0) \mathcal{O}_a(u, v) | M \rangle \end{aligned} \tag{4.4}$$

For 2D CFTs at large  $c$ , it has been argued in [32] that the matrix elements (4.4) are dominated by the identity conformal block (which for  $G^+(u, v)$  is given by the term with  $h = 0$  on the left in figure 2.) For large  $c$ , this identity block is well approximated by the thermal 2-point function on an infinite 1D space

$$G_{ab}^\pm(u, v) \simeq \delta_{ab} \left( \frac{\pi/\beta}{\sinh\left(\frac{\pi}{\beta}(u \pm i\epsilon)\right)} \right)^{2h} \left( \frac{\pi/\beta}{\sinh\left(\frac{\pi}{\beta}(v \pm i\epsilon)\right)} \right)^{2h}. \tag{4.5}$$

with  $\beta = \pi\sqrt{c/6M}$ . This is a useful result, that supports both the ETH and the dual identification of the two point function as the boundary-to-boundary propagator of a bulk field in a BTZ black hole background.

The validity of equations (4.5) is somewhat limited, however. It only holds for spatial separations that are small compared to the size of the spatial circle, and for the OTO two-point function, the time difference must be short compared to the scrambling time, since otherwise one enters the Lyapunov regime. On the gravity side, the perturbation  $\mathcal{O}_b$  creates an incoming wave that may collide with the outgoing wave detected by  $\mathcal{O}_a$ , and thereby substantially affect its future trajectory. This gravitational effect will show up as a modification of the OTO two-point function  $G^-(u, v)$ , and was studied in section 2. Here we will focus on the late time behavior of the time ordered 2-point function  $G^+(u, v)$ .

The incoming wave deforms the black hole horizon state. The subsequent ring down of the black hole towards equilibrium is the dual of the thermalization process of the CFT. Both processes are governed by an infinite set of resonances. On the gravity side, these resonances are the quasi-normal modes. These can be analyzed perturbatively, by considering small fluctuations of fields propagating in the neighborhood of the black hole horizon. This resonant quasi-normal frequencies are an infinite series of complex numbers, labeled by a non-negative integer  $n$  via [21]

$$\omega = \pm k - i \frac{4\pi}{\beta} (n + h). \tag{4.6}$$

with  $k$  is the momentum of the infalling mode and  $h$  the conformal dimension of the fluctuating field. This result was derived using the Poincaré patch, corresponding with a CFT on an infinite line, and with vanishing Dirichlet boundary conditions at infinity [21].<sup>8</sup> It is reasonable to assume that the result generalizes to black holes in global AdS, with a periodic spatial boundary, by replacing the momentum  $k$  by an integer angular momentum  $\ell$ .

In the CFT, the quasi-normal modes manifest themselves as Ruelle resonances, that appear as poles in the Fourier transform of the retarded thermal Green's function (4.3)

$$G_{ab}^{\text{ret}}(\omega, \ell) = \int du \int dv e^{i\frac{1}{2}(\omega+\ell)u} e^{i\frac{1}{2}(\omega-\ell)v} G_{ab}^{\text{ret}}(u, v), \tag{4.7}$$

which via equation (4.5) yields a spectrum that matches with the gravity prediction (4.6). Our goal in this section is to use the presence of these Ruelle poles to extract useful information about the OPE coefficients of the CFT. Earlier paper with results that overlap with this section are [32, 39].

## 4.2 Resonances and OPE coefficients

As a preparation, let us look at the different conformal block expansions of the matrix elements (4.4), as shown schematically in figure 2. The first equal sign of these identities represents the crossing symmetry relation<sup>9</sup>

$$G_{ab}(z) = \sum_{\omega} C_{aMM+\omega} C_{Mb}^{M+\omega} |\mathcal{F}_{M+\omega} \left[ \begin{matrix} MM \\ a \ b \end{matrix} \right] (z)|^2 = \sum_h C_{MM}^h C_{ahb} |\mathcal{F}_h \left[ \begin{matrix} M \ a \\ L \ M \ b \end{matrix} \right] (1-z)|^2 \tag{4.8}$$

where  $\mathcal{F}_h \left[ \begin{matrix} MM \\ a \ b \end{matrix} \right] (z)$  represents the Virasoro block shown on the left in figure 2. We see that crossing symmetry relates the ‘t-channel block’ with heavy intermediate channel (labeled by  $M+\omega$ ) to the ‘s-channel block’ with a light intermediate channel (labeled by  $h$ ).


The second relation in figure 2 is the exchange algebra relation,

$$\sum_{\omega} C_{aMM+\omega} C_{Mb}^{M+\omega} |\mathcal{F}_{M+\omega} \left[ \begin{matrix} MM \\ a \ b \end{matrix} \right] (z)|^2 = \sum_{\omega'} C_{bMM+\omega'} C_{Ma}^{M+\omega'} |\mathcal{F}_{M+\omega'} \left[ \begin{matrix} MM \\ b \ a \end{matrix} \right] (1/z)|^2, \tag{4.9}$$

---

<sup>8</sup>Notice that this Dirichlet boundary condition eliminates all gravitational excitations corresponding to the Virasoro descendants in the CFT. This restriction will become relevant later.

<sup>9</sup>Here we temporarily rotate to euclidean signature and set  $(u, v) = (z, \bar{z})$ .



**Figure 2.** Diagrammatic representation of crossing symmetry (given by the first identity) and the exchange algebra (given by the second) of the CFT correlation function of two heavy operators, labeled by  $M$ , and two light ones, labeled by  $a$  and  $b$ .

that imposes locality in the Euclidean region. In Lorentzian language, it implies that the R-matrix  $R_{\omega,\omega'}$  that relates the chiral time-ordered conformal block (labeled by  $M+\omega$ ) to the out-of-time-ordered conformal block (labeled by  $M+\omega'$ ) is an appropriate unitary transformation, so that in the euclidean region, it cancels out between the left- and right-movers of the complete CFT four-point function. After rotating to Lorentz signature, the R-matrix does show up in a non-trivial way, in the relation between the time-ordered Green's function  $G_{ab}^+(u, v)$  and the OTO Green's function  $G_{ab}^-(u, v)$  [3].

We wish to extract information regarding the Fourier transform of  $G_{ab}$  from its expansion (4.8) in conformal blocks, in the channel shown in the middle of figure 2. This is not directly possible, since no explicit expression for the Virasoro conformal blocks is known. So let us take a step back and write the crossing symmetry formula as a sum over primary operators and descendants. Let  $\gamma_n^\omega$  denote the  $n$ -th coefficient of the Laurent expansion of the Virasoro conformal block  $\mathcal{F}_{M+\omega}^{\left[ \begin{smallmatrix} MM \\ ab \end{smallmatrix} \right]}(z)$ . From now on we focus on the diagonal part of the two point function  $G_{ab}(u, v) = G(u, v)\delta_{ab}$ . It has the following expansion

$$G(u, v) = \sum_{|i\rangle} G_{\omega_{i,L}}(u) G_{\omega_{i,R}}(v) \quad (4.10)$$

$$G_{\omega_L}(u) = \sum_{n_L} C_{aMM+\omega_L} \gamma_{n_L}^{\omega_L} e^{i(\omega_L+n_L)u} \quad (4.11)$$

and a similar formula holds for  $G_{\omega_R}(v)$ . Here  $|i\rangle$  runs over all conformal primary states of the CFT in the neighborhood of the high energy state  $|M\rangle$ . In the sum we allowed all states with different left- and right conformal dimension  $(\Delta_{i,L}, \Delta_{i,R}) = (M + \omega_{i,L}, M + \omega_{i,R})$ .

We want to take the Fourier transform (4.7) with respect to both light-cone coordinates. It is useful to introduce the spectral density of CFT primary states

$$\rho(\omega_L, \omega_R) = \sum_{|i\rangle} \delta(M + \omega_L - \Delta_{i,L}) \delta(M + \omega_R - \Delta_{i,R}). \quad (4.12)$$

We then have

$$G^{\text{ret}}(\omega, \ell) = \int \frac{d\omega'}{2\pi} \frac{G(\omega', \ell)}{\omega' - \omega - i\epsilon} \quad (4.13)$$

$$G(\omega, \ell) = \sum_{n_L, n_R} C_{aMM+\omega_L-n_L} C_{aMM+\omega_R-n_R} \gamma_{n_L}^{\omega_L} \gamma_{n_R}^{\omega_R} \rho(\omega_L - n_L, \omega_R - n_R), \quad (4.14)$$

with  $\omega_L = \frac{1}{2}(\omega + \ell)$  and  $\omega_R = \frac{1}{2}(\omega - \ell)$ . For a given CFT,  $G^{\text{ret}}(\omega, \ell)$  contains exact information about the spectrum of primary fields, in the form of a dense set of poles along the real axis, with residues equal to the corresponding OPE coefficient. The Ruelle resonances appear as a series of poles in  $G^{\text{ret}}(\omega, \ell)$  located off the real axis. Based on equation (4.5) and the results of [32] and [21], we expect that their location should match with the quasi-normal frequencies (4.6).

The spectrum of an irrational CFT at large  $c$  becomes very dense in the Cardy regime. In this type of situation, it is customary to treat the spectrum as a continuum with spectral density given by the Cardy formula, and elevate the OPE coefficients to continuous functions of the conformal weights. The Ruelle resonances are then expected to arise as poles in the analytic continuation of the OPE coefficients.<sup>10</sup>

Let us summarize. The OPE coefficients between light and heavy operators satisfy several non-trivial compatibility conditions: they solve the CFT bootstrap equations (4.8) and (4.9), and must be compatible with the known location (4.6) of the Ruelle resonances. The question is: do these conditions uniquely fix the form of the OPE coefficients, in the universal high energy regime in which the CFT spectrum is governed by the Cardy formula? Do we know of any solutions to these conditions?

### 4.3 Ruelle from Liouville

The answer to the last question is affirmative: Liouville theory solves both conditions. The bootstrap program of Liouville CFT is by now on firm footing [40]. Our new observation is that the OPE coefficients of Liouville CFT, given by the famous DOZZ formula [41, 42], indeed exhibit a series of poles that precisely match with the quasi-normal frequencies (4.6) of the BTZ black hole. This observation gives extra support to the proposal that Liouville theory should be viewed as the effective CFT that captures universal high energy behavior of holographic CFTs. As we will discuss in the concluding section, this result also sheds light on whether the lattice model of section 3 has ergodic dynamics or not.

Liouville CFT has a continuous spectrum labeled by the momentum variable  $\alpha$  via  $\Delta_\alpha = \alpha(Q - \alpha)$  with  $c = 1 + 6Q^2$  and  $Q = b + b^{-1}$ . In appendix A we review the expression for the three point function  $C(\alpha_1, \alpha_2, \alpha_3)$  for a light operator, labeled by  $\alpha_1$ , and two heavy operators, labeled by  $\alpha_2$  and  $\alpha_3$ . Denoting the conformal dimensions as  $\Delta_1 = h$ ,  $\Delta_2 = M$  and  $\Delta_3 = M + \omega$ , the corresponding Liouville momenta are

$$\alpha_1 \simeq bh, \quad \alpha_3 - \alpha_2 \simeq i \frac{\omega}{2\sqrt{M}} = ib \frac{\beta}{4\pi} \omega \tag{4.15}$$

where  $\beta = 2\pi/b\sqrt{M}$  is the inverse temperature associated with the state  $M$ .

The DOZZ three-point function  $C(\alpha_1, \alpha_2, \alpha_3)$  has a rich pole structure. As explained in appendix A, the series of poles that are relevant to our physical situation are located at

$$\alpha_1 + \alpha_3 - \alpha_2 = nb, \quad n \in \mathbb{Z}, \tag{4.16}$$

---

<sup>10</sup>Evidently, the Ruelle poles do not arise from the density of states. The Laurent coefficients  $\gamma_n^\omega$  are fixed by conformal symmetry and they only exhibit poles for frequencies associated to degenerate states. The degenerate states appear at different locations than the quasi-normal modes.



which via equations (4.15) and (4.16) tells us that  $C_{\text{hMM}+\omega}$  has poles at

$$\omega = -i\frac{4\pi}{\beta}(n+h). \tag{4.17}$$

Plugging this into (4.14), and doing the integral (4.13), we learn that the retarded Green's function  $G^{\text{ret}}(\omega, \ell)$  has poles for

$$\begin{aligned} \omega &= -\ell + 2n_L - i\frac{4\pi}{\beta}(n+h), \\ \omega &= \ell + 2n_R - i\frac{4\pi}{\beta}(n+h). \end{aligned} \quad n \in \mathbb{N}. \tag{4.18}$$

These are the Ruelle resonances that govern the thermalization dynamics of Liouville CFT. Notice that relative to the list (4.6) of BTZ quasi-normal modes, the series (4.18) reveals additional poles shifted by the excitation numbers  $n_L$  and  $n_R$  of the left- and right-moving Virasoro descendants. These additional poles arise because in our CFT calculation, we did not exclude the possibility that the incoming wave created by  $\mathcal{O}_a$  also excites boundary gravitons. If we ignore the energy stored in the boundary gravitons, we recover the expected BTZ result (4.6).

## 5 Conclusions

In this note we have made three observations that clarify the geometric origin of chaotic behavior in irrational 2D CFTs. We argued that in holographic CFTs at finite temperature, conformal symmetry is non-linearly realized by means of universal Goldstone-like fields  $\xi(u)$  and  $\eta(v)$ , that describe the near-horizon gravitational dynamics of the dual theory. The effective field theory is weakly coupled and its maximal Lyapunov behavior can be demonstrated at the semi-classical level. We expect this mode to dominate in irrational CFT with large central charge and a dense spectrum. Understanding this behavior from first principles is still an open question.

We used this insight to propose a new toy model for quantum chaos in the form of the FKV lattice model, with an integrable equation of motion given by a Y-system. Integrability may seem unhelpful for generating ergodic behavior. Indeed, integrable systems are seen as prototypical counter-examples for the ETH: their single state microcanonical ensemble is understood to be described by the generalized Gibbs ensemble (GGE), which has many chemical potentials, one for each conserved quantity [47]. However, this reasoning assumes that the state that defines the microcanonical ensemble is an (approximate) eigenstate of many or all conserved quantities. Instead, if we choose an energy eigenstate that otherwise is a random linear superposition of eigenstates of all other conserved quantities, then the usual ETH can still apply. The conserved quantities in the FKV lattice model are highly non-local, and with respect to local observables the dynamics still looks random and thermalizing. As discussed in the introduction, this random dynamics can be reinforced by introducing some degree of disorder.

Indeed the discrete model seems particularly useful for studying propagation of entanglement, and even though the continuum limit is expected to be described by a CFT,

the entanglement propagation generated by the Y-system rule (1.3) is non-ballistic and mixes left- and right-moving signals. Our conjecture that the lattice dynamics is ergodic is further supported by the fact that the continuum limit of the model is expected to be described by Liouville theory, which via the observation of section 4 has Ruelle resonances that prescribe the approach towards thermal equilibrium.

Of course, underlying all three observations in this note, is the idea that the bulk gravitational dynamics of holographic 2D CFTs is accurately captured by 2D Liouville CFT [3]. This emergent Liouville field can be viewed as encoding the dynamical interplay between geometric entanglement and energy flow. This interpretation combines the idea of kinematic space [43, 44], that the entanglement entropy  $S(u, v)$  of an interval  $[u, v]$  between two space-like separated points  $x = u$  and  $x = v$  describes a metric on a 2D hyperbolic space via

$$ds^2 = \partial_u \partial_v S(u, v) du dv, \tag{5.1}$$

with the first law of entanglement thermodynamics

$$\delta S(u, v) = \delta K(u, v) = \int_u^v dx P_{[u,v]}(x) \delta T_{00}(x) \tag{5.2}$$

with  $P_{[u,v]}(x)$  the conformal Killing vector associated with the Rindler Hamiltonian  $K(u, v)$  of the interval  $[u, v]$ . Equation (5.2) can be integrated [45] into an expression for the energy-momentum tensor  $T_{\alpha\beta}$  in terms of the entanglement entropy  $S(u, v)$ , which looks exactly like the Liouville energy momentum tensor, via the identification

$$\phi(u, v) = S(u, v) \tag{5.3}$$

of the Liouville field with the entanglement entropy. Note that both quantities define locally constant curvature metrics, and both transform inhomogeneously under coordinate transformations. Hence the dynamics of kinematic space seems intimately connected with the emergence of an effective Liouville field in holographic 2D CFT.

## Acknowledgments

We thank Nele Callebaut, Bruno le Floch, Aitor Lewkowycz, Lauren McGough, Mark Mezei, Juan Maldacena, Eric Perlmutter, Eliezer Rabinovici and Douglas Stanford for helpful discussions. This research of H.V. is supported by NSF grant PHY-1620059.

## A DOZZ three point function

In this appendix we summarize the DOZZ formula for the OPE coefficients of Liouville theory [41, 42]. A nice review can be found in [46]. After introducing the formula we will study its analytic properties which are relevant for the application we consider in the main text.

The DOZZ formula computes the OPE coefficients between three primary operators of Liouville theory. These operators are labeled by a complex parameter  $\alpha$  and can be written in terms of the Liouville field  $\varphi(z, \bar{z})$  in the following way

$$V_{\alpha_j}(z, \bar{z}) = e^{2\alpha_j \varphi(z, \bar{z})}, \quad j = 1, 2, 3. \tag{A.1}$$

The dimension of the state in term of its label is  $\Delta_\alpha = \bar{\Delta}_\alpha = \alpha(Q - \alpha)$ . As usual the central charge is  $c = 1 + 6Q^2$  and  $Q = b + \frac{1}{b}$ , where  $b$  is a positive real parameter. The semiclassical limit  $c \gg 1$  corresponds to  $b \ll 1$ . That is the limit we are interested in, although the result for Liouville theory is supposed to be valid more generally. Now we can state the DOZZ formula which for generic  $\alpha_{1,2,3}$  and  $b$  is given by

$$C(\alpha_1, \alpha_2, \alpha_3) = \frac{\left[ \pi \mu \gamma(b^2) b^{2-2b^2} \right]^{(Q - \sum_i \alpha_i)/b} \Upsilon_0 \Upsilon_b(2\alpha_1) \Upsilon_b(2\alpha_2) \Upsilon_b(2\alpha_3)}{\Upsilon_b(\sum_i \alpha_i - Q) \Upsilon_b(\alpha_1 + \alpha_2 - \alpha_3) \Upsilon_b(\alpha_1 - \alpha_3 - \alpha_2) \Upsilon_b(\alpha_2 + \alpha_3 - \alpha_1)}, \quad (\text{A.2})$$

where  $\mu$  is the cosmological constant,  $\gamma(x) \equiv \Gamma(x)/\Gamma(1-x)$  and  $\Upsilon_0 = \left. \frac{d\Upsilon_b(x)}{dx} \right|_0$ .  $\Upsilon_b(x)$  is an entire function. It is usually defined by analytic continuation of an integral representation valid for  $0 < \text{Re}(x) < Q$  which can be found in [41, 42]. Here we will not need more information about this function other than its zeros since it gives the position of the poles in the DOZZ formula in terms of the  $\alpha$ 's. Specifically, they are located at

$$x = -\frac{m}{b} - bn, \quad x = \frac{m' + 1}{b} + b(n' + 1), \quad m, m', n, n' \in \mathbb{Z}^+. \quad (\text{A.3})$$

Looking at the formula (A.2) we see all the poles are located in terms of the labels  $\alpha$  at

$$\alpha_1 + \alpha_2 + \alpha_3 - Q = -\frac{m}{b} - bn, \quad \text{or} \quad \frac{m' + 1}{b} + (b + 1)n', \quad (\text{A.4})$$

$$\alpha_1 + \alpha_2 - \alpha_3 = -\frac{m}{b} - bn, \quad \text{or} \quad \frac{m' + 1}{b} + (b + 1)n', \quad (\text{A.5})$$

$$\alpha_1 - \alpha_2 + \alpha_3 = -\frac{m}{b} - bn, \quad \text{or} \quad \frac{m' + 1}{b} + (b + 1)n', \quad (\text{A.6})$$

$$\alpha_2 + \alpha_3 - \alpha_1 = -\frac{m}{b} - bn, \quad \text{or} \quad \frac{m' + 1}{b} + (b + 1)n'. \quad (\text{A.7})$$

These are all the poles of the OPE coefficients. Now we will use the semiclassical limit to identify the poles that are physically relevant for the discussion in the main text, i.e. the ones that survive the  $b \rightarrow 0$  limit. The external operators that we are interested in are such that one is light,  $\alpha_1$ , two are heavy,  $\alpha_2$  and  $\alpha_3$ , and the difference between the two heavy operators is small. This means we fix the scaling with  $b$  in the  $b \rightarrow 0$  limit such that  $\alpha_1 \sim b$ ,  $\alpha_2 \sim \alpha_3 \sim b^{-1}$  and  $\alpha_3 - \alpha_2 \sim b$ . Then it is clear that the relevant poles to retain are the ones in equation (A.5) and (A.6) for only  $n$  non zero. These two sets of pole for  $n > 0$  can be combined into a single formula

$$\alpha_1 + \alpha_3 - \alpha_2 = bn, \quad n \in \mathbb{Z}, \quad (\text{A.8})$$

where now  $n$  runs over all the integers. We see this has the right scaling since both the left and right hand side scale as  $b$  in the semiclassical limit. All the rest of the poles disappear in the heavy-heavy-light limit we are interested in here.

**Open Access.** This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

## References

- [1] S.H. Shenker and D. Stanford, *Black holes and the butterfly effect*, *JHEP* **03** (2014) 067 [[arXiv:1306.0622](#)] [[INSPIRE](#)].
- [2] S.H. Shenker and D. Stanford, *Multiple shocks*, *JHEP* **12** (2014) 046 [[arXiv:1312.3296](#)] [[INSPIRE](#)].
- [3] S. Jackson, L. McGough and H. Verlinde, *Conformal bootstrap, universality and gravitational scattering*, *Nucl. Phys. B* **901** (2015) 382 [[arXiv:1412.5205](#)] [[INSPIRE](#)].
- [4] J. Maldacena, S.H. Shenker and D. Stanford, *A bound on chaos*, *JHEP* **08** (2016) 106 [[arXiv:1503.01409](#)] [[INSPIRE](#)].
- [5] A. Kitaev, *A simple model of quantum holography (part 1)*, talk at *KITP*, U.S.A. April 7 2015.
- [6] A. Kitaev, *A simple model of quantum holography (part 2)*, talk at *KITP*, U.S.A. May 27 2015.
- [7] S. Sachdev and J. Ye, *Gapless spin fluid ground state in a random, quantum Heisenberg magnet*, *Phys. Rev. Lett.* **70** (1993) 3339 [[cond-mat/9212030](#)] [[INSPIRE](#)].
- [8] J. Polchinski and V. Rosenhaus, *The spectrum in the Sachdev-Ye-Kitaev model*, *JHEP* **04** (2016) 001 [[arXiv:1601.06768](#)] [[INSPIRE](#)].
- [9] J. Polchinski, *Chaos in the black hole S-matrix*, [arXiv:1505.08108](#) [[INSPIRE](#)].
- [10] G. Gur-Ari, M. Hanada and S.H. Shenker, *Chaos in classical D0-brane mechanics*, *JHEP* **02** (2016) 091 [[arXiv:1512.00019](#)] [[INSPIRE](#)].
- [11] D. Stanford, *Many-body chaos at weak coupling*, *JHEP* **10** (2016) 009 [[arXiv:1512.07687](#)] [[INSPIRE](#)].
- [12] P. Caputa, T. Numasawa and A. Veliz-Osorio, *Out-of-time-ordered correlators and purity in rational conformal field theories*, *Prog. Theor. Exp. Phys.* **2016** (2016) 113B06 [[arXiv:1602.06542](#)] [[INSPIRE](#)].
- [13] Y. Gu and X.-L. Qi, *Fractional statistics and the butterfly effect*, *JHEP* **08** (2016) 129 [[arXiv:1602.06543](#)] [[INSPIRE](#)].
- [14] B. Michel, J. Polchinski, V. Rosenhaus and S.J. Suh, *Four-point function in the IOP matrix model*, *JHEP* **05** (2016) 048 [[arXiv:1602.06422](#)] [[INSPIRE](#)].
- [15] E. Perlmutter, *Bounding the space of holographic CFTs with chaos*, *JHEP* **10** (2016) 069 [[arXiv:1602.08272](#)] [[INSPIRE](#)].
- [16] L.D. Faddeev and A. Yu. Volkov, *Hirota equation as an example of integrable symplectic map*, *Lett. Math. Phys.* **32** (1994) 125 [[hep-th/9405087](#)] [[INSPIRE](#)].
- [17] L.D. Faddeev, R.M. Kashaev and A. Yu. Volkov, *Strongly coupled quantum discrete Liouville theory. 1. Algebraic approach and duality*, *Commun. Math. Phys.* **219** (2001) 199 [[hep-th/0006156](#)] [[INSPIRE](#)].
- [18] L.D. Faddeev and R.M. Kashaev, *Strongly coupled quantum discrete Liouville theory. 2. Geometric interpretation of the evolution operator*, *J. Phys. A* **35** (2002) 4043 [[hep-th/0201049](#)] [[INSPIRE](#)].
- [19] L.D. Faddeev and L.A. Takhtajan, *Liouville model on the lattice*, *Lect. Notes Phys.* **246** (1986) 166 [[INSPIRE](#)].

- [20] A. Bytsko and J. Teschner, *The integrable structure of nonrational conformal field theory*, *Adv. Theor. Math. Phys.* **17** (2013) 701 [[arXiv:0902.4825](#)] [[INSPIRE](#)].
- [21] D. Birmingham, I. Sachs and S.N. Solodukhin, *Conformal field theory interpretation of black hole quasinormal modes*, *Phys. Rev. Lett.* **88** (2002) 151301 [[hep-th/0112055](#)] [[INSPIRE](#)].
- [22] A. Alekseev and S.L. Shatashvili, *From geometric quantization to conformal field theory*, *Commun. Math. Phys.* **128** (1990) 197 [[INSPIRE](#)].
- [23] Y. Kiem, H.L. Verlinde and E.P. Verlinde, *Black hole horizons and complementarity*, *Phys. Rev. D* **52** (1995) 7053 [[hep-th/9502074](#)] [[INSPIRE](#)].
- [24] T.D. Chung and H.L. Verlinde, *Dynamical moving mirrors and black holes*, *Nucl. Phys. B* **418** (1994) 305 [[hep-th/9311007](#)] [[INSPIRE](#)].
- [25] D. Ruelle, *Resonances of chaotic dynamical systems*, *Phys. Rev. Lett.* **56** (1986) 405.
- [26] A. Peres, *Ergodicity and mixing in quantum theory. I*, *Phys. Rev. A* **30** (1984) 504 [[INSPIRE](#)].
- [27] M. Feingold and A. Peres, *Distribution of matrix elements of chaotic systems*, *Phys. Rev. A* **34** (1986) 591 [[INSPIRE](#)].
- [28] J.M. Deutsch, *Quantum statistical mechanics in a closed system*, *Phys. Rev. A* **43** (1991) 2046.
- [29] M. Srednicki, *Chaos and quantum thermalization*, *Phys. Rev. E* **50** (1994) 888 [[cond-mat/9403051](#)].
- [30] M. Srednicki, *The approach to thermal equilibrium in quantized chaotic systems*, *J. Phys. A* **32** (1999) 1163 [[cond-mat/9809360](#)].
- [31] A.L. Fitzpatrick, J. Kaplan and M.T. Walters, *Universality of long-distance AdS physics from the CFT bootstrap*, *JHEP* **08** (2014) 145 [[arXiv:1403.6829](#)] [[INSPIRE](#)].
- [32] A.L. Fitzpatrick, J. Kaplan and M.T. Walters, *Virasoro conformal blocks and thermality from classical background fields*, *JHEP* **11** (2015) 200 [[arXiv:1501.05315](#)] [[INSPIRE](#)].
- [33] J.A. Teschner, *On quantization of Liouville theory and related conformal field theories*, Ph.D. thesis, DESY-95-118, Hamburg University, Hamburg Germany (1995) [[INSPIRE](#)].
- [34] H. Dorn and G. Jorjadze, *Boundary Liouville theory: Hamiltonian description and quantization*, *SIGMA* **3** (2007) 012 [[hep-th/0610197](#)] [[INSPIRE](#)].
- [35] J. Balog, L. Feher and L. Palla, *Coadjoint orbits of the Virasoro algebra and the global Liouville equation*, *Int. J. Mod. Phys. A* **13** (1998) 315 [[hep-th/9703045](#)] [[INSPIRE](#)].
- [36] J.-L. Gervais and A. Neveu, *The dual string spectrum in Polyakov's quantization. 1*, *Nucl. Phys. B* **199** (1982) 59 [[INSPIRE](#)].
- [37] J.-L. Gervais and A. Neveu, *Dual string spectrum in Polyakov's quantization. 2. Mode separation*, *Nucl. Phys. B* **209** (1982) 125 [[INSPIRE](#)].
- [38] J.-L. Gervais and A. Neveu, *New quantum treatment of Liouville field theory*, *Nucl. Phys. B* **224** (1983) 329 [[INSPIRE](#)].
- [39] K. Krasnov and S.N. Solodukhin, *Effective stringy description of Schwarzschild black holes*, *Adv. Theor. Math. Phys.* **8** (2004) 421 [[hep-th/0403046](#)] [[INSPIRE](#)].
- [40] J. Teschner, *Liouville theory revisited*, *Class. Quant. Grav.* **18** (2001) R153 [[hep-th/0104158](#)] [[INSPIRE](#)].

- [41] H. Dorn and H.J. Otto, *Two and three point functions in Liouville theory*, *Nucl. Phys. B* **429** (1994) 375 [[hep-th/9403141](#)] [[INSPIRE](#)].
- [42] A.B. Zamolodchikov and A.B. Zamolodchikov, *Structure constants and conformal bootstrap in Liouville field theory*, *Nucl. Phys. B* **477** (1996) 577 [[hep-th/9506136](#)] [[INSPIRE](#)].
- [43] B. Czech, L. Lamprou, S. McCandlish and J. Sully, *Integral geometry and holography*, *JHEP* **10** (2015) 175 [[arXiv:1505.05515](#)] [[INSPIRE](#)].
- [44] B. Czech, L. Lamprou, S. McCandlish and J. Sully, *Tensor networks from kinematic space*, *JHEP* **07** (2016) 100 [[arXiv:1512.01548](#)] [[INSPIRE](#)].
- [45] N. Callebaut and H. Verlinde, in preparation.
- [46] D. Harlow, J. Maltz and E. Witten, *Analytic continuation of Liouville theory*, *JHEP* **12** (2011) 071 [[arXiv:1108.4417](#)] [[INSPIRE](#)].
- [47] M. Rigol, V. Dunjko and M. Olshanii, *Thermalization and its mechanism for generic isolated quantum systems*, *Nature* **452** (2008) 854 [[arXiv:0708.1324](#)].