

RECEIVED: November 9, 2016

REVISED: December 8, 2016

ACCEPTED: December 21, 2016

PUBLISHED: December 28, 2016

Asymptotically safe grand unification

Borut Bajc^a and Francesco Sannino^{b,c}

^a*J. Stefan Institute,
1000 Ljubljana, Slovenia*

^b*CP³-Origins & the Danish IAS, University of Southern Denmark,
Campusvej 55, DK-5230 Odense M, Denmark*

^c*Université de Lyon, France, Université Lyon 1, CNRS/IN2P3, UMR5822 IPNL,
F-69622 Villeurbanne Cedex, France*

E-mail: borut.bajc@ijs.si, sannino@cp3.dias.sdu.dk

ABSTRACT: Phenomenologically appealing supersymmetric grand unified theories have large gauge representations and thus are not asymptotically free. Their ultraviolet validity is limited by the appearance of a Landau pole well before the Planck scale. One could hope that these theories save themselves, before the inclusion of gravity, by generating an interacting ultraviolet fixed point, similar to the one recently discovered in non-supersymmetric gauge-Yukawa theories. Employing a-maximization, a-theorem, unitarity bounds, as well as positivity of other central charges we nonperturbatively rule out this possibility for a broad class of prime candidates of phenomenologically relevant supersymmetric grand unified theories. We also uncover candidates passing these tests, which have either exotic matter or contain one field decoupled from the superpotential. The latter class of theories contains a model with the minimal matter content required by phenomenology.

KEYWORDS: GUT, Supersymmetric gauge theory

ARXIV EPRINT: [1610.09681](https://arxiv.org/abs/1610.09681)

Contents

1	Introduction	1
2	Consistency checks and constraints	3
2.1	Unitary constraints	3
2.2	Central charges and their positivity	3
2.3	a -maximization	4
2.4	Beta functions	5
2.5	a , b and c -theorems	5
2.6	Tracking the R -charge without the superpotential	6
3	Can SO(10) GUT be asymptotically safe?	7
3.1	Gaining R parity by loosing asymptotic freedom	7
3.2	SO(10) GUT without superpotential is unsafe	8
3.3	Minimal model with a superpotential	10
3.4	Gauge invariant fields becoming free	11
3.5	On the doublet-triplet splitting problem	12
4	Outlook and conclusions	13

1 Introduction

Theories of grand unification continue to play an important role as guiding principle when searching for extensions of the Standard Model. They offer a natural explanation for the observed quantization of the electric charge [1, 2], and predict the unification of the distinct SM gauge couplings at high energy [3]. The latter occurs when adding to the SM specific matter transforming according to incomplete representations of the grand unified theory (GUT).

Supersymmetry is a natural playground for the unification scenario since it almost automatically predicts the correct low energy spectrum that allows for one step-unification of the 3 gauge couplings. These meet at approximately 2×10^{16} GeV [4–7]. Moreover the different low energy matter fields, of a given generation, also unify in a larger representation of the gauge group, i.e. the 16 (27) of SO(10) [8, 9] (or E_6 [10]). This also, in turn, predicts new states such as the occurrence of a right-handed neutrino (plus extra vector-like matter in E_6) that fits naturally in the see-saw mechanism [11–15].

It is a fact, however, that asymptotic freedom is not always respected in supersymmetric GUTs such as the ones that predict exact R-parity conservation [16–18] at low

energy [19–21]. The reason being that one needs large matter representations [22–25] under $SO(10)$. This means that the coefficient of the one-loop gauge beta function

$$\beta_{1\text{-loop}} = 3T(G) - \sum_i T(R_i) \tag{1.1}$$

is strongly negative leading to a Landau pole typically just above the GUT scale but comfortably below the canonical gravity scale (for a 2-loop study see [26]). Embedding $SO(10)$ in larger gauge groups, for example E_6 cannot help [27, 28] because the resulting theory is even less asymptotically free.

One could envision different ways to go around this issue, for example one could push the unification scale closer to the gravity one via specific threshold corrections and hope that gravity will work its magic.

Another appealing possibility is that these theories save themselves, before gravity sets in, by developing an ultraviolet interacting fixed point in all couplings.

The hope for such a possibility stems from the discovery [29] that vector-like non asymptotically free gauge-Yukawa theories can indeed be fundamental theories at all scales.¹ The situation changes when considering the supersymmetric cousins of the theory investigated in [29]. It was, in fact, demonstrated in [36] that these supersymmetric cousins are unsafe, along with a much broader class of supersymmetric theories, further extending the one in [37]. The first study of asymptotically safe chiral gauge theories, some of which resembling GUT-like non-supersymmetric theories, appeared in [38] while semi-simple gauge groups in [39]. Asymptotic safety has been invoked by Weinberg [40] to tame quantum gravity [41–45].²

It is therefore timely and relevant to investigate the ultraviolet fate of a broad class of supersymmetric GUTs in which asymptotic freedom is lost.

We start with a pedagogical introduction and description of the tools that we will use to uncover the dynamics of these theories. In particular we will investigate non asymptotically free $SO(10)$ theories with different matter representations and with(out) superpotentials. Although we will show that a wide class of theories cannot abide all the constraints simultaneously we do find exotic theories featuring extremely large numbers of matter fields passing the tests. Besides the exotic models we also uncover a minimal model, with just 3 copies of 16’s, as well as one representative for each of the 10, 210, 126 and $\overline{126}$ multiplets of $SO(10)$ that can still be asymptotically safe.

We structure our paper as follows: in section 2 we briefly review, for the benefit of the reader, the rationale behind the full set of field-theoretical constraints we will use to discriminate among the possible candidate fixed points we will analyse. Section 3 opens with a brief self-contained introduction and justification of $SO(10)$ grand unified models

¹An important aspect of asymptotic safety in perturbative gauge-Yukawa theories is that scalars are required to tame the gauge fluctuations [29–31]. Earlier investigations of perturbative IR and UV interacting fixed points for gauge-Yukawa theories [32] were instrumental for the discovery in [29]. Asymptotic safety might occur also without elementary scalars [33, 34] but it would require a phase transition in the number of matter fields [35].

²UV conformal extensions of the standard model with and without gravity have also been discussed in the literature [46–86].

featuring several matter representations. The rest of the section is devoted to the analysis of a broad class of SO(10) grand unified models with and without superpotentials. We finally offer our conclusions in section 4.

2 Consistency checks and constraints

Since the grand unified theories investigated here supersymmetric we have a number of consistency checks and general constraints at our disposal to analyse the potential existence of any RG fixed point. If such a RG fixed points exists in an $\mathcal{N} = 1$ superconformal field theory (SCFT) it will necessarily possess a conserved $U(1)_R$ global symmetry. Furthermore the $U(1)_R$ current is in the same supermultiplet [87] as the energy-momentum tensor and the supercharge currents; this leads to several exact relations and constraints that we briefly review in this section.

2.1 Unitary constraints

For a unitary theory, the operators form unitary representations of the superconformal group, which implies that operator dimensions have various lower bounds. For example, regardless of supersymmetry, all gauge invariant spin $j = \bar{j} = 0$ operators have the lower bound (generators act with implicit commutators) [88] (see also e.g. [89])

$$D(\mathcal{O}) \geq 1, \quad D(\mathcal{O}) = 1 \leftrightarrow P_\mu P^\mu(\mathcal{O}) = 0, \quad (2.1)$$

so the bound is saturated if and only if the operator \mathcal{O} is a decoupled, free field. Chiral primary operators have dimension, D , and superconformal $U(1)_R$ charge, R , related by

$$D(\mathcal{O}) = \frac{3}{2}R(\mathcal{O}). \quad (2.2)$$

Using (2.2) for the matter chiral superfields Q_i one can relate the matter anomalous dimensions γ_i to their superconformal $U(1)_R$ charge.

$$D(Q_i) \equiv 1 + \frac{1}{2}\gamma_i(g) = \frac{3}{2}R(Q_i) \equiv \frac{3}{2}R_i. \quad (2.3)$$

2.2 Central charges and their positivity

We summarise here the constraints due to the positivity of the coefficients related to the stress-energy trace anomaly. These have been derived by considering the effects of an external supergravity background for theories with sources for conserved flavor currents stemming from trace anomaly and proportional to the square of the dual of the Riemann curvature, the square of the Weyl tensor, as well as the square of the flavor symmetry field strength. These functions of the R charge are indicated respectively with $a(R)$, $c(R)$ and $b(R)$ [90, 91].

The conformal anomaly a of the SCFT is exactly given by the superconformal $U(1)_R$ 't Hooft anomalies [90, 91] (we rescale the overall normalization factor of 3/32 for convenience)

$$a(R) = 3\text{Tr}U(1)_R^3 - \text{Tr}U(1)_R. \quad (2.4)$$

Let's determine this function for a gauge theory with gauge group, G , and matter fields Q_i , in representations r_i of G , the 't Hooft anomalies evaluate to

$$a(R) = |G| [3R_V^3 - R_V] + \sum_i |r_i| [3(R_i - 1)^3 - (R_i - 1)] = 2|G| + \sum_i |r_i| a_1(R_i) \quad (2.5)$$

where $|G| = r_{\text{adjoint}}$ is the number of generators in the adjoint representation and $|r_i|$ is the dimension of the representation r_i . We must use in $a(R)$ the fermion R charges that for the gluino is exactly $R_V = R(V) = 1$ with V the vector chiral superfield, while for each chiral superfield $Q = \phi_Q + \sqrt{2}q\theta + \theta^2 F_Q$ we have $R(q) = R(Q) - 1$ because $R(\theta) = -1$, and we define the function

$$a_1(R) \equiv 3(R - 1)^3 - (R - 1) = (1 - R) [1 - 3(1 - R)^2]. \quad (2.6)$$

The c-function reads [90, 91]

$$c(R) = 9\text{TrU}(1)_R^3 - 5\text{TrU}(1)_R. \quad (2.7)$$

For a generic gauge theory we have:

$$c(R) = |G|(9 - 5) + \sum_i |r_i| [9(R_i - 1)^3 - 5(R_i - 1)] = 4|G| + \sum_i |r_i|(1 - R_i) [5 - 9(1 - R_i)^2], \quad (2.8)$$

and we dropped the overall normalization factor of $1/32$.

The flavor b-function reads [90, 91]

$$b(R) = \text{TrU}(1)_R F^2 = \sum_i |r_i|(1 - R_i) F_i^2. \quad (2.9)$$

We have dropped the overall normalization factor of 3 and F_i are the flavor charges for each representation.

2.3 a -maximization

Among all possible, conserved $U(1)_R$ symmetries, the superconformal $U(1)_R$ is the one maximizing $a(R)$ pioneered in [92]. For example, for a chiral superfield X of charge $R(X) = R$ (so $R(\psi_X) = R - 1$), the function is $a(R) = a_1(R)$ in (2.6). The function $a_1(R)$ has a local maximum at the free-field value, $R = \frac{2}{3}$, and a local minimum at $R = \frac{4}{3}$, see figure 1. In addition $a_1(R)$ is below the local maximum, $a_1(R) < a_1(R = 2/3)$ for $R < 5/3$. (see [93] for a further related phase diagnostic). We maximize the function (2.6) for unconstrained, i.e. free chiral superfields and obtain $R_* = 2/3$, which is the free-field value of the R-charge, corresponding to $D(X) = 1$. When interactions are present, we maximize $a(R)$ requiring the interactions to preserve the R-symmetry. Accidental symmetries, if present, affect a-maximization [94, 95] yielding a larger value of a .




Figure 1. The function $a_1(R)$.

2.4 Beta functions

Beta functions are proportional to how the couplings break the superconformal $U(1)_R$ when going away from the fixed point. The supersymmetric gauge coupling beta function embodies a remarkable property, it is proportional to the ABJ triangle anomaly of the $U(1)_R$ current with two G gauge fields, i.e. $\text{Tr } G^2 U(1)_R$:

$$\beta(g) = -\frac{3g^3}{16\pi^2} f(g^2) \text{Tr } G^2 U(1)_R, \quad \text{Tr } G^2 U(1)_R = T(G) + \sum_i T(r_i)(R_i - 1). \quad (2.10)$$

Our normalization for the quadratic Casimir of the adjoint $T(G)$ is $T(\text{SU}(N_c)) = N_c$, so that the fundamental representation of $\text{SU}(N)$ has $T(r_{\text{fund}}) = \frac{1}{2}$. The function $f(g^2) = 1 + \mathcal{O}(g^2)$ is scheme dependent (and presumed positive). The above (2.3) is the NSVZ exact beta function [96], in which a specific scheme is employed for $f(g^2)$:

$$\beta(8\pi^2 g^{-2}) = f(g^2) \left(3T(G) - \sum_i T(r_i)(1 - \gamma_i(g)) \right). \quad (2.11)$$

For superpotential terms with trilinear interactions W_y , the beta function for the holomorphic coupling y reads

$$\beta(y) = \frac{3}{2} y (R(W_y) - 2). \quad (2.12)$$

2.5 a , b and c -theorems

For any super CFT not only these coefficients must be positive [90, 91] but it is also expected, following Cardy's conjecture, a 4d version of the a -theorem [97–102], that reads

$$\Delta a \equiv a_{\text{UV}} - a_{\text{IR}} > 0. \quad (2.13)$$

For free theories these coefficients are automatically positive. This implies that for asymptotically free theories they are automatically positive at the trivial UV fixed point while for asymptotically safe theories they are automatically positive in the infrared. In fact, the

free value for gauge theories reads:

$$a_{\text{free}} = 2|G| + \frac{2}{9} \sum_i |r_i|, \quad c_{\text{free}} = 4|G| + \frac{4}{3} \sum_i |r_i|, \quad b_{\text{free}} = \frac{1}{3} \sum_i |r_i| F_i^2 \quad (2.14)$$

where $R_i = 2/3$ because all the chiral superfields are free, i.e. $D_i = 1$, as it should be for a non-interacting field. It is worth mentioning that the physical dimension of the vector chiral superfield is always (also in the interacting theory) the free one since the R charge of the gluino is fixed and $D(V) = 3/2R(V) = 3/2 = D(\lambda)$ with λ the gluino. For the free theory we do not care about anomaly free value for R charges because there are no interactions.

Interesting constraints emerge when requiring positivity of $b_{\text{IR(UV)}}$ and $c_{\text{IR(UV)}}$ for the interacting IR(UV) fixed point in asymptotically free (safe) field theories along with the $\Delta a > 0$ condition. At the interacting fixed point the only R charges that matter are the ones that allow for the interacting field theory to be consistent, including 't Hooft anomaly (free) conditions and superpotential constraints. The above implies:

$$c_{\text{FP}} = 4|G| + \sum_i |r_i|(1-R_i) [5-9(1-R_i)^2] > 0, \quad b_{\text{FP}} = \sum_i |r_i|(1-R_i) F_i^2 > 0, \quad (2.15)$$

and

$$\Delta a = a_{\text{UV}} - a_{\text{IR}} = \pm \left(2|G| + \sum_i |r_i| a_1(R_i) - a_{\text{free}} \right) = \pm \frac{1}{9} \sum_i |r_i| [(3R_i - 2)^2 (3R_i - 5)] > 0, \quad (2.16)$$

where the plus(minus) sign corresponds to the asymptotically safe(free) interacting fixed point. The constraint in (2.16) is stronger for asymptotically safe theories [36, 37] since it requires at least one chiral superfield to have a quite sizable R charge larger than $5/3$. Assuming, for example, the presence of an asymptotically safe fixed point for super QCD once asymptotic freedom is lost, i.e. $N_f > 3N_c$, one discovers that $R_Q = R_{\tilde{Q}} = 1 - N_c/N_f$ assume values between 1 and $2/3$ and therefore the theory violates the constraint in (2.16) [36, 37] while it still respects positivity of the remaining constraints. An IR interacting fixed point, relevant for super QCD conformal window [103, 104], in asymptotically free field theories, on the other hand, requires the milder condition $R_i < 5/3$.

2.6 Tracking the R -charge without the superpotential

With vanishing superpotential, the a -function is defined as [105, 106]

$$a(R_i, \lambda_G) = 2|G| + \sum_i |r_i| (3(R_i - 1)^3 - (R_i - 1)) + \lambda_G \left(T(G) + \sum_i T(r_i)(R_i - 1) \right) \quad (2.17)$$

where λ_G is the Lagrange multiplier which enforces the vanishing of the NSVZ β -function at the superconformal fixed point. From

$$\frac{\partial a(R_i, \lambda_G)}{\partial R_i} = 0 \quad (2.18)$$

one finds

$$R_i = 1 - \frac{\epsilon_i}{3} \sqrt{1 - \frac{T(r_i)\lambda_G}{|r_i|}} \quad (2.19)$$

with $\epsilon_i^2 = 1$. Reality of R_i requires

$$\lambda_G \leq \lambda_G^{\max} \equiv \min_i \left(\frac{|r_i|}{T(r_i)} \right) \quad (2.20)$$

We will also presume that the interacting fixed point is smoothly connected to the non-interacting fixed point when the coupling vanishes. This allows to enforce continuity of the R charges.

We are now ready to investigate the dynamics of grand unified theories that are not asymptotically free.

3 Can SO(10) GUT be asymptotically safe?

We will now use the above machinery to investigate whether SO(10) GUT theories can be asymptotically safe rather than free. We first summarize how and why the loss of asymptotic freedom appears when trying to construct models that automatically embody R -parity. We then analyze whether these theories can nonperturbatively flow to an UV fixed point by applying the above tests.

3.1 Gaining R parity by loosing asymptotic freedom

We mentioned in the introduction that non asymptotically free grand unifications can provide a rationale for the existence of low energy R parity [23–25]. The latter stems from the SO(10) Cartan subalgebra generator $B - L$ through

$$R = (-1)^{3(B-L)+2S} = M(-1)^{2S} \quad \text{with} \quad M = (-1)^{3(B-L)}. \quad (3.1)$$

We see that R parity, up to the spin S , identifies with the matter parity M . An elegant way to break the rank of SO(10) without breaking spontaneously the R -parity is to introduce a Higgs sector transforming according to the $126 + \overline{126}$ dimensional representation³ of SO(10). Indeed in the 126 ($\overline{126}$) the only possible SM and SU(5) singlet has $B - L = -2(2)$ that preserves R -parity.

Since in SO(10)

$$16 \times 16 = 10 + 126 + 120, \quad (3.2)$$

the Yukawa couplings (and thus all SM fermion masses) could arise via the following linear combination:

$$W_{\text{Yukawa}} = 16_a \left(Y_{10}^{ab} 10 + Y_{126}^{ab} \overline{126} + Y_{120}^{ab} 120 \right) 16_b, \quad (3.3)$$

with $a, b = 1, 2, 3$ running over the generations. From SO(10) one can show that

$$Y_{10,126} = +Y_{10,126}^T, \quad Y_{120} = -Y_{120}^T. \quad (3.4)$$

³We need simultaneously 126 and $\overline{126}$ to cancel the D-terms.

In fact a minimal choice to generate realistic mixings among the generations, i.e. the physical V_{CKM} and V_{PMNS} matrices, is to add to the already present $\overline{126}$ the 10 dimensional representation alone.⁴

For successful model building two requirements must still be met: first, we need to break $\text{SO}(10)$ down to the SM gauge group which cannot be accomplished by the $\overline{126}$ that at most can break it to $\text{SU}(5)$; second, the MSSM Higgses must be contained in both the 10 and $\overline{126}$. Both problems can be addressed by introducing the 210 representation. The reasons being that: the 3 SM singlets of 210 are enough to further break $\text{SU}(5)$ to the SM group; the renormalizable operator $210\ 10\ \overline{126}$ sources via a nonzero doublet vev in 10 a related vev in $\overline{126}$.

To summarize, the minimal $\text{SO}(10)$ model we will consider is composed of

$$3 \times 16 + 126 + \overline{126} + 10 + 210, \tag{3.5}$$

that in the end yields a non-asymptotically free theory with the following extremely large coefficient of the one-loop beta function

$$\beta_{1\text{-loop}} = -109, \tag{3.6}$$

implying that a Landau pole is reached very quickly and below the Planck scale. The emergence of an interacting ultraviolet fixed point could save the theory. We will therefore investigate such a possibility in the next session.

3.2 $\text{SO}(10)$ GUT without superpotential is unsafe

We commence our analysis by demonstrating that: *Minimal $\text{SO}(10)$ with $3 \times 16 + 126 + \overline{126} + 10 + 210$ matter content and vanishing superpotential does not have a UV fixed point.*

To prove this we start with the NSVZ beta function

$$\beta_{\text{NSVZ}}(\lambda_G) \equiv T(G) + \sum_i T(r_i)(R_i(\lambda_G) - 1), \tag{3.7}$$

in which $R_i(\lambda_G)$ is given by (2.19). We, of course, reproduce in the non-interacting IR limit ($\lambda_G = 0$ and $\epsilon_i = +1$)

$$\beta_{\text{NSVZ}}(0) = \beta_{1\text{-loop}}/3 = -109/3. \tag{3.8}$$

We use the Dynkin indices from [107, 108] summarized, for reader's convenience, for the lowest dimensional representations in $\text{SO}(10)$ in table 1.

Now, let's assume that an UV fixed point occurs nonperturbatively in the theory. Because we require $a_{\text{UV}} > a_{\text{IR}}$ at least one $\epsilon_i = -1$ must be negative implying that by continuity in λ_G we need to reach the point where

$$\lambda_G^{\text{max}} = \min_i \left(\frac{|r_i|}{T(r_i)} \right) = \frac{|126|}{T(126)} = \frac{|\overline{126}|}{T(\overline{126})} = \frac{126}{35}. \tag{3.9}$$

⁴The other choice, i.e. $\overline{126} + 120$, does not have enough parameters to reproduce the physical results. The reason being the antisymmetric nature of the matrix Y_{120} .

ϕ_i	10	16	45	54	120	126	144	210
$T(r_i)$	1	2	8	12	28	35	34	56

Table 1. Dynkin indices $T(r_i)$ for some low-dimensional representations ϕ_i of $\text{SO}(10)$.

However for this value of λ_G we find

$$\beta_{\text{NSVZ}}(\lambda_G^{\text{max}}) = 4 - \sqrt{\frac{11}{5}} > 0, \quad (3.10)$$

and therefore the $\beta_{\text{NSVZ}}(\lambda_G)$ must have changed sign between $\lambda_G = 0$ in the infrared and $\lambda_G = \lambda_G^{\text{max}}$. Assuming continuity an apparent fixed point exists $\lambda_G^* < \lambda_G^{\text{max}}$ for which⁵

$$\beta_{\text{NSVZ}}(\lambda_G^*) = 0, \quad \text{and} \quad \lambda_G^* = 3.57. \quad (3.11)$$

However for this value we find

$$a_{\text{UV}} = a(R_i(\lambda_G^*)) = 125 < 206 = a(R_i(0)) = a_{\text{IR}}, \quad (3.12)$$

showing that the alleged fixed point violates the a-theorem constraint expressed in (2.16) and therefore cannot be physical.

We now move to consider a more general matter field content of $\text{SO}(10)$ without superpotential and test whether one can achieve an acceptable UV fixed point. In practice we require:

- 1) No zero to appear in the $\beta_{\text{NSVZ}}(\lambda_G)$ for the branch connected to the perturbative IR region (i.e. with all $\epsilon_i = +1$) with $\lambda_G \leq \lambda_G^{\text{max}}$. Notice that λ_G^{max} differs for different theories;
- 2) A possible UV zero in the $\beta_{\text{NSVZ}}(\lambda_G)$ to occur for, at least, some negative ϵ 's, i.e. $\epsilon_k = -1$ with $\lambda_G \leq \lambda_G^{\text{max}} \equiv \min_i(|r_i|/T(r_i)) = |r_k|/T(r_k)$;
- 3) This solution must either satisfy $a_{\text{UV}} > a_{\text{IR}}$, or develop at least one non-interacting gauge invariant operator (GIO) and thus by eliminating the operator has a chance for a modified a .

We now perform a scan over the following two families of theories

- i) We first consider the same type of matter fields 10, 16 and/or $\overline{16}$, 126 and/or $\overline{126}$ and 210 but scan over the theories featuring from 0 up to 3 copies of each field. The total number of cases is therefore $4^4 - 1 = 255$. Only 240 of these combinations have a negative 1-loop β -function and are thus interesting to investigate. Clearly our first example, the one discussed in detail at the beginning of this section, corresponds to a special case of this family of theories, i.e. 1 multiplet of 10, 3 multiplets 16, 2 multiplets 126 (or $\overline{126}$) and one multiplet 210. Among these 240 theories we found that 37 of them satisfy point 1) and 2) above.

⁵Here and in the following we round all real numbers to 3 digits.

- ii) In the second example we consider the fields 10, 16 (or $\overline{16}$), 45, 54, 120 126 (or $\overline{126}$), 144 (or $\overline{144}$), and 210. We scan over all possibilities of having or not having each of these fields once. This means we consider $2^8 - 1 = 255$ different theories, of which again 240 have a negative 1-loop β -function. Of these theories 23 are found to satisfy point 1) and 2) above.

We were unable to find acceptable asymptotically safe solutions for any of the $2 \times 240 = 480$ different models above satisfying simultaneously the conditions 1), 2) and 3). These findings extend the results of [36]. Nevertheless exotic theories exist passing these tests such as the theory with 274909 generations of 10, and 5161 generations of 126 (part or all of them can be $\overline{126}$). The would be UV fixed point seems to occur for

$$\lambda_G^* = -28.5, \tag{3.13}$$

for which

$$a_{\text{UV}} - a_{\text{IR}} = 1.17 \times 10^4, \tag{3.14}$$

$$(R_{10}, R_{126}) = (0.346, 2.00), \tag{3.15}$$

$$b_{\text{UV}} = 2.99 \times 10^5 \times F_{10}^2, \tag{3.16}$$

$$c_{\text{UV}} = 4.60 \times 10^6. \tag{3.17}$$

All constraints are met and no GIO becomes non-interacting. There are other exotic solutions of this type, with some containing 3 generations of the 16 matter. These solutions are far from phenomenologically viable while help elucidating the difficulty in constructing asymptotically safe supersymmetric quantum field theories.

3.3 Minimal model with a superpotential

We now extend the analysis above to the case of a non-vanishing superpotential. We shall use here as well the continuity of the R-charges R_i as functions of the Lagrange multiplier λ_G , and further add Lagrange multipliers λ_a stemming from each new interaction in the superpotential W .

Let's therefore consider all the permitted trilinear terms in the superpotential [22–25]:

$$W = y_1 210^3 + y_2 210 126 \overline{126} + y_3 210 126 10 + y_4 210 \overline{126} 10 + \sum_{a,b=1,2,3} 16_a 16_b (y_{5,ab} 10 + y_{6,ab} \overline{126}) \tag{3.18}$$

The function a assumes the form

$$a = 2|G| + \sum_i |r_i| a_1(R_i) + \lambda_G \left(T(G) + \sum_i T(r_i) (R_i - 1) \right) + \lambda_1 (2 - 3R_{210}) + \lambda_2 (2 - R_{210} - R_{126} - R_{\overline{126}}) + \lambda_3 (2 - R_{210} - R_{126} - R_{10}) + \lambda_4 (2 - R_{210} - R_{\overline{126}} - R_{10}) + \sum_{a,b=1,2,3} \lambda_{5,ab} (2 - R_{10} - R_{16_a} - R_{16_b}) + \sum_{a,b=1,2,3} \lambda_{6,ab} (2 - R_{\overline{126}} - R_{16_a} - R_{16_b}). \tag{3.19}$$

If all the trilinear terms are present then the matter R -charges are constrained to be the free ones, i.e. $R_i = 2/3$ for any i forbidding a zero in the NSVZ beta function. A minimal approach is to remove just one field from the superpotential. It turns out that the best choice is one of the 16 fields which we choose to be 16_1 and the sum over a, b in (3.18) and (3.19) go over 2 and 3.

By extremizing the a -function we now obtain $R_{16_1} = 113/6$ and all the others fields still possess $R = 2/3$. The positivity requirements are satisfied, since

$$a_{UV} - a_{IR} = 2.72 \times 10^5 > 0, \quad \text{and} \quad c_{UV} = 8.16 \times 10^5 > 0. \quad (3.20)$$

while there are no extra flavor symmetries.

Therefore the present solution passes all known constraints needed by a superconformal fixed point. By construction our solution describes a world with a decoupled massless generation.

Our solution corresponds to a manifold of UV fixed points. In fact the 11 equations (we consider here the sums over $a, b = 2, 3$) $\partial a / \partial \lambda = 0$, which extremize the a -function (3.19), are expressed with only 7 combinations $R_i(\lambda)$ (got previously from $\partial a / \partial R = 0$). This means that from the numerical values of R_i in the UV limit we can determine

$$\lambda_G = -2.29 \times 10^4 \quad (3.21)$$

$$\lambda_1 = -1.64 \times 10^5 \quad (3.22)$$

$$\lambda_4 = 1.14 \times 10^4 \quad (3.23)$$

and the following linear combinations

$$\lambda_2 + \lambda_3 = -8.01 \times 10^5 \quad (3.24)$$

$$(\lambda_{5,22} + \lambda_{5,23} + \lambda_{5,33}) - \lambda_2 = 7.67 \times 10^5 \quad (3.25)$$

$$(\lambda_{6,22} + \lambda_{6,23} + \lambda_{6,33}) + \lambda_2 = -8.13 \times 10^5 \quad (3.26)$$

$$(\lambda_{5,33} + \lambda_{6,33}) - (\lambda_{5,22} + \lambda_{6,22}) = 0 \quad (3.27)$$

The solution we found is thus not a fixed point, but a manifold of fixed points, somehow reminiscent (although with the role of UV and IR inverted) of the cases considered in [109].

3.4 Gauge invariant fields becoming free

If a singlet scalar gauge invariant operator of the chiral ring

$$\mathcal{O}_\alpha = \prod_i \phi_i^{q_{\alpha i}}, \quad (3.28)$$

at the fixed point acquires

$$R_\alpha = \sum_i q_{\alpha i} R_i < 2/3, \quad (3.29)$$

unitarity is violated unless it becomes free. If this occurs one needs to modify the a function accordingly [94]

$$a(R_i) \rightarrow a(R_i) + \sum_\alpha (a_1(2/3) - a_1(R_\alpha)). \quad (3.30)$$

The maximization of the modified a function is subject to the same constraints discussed earlier. Note that these additional terms to the a -function naively tend to increase its value. This means that a candidate fixed point with $a_{UV} < a_{IR}$ can in principle turn into a candidate fixed point with $a_{UV} > a_{IR}$ once the contribution of all free GIOs are subtracted from a . For this reason rather than imposing $a_{UV} - a_{IR} > 0$ from the beginning it is better to consider every real solution stemming from maximizing the a -function. Here we limit the analysis to the positive R_i case with at least one $R_i < 1/3$ but without enforcing the bounds $b < 0$ and $a_{UV} - a_{IR} < 0$ or $c < 0$. This is because we expect that if one of the $R_i < 1/3$ some GIO can become free and the minimization analysis needs to be redone.

However, even the enlarged analysis, didn't return potentially relevant asymptotically safe candidates. The main reason is that in all cases only few GIOs can become free making it difficult to return a positive variation of the a -function. Specifically, in most of the cases, the smallest R_i is for the 10 chiral superfield that has only 10 10 as a GIO. In few cases also other fields like 126 or $\overline{126}$ or 16 have $R < 2/3$, but typically not very small, therefore it is hard to construct singlet operators with R -charges less than $2/3$, while in combination with 10 it is hard to get many more invariants. For example the invariant $126 10^5$ is antisymmetric in 10 so with one 10 only it vanishes.

Let's give an explicit example, i.e. the theory with the maximum among the negative $a_{UV} - a_{IR}$ we were able to find for all $R_i > 0$. This corresponds to the theory with superpotential

$$W = y_3 10 210 126 + y_4 10 210 \overline{126}. \tag{3.31}$$

We find upon maximization of a

$$R = (0.893, 1.00, 1.00, 0.104, 0.781, 0.781, 0.781), \tag{3.32}$$

$$b_{UV} = (1.37 \times 10^4, 27.4, 7.02, 7.00), \tag{3.33}$$

$$a_{UV} - a_{IR} = -98.2, \quad c_{UV} = 315.. \tag{3.34}$$

Only the 126 and $\overline{126}$ have $R > 1$ and thus $\epsilon = -1$. We find that only $R_{10} = 0.104$ is smaller than $1/3$ and that there is only one GIO, i.e. 10 10, with the correction $\Delta(a) = a_0(2/3) - a_0(2R_{10})$ to be added to the previous a . The new maximization yields (for the solution with all $b > 0$)

$$R = (0.897, 1.00, 1.00, 0.103, 0.777, 0.777, 0.777), \tag{3.35}$$

$$b_{UV} = (1.40 \times 10^4, 26.8, 7.15, 7.13), \tag{3.36}$$

$$a_{UV} - a_{IR} = -97.3, \quad c_{UV} = 317.. \tag{3.37}$$

Although $a_{UV} - a_{IR}$ is slightly larger it is still negative and the fixed point is excluded. Of course, also c and b receive small corrections.

So far we have investigated the case in which all R were positive. However one could have one or more negative R -charges. This interesting case will be investigated elsewhere.

3.5 On the doublet-triplet splitting problem

Grand unified theories require the SM Higgs to arise from representations of the unified group. These contain, besides the SM Higgs weak doublet, also other states that include

color triplets. In supersymmetric theories color triplet Higgses can induce dimension five supersymmetric operators mediating proton decay. Consequently one needs to keep the color triplet very heavy, typically heavier than the GUT scale, while keeping the doublet light for the theory to be viable. The doublet-triplet splitting (DT) problem begs the question: what keeps the doublets light and the triplets heavy? To ameliorate the severity of the problem several proposals have been made in the literature, such as the Dimopoulos-Wilczek mechanism that requires the introduction of an adjoint $SO(10)$ chiral field. The field can acquire, due to its antisymmetric nature, two independent vacuum expectation values: a non-vanishing one for the color triplet and a vanishing-one for the doublet [110]. This is also known as the missing VEV mechanism [110, 111]. Other possibilities are the missing partner [112–116] and the orbifold construction [117–124]. These mechanisms require another layer of model building in $SO(10)$ with the addition of extra fields such as the aforementioned 45 (missing VEV), additional $126 + \overline{126}$ (missing partner) or extra Kaluza-Klein states (orbifold).

Although a more thorough analysis of the possible occurrence of an UV fixed point in models directly addressing the DT problem and low energy R-symmetry will be performed elsewhere we can already discuss a special case here. By using the solution, found in section 3.3, where all chiral fields, including the extra ones needed for the DT splitting, are constrained to have R-charge equal to $2/3$ except for the 16_1 which is allowed to have a very large R-charge we can argue that this solution is a plausible candidate for an UV finite GUT theory.

4 Outlook and conclusions

We investigated the possibility for phenomenologically motivated supersymmetric grand unified theories to feature an interacting ultraviolet fixed point before reaching the gravity transition scale. Using a set of nonperturbative tools ranging from a-maximization to the positivity of relevant central charges we nonperturbatively rule out this possibility for a broad class of prime candidates. We have also discovered a less exotic candidate theory, passing these tests that, although features the physically relevant fields, is not yet phenomenologically viable. Nevertheless the exotic candidates simultaneously elucidate the challenges and hint to the required underlying structure of potentially viable asymptotically safe grand unified theories.

We focussed in this initial work on grand unified theories on the $SO(10)$ theory but we plan to extend the analysis to similar E_6 realizations [27, 28] as well as $SU(5)$ [112–114].

Acknowledgments

BB would like to thank CP3-Origins & the Danish IAS for hospitality, during which this work started. FS thanks the hospitality of the Lyon high energy group as *Professeur mois invite* where this work was finalized. The work of BB has been supported by the Slovenian Research Agency. FS acknowledges the support of the Danish National Research Foundation under the grant number DNR90.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] J.C. Pati and A. Salam, *Lepton Number as the Fourth Color*, *Phys. Rev. D* **10** (1974) 275 [*Erratum ibid.* **D 11** (1975) 703] [[INSPIRE](#)].
- [2] H. Georgi and S.L. Glashow, *Unity of All Elementary Particle Forces*, *Phys. Rev. Lett.* **32** (1974) 438 [[INSPIRE](#)].
- [3] H. Georgi, H.R. Quinn and S. Weinberg, *Hierarchy of Interactions in Unified Gauge Theories*, *Phys. Rev. Lett.* **33** (1974) 451 [[INSPIRE](#)].
- [4] S. Dimopoulos, S. Raby and F. Wilczek, *Supersymmetry and the Scale of Unification*, *Phys. Rev. D* **24** (1981) 1681 [[INSPIRE](#)].
- [5] L.E. Ibáñez and G.G. Ross, *Low-Energy Predictions in Supersymmetric Grand Unified Theories*, *Phys. Lett. B* **105** (1981) 439 [[INSPIRE](#)].
- [6] M.B. Einhorn and D.R.T. Jones, *The Weak Mixing Angle and Unification Mass in Supersymmetric SU(5)*, *Nucl. Phys. B* **196** (1982) 475 [[INSPIRE](#)].
- [7] W.J. Marciano and G. Senjanović, *Predictions of Supersymmetric Grand Unified Theories*, *Phys. Rev. D* **25** (1982) 3092 [[INSPIRE](#)].
- [8] H. Georgi, *The State of the Art — Gauge Theories*, *AIP Conf. Proc.* **23** (1975) 575 [[INSPIRE](#)].
- [9] H. Fritzsch and P. Minkowski, *Unified Interactions of Leptons and Hadrons*, *Annals Phys.* **93** (1975) 193 [[INSPIRE](#)].
- [10] F. Gursev, P. Ramond and P. Sikivie, *A Universal Gauge Theory Model Based on E6*, *Phys. Lett. B* **60** (1976) 177 [[INSPIRE](#)].
- [11] P. Minkowski, $\mu \rightarrow e\gamma$ at a Rate of One Out of 10^9 Muon Decays?, *Phys. Lett. B* **67** (1977) 421 [[INSPIRE](#)].
- [12] M. Gell-Mann, P. Ramond and R. Slansky, *Complex Spinors and Unified Theories*, in *Supergravity*, P. van Nieuwenhuizen and D.Z. Freedman eds., North Holland Publ. Co. (1979), pg. 315, *Conf. Proc. C* **790927** (1979) 315 [[arXiv:1306.4669](#)] [[INSPIRE](#)].
- [13] T. Yanagida, *Horizontal Gauge Symmetry And Masses Of Neutrinos*, in *Proceedings of the Workshop on the Baryon Number of the Universe and Unified Theories*, O. Sawada and A. Sugamoto eds., Tsukuba, Japan, 13–14 February 1979, pg. 95 [[INSPIRE](#)].
- [14] S.L. Glashow, *The Future Of Elementary Particle Physics*, in *Quarks and Leptons*, Cargèse Lectures, 9–29 July 1979, Plenum, New York (1980), pg. 687, *NATO Sci. Ser. B* **61** (1980) 687 [[INSPIRE](#)].
- [15] R.N. Mohapatra and G. Senjanović, *Neutrino Mass and Spontaneous Parity Violation*, *Phys. Rev. Lett.* **44** (1980) 912 [[INSPIRE](#)].
- [16] R.N. Mohapatra, *New Contributions to Neutrinoless Double beta Decay in Supersymmetric Theories*, *Phys. Rev. D* **34** (1986) 3457 [[INSPIRE](#)].

- [17] A. Font, L.E. Ibáñez and F. Quevedo, *Does Proton Stability Imply the Existence of an Extra Z_0 ?*, *Phys. Lett. B* **228** (1989) 79 [INSPIRE].
- [18] S.P. Martin, *Some simple criteria for gauged R-parity*, *Phys. Rev. D* **46** (1992) R2769 [hep-ph/9207218] [INSPIRE].
- [19] C.S. Aulakh, K. Benakli and G. Senjanović, *Reconciling supersymmetry and left-right symmetry*, *Phys. Rev. Lett.* **79** (1997) 2188 [hep-ph/9703434] [INSPIRE].
- [20] C.S. Aulakh, A. Melfo and G. Senjanović, *Minimal supersymmetric left-right model*, *Phys. Rev. D* **57** (1998) 4174 [hep-ph/9707256] [INSPIRE].
- [21] C.S. Aulakh, A. Melfo, A. Rasin and G. Senjanović, *Seesaw and supersymmetry or exact R-parity*, *Phys. Lett. B* **459** (1999) 557 [hep-ph/9902409] [INSPIRE].
- [22] C.S. Aulakh and R.N. Mohapatra, *Implications of Supersymmetric SO(10) Grand Unification*, *Phys. Rev. D* **28** (1983) 217 [INSPIRE].
- [23] T.E. Clark, T.-K. Kuo and N. Nakagawa, *A SO(10) Supersymmetric Grand Unified Theory*, *Phys. Lett. B* **115** (1982) 26 [INSPIRE].
- [24] K.S. Babu and R.N. Mohapatra, *Predictive neutrino spectrum in minimal SO(10) grand unification*, *Phys. Rev. Lett.* **70** (1993) 2845 [hep-ph/9209215] [INSPIRE].
- [25] C.S. Aulakh, B. Bajc, A. Melfo, G. Senjanović and F. Vissani, *The minimal supersymmetric grand unified theory*, *Phys. Lett. B* **588** (2004) 196 [hep-ph/0306242] [INSPIRE].
- [26] C.S. Aulakh, I. Garg and C.K. Khosa, *NMSGUT emergence and Trans-Unification RG flows*, [arXiv:1509.00422](https://arxiv.org/abs/1509.00422) [INSPIRE].
- [27] B. Bajc and V. Susič, *Towards the minimal renormalizable supersymmetric E_6 model*, *JHEP* **02** (2014) 058 [[arXiv:1311.0775](https://arxiv.org/abs/1311.0775)] [INSPIRE].
- [28] K.S. Babu, B. Bajc and V. Susič, *A minimal supersymmetric E_6 unified theory*, *JHEP* **05** (2015) 108 [[arXiv:1504.00904](https://arxiv.org/abs/1504.00904)] [INSPIRE].
- [29] D.F. Litim and F. Sannino, *Asymptotic safety guaranteed*, *JHEP* **12** (2014) 178 [[arXiv:1406.2337](https://arxiv.org/abs/1406.2337)] [INSPIRE].
- [30] D.F. Litim, M. Mojaza and F. Sannino, *Vacuum stability of asymptotically safe gauge-Yukawa theories*, *JHEP* **01** (2016) 081 [[arXiv:1501.03061](https://arxiv.org/abs/1501.03061)] [INSPIRE].
- [31] A.D. Bond and D.F. Litim, *Theorems for Asymptotic Safety of Gauge Theories*, [arXiv:1608.00519](https://arxiv.org/abs/1608.00519) [INSPIRE].
- [32] O. Antipin, M. Gillioz, E. Mølgaard and F. Sannino, *The a theorem for gauge-Yukawa theories beyond Banks-Zaks fixed point*, *Phys. Rev. D* **87** (2013) 125017 [[arXiv:1303.1525](https://arxiv.org/abs/1303.1525)] [INSPIRE].
- [33] C. Pica and F. Sannino, *UV and IR Zeros of Gauge Theories at The Four Loop Order and Beyond*, *Phys. Rev. D* **83** (2011) 035013 [[arXiv:1011.5917](https://arxiv.org/abs/1011.5917)] [INSPIRE].
- [34] B. Holdom, *Large- N flavor β -functions: a recap*, *Phys. Lett. B* **694** (2011) 74 [[arXiv:1006.2119](https://arxiv.org/abs/1006.2119)] [INSPIRE].
- [35] F. Sannino, *Exact results for (Super) Asymptotically Safe QFT*, plenary talk held for the 8th International Conference on the Exact Renormalization Group (ERG2016), Trieste, Italy, 19–23 September 2016, the slides of the presentation can be found at <http://indico.ictp.it/event/7608/session/3/contribution/10/material/slides/0.pdf>.

- [36] K. Intriligator and F. Sannino, *Supersymmetric asymptotic safety is not guaranteed*, *JHEP* **11** (2015) 023 [[arXiv:1508.07411](#)] [[INSPIRE](#)].
- [37] S.P. Martin and J.D. Wells, *Constraints on ultraviolet stable fixed points in supersymmetric gauge theories*, *Phys. Rev. D* **64** (2001) 036010 [[hep-ph/0011382](#)] [[INSPIRE](#)].
- [38] E. Molgaard and F. Sannino, *Phases of (Asymptotically) Safe Chiral Theories with(out) Scalars*, [arXiv:1610.03130](#) [[INSPIRE](#)].
- [39] J.K. Esbensen, T.A. Rytlov and F. Sannino, *Quantum critical behavior of semisimple gauge theories*, *Phys. Rev. D* **93** (2016) 045009 [[arXiv:1512.04402](#)] [[INSPIRE](#)].
- [40] S. Weinberg, *Ultraviolet Divergences In Quantum Theories Of Gravitation*, in *General Relativity: An Einstein centenary survey*, S.W. Hawking and W. Israel eds., Cambridge University Press (1979), pg. 790–831 [[INSPIRE](#)].
- [41] M. Niedermaier, *The asymptotic safety scenario in quantum gravity: An introduction*, *Class. Quant. Grav.* **24** (2007) R171 [[gr-qc/0610018](#)] [[INSPIRE](#)].
- [42] M. Niedermaier and M. Reuter, *The Asymptotic Safety Scenario in Quantum Gravity*, *Living Rev. Rel.* **9** (2006) 5.
- [43] R. Percacci, *Asymptotic Safety*, [arXiv:0709.3851](#) [[INSPIRE](#)].
- [44] M. Reuter and F. Saueressig, *Quantum Einstein Gravity*, *New J. Phys.* **14** (2012) 055022 [[arXiv:1202.2274](#)] [[INSPIRE](#)].
- [45] D.F. Litim, *Renormalisation group and the Planck scale*, *Phil. Trans. Roy. Soc. Lond. A* **369** (2011) 2759 [[arXiv:1102.4624](#)] [[INSPIRE](#)].
- [46] D.I. Kazakov, *Ultraviolet fixed points in gauge and SUSY field theories in extra dimensions*, *JHEP* **03** (2003) 020 [[hep-th/0209100](#)] [[INSPIRE](#)].
- [47] H. Gies, J. Jaeckel and C. Wetterich, *Towards a renormalizable standard model without fundamental Higgs scalar*, *Phys. Rev. D* **69** (2004) 105008 [[hep-ph/0312034](#)] [[INSPIRE](#)].
- [48] T.R. Morris, *Renormalizable extra-dimensional models*, *JHEP* **01** (2005) 002 [[hep-ph/0410142](#)] [[INSPIRE](#)].
- [49] P. Fischer and D.F. Litim, *Fixed points of quantum gravity in extra dimensions*, *Phys. Lett. B* **638** (2006) 497 [[hep-th/0602203](#)] [[INSPIRE](#)].
- [50] P. Fischer and D.F. Litim, *Fixed points of quantum gravity in higher dimensions*, *AIP Conf. Proc.* **861** (2006) 336 [[hep-th/0606135](#)] [[INSPIRE](#)].
- [51] D.I. Kazakov and G.S. Vartanov, *Renormalizable $1/N_f$ Expansion for Field Theories in Extra Dimensions*, *JHEP* **06** (2007) 081 [[arXiv:0707.2564](#)] [[INSPIRE](#)].
- [52] O. Zanusso, L. Zambelli, G.P. Vacca and R. Percacci, *Gravitational corrections to Yukawa systems*, *Phys. Lett. B* **689** (2010) 90 [[arXiv:0904.0938](#)] [[INSPIRE](#)].
- [53] H. Gies, S. Rechenberger and M.M. Scherer, *Towards an Asymptotic-Safety Scenario for Chiral Yukawa Systems*, *Eur. Phys. J. C* **66** (2010) 403 [[arXiv:0907.0327](#)] [[INSPIRE](#)].
- [54] J.-E. Daum, U. Harst and M. Reuter, *Running Gauge Coupling in Asymptotically Safe Quantum Gravity*, *JHEP* **01** (2010) 084 [[arXiv:0910.4938](#)] [[INSPIRE](#)].
- [55] G.P. Vacca and O. Zanusso, *Asymptotic Safety in Einstein Gravity and Scalar-Fermion Matter*, *Phys. Rev. Lett.* **105** (2010) 231601 [[arXiv:1009.1735](#)] [[INSPIRE](#)].

- [56] S. Folkerts, D.F. Litim and J.M. Pawłowski, *Asymptotic freedom of Yang-Mills theory with gravity*, *Phys. Lett. B* **709** (2012) 234 [[arXiv:1101.5552](#)] [[INSPIRE](#)].
- [57] F. Bazzocchi, M. Fabbrichesi, R. Percacci, A. Tonero and L. Vecchi, *Fermions and Goldstone bosons in an asymptotically safe model*, *Phys. Lett. B* **705** (2011) 388 [[arXiv:1105.1968](#)] [[INSPIRE](#)].
- [58] H. Gies, S. Rechenberger, M.M. Scherer and L. Zambelli, *An asymptotic safety scenario for gauged chiral Higgs-Yukawa models*, *Eur. Phys. J. C* **73** (2013) 2652 [[arXiv:1306.6508](#)] [[INSPIRE](#)].
- [59] O. Antipin, M. Mojaza and F. Sannino, *Conformal Extensions of the Standard Model with Veltman Conditions*, *Phys. Rev. D* **89** (2014) 085015 [[arXiv:1310.0957](#)] [[INSPIRE](#)].
- [60] P. Donà, A. Eichhorn and R. Percacci, *Matter matters in asymptotically safe quantum gravity*, *Phys. Rev. D* **89** (2014) 084035 [[arXiv:1311.2898](#)] [[INSPIRE](#)].
- [61] A. Bonanno and M. Reuter, *Cosmology of the Planck era from a renormalization group for quantum gravity*, *Phys. Rev. D* **65** (2002) 043508 [[hep-th/0106133](#)] [[INSPIRE](#)].
- [62] K.A. Meissner and H. Nicolai, *Conformal Symmetry and the Standard Model*, *Phys. Lett. B* **648** (2007) 312 [[hep-th/0612165](#)] [[INSPIRE](#)].
- [63] R. Foot, A. Kobakhidze, K.L. McDonald and R.R. Volkas, *A solution to the hierarchy problem from an almost decoupled hidden sector within a classically scale invariant theory*, *Phys. Rev. D* **77** (2008) 035006 [[arXiv:0709.2750](#)] [[INSPIRE](#)].
- [64] J. Hewett and T. Rizzo, *Collider Signals of Gravitational Fixed Points*, *JHEP* **12** (2007) 009 [[arXiv:0707.3182](#)] [[INSPIRE](#)].
- [65] D.F. Litim and T. Plehn, *Signatures of gravitational fixed points at the LHC*, *Phys. Rev. Lett.* **100** (2008) 131301 [[arXiv:0707.3983](#)] [[INSPIRE](#)].
- [66] M. Shaposhnikov and D. Zenhausern, *Quantum scale invariance, cosmological constant and hierarchy problem*, *Phys. Lett. B* **671** (2009) 162 [[arXiv:0809.3406](#)] [[INSPIRE](#)].
- [67] M. Shaposhnikov and D. Zenhausern, *Scale invariance, unimodular gravity and dark energy*, *Phys. Lett. B* **671** (2009) 187 [[arXiv:0809.3395](#)] [[INSPIRE](#)].
- [68] M. Shaposhnikov and C. Wetterich, *Asymptotic safety of gravity and the Higgs boson mass*, *Phys. Lett. B* **683** (2010) 196 [[arXiv:0912.0208](#)] [[INSPIRE](#)].
- [69] S. Weinberg, *Asymptotically Safe Inflation*, *Phys. Rev. D* **81** (2010) 083535 [[arXiv:0911.3165](#)] [[INSPIRE](#)].
- [70] G. 't Hooft, *Probing the small distance structure of canonical quantum gravity using the conformal group*, [arXiv:1009.0669](#) [[INSPIRE](#)].
- [71] M. Hindmarsh, D. Litim and C. Rahmede, *Asymptotically Safe Cosmology*, *JCAP* **07** (2011) 019 [[arXiv:1101.5401](#)] [[INSPIRE](#)].
- [72] T. Hur and P. Ko, *Scale invariant extension of the standard model with strongly interacting hidden sector*, *Phys. Rev. Lett.* **106** (2011) 141802 [[arXiv:1103.2571](#)] [[INSPIRE](#)].
- [73] B. Dobrich and A. Eichhorn, *Can we see quantum gravity? Photons in the asymptotic-safety scenario*, *JHEP* **06** (2012) 156 [[arXiv:1203.6366](#)] [[INSPIRE](#)].
- [74] G. Marques Tavares, M. Schmaltz and W. Skiba, *Higgs mass naturalness and scale invariance in the UV*, *Phys. Rev. D* **89** (2014) 015009 [[arXiv:1308.0025](#)] [[INSPIRE](#)].

- [75] C. Tamarit, *Running couplings with a vanishing scale anomaly*, *JHEP* **12** (2013) 098 [[arXiv:1309.0913](#)] [[INSPIRE](#)].
- [76] S. Abel and A. Mariotti, *Novel Higgs Potentials from Gauge Mediation of Exact Scale Breaking*, *Phys. Rev. D* **89** (2014) 125018 [[arXiv:1312.5335](#)] [[INSPIRE](#)].
- [77] O. Antipin, J. Krog, M. Mojaza and F. Sannino, *Stable $E\chi$ tensions with(out) gravity*, *Nucl. Phys. B* **886** (2014) 125 [[arXiv:1311.1092](#)] [[INSPIRE](#)].
- [78] M. Heikinheimo, A. Racioppi, M. Raidal, C. Spethmann and K. Tuominen, *Physical Naturalness and Dynamical Breaking of Classical Scale Invariance*, *Mod. Phys. Lett. A* **29** (2014) 1450077 [[arXiv:1304.7006](#)] [[INSPIRE](#)].
- [79] E. Gabrielli, M. Heikinheimo, K. Kannike, A. Racioppi, M. Raidal and C. Spethmann, *Towards Completing the Standard Model: Vacuum Stability, EWSB and Dark Matter*, *Phys. Rev. D* **89** (2014) 015017 [[arXiv:1309.6632](#)] [[INSPIRE](#)].
- [80] M. Holthausen, J. Kubo, K.S. Lim and M. Lindner, *Electroweak and Conformal Symmetry Breaking by a Strongly Coupled Hidden Sector*, *JHEP* **12** (2013) 076 [[arXiv:1310.4423](#)] [[INSPIRE](#)].
- [81] G.C. Dorsch, S.J. Huber and J.M. No, *Cosmological Signatures of a UV-Conformal Standard Model*, *Phys. Rev. Lett.* **113** (2014) 121801 [[arXiv:1403.5583](#)] [[INSPIRE](#)].
- [82] A. Eichhorn and M.M. Scherer, *Planck scale, Higgs mass and scalar dark matter*, *Phys. Rev. D* **90** (2014) 025023 [[arXiv:1404.5962](#)] [[INSPIRE](#)].
- [83] F. Sannino and I.M. Shoemaker, *Asymptotically Safe Dark Matter*, *Phys. Rev. D* **92** (2015) 043518 [[arXiv:1412.8034](#)] [[INSPIRE](#)].
- [84] N.G. Nielsen, F. Sannino and O. Svendsen, *Inflation from Asymptotically Safe Theories*, *Phys. Rev. D* **91** (2015) 103521 [[arXiv:1503.00702](#)] [[INSPIRE](#)].
- [85] A. Codello, K. Langæble, D.F. Litim and F. Sannino, *Conformal Gauge-Yukawa Theories away From Four Dimensions*, *JHEP* **07** (2016) 118 [[arXiv:1603.03462](#)] [[INSPIRE](#)].
- [86] J. Meibohm, J.M. Pawłowski and M. Reichert, *Asymptotic safety of gravity-matter systems*, *Phys. Rev. D* **93** (2016) 084035 [[arXiv:1510.07018](#)] [[INSPIRE](#)].
- [87] S. Ferrara and B. Zumino, *Transformation Properties of the Supercurrent*, *Nucl. Phys. B* **87** (1975) 207 [[INSPIRE](#)].
- [88] G. Mack, *All Unitary Ray Representations of the Conformal Group $SU(2,2)$ with Positive Energy*, *Commun. Math. Phys.* **55** (1977) 1 [[INSPIRE](#)].
- [89] B. Grinstein, K.A. Intriligator and I.Z. Rothstein, *Comments on Unparticles*, *Phys. Lett. B* **662** (2008) 367 [[arXiv:0801.1140](#)] [[INSPIRE](#)].
- [90] D. Anselmi, D.Z. Freedman, M.T. Grisaru and A.A. Johansen, *Nonperturbative formulas for central functions of supersymmetric gauge theories*, *Nucl. Phys. B* **526** (1998) 543 [[hep-th/9708042](#)] [[INSPIRE](#)].
- [91] D. Anselmi, J. Erlich, D.Z. Freedman and A.A. Johansen, *Positivity constraints on anomalies in supersymmetric gauge theories*, *Phys. Rev. D* **57** (1998) 7570 [[hep-th/9711035](#)] [[INSPIRE](#)].
- [92] K.A. Intriligator and B. Wecht, *The exact superconformal R symmetry maximizes a* , *Nucl. Phys. B* **667** (2003) 183 [[hep-th/0304128](#)] [[INSPIRE](#)].

- [93] K.A. Intriligator, *IR free or interacting? A proposed diagnostic*, *Nucl. Phys. B* **730** (2005) 239 [[hep-th/0509085](#)] [[INSPIRE](#)].
- [94] D. Kutasov, A. Parnachev and D.A. Sahakyan, *Central charges and $U(1)_R$ symmetries in $N = 1$ super Yang-Mills*, *JHEP* **11** (2003) 013 [[hep-th/0308071](#)] [[INSPIRE](#)].
- [95] K.A. Intriligator and B. Wecht, *RG fixed points and flows in SQCD with adjoints*, *Nucl. Phys. B* **677** (2004) 223 [[hep-th/0309201](#)] [[INSPIRE](#)].
- [96] V.A. Novikov, M.A. Shifman, A.I. Vainshtein and V.I. Zakharov, *Exact Gell-Mann-Low Function of Supersymmetric Yang-Mills Theories from Instanton Calculus*, *Nucl. Phys. B* **229** (1983) 381 [[INSPIRE](#)].
- [97] J.L. Cardy, *Is There a c Theorem in Four-Dimensions?*, *Phys. Lett. B* **215** (1988) 749 [[INSPIRE](#)].
- [98] Z. Komargodski and A. Schwimmer, *On Renormalization Group Flows in Four Dimensions*, *JHEP* **12** (2011) 099 [[arXiv:1107.3987](#)] [[INSPIRE](#)].
- [99] Z. Komargodski, *The Constraints of Conformal Symmetry on RG Flows*, *JHEP* **07** (2012) 069 [[arXiv:1112.4538](#)] [[INSPIRE](#)].
- [100] A.B. Zamolodchikov, *Irreversibility of the Flux of the Renormalization Group in a 2D Field Theory*, *JETP Lett.* **43** (1986) 730 [[INSPIRE](#)].
- [101] H. Osborn, *Derivation of a Four-dimensional c Theorem*, *Phys. Lett. B* **222** (1989) 97 [[INSPIRE](#)].
- [102] I. Jack and H. Osborn, *Analogs for the c Theorem for Four-dimensional Renormalizable Field Theories*, *Nucl. Phys. B* **343** (1990) 647 [[INSPIRE](#)].
- [103] N. Seiberg, *Electric-magnetic duality in supersymmetric nonAbelian gauge theories*, *Nucl. Phys. B* **435** (1995) 129 [[hep-th/9411149](#)] [[INSPIRE](#)].
- [104] K.A. Intriligator and N. Seiberg, *Lectures on supersymmetric gauge theories and electric-magnetic duality*, *Nucl. Phys. Proc. Suppl.* **45BC** (1996) 1 [[hep-th/9509066](#)] [[INSPIRE](#)].
- [105] D. Kutasov, *New results on the ‘a theorem’ in four-dimensional supersymmetric field theory*, [hep-th/0312098](#) [[INSPIRE](#)].
- [106] D. Kutasov and A. Schwimmer, *Lagrange multipliers and couplings in supersymmetric field theory*, *Nucl. Phys. B* **702** (2004) 369 [[hep-th/0409029](#)] [[INSPIRE](#)].
- [107] R. Slansky, *Group Theory for Unified Model Building*, *Phys. Rept.* **79** (1981) 1 [[INSPIRE](#)].
- [108] R. Feger and T.W. Kephart, *LieART — A Mathematica application for Lie algebras and representation theory*, *Comput. Phys. Commun.* **192** (2015) 166 [[arXiv:1206.6379](#)] [[INSPIRE](#)].
- [109] R.G. Leigh and M.J. Strassler, *Exactly marginal operators and duality in four-dimensional $N = 1$ supersymmetric gauge theory*, *Nucl. Phys. B* **447** (1995) 95 [[hep-th/9503121](#)] [[INSPIRE](#)].
- [110] S. Dimopoulos and F. Wilczek, *Incomplete multiplets in supersymmetric unified models*, [NSF-ITP-82-07](#) (1981).
- [111] K.S. Babu and S.M. Barr, *Natural suppression of Higgsino mediated proton decay in supersymmetric $SO(10)$* , *Phys. Rev. D* **48** (1993) 5354 [[hep-ph/9306242](#)] [[INSPIRE](#)].

- [112] B. Grinstein, *A Supersymmetric SU(5) Gauge Theory with No Gauge Hierarchy Problem*, *Nucl. Phys. B* **206** (1982) 387 [[INSPIRE](#)].
- [113] A. Masiero, D.V. Nanopoulos, K. Tamvakis and T. Yanagida, *Naturally Massless Higgs Doublets in Supersymmetric SU(5)*, *Phys. Lett. B* **115** (1982) 380 [[INSPIRE](#)].
- [114] J. Hisano, T. Moroi, K. Tobe and T. Yanagida, *Suppression of proton decay in the missing partner model for supersymmetric SU(5) GUT*, *Phys. Lett. B* **342** (1995) 138 [[hep-ph/9406417](#)] [[INSPIRE](#)].
- [115] K.S. Babu, I. Gogoladze and Z. Tavartkiladze, *Missing Partner Mechanism in SO(10) Grand Unification*, *Phys. Lett. B* **650** (2007) 49 [[hep-ph/0612315](#)] [[INSPIRE](#)].
- [116] K.S. Babu, I. Gogoladze, P. Nath and R.M. Syed, *Variety of SO(10) GUTs with Natural Doublet-Triplet Splitting via the Missing Partner Mechanism*, *Phys. Rev. D* **85** (2012) 075002 [[arXiv:1112.5387](#)] [[INSPIRE](#)].
- [117] Y. Kawamura, *Triplet doublet splitting, proton stability and extra dimension*, *Prog. Theor. Phys.* **105** (2001) 999 [[hep-ph/0012125](#)] [[INSPIRE](#)].
- [118] G. Altarelli and F. Feruglio, *SU(5) grand unification in extra dimensions and proton decay*, *Phys. Lett. B* **511** (2001) 257 [[hep-ph/0102301](#)] [[INSPIRE](#)].
- [119] L.J. Hall and Y. Nomura, *Gauge unification in higher dimensions*, *Phys. Rev. D* **64** (2001) 055003 [[hep-ph/0103125](#)] [[INSPIRE](#)].
- [120] A. Hebecker and J. March-Russell, *A minimal $S^1/(Z_2 \times Z_2)$ orbifold GUT*, *Nucl. Phys. B* **613** (2001) 3 [[hep-ph/0106166](#)] [[INSPIRE](#)].
- [121] A. Hebecker and J. March-Russell, *The structure of GUT breaking by orbifolding*, *Nucl. Phys. B* **625** (2002) 128 [[hep-ph/0107039](#)] [[INSPIRE](#)].
- [122] T. Asaka, W. Buchmüller and L. Covi, *Gauge unification in six-dimensions*, *Phys. Lett. B* **523** (2001) 199 [[hep-ph/0108021](#)] [[INSPIRE](#)].
- [123] L.J. Hall, Y. Nomura, T. Okui and D. Tucker-Smith, *SO(10) unified theories in six-dimensions*, *Phys. Rev. D* **65** (2002) 035008 [[hep-ph/0108071](#)] [[INSPIRE](#)].
- [124] R. Dermisek and A. Mafi, *SO(10) grand unification in five-dimensions: Proton decay and the mu problem*, *Phys. Rev. D* **65** (2002) 055002 [[hep-ph/0108139](#)] [[INSPIRE](#)].