

Displaced heavy neutrinos from Z' decays at the LHC

Cheng-Wei Chiang,^a Giovanna Cottin,^{a,b} Arindam Das^c and Sanjoy Mandal^{d,e}

^a*Department of Physics, National Taiwan University,
Taipei 10617, Taiwan*

^b*Instituto de Física, Pontificia Universidad Católica de Chile,
Avenida Vicuña Mackenna 4860, Santiago, Chile*

^c*Department of Physics, Osaka University,
Toyonaka, Osaka 560-0043, Japan*

^d*Institute of Physics, Sachivalaya Marg,
Bhubaneswar 751005, India*

^e*Homi Bhabha National Institute, BARC Training School Complex,
Anushakti Nagar, Mumbai 400094, India*

E-mail: chengwei@phys.ntu.edu.tw, gfcottin@uc.cl,
arindam.das@hetmail.phys.sci.osaka-u.ac.jp, smandal@iopb.res.in

ABSTRACT: We study the LHC sensitivity to probe a long-lived heavy neutrino N in the context of Z' models. We focus on displaced vertex signatures of N when pair produced via a Z' , decaying to leptons and jets inside the inner trackers of the LHC experiments. We explore the LHC reach with current long-lived particle search strategies for either one or two displaced vertices in association with hadronic tracks or jets. We focus on two well-motivated models, namely, the minimal $U(1)_{B-L}$ scenario and its $U(1)_X$ extension. We find that searches for at least one displaced vertex can cover a significant portion of the parameter space, with light-heavy neutrino mixings as low as $|V_{lN}|^2 \approx 10^{-17}$, and $l = e, \mu$ accessible across GeV scale heavy neutrino masses.

KEYWORDS: Beyond Standard Model, Neutrino Physics

ARXIV EPRINT: [1908.09838](https://arxiv.org/abs/1908.09838)

Contents

1	Introduction	1
2	The model	2
3	LHC sensitivity with displaced vertex searches	6
3.1	Reach with ATLAS 1DV ID	7
3.2	Reach with CMS 2DV + jets	9
4	Conclusions	12
A	Useful formulas	15

1 Introduction

The observed light mass of neutrinos in the Standard Model (SM) begs for new physics explanations. The effects of such new physics are being actively looked for at the Large Hadron Collider (LHC) and several other experimental facilities worldwide. Such light neutrino masses can be explained by employing the so-called seesaw mechanism [1–3] that in its simplest form introduces new, heavy right-handed neutrinos, which can mix with the light neutrinos in the SM. For small enough values of the mixing, the heavy neutrinos can be long-lived, leading to macroscopic decays in the LHC experiments. Such decays can be reconstructed as displaced vertices (DVs) inside the inner trackers of the LHC detectors. Growing attention to these signatures has taken place over recent years (for a recent state-of-art review of long-lived particle searches at the LHC, see ref. [4]), as null results at the LHC may point to the possibilities that the new physics has more complex decay patterns, and that its effects may have been overlooked or misidentified by standard searches. New physics may become evident in these spectacular displaced signatures, as the SM can hardly mimic them. Their very small backgrounds will continue to make them attractive and their current study is of top importance for the high luminosity run of the LHC and future experimental facilities [5–7].

Heavy neutrinos are predicted in several models of new physics beyond the SM. Of particular interest are the $B - L$ extensions of the SM, which have an extended gauge symmetry $U(1)_{B-L}$ [8] and an associated new heavy Z' vector boson. The $U(1)_{B-L}$ symmetry can be broken spontaneously by the addition of a new SM-singlet Higgs field that attains a vacuum expectation value (VEV), and the theory includes three-generations of right-handed (or sterile) heavy neutrinos N^i , enabling the seesaw mechanism of light neutrino mass generation. The N^i can be produced from a SM-like Higgs associated with the $B - L$ breaking, or pair produced at colliders via a Z' . It can further decay with a displaced vertex (DV) depending on its couplings and mass.

Dedicated experimental searches for a Z' decaying to lepton pairs by the CMS collaboration placed a bound on the Z' mass to be $m_{Z'} > 4.5$ TeV [9] (assuming a SM-like gauge coupling). Recently the ATLAS collaboration analyzed the full Run 2 dataset [10], excluding a Z' just below 5 TeV. For a broad review on Z' models and early LHC strategies, see ref. [11] and references therein (see also refs. [12–17] for other collider studies in the $B-L$ case). ATLAS has now implemented a search that targets at displaced heavy neutrinos as benchmark, in the scenario where only one extra right-handed neutrino (produced in W boson decays) is added to the SM [18]. Current and proposed displaced strategies in several other heavy neutrino models are an attractive focus of research in recent years [19–42].

LHC constraints for the minimal $U(1)_{B-L}$ model were addressed through a global fit in ref. [43] for several choices of model parameters, but signatures involving displaced heavy neutrinos were not considered. Recent works on displaced neutrinos in $U(1)_{B-L}$ models have focused on displaced signatures coming from Higgs bosons due to a higher production cross section [44, 45]. For this reason, production via a Z' has had less attention. Early displaced strategies for a simplified model were recast in ref. [35], with focus on a benchmark scenario with relatively unboosted N . Recently, the authors in ref. [46] estimated the reach of future lifetime frontier experiments (like FASER [6, 47] and MATHUSLA [5]) on a rather light N and Z' , of $\mathcal{O}(\text{GeV})$ masses. In this work, we focus on higher masses and the LHC capabilities running at high luminosity, by reinterpreting ongoing DV searches at ATLAS and CMS. We also investigate prospects in a more general scenario than the $B-L$ model, the so-called non-exotic $U(1)_X$ extension of the SM [48], as it has been shown in refs. [15, 49] that an enhancement in the Z' production is possible, providing increased sensitivity to more complex scenarios in the search for displaced heavy neutrinos when they come from a Z' . The lifetime of N as a function of the lightest neutrino mass under the general $U(1)_X$ -extended models have been studied in refs. [17, 50].

The paper is organized as follows. We summarize the model under study in the section 2. We also extract the exclusion region in the parameter space of new gauge coupling and Z' mass using the Drell-Yan processes measured by both ATLAS and CMS. In section 3, we discuss the ATLAS and CMS displaced searches, reinterpret their results, and identify discovery prospects at the high luminosity LHC. We summarize and conclude in section 4.

2 The model

We consider an extension of the SM to have the gauge group $SU(3)_c \times SU(2)_L \times U(1)_Y \times U(1)_X$, where $U(1)_X$ is realized as a linear combination of the SM $U(1)_Y$ and the $U(1)_{B-L}$ symmetries [51], known as the non-exotic $U(1)_X$ extension of the SM [48]. The model is free from all the gauge and mixed gauge-gravity anomalies due to the presence of three generations of right-hand neutrinos (RHNs) N^i (with $i = 1, 2, 3$) [52, 53]. A new scalar field Φ is also introduced to break the $U(1)_X$ symmetry by attaining a VEV. The particle content is given in table 1.

	SU(3) _c	SU(2) _L	U(1) _Y	U(1) _X
q_L^i	3	2	+1/6	$\frac{1}{6}x_H + \frac{1}{3}x_\Phi$
u_R^i	3	1	+2/3	$\frac{2}{3}x_H + \frac{1}{3}x_\Phi$
d_R^i	3	1	-1/3	$-\frac{1}{3}x_H + \frac{1}{3}x_\Phi$
ℓ_L^i	1	2	-1/2	$-\frac{1}{2}x_H - x_\Phi$
e_R^i	1	1	-1	$-x_H - x_\Phi$
H	1	2	+1/2	$\frac{1}{2}x_H$
N^i	1	1	0	$-x_\Phi$
Φ	1	1	0	$2x_\Phi$

Table 1. The particle content and gauge charges of the non-exotic U(1)_X model [48]. The choice of $x_H = 0$ and $x_\Phi = 1$ corresponds to the $B - L$ case.

The charges of the particles are controlled by two parameters only, x_H and x_Φ , as seen in table 1. As the U(1)_X gauge group can be defined as a linear combination of the SM U(1)_Y and the U(1)_{B-L}, by setting $x_H = 0$ and $x_\Phi = 1$, we recover the minimal $B - L$ scenario [51, 54]. Therefore, without loss of generality, we fix $x_\Phi = 1$ in our analysis throughout the paper.

The Yukawa sector of the model can be written in a gauge-invariant way as

$$\begin{aligned}
 \mathcal{L}_Y = & - \sum_{\alpha, \beta=1}^3 Y_u^{\alpha\beta} \bar{q}_L^\alpha \tilde{H} u_R^\beta - \sum_{\alpha, \beta=1}^3 Y_d^{\alpha\beta} \bar{q}_L^\alpha H d_R^\beta - \sum_{\alpha, \beta=1}^3 Y_e^{\alpha\beta} \bar{\ell}_L^\alpha H e_R^\beta \\
 & - \sum_{\alpha, \beta=1}^3 Y_D^{\alpha\beta} \bar{\ell}_L^\alpha \tilde{H} N^\beta - \sum_{\alpha=1}^3 Y_N^\alpha \Phi \overline{N^{\alpha C}} N^\alpha + \text{h.c.}, \tag{2.1}
 \end{aligned}$$

where α, β are generation indices, $\tilde{H} \equiv i\tau^2 H^*$, and C denotes the charge conjugation. The fourth and fifth terms in eq. (2.1) are the Dirac and Majorana Yukawa terms. Without loss of generality, we use a diagonal basis for the Majorana Yukawa matrix. After the breaking of the U(1)_X and the electroweak symmetries, the U(1)_X gauge boson Z' mass, the Majorana and neutrino Dirac masses are generated, given by

$$\begin{aligned}
 M_{Z'} &= g'_1 \sqrt{4v_\Phi^2 + \frac{1}{4}x_H^2 v_{\text{SM}}^2} \simeq 2g'_1 v_\Phi, \\
 M_{N^\alpha} &= \frac{Y_N^\alpha}{\sqrt{2}} v_\Phi, \\
 M_D^{\alpha\beta} &= \frac{Y_D^{\alpha\beta}}{\sqrt{2}} v_{\text{SM}}, \tag{2.2}
 \end{aligned}$$

where g'_1 is the U(1)_X gauge coupling, v_Φ is the VEV of Φ , and $v_{\text{SM}} = 246 \text{ GeV}$ is the SM Higgs VEV. To be in agreement with LEP constraints, we take $v_\Phi \gg v_{\text{SM}}$ [55, 56]. LHC constraints will be discussed shortly.

In this model, through the $U(1)_X$ symmetry breaking, the Majorana mass terms for the RHNs are generated, and the light neutrinos acquire masses via the seesaw mechanism [1, 3]. The neutrino mass matrix is given by

$$m_\nu = \begin{pmatrix} 0 & M_D \\ M_D^T & M_N \end{pmatrix}. \quad (2.3)$$

Considering $|M_D^{\alpha\beta}/M_N^\alpha| \ll 1$ and diagonalizing the neutrino mass matrix in eq. (2.3), we obtain the light neutrino mass eigenvalue matrix as

$$m_\nu \simeq -M_D M_N^{-1} M_D^T. \quad (2.4)$$

The heavy neutrinos are SM gauge singlets. They interact with the W and Z bosons via mixing with the SM neutrinos. As a result, the SM neutrino flavor eigenstate (ν) can be expressed as a linear combination of the light mass eigenstates (ν^m) and heavy mass eigenstates (N^n):

$$\nu_\ell \simeq U_{\ell m} \nu^m + V_{\ell n} N^n, \quad (2.5)$$

where ℓ and m are generation indices, $U_{\ell m}$ is the 3×3 light neutrino mixing matrix and it is identical to the PMNS matrix at the leading order (ignoring effects of non-unitarity), and

$$V_{\ell n} \simeq M_D M_N^{-1} \quad (2.6)$$

is the mixing between the SM neutrinos and the heavy neutrinos assumed to be much less than 1.

The charged-current interactions can be expressed in terms of the neutrino mass eigenstates as

$$\mathcal{L}_{CC} \supset -\frac{g}{\sqrt{2}} W_\mu \bar{e} \gamma^\mu P_L V_{\ell n} N^n + \text{h.c.}, \quad (2.7)$$

where e represents the three generations of the charged leptons, and $P_L = \frac{1}{2}(1 - \gamma_5)$ is the projection operator. Similarly, in terms of the mass eigenstates, the neutral-current interactions are written as

$$\mathcal{L}_{NC} \supset -\frac{g}{2c_w} Z_\mu \left[\bar{N}^m \gamma^\mu P_L (V^\dagger V)_{mn} N^n + \left\{ \bar{\nu}^m \gamma^\mu P_L (U^\dagger V)_{mn} N^n + \text{h.c.} \right\} \right], \quad (2.8)$$

where $c_w \equiv \cos \theta_w$ with θ_w being the weak mixing angle.

Due to the nonzero $U(1)_X$ charges, the Z' boson interacts with the particles in the same way as it does in the $B - L$ scenario. However, the couplings of such interactions will depend upon the x_H and x_Φ parameters. As we have already fixed $x_\Phi = 1$, the corresponding partial decay widths of Z' into fermions, which are of interest in this work, will depend upon the choice of x_H . The expressions are given in the appendix.

The interaction between the Z' and the quarks can be written as

$$\mathcal{L}_{\text{int}} = -g'_1 \left(\bar{q}_L \gamma_\mu Q_{x_L}^q q_L + \bar{q}_R \gamma_\mu Q_{x_R}^q q_R \right) Z'_\mu, \quad (2.9)$$




Figure 1. Reinterpreted 95% CL exclusion limits based on the ATLAS [10] and CMS [9] results in the search for a heavy resonance decaying to lepton pairs, in the context of the $U(1)_{B-L}$ (left) and $U(1)_X$ (right) models.

where q_L (q_R) is the left- (right-) handed quark and $Q_{x_L}^q$ ($Q_{x_R}^q$) is the corresponding $U(1)_X$ charge. The corresponding interaction between the lepton sector and Z' can be written as

$$\mathcal{L}_{\text{int}} = -g'_1 \left(\bar{\ell}_L \gamma_\mu Q_{x_L}^\ell \ell_L + \bar{e}_R \gamma_\mu Q_{x_R}^\ell e_R \right) Z'_\mu, \quad (2.10)$$

where ℓ_L (e_R) is the left- (right-) handed lepton and the $Q_{x_L}^\ell$ ($Q_{x_R}^\ell$) is the corresponding $U(1)_X$ charge. All these charges are given in table 1 as a function of x_H and x_Φ .

For the rest of the paper, we choose two values for x_H : (a) $x_H = 0$, that corresponds to the $B - L$ case, or (b) $x_H = -1.2$, which is the value found to maximize the branching ratio of Z' to RHNs [49]. These charge assignments determine our labels for the $U(1)_{B-L}$ and $U(1)_X$ models in what follows. We also set the mixing between the SM Higgs and the new scalar Φ to zero, as this ensures pair production of the RHNs through a Z' .

With the above choices, we implement both the $U(1)_X$ and $U(1)_{B-L}$ models in the UFO format [57]. We first study the LHC process $pp \rightarrow Z' \rightarrow \ell^+ \ell^-$ for $\ell = e, \mu$. We reinterpret the latest ATLAS [10] and CMS [9] limits on both models, after validating the exclusions for the sequential Z'_{SSM} model the experiments present. For CMS, we consider the last 13-TeV result available, corresponding to an integrated luminosity of 36/fb. For ATLAS we make use of the latest full Run-II result with 139/fb.

Events are generated for different $(m_{Z'}, g'_1)$ values for both models with MADGRAPH5 [58] at the leading order. By comparing with the combined experimental upper limits (figure 3b of ref. [10] at $\Gamma/m = 0.5\%$, and figure 3c of ref. [9]), we can obtain lower (upper) bounds on $m_{Z'}$ (g'_1). Figure 1 shows the excluded region at 95% CL. Our ATLAS result is consistent with figure 3a of ref. [43], performed with their previous dataset.




Figure 2. Heavy neutrino N production and decay at the LHC in the Z' models considered in this work. Production proceeds through a Z' , which can be either on-shell or off-shell. The neutrino then decays to a lepton (either e or μ) and jets, giving visible charged decay products coming from the DV. The DV positions are marked by purple circles.

We use these bounds to fix a benchmark scenario of $(m_{Z'}, g'_1) = (6 \text{ TeV}, 0.8)$ for the rest of the paper, consistent with current resonance searches at the LHC.

We now focus on heavy neutrino production and decay at the LHC via a Z' . We scan m_N from 20 GeV to 1200 GeV, and $|V_{lN}|^2$ from 10^{-5} to 10^{-20} . Existing constraints in the light-heavy neutrino mixing and mass plane for $m_N \geq 20 \text{ GeV}$ exclude $|V_{lN}|^2 > 10^{-6}$. LEP data excludes $|V_{lN}|^2 > 10^{-5}$ [59], where DELPHI places limits on right-handed states produced in Z boson decays. Prompt collider LHC searches for three leptons in the final state exclude mixings $|V_{\mu N}|^2 > 10^{-5}$ [18, 60]. Also, the ATLAS displaced search for sterile neutrinos places bounds on $|V_{\mu N}|^2$ to be less than 10^{-6} [18] for masses of tens of GeV. Neutrinoless double beta decay ($0\nu\beta\beta$) experiments such as GERDA can also constrain $|V_{eN}|^2$. In ref. [27], an updated limit in the case where the SM is extended with only one right-handed neutrino produced in the decay of W boson excludes values above 10^{-7} . We will show in the next section that strategies presented in this work are more stringent than all of these constraints.

A diagram showing the N production and displaced decay considered here is presented in figure 2. The cross section of heavy neutrino pair production and that with heavy neutrino decays are shown in figure 3 for both models.

3 LHC sensitivity with displaced vertex searches

We focus on light-heavy neutrino mixing in the electron and muon sector separately (V_{lN} , with $l = e, \mu$); so only one N^i will couple to either electrons or muons. In both cases, only one neutrino, now defined as $N^i = N$, is in the kinematical region of interest. We adapt




Figure 3. Pair production cross section of right handed neutrinos (left) and that including their further decays to leptons and jets (right) as a function of the heavy neutrino mass m_N , for $(m_{Z'}, g'_1) = (6 \text{ TeV}, 0.8)$. Curves for both the $U(1)_X$ and $U(1)_{B-L}$ models are shown.

the UFOs so that the light-heavy neutrino mixing (V_{lN}) and the sterile neutrino masses (m_N) are treated as independent parameters.

We study the production process $pp \rightarrow Z' \rightarrow NN$, with each heavy neutrino decaying via $N \rightarrow ljj$ and $l = e$ or μ . Event generation is performed with MADGRAPH5 [58] at the leading order and the decay of N is processed with MADSPIN [61]. We save the decay information by setting the `time_of_flight` variable inside MADGRAPH5's `run_card`. The generated events are then interfaced to PYTHIA8 v2.3 [62] for showering and hadronization. Plots are generated with `matplotlib` [63].

We first reconstruct objects at the truth level and then apply efficiency corrections depending on the displaced search strategy of interest. We consider two complementary searches, each sensitive to a particular proper lifetime of the heavy neutrino. In figure 4 we show the naive reach in the proper lifetime and mass plane, for fixed mixing. We label the searches that we recast as: the “ATLAS 1DV ID” search for at least one DV in the ATLAS Inner Detector [64] and the “CMS 2DV+jets” [65] search for DV pairs in association with jets. The former is a proposal inspired by searches for multitrack DVs at ATLAS [64, 66]. The latter is a direct reinterpretation of the CMS search for two DVs in the inner tracker and jets [65].

3.1 Reach with ATLAS 1DV ID

The 8-TeV ATLAS 1DV ID search [64] looks for high track-multiplicity DVs in events possessing at least one DV and either leptons, jets or missing transverse momenta. The 13-TeV version of the search in ref. [66] only triggers on missing transverse momenta, but provides a prescription to implement parametrized efficiencies for DVs as a function of the




Figure 4. Proper decay distance $c\tau_N$ as a function of N mass and light-heavy neutrino mixing $|V_{lN}|^2$. The shaded region represents roughly the region that can be accessed with current DV searches at CMS [65] and the ATLAS inner tracker [66].

vertex invariant mass and number of charged tracks. Therefore, in this work we propose a 13-TeV search with the same displaced reconstruction but with a lepton trigger (as in the 8-TeV search).

The DV reconstruction in both 8- and 13-TeV searches is very similar to the experiment’s recent search for displaced heavy neutrinos in ref. [18]. Tunes of the former ATLAS multitrack search to target at displaced neutrinos were proposed in ref. [28], matching closely the actual experimental cuts in ref. [18]. Our vertex implementation in this work is basically the same as done in ref. [28].

The DV is reconstructed inside the ATLAS inner tracker from all charged particle tracks. The search triggers on a lepton, which is required to be associated with the DV. Cuts on the invariant mass of the vertex m_{DV} and the amount of charged particle tracks N_{trk} coming from it ensure this search to be free of backgrounds.¹ We follow the same optimized cuts for m_{DV} and N_{trk} justified in ref. [28].

We have to identify electrons, muons, tracks and DVs. For muons (electrons), we use a flat identification efficiency of 90% (70%). We apply vertex-level efficiencies as a function of DV distance, mass and number of tracks, using the parametrized selection efficiencies from the ATLAS 13-TeV DV search [66]. A validation study for these efficiencies was performed for the Les Houches PhysTeV 2017 proceedings in ref. [67].

¹Backgrounds to this search are mostly instrumental, from random crossings of tracks that fake a DV, and the background from heavy flavour, which are mostly B hadrons that merge to fake vertices, or are crossed by a random track, making a high mass “fake” vertex.

Trigger	Muon: $ \eta < 1.07$ and $p_T > 55$ GeV Electron: $ \eta < 2.47$ and $p_T > 120$ GeV
DV region	DV within $4 \text{ mm} < r_{DV} < 300 \text{ mm}$ and $ z_{DV} < 300 \text{ mm}$
DV selection	Made from tracks with $ d_0 > 2 \text{ mm}$ and with $p_T > 1 \text{ GeV}$ DV track multiplicity $N_{trk} \geq 4$ and invariant mass $m_{DV} \geq 5 \text{ GeV}$

Table 2. Cuts for the ATLAS 1DV ID proposed search. These are optimized as in ref. [27], and are inspired by the ATLAS search [64].

Events are first selected by triggering on a lepton. For electrons we require $p_T > 120 \text{ GeV}$ within $|\eta| < 2.47$. For muons, we demand $p_T > 55 \text{ GeV}$ and within $|\eta| < 1.07$. DVs are selected by reconstructing tracks with a large transverse impact parameter,² $|d_0| > 2 \text{ mm}$ and with $p_T > 1 \text{ GeV}$. The vertex position must be between 4 mm and 300 mm and must have at least 4 tracks, $N_{trk} \geq 4$. The lepton that fires the trigger must be associated with the DV. We account for this by truth-matching the lepton index with one of the displaced tracks. Note that no isolation requirement is applied on leptons. Finally, the invariant mass of the DV, m_{DV} , is calculated assuming all tracks have the mass of the pion and it is required to be larger than 5 GeV. A summary of all selections can be found in table 2.

The event level efficiency of this strategy, after all selections, is shown in figure 5 for the $U(1)_X$ model. The efficiency has a cigar-like shape, and is bounded by the case when the neutrinos are decaying either too promptly or too far away, outside of the detector’s acceptance. For a fixed mass, such as $m_N = 100 \text{ GeV}$, and mixings bigger than $\sim 10^{-10}$, the neutrino already decays too promptly. The efficiency in this case goes down with increasing mixing for a fixed heavy neutrino mass. For fixed mixing and smaller masses, the neutrinos are decaying outside of the tracker’s acceptance.

We show in figure 6 the estimated number of signal events at 13 TeV and 3000 fb^{-1} for both electron and muon mixing, for the $U(1)_{B-L}$ model. One can set a 95%CL exclusion region with at least 3 signal events, which is reasonable to set as a requirement for discovery in the absence of background. Analogous plots for the $U(1)_X$ model are shown in figure 7. A larger parameter region can be excluded in the $U(1)_X$ model due to its higher cross section, although the strategy is sensitive to both models. Mixings as low as $\sim 10^{-16}$ can be accessed for $m_N \sim 500 \text{ GeV}$ in the $U(1)_{B-L}$ model, and $\sim 10^{-17}$ for $m_N \sim 1 \text{ TeV}$ in the $U(1)_X$ model.

3.2 Reach with CMS 2DV + jets

The above search required at least one DV reconstructed in the inner tracker. Now we explore the reach with a search requiring exactly two DVs. This has the immediate advantage of being a search free from backgrounds. In addition, with the identification of two DVs we can access the mass of the Z' boson, when combining the kinematics with DV information [68].

²The transverse impact parameter is defined as $d_0 = r \times \Delta\phi$, with r being the transverse distance of the track and $\Delta\phi$ being the azimuthal angle between the decay product and the trajectory of the long-lived neutrino.




Figure 5. Representative event level efficiency of the ATLAS 1DV ID search as a function of m_N and $|V_{\mu N}|^2$.




Figure 6. Number of signal events for the $U(1)_{B-L}$ model at $\sqrt{s} = 13$ TeV expected in $\mathcal{L} = 3000 \text{ fb}^{-1}$ with the ATLAS 1DV ID search. Reach for mixings in the electron (left) and muon (right) sector are shown.

The CMS search in ref. [65] looked for two DVs in the CMS inner tracker in addition to jets in the final state. The original search targeted at supersymmetric models, but the collaboration provided a reinterpretation method for extending the results to other models with pair-produced long-lived particles. Here we validate this method,³ which is based on generator-level selections that approximately replicate the vertex-reconstruction efficiency.

³Another reinterpretation of this procedure in the context of Twin Higgs Doublet models was done recently in ref. [69].




Figure 7. Number of signal events for the $U(1)_X$ model at $\sqrt{s} = 13$ TeV expected in $\mathcal{L} = 3000 \text{ fb}^{-1}$ with the ATLAS 1DV ID search. Reach for mixings in the electron (left) and muon (right) sector are shown.

Trigger	$H_T > 1000 \text{ GeV}$
Jet selection	At least 4 jets with $p_T > 20 \text{ GeV}$ and $ \eta < 2.5$
DV region	2 DVs within $0.1 \text{ mm} < r_{DV} < 20 \text{ mm}$ and $d_{VV} > 0.4 \text{ mm}$
DV selection	Made from tracks with $ d_0 \geq 0.1 \text{ mm}$, $p_T > 20 \text{ GeV}$ and $ \eta < 2.5$. $\sum p_T \geq 350 \text{ GeV}$, correcting for b quarks.

Table 3. Cuts for the CMS 2DV + jets search following the reinterpretation procedure in [65].

We reconstruct jets with FASTJET 3.1.3 [70] using the anti- k_t clustering algorithm with jet radius parameter $R = 0.4$. At least four jets are required with $p_T > 20 \text{ GeV}$ and $|\eta| < 2.5$. The variable H_T , which is the scalar sum of the p_T of all generated jets with $p_T > 40 \text{ GeV}$, is computed and events must have $H_T > 1000 \text{ GeV}$.

Two DVs are required, both within a transverse distance between 0.1 and 20 mm (near the CMS beampipe). The two DVs must also be separated from each other in the transverse plane, with distance $d_{VV} > 0.4 \text{ mm}$. All daughter particles coming from the DVs (namely, u, d, s, c, b quarks and electrons, muons and tau leptons) must satisfy $p_T > 20 \text{ GeV}$ and $|\eta| < 2.5$, and have impact parameter $|d_0| \geq 0.1 \text{ mm}$. In addition, the sum of the p_T of the daughter particles has to be at least 350 GeV. When calculating this value, we multiply the p_T of b quarks by 0.65 to account for lower reconstruction efficiency due to the long lifetime of B hadrons, as instructed in ref. [65]. All selections are summarized in table 3.

As no prior validation has been done before, we validate this search on a minimal flavour violating model of R -parity violating supersymmetry (MFV RPV SUSY), same as the CMS benchmark signal model. We generate events in PYTHIA8 [62] for pair production of long-lived neutralinos $\tilde{\chi}_1^0$ by quark-antiquark annihilation. The neutralino then decays into a top anti-quark and a virtual top squark, and the virtual top squark decays into strange and bottom anti-quarks. The model spectrum is generated with SOFTSUSY 3.6.1 [71], and is read as input to PYTHIA8 using the SLHA structure [72].

Figure 8 shows our recast 95% CL limit taken with zero background and 3 signal events, against the CMS exclusion for three different benchmarks.




Figure 8. Validation of the 95% CL observed CMS upper limits [65] for three mass points in the MFV RPV SUSY model: $m_{\tilde{\chi}_1^0} = 800, 1600, 2400$ GeV.

The event-level efficiency is shown in figure 9 for the $U(1)_X$ model. We note the same pattern as with the ATLAS 1DV ID search, although the sensitivity here is affected by the mass ratio between m_N and $m_{Z'}$, as the angular opening of the decay products of the long-lived neutrino is proportional to its boost. This means that for neutrino masses roughly around 100 GeV and below, it is harder for the decay products to be effectively resolved into our reconstructed jets, failing the selection criteria. The additional condition that both neutrinos must decay well separated near the CMS beampipe imposes another strict requirement on our displaced events.

When estimating the number of signal events, we note nearly no sensitivity to the $B - L$ scenario, as shown in figure 10. The situation improves in the $U(1)_X$ model shown in figure 11, although the parameter space of the models that can be accessed can also be covered by the ATLAS 1DV search, except for a small region at small mixings and TeV masses. We can reach mixings down to $\approx 10^{-17}$ for $m_N = 1200$ GeV in the electron case. The mass reach in m_N is limited by its proper lifetime. As for mixings above $\approx 10^{-15}$ and masses higher than ≈ 1 TeV, the heavy neutrino is already decaying too promptly.

4 Conclusions

We have studied discovery prospects at the LHC of a new right-handed neutrino, which can be pair produced via a Z' gauge boson. The neutrino has a macroscopic lifetime to be able to decay inside the inner trackers of the LHC detectors to leptons and jets, enabling the reconstruction of displaced vertices from its charged tracks.




Figure 9. Representative event level efficiency of the CMS 2DV + jets search as a function of m_N and $|V_{\mu N}|^2$.




Figure 10. Number of signal events for the $U(1)_{B-L}$ model at $\sqrt{s} = 13$ TeV expected for $\mathcal{L} = 3000 \text{ fb}^{-1}$ with the CMS 2DV+jets strategy. Reach for mixings in the electron (left) and muon (right) sector are shown.

We focus on two models, namely the minimal $U(1)_{B-L}$ model and its $U(1)_X$ extension, allowing a more general quantum charge assignments for an increased Z' production cross section. We examine discovery prospects on a benchmark scenario near the limit of current lepton resonance searches at the LHC, with the explicit parameters $(m_{Z'}, g'_1) = (6 \text{ TeV}, 0.8)$. We study production and decay of the heavy displaced neutrino to leptons and jets, and reinterpret existing searches for displaced vertices. Our focus is on the current inner tracker DV searches: an ATLAS search for at least one DV (ATLAS 1DV ID) and a CMS search for exactly two DVs in multi-jet events (CMS 2DV+jets). We find that the ATLAS 1DV ID is the most sensitive across a wide range in the light-heavy neutrino mixing and mass




Figure 11. Number of signal events for the $U(1)_X$ model at $\sqrt{s} = 13$ TeV expected for $\mathcal{L} = 3000 \text{ fb}^{-1}$ with the CMS 2DV+jets strategy. Reach for mixings in the electron (left) and muon (right) sector are shown.

space. Mixings as low as $\approx 10^{-17}$ can be accessible for a heavy neutrino of mass around hundreds of GeV.

By considering displaced vertices reconstructed inside the muon spectrometer (MS) [66], one should be able to further constrain the parameter space of these heavy neutrino models. The scenario with heavy neutrino masses lower than the ones considered in this work was studied first in ref. [35] and later in ref. [45], the latter focusing on neutrinos which were pair produced from decays of the Higgs boson. The authors show that mixings with muons as small as $\approx 10^{-14}$ can be probed at the LHC for heavy neutrino masses below 60 GeV. In our case, the MS searches would be sensitive to even smaller values of the mixing, below $\approx 10^{-20}$ (see figure 4). In view of the challenges in simulating vertex reconstruction in the muon spectrometer, estimation of displaced efficiencies and proper treatment of backgrounds, we leave this study for a future work.

Acknowledgments

G.C. would like to thank the Korea Institute of Advance Studies (KIAS) for hospitality offered (at the time of ICHEP 2018) when this work was initiated, Brian Shuve for useful comments and Gabriel Torrealba for technical advice with python. The research of C.-W.C. was supported in part by the Ministry of Science and Technology (MOST) of Taiwan under Grant No. MOST-104-2628-M-002-014-MY4 and MOST-108-2112-M-002-005-MY3. G.C. acknowledges support by Grant No. MOST-107-2811-M-002-3120 and CONICYT-Chile FONDECYT Grant No. 3190051. The work of A.D. is supported by the Japan Society for the Promotion of Science (JSPS) Post- doctoral Fellowship for Research in Japan.

A Useful formulas

We write the partial decay widths of the Z' as a function of x_H and g'_1 taking $x_\Phi = 1$. The partial decay width of the Z' into a pair of light neutrinos:

$$\Gamma(Z' \rightarrow 2\nu) = \frac{m_{Z'}}{24\pi} g_L^\nu [g'_1, x_H]^2, \quad (\text{A.1})$$

with $g_L^\nu [g'_1, x_H] \equiv \left(-\frac{1}{2}x_H + (-1)\right)g'_1$. The partial decay width of the Z' into a pair of charged leptons:

$$\Gamma(Z' \rightarrow 2\ell) = \frac{m_{Z'}}{24\pi} \left(g_L^e [g'_1, x_H]^2 + g_R^e [g'_1, x_H]^2 \right), \quad (\text{A.2})$$

with $g_L^e = \left(-\frac{1}{2}x_H + (-1)\right)g'_1$ and $g_R^e = \left(-1x_H + (-1)\right)g'_1$. The partial decay width of the Z' into a pair of up-type quarks:

$$\Gamma(Z' \rightarrow 2u) = \frac{m_{Z'}}{24\pi} \left(g_L^u [g'_1, x_H]^2 + g_R^u [g'_1, x_H]^2 \right), \quad (\text{A.3})$$

with $g_L^u = \left(\frac{1}{6}x_H + \frac{1}{3}\right)g'_1$ and $g_R^u = \left(\frac{2}{3}x_H + \frac{1}{3}\right)g'_1$. The partial decay width of the Z' into a pair of down-type quarks:

$$\Gamma(Z' \rightarrow 2d) = \frac{m_{Z'}}{24\pi} \left(g_L^d [g'_1, x_H]^2 + g_R^d [g'_1, x_H]^2 \right), \quad (\text{A.4})$$

with $g_L^d = \left(\frac{1}{6}x_H + \frac{1}{3}\right)g'_1$ and $g_R^d = \left(-\frac{1}{3}x_H + \frac{1}{3}\right)g'_1$. In these partial decay widths, terms involving the fermion masses are negligible because they are suppressed by $m_{Z'}$, which we take to be 6 TeV in this analysis. The partial decay width of the Z' into a pair of Majorana heavy neutrinos for $m_{Z'} > 2m_N$ (at least) is

$$\Gamma(Z' \rightarrow 2N) = \frac{m_{Z'}}{24\pi} g_R^N [g'_1, x_H]^2 \left(1 - 4 \frac{m_N^2}{m_{Z'}^2} \right)^{\frac{3}{2}}, \quad (\text{A.5})$$

with $g_R^N [g'_1, x_H] = \left(0x_H + (-1)\right)g'_1$. The partial decay widths of the heavy neutrino can be found in ref. [17] for different masses, lighter and heavier than the SM gauge bosons.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] P. Minkowski, $\mu \rightarrow e\gamma$ at a Rate of One Out of 10^9 Muon Decays?, *Phys. Lett.* **67B** (1977) 421 [[INSPIRE](#)].
- [2] R.N. Mohapatra and G. Senjanović, Neutrino Mass and Spontaneous Parity Nonconservation, *Phys. Rev. Lett.* **44** (1980) 912 [[INSPIRE](#)].

- [3] J. Schechter and J.W.F. Valle, *Neutrino Masses in $SU(2) \times U(1)$ Theories*, *Phys. Rev. D* **22** (1980) 2227 [INSPIRE].
- [4] J. Alimena et al., *Searching for Long-Lived Particles beyond the Standard Model at the Large Hadron Collider*, [arXiv:1903.04497](#) [INSPIRE].
- [5] D. Curtin et al., *Long-Lived Particles at the Energy Frontier: The MATHUSLA Physics Case*, *Rept. Prog. Phys.* **82** (2019) 116201 [[arXiv:1806.07396](#)] [INSPIRE].
- [6] FASER collaboration, *FASER's physics reach for long-lived particles*, *Phys. Rev. D* **99** (2019) 095011 [[arXiv:1811.12522](#)] [INSPIRE].
- [7] S. Alekhin et al., *A facility to Search for Hidden Particles at the CERN SPS: the SHiP physics case*, *Rept. Prog. Phys.* **79** (2016) 124201 [[arXiv:1504.04855](#)] [INSPIRE].
- [8] L. Basso, A. Belyaev, S. Moretti and C.H. Shepherd-Themistocleous, *Phenomenology of the minimal B-L extension of the Standard model: Z' and neutrinos*, *Phys. Rev. D* **80** (2009) 055030 [[arXiv:0812.4313](#)] [INSPIRE].
- [9] CMS collaboration, *Search for high-mass resonances in dilepton final states in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *JHEP* **06** (2018) 120 [[arXiv:1803.06292](#)] [INSPIRE].
- [10] ATLAS collaboration, *Search for high-mass dilepton resonances using 139 fb^{-1} of pp collision data collected at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Phys. Lett. B* **796** (2019) 68 [[arXiv:1903.06248](#)] [INSPIRE].
- [11] E. Salvioni, G. Villadoro and F. Zwirner, *Minimal Z' models: Present bounds and early LHC reach*, *JHEP* **11** (2009) 068 [[arXiv:0909.1320](#)] [INSPIRE].
- [12] L. Basso, A. Belyaev, S. Moretti and G.M. Pruna, *Probing the Z-prime sector of the minimal B-L model at future Linear Colliders in the $e^+e^- \rightarrow \mu^+\mu^-$ process*, *JHEP* **10** (2009) 006 [[arXiv:0903.4777](#)] [INSPIRE].
- [13] L. Basso, A. Belyaev, S. Moretti, G.M. Pruna and C.H. Shepherd-Themistocleous, *Z' discovery potential at the LHC in the minimal B – L extension of the Standard Model*, *Eur. Phys. J. C* **71** (2011) 1613 [[arXiv:1002.3586](#)] [INSPIRE].
- [14] F.F. Deppisch, N. Desai and J.W.F. Valle, *Is charged lepton flavor violation a high energy phenomenon?*, *Phys. Rev. D* **89** (2014) 051302 [[arXiv:1308.6789](#)] [INSPIRE].
- [15] A. Das, N. Okada and D. Raut, *Heavy Majorana neutrino pair productions at the LHC in minimal U(1) extended Standard Model*, *Eur. Phys. J. C* **78** (2018) 696 [[arXiv:1711.09896](#)] [INSPIRE].
- [16] S. Jana, N. Okada and D. Raut, *Displaced vertex signature of type-I seesaw model*, *Phys. Rev. D* **98** (2018) 035023 [[arXiv:1804.06828](#)] [INSPIRE].
- [17] A. Das, N. Okada, S. Okada and D. Raut, *Probing the seesaw mechanism at the 250 GeV ILC*, *Phys. Lett. B* **797** (2019) 134849 [[arXiv:1812.11931](#)] [INSPIRE].
- [18] ATLAS collaboration, *Search for heavy neutral leptons in decays of W bosons produced in 13 TeV pp collisions using prompt and displaced signatures with the ATLAS detector*, *JHEP* **10** (2019) 265 [[arXiv:1905.09787](#)] [INSPIRE].
- [19] M. Drewes and J. Hajer, *Heavy Neutrinos in displaced vertex searches at the LHC and HL-LHC*, [arXiv:1903.06100](#) [INSPIRE].
- [20] I. Boiarska et al., *Probing baryon asymmetry of the Universe at LHC and SHiP*, [arXiv:1902.04535](#) [INSPIRE].

- [21] G. Cottin, J.C. Helo, M. Hirsch and D. Silva, *Revisiting the LHC reach in the displaced region of the minimal left-right symmetric model*, *Phys. Rev. D* **99** (2019) 115013 [[arXiv:1902.05673](#)] [[INSPIRE](#)].
- [22] SHiP collaboration, *Sensitivity of the SHiP experiment to Heavy Neutral Leptons*, *JHEP* **04** (2019) 077 [[arXiv:1811.00930](#)] [[INSPIRE](#)].
- [23] A. Abada, N. Bernal, M. Losada and X. Marcano, *Inclusive Displaced Vertex Searches for Heavy Neutral Leptons at the LHC*, *JHEP* **01** (2019) 093 [[arXiv:1807.10024](#)] [[INSPIRE](#)].
- [24] D. Dercks, H.K. Dreiner, M. Hirsch and Z.S. Wang, *Long-Lived Fermions at AL3X*, *Phys. Rev. D* **99** (2019) 055020 [[arXiv:1811.01995](#)] [[INSPIRE](#)].
- [25] J.C. Helo, M. Hirsch and Z.S. Wang, *Heavy neutral fermions at the high-luminosity LHC*, *JHEP* **07** (2018) 056 [[arXiv:1803.02212](#)] [[INSPIRE](#)].
- [26] M. Nemešek, F. Nesti and G. Popara, *Keung-Senjanović process at the LHC: From lepton number violation to displaced vertices to invisible decays*, *Phys. Rev. D* **97** (2018) 115018 [[arXiv:1801.05813](#)] [[INSPIRE](#)].
- [27] G. Cottin, J.C. Helo and M. Hirsch, *Displaced vertices as probes of sterile neutrino mixing at the LHC*, *Phys. Rev. D* **98** (2018) 035012 [[arXiv:1806.05191](#)] [[INSPIRE](#)].
- [28] G. Cottin, J.C. Helo and M. Hirsch, *Searches for light sterile neutrinos with multitrack displaced vertices*, *Phys. Rev. D* **97** (2018) 055025 [[arXiv:1801.02734](#)] [[INSPIRE](#)].
- [29] S. Dube, D. Gadkari and A.M. Thalapillil, *Lepton-Jets and Low-Mass Sterile Neutrinos at Hadron Colliders*, *Phys. Rev. D* **96** (2017) 055031 [[arXiv:1707.00008](#)] [[INSPIRE](#)].
- [30] P.S.B. Dev, R.N. Mohapatra and Y. Zhang, *Long Lived Light Scalars as Probe of Low Scale Seesaw Models*, *Nucl. Phys. B* **923** (2017) 179 [[arXiv:1703.02471](#)] [[INSPIRE](#)].
- [31] E. Accomando, L. Delle Rose, S. Moretti, E. Olaiya and C.H. Shepherd-Themistocleous, *Extra Higgs boson and Z' as portals to signatures of heavy neutrinos at the LHC*, *JHEP* **02** (2018) 109 [[arXiv:1708.03650](#)] [[INSPIRE](#)].
- [32] A. Caputo, P. Hernández, J. Lopez-Pavon and J. Salvado, *The seesaw portal in testable models of neutrino masses*, *JHEP* **06** (2017) 112 [[arXiv:1704.08721](#)] [[INSPIRE](#)].
- [33] A. Caputo, P. Hernández, M. Kekic, J. López-Pavón and J. Salvado, *The seesaw path to leptonic CP-violation*, *Eur. Phys. J. C* **77** (2017) 258 [[arXiv:1611.05000](#)] [[INSPIRE](#)].
- [34] M. Mitra, R. Ruiz, D.J. Scott and M. Spannowsky, *Neutrino Jets from High-Mass W_R Gauge Bosons in TeV-Scale Left-Right Symmetric Models*, *Phys. Rev. D* **94** (2016) 095016 [[arXiv:1607.03504](#)] [[INSPIRE](#)].
- [35] B. Batell, M. Pospelov and B. Shuve, *Shedding Light on Neutrino Masses with Dark Forces*, *JHEP* **08** (2016) 052 [[arXiv:1604.06099](#)] [[INSPIRE](#)].
- [36] B. Shuve and M.E. Peskin, *Revision of the LHCb Limit on Majorana Neutrinos*, *Phys. Rev. D* **94** (2016) 113007 [[arXiv:1607.04258](#)] [[INSPIRE](#)].
- [37] S. Antusch, E. Cazzato and O. Fischer, *Displaced vertex searches for sterile neutrinos at future lepton colliders*, *JHEP* **12** (2016) 007 [[arXiv:1604.02420](#)] [[INSPIRE](#)].
- [38] S. Antusch, E. Cazzato and O. Fischer, *Sterile neutrino searches at future e^-e^+ , pp and e^-p colliders*, *Int. J. Mod. Phys. A* **32** (2017) 1750078 [[arXiv:1612.02728](#)] [[INSPIRE](#)].
- [39] E. Izaguirre and B. Shuve, *Multilepton and Lepton Jet Probes of Sub-Weak-Scale Right-Handed Neutrinos*, *Phys. Rev. D* **91** (2015) 093010 [[arXiv:1504.02470](#)] [[INSPIRE](#)].

- [40] A.M. Gago, P. Hernández, J. Jones-Pérez, M. Losada and A. Moreno Briceño, *Probing the Type I Seesaw Mechanism with Displaced Vertices at the LHC*, *Eur. Phys. J. C* **75** (2015) 470 [[arXiv:1505.05880](#)] [[INSPIRE](#)].
- [41] S. Eijima, M. Shaposhnikov and I. Timiryasov, *Parameter space of baryogenesis in the ν MSM*, *JHEP* **07** (2019) 077 [[arXiv:1808.10833](#)] [[INSPIRE](#)].
- [42] A. Ibarra, E. Molinaro and S. Vogl, *Potential for probing three-body decays of Long-Lived Particles with MATHUSLA*, *Phys. Lett. B* **789** (2019) 127 [[arXiv:1809.01683](#)] [[INSPIRE](#)].
- [43] S. Amrith, J.M. Butterworth, F.F. Deppisch, W. Liu, A. Varma and D. Yallup, *LHC constraints on a $B - L$ gauge model using Contur*, *JHEP* **05** (2019) 154 [[arXiv:1811.11452](#)] [[INSPIRE](#)].
- [44] E. Accomando, L. Delle Rose, S. Moretti, E. Olaiya and C.H. Shepherd-Themistocleous, *Novel SM-like Higgs decay into displaced heavy neutrino pairs in $U(1)'$ models*, *JHEP* **04** (2017) 081 [[arXiv:1612.05977](#)] [[INSPIRE](#)].
- [45] F.F. Deppisch, W. Liu and M. Mitra, *Long-lived Heavy Neutrinos from Higgs Decays*, *JHEP* **08** (2018) 181 [[arXiv:1804.04075](#)] [[INSPIRE](#)].
- [46] F. Deppisch, S. Kulkarni and W. Liu, *Heavy neutrino production via Z' at the lifetime frontier*, *Phys. Rev. D* **100** (2019) 035005 [[arXiv:1905.11889](#)] [[INSPIRE](#)].
- [47] F. Kling and S. Trojanowski, *Heavy Neutral Leptons at FASER*, *Phys. Rev. D* **97** (2018) 095016 [[arXiv:1801.08947](#)] [[INSPIRE](#)].
- [48] T. Appelquist, B.A. Dobrescu and A.R. Hopper, *Nonexotic Neutral Gauge Bosons*, *Phys. Rev. D* **68** (2003) 035012 [[hep-ph/0212073](#)] [[INSPIRE](#)].
- [49] A. Das, N. Okada and D. Raut, *Enhanced pair production of heavy Majorana neutrinos at the LHC*, *Phys. Rev. D* **97** (2018) 115023 [[arXiv:1710.03377](#)] [[INSPIRE](#)].
- [50] A. Das, P.S.B. Dev and N. Okada, *Long-lived TeV-scale right-handed neutrino production at the LHC in gauged $U(1)_X$ model*, *Phys. Lett. B* **799** (2019) 135052 [[arXiv:1906.04132](#)] [[INSPIRE](#)].
- [51] R.N. Mohapatra and R.E. Marshak, *Local $B - L$ Symmetry of Electroweak Interactions, Majorana Neutrinos and Neutron Oscillations*, *Phys. Rev. Lett.* **44** (1980) 1316 [Erratum *ibid.* **44** (1980) 1643] [[INSPIRE](#)].
- [52] A. Das, S. Oda, N. Okada and D.-s. Takahashi, *Classically conformal $U(1)'$ extended standard model, electroweak vacuum stability and LHC Run-2 bounds*, *Phys. Rev. D* **93** (2016) 115038 [[arXiv:1605.01157](#)] [[INSPIRE](#)].
- [53] S. Oda, N. Okada and D.-s. Takahashi, *Classically conformal $U(1)'$ extended standard model and Higgs vacuum stability*, *Phys. Rev. D* **92** (2015) 015026 [[arXiv:1504.06291](#)] [[INSPIRE](#)].
- [54] A. Davidson, *$B - L$ as the Fourth Color, Quark-Lepton Correspondence and Natural Masslessness of Neutrinos Within a Generalized W_s Model*, *Phys. Rev. D* **20** (1979) 776 [[INSPIRE](#)].
- [55] M. Carena, A. Daleo, B.A. Dobrescu and T.M.P. Tait, *Z' gauge bosons at the Tevatron*, *Phys. Rev. D* **70** (2004) 093009 [[hep-ph/0408098](#)] [[INSPIRE](#)].
- [56] J. Heeck, *Unbroken $B - L$ symmetry*, *Phys. Lett. B* **739** (2014) 256 [[arXiv:1408.6845](#)] [[INSPIRE](#)].

- [57] C. Degrande, C. Duhr, B. Fuks, D. Grellscheid, O. Mattelaer and T. Reiter, *UFO — The Universal FeynRules Output*, *Comput. Phys. Commun.* **183** (2012) 1201 [[arXiv:1108.2040](#)] [[INSPIRE](#)].
- [58] J. Alwall et al., *The automated computation of tree-level and next-to-leading order differential cross sections and their matching to parton shower simulations*, *JHEP* **07** (2014) 079 [[arXiv:1405.0301](#)] [[INSPIRE](#)].
- [59] F.F. Deppisch, P.S. Bhupal Dev and A. Pilaftsis, *Neutrinos and Collider Physics*, *New J. Phys.* **17** (2015) 075019 [[arXiv:1502.06541](#)] [[INSPIRE](#)].
- [60] CMS collaboration, *Search for heavy neutral leptons in events with three charged leptons in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *Phys. Rev. Lett.* **120** (2018) 221801 [[arXiv:1802.02965](#)] [[INSPIRE](#)].
- [61] P. Artoisenet, R. Frederix, O. Mattelaer and R. Rietkerk, *Automatic spin-entangled decays of heavy resonances in Monte Carlo simulations*, *JHEP* **03** (2013) 015 [[arXiv:1212.3460](#)] [[INSPIRE](#)].
- [62] T. Sjöstrand et al., *An Introduction to PYTHIA 8.2*, *Comput. Phys. Commun.* **191** (2015) 159 [[arXiv:1410.3012](#)] [[INSPIRE](#)].
- [63] J.D. Hunter, *Matplotlib: A 2D Graphics Environment*, *Comput. Sci. Eng.* **9** (2007) 90 [[INSPIRE](#)].
- [64] ATLAS collaboration, *Search for massive, long-lived particles using multitrack displaced vertices or displaced lepton pairs in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector*, *Phys. Rev. D* **92** (2015) 072004 [[arXiv:1504.05162](#)] [[INSPIRE](#)].
- [65] CMS collaboration, *Search for long-lived particles with displaced vertices in multijet events in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *Phys. Rev. D* **98** (2018) 092011 [[arXiv:1808.03078](#)] [[INSPIRE](#)].
- [66] ATLAS collaboration, *Search for long-lived, massive particles in events with displaced vertices and missing transverse momentum in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector*, *Phys. Rev. D* **97** (2018) 052012 [[arXiv:1710.04901](#)] [[INSPIRE](#)].
- [67] G. Brooijmans et al., *Les Houches 2017: Physics at TeV Colliders New Physics Working Group Report*, [fermilab-conf-17-664](#) [[arXiv:1803.10379](#)].
- [68] G. Cottin, *Reconstructing particle masses in events with displaced vertices*, *JHEP* **03** (2018) 137 [[arXiv:1801.09671](#)] [[INSPIRE](#)].
- [69] S. Alipour-Fard, N. Craig, S. Gori, S. Koren and D. Redigolo, *The second Higgs at the lifetime frontier*, [arXiv:1812.09315](#) [[INSPIRE](#)].
- [70] M. Cacciari, G.P. Salam and G. Soyez, *FastJet User Manual*, *Eur. Phys. J. C* **72** (2012) 1896 [[arXiv:1111.6097](#)] [[INSPIRE](#)].
- [71] B.C. Allanach, *SOFTSUSY: a program for calculating supersymmetric spectra*, *Comput. Phys. Commun.* **143** (2002) 305 [[hep-ph/0104145](#)] [[INSPIRE](#)].
- [72] B.C. Allanach et al., *SUSY Les Houches Accord 2*, *Comput. Phys. Commun.* **180** (2009) 8 [[arXiv:0801.0045](#)] [[INSPIRE](#)].