Towards the classification of rank- r $\mathcal{N} = 2$ SCFTs. Part I. Twisted partition function and central charge formulae

Mario Martone

*Physics Department, University of Texas at Austin,
Austin TX 78712, U.S.A.*

E-mail: mariomartone@utexas.edu

ABSTRACT: We derive explicit formulae to compute the a and c central charges of four dimensional $\mathcal{N} = 2$ superconformal field theories (SCFTs) directly from Coulomb branch related quantities. The formulae apply at arbitrary rank. We also discover general properties of the low-energy limit behavior of the flavor symmetry of $\mathcal{N} = 2$ SCFTs which culminate with our $\mathcal{N} = 2$ *UV-IR simple flavor condition*. This is done by determining precisely the relation between the integrand of the partition function of the topologically twisted version of the 4d $\mathcal{N} = 2$ SCFTs and the singular locus of their Coulomb branches. The techniques developed here are extensively applied to many rank-2 SCFTs, including new ones, in a companion paper.

This manuscript is dedicated to the memory of Rayshard Brooks, George Floyd, Breonna Taylor and the countless black lives taken by US police forces and still awaiting justice. Our hearts are with our colleagues of color who suffer daily the consequences of this racist world.

KEYWORDS: Conformal Field Theory, Extended Supersymmetry, Supersymmetric Effective Theories

ARXIV EPRINT: [2006.16255](https://arxiv.org/abs/2006.16255)

Contents

1	Introduction	1
2	Discriminant locus, quantum discriminant, and physical discriminant	3
2.1	Structure of the singularity	3
2.2	Discriminant locus	4
2.3	Low-energy effective theory on the singularity	7
2.4	Quantum discriminant	8
2.5	Physical discriminant	12
3	Twisted partition function and central charge	12
3.1	U(1) anomaly and central charges	14
3.2	Final form of the partition function and central charge formulae	16
4	Flavor structure of general $\mathcal{N} = 2$ SCFTs	19
5	Conclusions	22
A	Coulomb branch generalities	23
A.1	Special Kahler stratification	24
A.2	Accidental U(1) _R near strata	25

1 Introduction

Since Witten’s seminal paper [1], the connection between topological and supersymmetric field theories has been fruitful. Beyond the deep insights that it brought into the study of four manifolds [2–4], see [5–7] and references therein for recent developments, the fact that supersymmetric theories contain a sub-sector which is topological has brought considerable progress to our understanding of supersymmetric field theories themselves, see e.g. [8–11]. The procedure of projecting onto the topological sector of a supersymmetric field theory is now standard and known as topological twist. The work presented here represents yet another instance in which the constrained picture of topological field theories is used to advance our understanding of the properties of the their non-topological cousins.

Our interest in studying topologically twisted $\mathcal{N} = 2$ theories, and in particular their partition functions, is motivated by our ambitious classification program of $\mathcal{N} = 2$ superconformal field theories (SCFTs) in four dimensions. This program, see [12] for a lightning review and a list of relevant references, is based on a systematic study of Coulomb branches

whose geometry appears to contain enough structure to nearly determine the space of allowed $\mathcal{N} = 2$ SCFTs.¹ The results of this paper give further evidence for this fact as we are able to derive general formulae which allow to straightforwardly compute, directly from Coulomb branch related quantities, the (a, c) central charges, as well as the flavor levels $k_{\mathfrak{f}}$, of any $\mathcal{N} = 2$ SCFT at arbitrary rank.

It is useful to summarize the main results of this paper right away:

- i.* We derive explicit formulae which express in terms of Coulomb branch quantities the (a, c) central charges and the level of any simple (i.e. non-abelian) flavor factor $\mathfrak{f}$, of an arbitrary $\mathcal{N} = 2$ SCFT:²

$$24a = 5r + h + 6 \left(\sum_{\ell=1}^r \Delta_{\mathbf{u}_\ell} - r \right) + \sum_{i \in I} \Delta_i^{\text{sing}} \frac{12c_i - 2 - h_i}{\Delta_i}, \quad (1.1a)$$

$$12c = 2r + h + \sum_{i \in I} \Delta_i^{\text{sing}} \frac{12c_i - 2 - h_i}{\Delta_i}, \quad (1.1b)$$

$$k_{\mathfrak{f}} = \sum_{i \in I_{\mathfrak{f}}} \frac{\Delta_i^{\text{sing}}}{d_i \Delta_i} (k^i - T(\mathbf{2}h_i)) + T(\mathbf{2}h). \quad (1.1c)$$

Here r is the rank of the SCFT, h is the quaternionic dimension of the theory's extended Coulomb branch, $\Delta_{\mathbf{u}_\ell}$ is the scaling dimension of the theory's ℓ -th Coulomb branch parameters, Δ_i^{sing} is defined in (2.3) and all the remaining quantities indexed by i refer to known quantities of rank-1 theories which are supported on special loci of the Coulomb branch (see section 2.3 below). We call these formulae *central charge formulae*.

The central charge formulae, which systematize and extend the beautiful work by Shapere and Tachikawa [10], are derived in section 3 and 4 using the well-known relation between the $(a, c, k_{\mathfrak{f}})$ central charges of an untwisted $\mathcal{N} = 2$ SCFT and the $U(1)_R$ symmetry anomaly of its twisted topological version.

- ii.* We formulate and prove the $\mathcal{N} = 2$ *UV-IR simple flavor condition*. This condition states that all mass deformations of a rank ≥ 2 $\mathcal{N} = 2$ SCFT modify the asymptotics of the CB and are realized as mass deformations of the same rank-1 theories which appear in their central charge formulae. This realization de-facto reduces the problem of analyzing mass deformations of higher rank theories to that of rank-1,³ providing a clear route towards a systematic understanding of the space of higher rank theories. The UV-IR simple flavor condition is discussed in section 4 and it is a straightforward consequence of the structure of the integrand of the partition function of the twisted version of the $\mathcal{N} = 2$ SCFT.

¹The qualification “nearly” refers to possible ambiguities which arise if we also allowed for discrete gauging [13].

²Our methods are extremely powerful when applied to simple factors of the flavor symmetry but are nearly insensitive to abelian ones (see section 4). Throughout the manuscript we will therefore neglect U_1 factors.

³Note that this result only applies to mass deformations, and not to the other class of relevant $\mathcal{N} = 2$ deformations known as chiral deformations.

- iii. We determine the relation between the integrand of the partition function of a topologically twisted $\mathcal{N} = 2$ SCFT and the structure of the singular locus of its Coulomb branch at arbitrary rank. This precise relation is used copiously to derive the results in *i.* and *ii.* and lead to a sharp definition and understanding of both the *quantum* and *physical* discriminant, see section 2.4 and 2.5.

It is, of course, not the first time that the partition function of topologically twisted $\mathcal{N} = 2$ theories with rank higher than one is studied in detail. The fact that the zeros and the singularities of the measure of the partition function could only lie along the singular locus, which is used here as a starting point, was noticed since the early days [14–19]. Here we point out that the orders of such zeros is subtle to determine but we are able to carry out this calculation in general. This is not just a mathematical curiosity. In fact our formulae above, precisely rely on a detailed determination of these exponents.

The paper is organized as follows. In the next section we will give a reminder of the geometric structure of Coulomb branches of $\mathcal{N} = 2$ theories at arbitrary rank and discuss the singular locus and the concept of the discriminant in detail. In section 3 we will set up the study of the partition function of a topologically twisted $\mathcal{N} = 2$ SCFT and derive our central charge formulae. Section 4 will instead be dedicated to a systematic study of the flavor structure of $\mathcal{N} = 2$ SCFTs and making many general remarks connecting the behavior of the flavor symmetry at high and low energies. We end the paper with our conclusions.

2 Discriminant locus, quantum discriminant, and physical discriminant

The central charge formulae (1.1a)–(1.1c) depend on a series quantities indexed by $i \in I$ which are associated to rank-1 theories appearing on special loci of the Coulomb branch (CB). We define these quantities here and explain our definitions in explicit examples. In doing so we clarify what we mean by discriminant of the CB a term which is often used in the literature in different contexts and with different meanings. Specifically we will give our definitions for three different but related objects: the *discriminant locus*, the *quantum discriminant*, or loosely the discriminant of the Seiberg-Witten (SW) curve, and the *physical discriminant* which instead enters the measure factor of the partition function of the twisted $\mathcal{N} = 2$ theory. These definitions will be needed in the next section. To understand this discussion, it might be useful to consult appendix A where the basics of CB geometry and elements of the *special Kahler stratification* are given.

2.1 Structure of the singularity

The low-energy theory on a generic point of the CB $\mathcal{C}$ is almost as boring as it gets; a free $\mathcal{N} = 2$ supersymmetric $U(1)^r$ gauge theory with no massless charged states. r is called the *rank* of the theory and coincides with the complex dimensionality of $\mathcal{C}$, $\dim_{\mathbb{C}} \mathcal{C} = r$. $\mathcal{C}$ is a singular space and its singular locus will be at the center of our discussion and denoted as $\overline{\mathcal{S}}$. $\overline{\mathcal{S}}$ stands both for “singular” and “stratum”, the reason for the bar is that $\overline{\mathcal{S}}$ is a closed subset of $\mathcal{C}$ and the smooth part of the CB is $\mathcal{C}_{\text{reg}} := \mathcal{C} \setminus \overline{\mathcal{S}}$. Thus $\mathcal{C}_{\text{reg}}$ is an open subset of $\mathcal{C}$.

Henceforth we will only discuss the special case when the $\mathcal{N} = 2$ theory is superconformal in which case the symmetry group includes an $\mathbb{R}^+ \times \text{U}(1)_R$ (we are neglecting the $\text{SU}(2)_R$ factor as it acts trivially on $\mathcal{C}$) which can be spontaneously broken, and so acts nontrivially on $\mathcal{C}$, and combines to give a $\mathbb{C}^*$ action on the CB. We will label the $\mathbb{C}^*$ weight or scaling dimension of a quantity $\cdot$ interchangeably as $[\cdot]$ or $\Delta(\cdot)$. The origin of $\mathcal{C}$ is the single vacuum of the entire moduli space which is invariant under such action and for this reason we will call it the superconformal vacuum. The entire structure of $\mathcal{C}$ has to be compatible with the $\mathbb{C}^*$ action and in particular $\bar{\mathcal{S}}$ and $\mathcal{C}_{\text{reg}}$ have to be closed under it. We will often refer to the set of constraints arising from the compatibility with the $\mathbb{C}^*$ action as the constraints coming from scale invariance. In this language we will only consider here scale invariant CB geometries.

Finally, the CB is both a complex space and a metric space, and so $\mathcal{C}$ can have singularities in each of these structures [20] and the singular locus is the union of the loci of the two types of singularities, $\bar{\mathcal{S}} = \bar{\mathcal{S}}_{\text{metr}} \cup \bar{\mathcal{S}}_{\text{cplx}}$. The physics interpretations of the two are remarkably different. $\bar{\mathcal{S}}_{\text{cplx}}$ is the locus of vacua for which the operators generating the corresponding CB chiral ring satisfy non-trivial relations. This means that the chiral ring is not freely generated at points in $\bar{\mathcal{S}}_{\text{cplx}}$. In the following we will make the assumption that $\bar{\mathcal{S}}_{\text{cplx}} = \emptyset$, therefore $\bar{\mathcal{S}} \equiv \bar{\mathcal{S}}_{\text{metr}}$. This condition can be equivalently stated as assuming that $\mathcal{C}$ admits a set of globally defined r complex coordinates which have definite weight under the $\mathbb{C}^*$ action and which we will collectively indicate as $\mathbf{u}$.

2.2 Discriminant locus

$\bar{\mathcal{S}}_{\text{metr}}$ is the locus of $\mathcal{C}$ where extra charged states become massless or, in other words, where the low-energy physics is not captured solely by a bunch of free $\mathcal{N} = 2$ vector multiplets. The central charge $|Z_Q|$, see (A.5), is a lower bound on the mass of a state with charge Q , therefore $Z_Q(\mathbf{u})$ vanishes for any $\mathbf{u} \in \bar{\mathcal{S}}_{\text{metr}}$. Assuming away some pathological behavior and carefully keeping track of the structure of the CB geometry, it is possible to prove that $\bar{\mathcal{S}}$ is an $r - 1$ complex dimensional algebraic subvariety of $\mathcal{C}$, which is the union of connected, irreducible, components $\bar{\mathcal{S}}_i$:

$$\bar{\mathcal{S}} := \bigcup_{i \in I} \bar{\mathcal{S}}_i, \quad \bar{\mathcal{S}}_i := \left\{ \mathbf{u} \in \mathcal{C} \mid Z_Q(\sigma(\mathbf{u})) = 0, \quad \forall Q \in \Lambda_i \right\}. \quad (2.1)$$

Each component $\bar{\mathcal{S}}_i$ is defined by the vanishing of the central charge for charges corresponding to the set of BPS states in the theory which become massless there. Since Z_Q is linear in the charges, what defines $\bar{\mathcal{S}}_i$ is the sublattice of charges Λ_i given by the integer linear span of such charges. By charge conservation, Λ_i is constant along $\bar{\mathcal{S}}_i$ irregardless of walls of marginal stability. The index i runs over some finite set I which corresponds to the range of the sum of the equations we have reported in the introduction (1.1a)–(1.1c). We will discuss in the next section how a rank-1 theory can be associated to each component $\bar{\mathcal{S}}_i$.

Since (2.1) is a complex co-dimension one algebraic subvariety of $\mathcal{C}$ it can be cut out by a single polynomial on the CB, which is a product of polynomials whose zero locus corresponds to each component $\bar{\mathcal{S}}_i$. If this polynomial is reduced, then it is unique up

to an overall constant factor. We then define the *discriminant locus* to be the following quantity:

$$D_x := \prod_{i \in I} P_i(\mathbf{u}), \quad \overline{\mathcal{S}}_i =: \left\{ \mathbf{u} \in \mathcal{C} \mid P_i(\mathbf{u}) = 0 \right\} \quad (2.2)$$

with $\overline{\mathcal{S}}_i$ as in (2.1) and the $P_i(\mathbf{u})$ are distinct and irreducible for all $i \in I$. By scale invariance, the $P_i(\mathbf{u})$ are weighted homogeneous polynomials in the $\mathbf{u}$, and we will call Δ_i^{sing} the weight of $P_i(\mathbf{u})$ under the $\mathbb{C}^*$ action:

$$\Delta_i^{\text{sing}} := \Delta(P_i(\mathbf{u})). \quad (2.3)$$

This quantity plays an important role in our central charge formulae (1.1a)–(1.1c).

Example: SCFTs with $\mathfrak{su}(3)$ gauge algebra. It might be useful to immediately follow this definition with a concrete example. Consider an $\mathfrak{su}(3)$ gauge theory with hypermultiplets in representations for which the beta function for the gauge coupling vanishes. It is a straightforward group theory exercise to show that there are three possibilities:

$$\mathfrak{su}(3) \mathcal{N} = 2 \text{ SCFTs} : \begin{cases} 1. & 6(\mathbf{3}) \\ 2. & 1(\mathbf{6}) \oplus 1(\mathbf{3}) \\ 3. & 1(\mathbf{8}) \end{cases} \quad (2.4)$$

entry 3 corresponds to the $\mathcal{N} = 4$ theory with $\mathfrak{su}(3)$ Lie-algebra,⁴ while entry 1 and 2 correspond to genuinely $\mathcal{N} = 2$ theories. These are all rank-2 theories and we will indicate the two complex coordinates which parametrize their CBs as (u, v) .

To derive the discriminant locus we can take advantage of the fact that the theory is lagrangian and directly study the values of (u, v) for which extra massless states enter the theory. To do that it is helpful to write the CB coordinates explicitly in terms of the eigenvalues of Φ , the scalar component of the $\mathfrak{su}(3)$ adjoint $\mathcal{N} = 2$ vector multiplet. First imposing F and D term conditions, constraints Φ to be gauge equivalent to the following form:

$$\Phi = \begin{pmatrix} a_1 & 0 & 0 \\ 0 & a_2 & 0 \\ 0 & 0 & -a_1 - a_2 \end{pmatrix} \quad (2.5)$$

and for $a_1 \neq a_2$, $\mathfrak{su}(3) \rightarrow \text{U}(1)^2$. A convenient choice for u and v is, explicitly:

$$u := \text{Tr}[\Phi^2] \sim a_1^2 + a_2^2 + a_1 a_2, \quad v := \text{Tr}[\Phi^3] \sim -a_1^2 a_2 - a_1 a_2^2 \quad (2.6)$$

where we have neglected inessential numerical coefficients. From (2.6), it follows straightforwardly that the coordinates of the CB of all three theories in (2.4) have scaling dimensions $\Delta_u = 2$ and $\Delta_v = 3$.

When two of the eigenvalues in (2.5) coincide, one of the two low energy $\text{U}(1) \rightarrow \mathfrak{su}(2)$ and the $\mathcal{N} = 2$ vector multiplets of the corresponding gauge bosons become massless giving

⁴It is well known that, for a given gauge algebra, there are multiple $\mathcal{N} = 4$ theories [21]. Here we are using the determinative article because we are neglecting all the subtleties connected with the global structure of the gauge group.

rise to singularities on the CB. It is easy for the reader to check that $a_1 = a_2$ corresponds to the following hypersurface in $\mathcal{C}$:

$$u^3 + \lambda v^2 = 0, \quad (2.7)$$

where $\lambda \in \mathbb{C}^*$ depends on the specific normalization of u and v .

Due to superpotential terms, the hypermultiplets are massive on a generic point of $\mathcal{C}$ but extra components of the singular locus might arise if special values of a_1 and a_2 exist for which some of the hypermultiplets components become massless. Here the result depends on the specific representations in (2.4). Straightforward representation theory shows that components of the hypermultiplets become massless at:

$$\left. \begin{array}{l} 1. \\ 2. \end{array} \right\} \quad a_1/a_2 = 0 \quad \Rightarrow \quad v = 0 \quad (2.8)$$

$$3. \quad a_1 = a_2 \quad \Rightarrow \quad u^3 + \lambda v^2 = 0. \quad (2.9)$$

The fact that components of the hypermultiplets for theory 3 become massless at (2.7) follows from the fact that for this theory the hypermultiplets transform in the same representation as the $\mathcal{N} = 2$ vector multiplets.

To improve on this semi-classical analysis and write down the discriminant locus, we notice that for theories 1. and 2. no hypermultiplets become massless at (2.7). The low energy effective theory there is then a $\mathcal{N} = 2$ $\mathfrak{su}(2)$ Yang-Mills and the well-known results of [22] imply that (2.7) has to “split” in two components accounting for the monopole and dyon singularities.⁵ No splitting instead happens for theory 3. because the extra massless hypermultiplets component make the low-energy theory on (2.7) superconformal as we will see explicitly below.

Bringing all together we find:

$$\left. \begin{array}{l} 1. \\ 2. \end{array} \right\} \quad D_x \sim v(u^3 + \lambda_1 v^2)(u^3 + \lambda_2 v^2) \quad (2.10a)$$

$$3. \quad D_x \sim (u^3 + \lambda v^2) \quad (2.10b)$$

and therefore we have the following for theories 1. and 2.:

$$\begin{aligned} \overline{\mathcal{S}}_1 &: \left\{ (u, v) \in \mathcal{C} \mid v = 0 \right\} & \Rightarrow & \Delta_1^{\text{sing}} = 3, \\ \overline{\mathcal{S}}_2 &: \left\{ (u, v) \in \mathcal{C} \mid u^3 + \lambda_1 v^2 = 0 \right\} & \Rightarrow & \Delta_2^{\text{sing}} = 6, \\ \overline{\mathcal{S}}_3 &: \left\{ (u, v) \in \mathcal{C} \mid u^3 + \lambda_2 v^2 = 0 \right\} & \Rightarrow & \Delta_3^{\text{sing}} = 6. \end{aligned} \quad (2.11)$$

while theory 3. only has a single component analogous to either $\overline{\mathcal{S}}_2$ or $\overline{\mathcal{S}}_3$. We will come back to these examples throughout the manuscript to elucidate the definitions that we make along the way.

⁵The closing of the singular locus under the $\mathbb{C}^*$ action implies that the separation between these two components increases as we move towards the asymptotics of the CB. This is consistent with a careful RG analysis and the original result [22].

2.3 Low-energy effective theory on the singularity

Let's see how we can naturally associate a rank-1 theory to each $i \in I$ indexing the singular components in (2.1). This will provide the missing information which we need to use (1.1a)–(1.1c) to compute the central charges of any $\mathcal{N} = 2$ SCFT.

Call the rank- r theory at the superconformal vacuum $\mathcal{T}$ and call $\mathcal{T}_{\mathbf{u}}$ the low-energy effective description of $\mathcal{T}$ at the generic point $\mathbf{u}$ of the CB $\mathcal{C}$ describing the massless degrees of freedom there. For example we have:

$$\mathcal{T}_{\mathbf{u}} \equiv \text{free } \mathcal{N} = 2 \text{ U}(1)^r, \quad \mathbf{u} \in \mathcal{C}_{\text{reg}}. \quad (2.12)$$

From what we just discussed, $\mathcal{T}_{\mathbf{u}}$ will include charged states if $\mathbf{u} \in \bar{\mathcal{S}}$. Each separate component $\bar{\mathcal{S}}_i$ is identified by a lattice of charges Λ_i which identifies the charges of these states (those becoming massless there and for which $Z_Q(\mathbf{u})|_{\bar{\mathcal{S}}_i}$ vanishes). To be precise, a given sublattice is associated to an open subset $\mathcal{S}_i \subset \bar{\mathcal{S}}_i$ and we will call $\mathcal{S}_i$ the stratum associated to $\bar{\mathcal{S}}_i$ (which is indeed the closure of $\mathcal{S}_i$). It is then natural to associate to each stratum the theory describing the massless degrees of freedom on the stratum itself. We will call this theory $\mathcal{T}_i$ and since $\mathcal{S}_i$ is complex co-dimension one, it can be shown that $\mathcal{T}_i$ has to be rank-1. See appendix A for clarification on this and other subtleties, like the difference between $\mathcal{S}_i$ and $\bar{\mathcal{S}}_i$. In summary:

$$\mathcal{T}_i \equiv \mathcal{T}_{\mathbf{u}}, \quad \mathbf{u} \in \mathcal{S}_i, \quad i \in I. \quad (2.13)$$

and the quantities indexed by $i \in I$, (c_i, k_i, h_i) , which enter the central charge formulae (1.1a)–(1.1c) refer to those of $\mathcal{T}_i$. We also use u_i to label the coordinate parametrizing the one complex dimensional CB $\mathcal{C}_i$ of $\mathcal{T}_i$ and define:

$$\Delta_i := \Delta(u_i) \quad (2.14)$$

which defines the last quantity entering the central charge formulae.

Example: SCFTs with $\mathfrak{su}(3)$ gauge algebra. To consolidate (2.13), let's see explicitly how this works in the examples (2.4). We have done most of the work above already. In this rank-2 case, the strata associated to the various components are straightforwardly obtained by subtraction of the origin of $\mathcal{C}$:

$$\mathcal{S}_i := \bar{\mathcal{S}}_i \setminus \{0\}. \quad (2.15)$$

We will clarify momentarily why the rank-1 theories are properly defined on $\mathcal{S}_i$ and not $\bar{\mathcal{S}}_i$.

Let's start from $\mathcal{S}_1$ in (2.11). For $\{v = 0\} \setminus \{0\}$, see (2.8), the unbroken gauge group is $\text{U}(1)^2$ with massless hypermultiplets corresponding to the weights associated to the vanishing eigenvalue. It is possible to choose the low-energy $\text{U}(1)$ s in such a way that the component of the massless hypermultiplets are only charged under one of the $\text{U}(1)$ factors. Given the $\mathfrak{su}(3)$ representations of the hypermultiplets, it is a straightforward group theory exercise to compute both the number and the $\text{U}(1)$ charges, q , of the massless components and we find:

$$\mathcal{T}_1 : \quad \begin{cases} 1 \rightarrow \text{U}(1) \text{ w/ 6 hypers with } q = 1 \\ 2 \rightarrow \text{U}(1) \text{ w/ 2 hypers with } q = 1 \oplus 1 \text{ hyper with } q = 2 \end{cases} \Rightarrow \Delta_1 = 1, \quad (2.16)$$

since the CB scaling dimension for $\mathcal{N} = 2$ U(1) gauge theories is one. The remaining components in (2.11) are associated to the monopole/dyon singularities of the $\mathcal{N} = 2$ $\mathfrak{su}(2)$ super Yang-Mills for both theory 1. and 2.. We can separately choose an electromagnetic duality basis for $\mathcal{S}_2$ and $\mathcal{S}_3$ which makes the massless degrees of freedom there electrically charged:

$$\mathcal{T}_{2,3} : \quad \rightarrow \quad \text{U}(1) \text{ w/ 1 hypers with } q = 1 \quad \Rightarrow \quad \Delta_{2,3} = 1, \quad (2.17)$$

This concludes the analysis of theories 1. and 2..

For the $\mathcal{N} = 4$ theory we only have to determine the low energy theory on the one locus where one of the low-energy U(1) factors enhances $\text{U}(1) \rightarrow \mathfrak{su}(2)$. Since the hypermultiplet of this theory transforms in the same representation of the $\mathcal{N} = 2$ vector multiplet, we expect the massless states coming from the hyper to fill an adjoint representation of $\mathfrak{su}(2)$ giving in the low-energy limit a rank-1 $\mathcal{N} = 4$ theory:

$$\mathcal{T}_{\mathcal{N}=4} : \quad \rightarrow \quad \mathcal{N} = 4 \mathfrak{su}(2) \quad \Rightarrow \quad \Delta_{\mathcal{N}=4} = 2. \quad (2.18)$$

The last equation follows from the fact that the CB parameter of an $\mathfrak{su}(2)$ theory is identified with $u \sim \langle \text{Tr} \Phi^2 \rangle$ and therefore $\Delta = 2$.

Now that we have identified the rank-1 theories supported on the various complex co-dimension strata $\mathcal{S}_i$, we can use the information (2.16)–(2.18) and compute the quantities which enter the central charge formulae for the $\mathfrak{su}(3)$ theories in (2.4) and which will be used in the next section:

Theory	$12c_i$	Δ_i	$\mathfrak{f}_{k_i}$	h_i	$T(\mathbf{2h})$
$\mathcal{T}_1 \left\{ \begin{array}{l} 1. \\ 2. \end{array} \right.$	8	1	$\mathfrak{su}(6)_2$	0	0
	5	1	$\emptyset$	0	0
$\mathcal{T}_{2,3}$	3	1	$\emptyset$	0	0
$\mathcal{T}_{\mathcal{N}=4}$	9	2	$\mathfrak{sp}(1)_3$	1	1

(2.19)

Here $\mathfrak{sp}(1) \cong \mathfrak{su}(2)$ and we have only reported the simple flavor factors of the flavor symmetries. The formula for computing $12c$ for lagrangian theories is standard but it is also recalled below, see (3.10). Finally h_i here is the quaternionic dimension of the ECB of the theory, that is the number of free hypermultiplets at the generic point of the theory's CB.

Let's conclude with a clarification of why rank-1 theories are associated to the $\overline{\mathcal{S}}_i \setminus \{0\}$. Setting both $u = v = 0$ fully restores the $\mathfrak{su}(3)$ gauge algebra. The superconformal vacuum is then associated to a rank-2 theory (the origin is also the only complex co-dimension higher than 1 stratum in this case, see appendix A). Therefore a non-trivial rank-1 factor is only defined away from $\{0\}$, that is strictly on $\mathcal{S}_i$.

2.4 Quantum discriminant

The discriminant locus (2.2) provides a nice algebraic description of $\overline{\mathcal{S}}$ but tells no information about the low-energy theory $\mathcal{T}_i$ supported on its strata, as we saw in the various examples. If a rank r $\mathcal{N} = 2$ SCFT admits a formulation of special Kahler geometry of

its CB in terms of a family of hyperelliptic curves, the Seiberg-Witten (SW) curve Σ , and one forms, the SW one-form λ_{SW} , we can improve on the discriminant locus and define *the quantum discriminant*. Even though we will not review here this formulation of SW theory,⁶ we will still require some technical definitions to arrive at the formulation of the quantum discriminant. The content of this subsection is important to understand the subtleties of how the geometry of the CB can be leveraged to understand the singular locus but will not be needed to understand the rest of the paper.

Bring the curve Σ in the following form:

$$\Sigma : y^2 = \mathcal{P}(x, \mathbf{u}) \quad (2.20)$$

where $\mathcal{P}(x, \mathbf{u})$ is a polynomial of degree, at most, $2r + 2$ in x with coefficients which in general depend on $\mathbf{u}$. Therefore for fixed value $\mathbf{u} \in \mathcal{C}$, (2.20) defines a hyper-elliptic curve $\Sigma_{\mathbf{u}}$ which degenerates if some of the $2r + 2$ roots of $\mathcal{P}(x, \mathbf{u})$ coincide. The loci $\mathbf{u}$ for which $\Sigma_{\mathbf{u}}$ degenerates corresponds to the singular locus $\overline{\mathcal{S}}$ and is therefore reasonable to define *the quantum discriminant* as the x discriminant of the r.h.s. of (2.20):

$$D_x^\Lambda := \text{Disc}_x \mathcal{P}(x, \mathbf{u}). \quad (2.21)$$

The zeros of the quantum discriminant should coincide with the discriminant locus but eq. (2.21) will be in general no longer reduced:

$$D_x^\Lambda \sim \prod_{i \in I} [P_i(\mathbf{u})]^{\ell_i}, \quad \overline{\mathcal{S}}_i =: \left\{ \mathbf{u} \in \mathcal{C} \mid P_i(\mathbf{u}) = 0 \right\} \quad (2.22)$$

The $\ell_i \in \mathbb{Z}$ are then the order of vanishing of D_x^Λ near the zeros of $P_i(\mathbf{u})$.

In a rank-1 theory the curve Σ is elliptic and it can always be brought in its Weierstrass form before taking its x discriminant. This is how we define the quantum discriminant of a rank-1 theory. Since the singular locus in this case is just the origin, the quantum discriminant will be proportional to u^ℓ , u being the global coordinate describing the CB of the rank-1 SCFT. It is known that in this case the order of vanishing ℓ strongly constrains the scaling dimension $\Delta(u)$ as can be seen looking at the third column of table 1. For instance, if $\ell = 1$ then the CB geometry has to be an I_1 and $\Delta(u) = 1$. If $\ell = 10$ there are instead three possibilities II^* , I_4^* or I_{10} , which correspond to $\Delta(u) = 6, 2, 1$, respectively. There is a natural generalization of this fact for the quantum discriminant of higher rank theories and the ℓ_i in (2.22) can be identified with the order of vanishing of the quantum discriminant of the rank-1 theory $\mathcal{T}_i$ supported on $\mathcal{S}_i$ and therefore strongly constrains the Δ_i in (2.14).

Example: SCFTs with $\mathfrak{su}(3)$ gauge algebra. Let's go back to our $\mathfrak{su}(3)$ theories and consider first theories 1 and 2. The SW curve for the two theories in terms of a two-parameter family of hyperelliptic curve was derived in [27, 28]. When the masses are turned off the two curves coincide:

$$y^2 = x^6 + (v + ux)^2 + x^3(v + ux)\tau \quad (2.23)$$

⁶Good references are the original papers [22, 23] or some of the reviews of SW geometry e.g. [24–26].

Scaling behaviors near rank 1 singularities			
Name	planar SW curve	$\text{ord}_0(D_x^\Lambda)$	$\Delta(u)$
II^*	$y^2 = x^3 + u^5$	10	6
III^*	$y^2 = x^3 + u^3x$	9	4
IV^*	$y^2 = x^3 + u^4$	8	3
I_0^*	$y^2 = \prod_{i=1}^3 (x - e_i(\tau)u)$	6	2
IV	$y^2 = x^3 + u^2$	4	$3/2$
III	$y^2 = x^3 + ux$	3	$4/3$
II	$y^2 = x^3 + u$	2	$6/5$
$I_n^* \ (n>0)$	$y^2 = x^3 + ux^2 + \Lambda^{-2n}u^{n+3}$	$n+6$	2
$I_n \ (n>0)$	$y^2 = (x-1)(x^2 + \Lambda^{-n}u^n)$	n	1

Table 1. Scaling forms of rank 1 planar special Kahler singularities, labeled by their Kodaira type (column 1), a representative family of elliptic curves with singularity at $u = 0$ (column 2), order of vanishing of the discriminant of the curve at $u = 0$ (column 3), mass dimension of u (column 4), a representative of the $SL(2, \mathbb{Z})$ conjugacy class of the monodromy around $u = 0$ (column 5), the deficit angle of the associated conical geometry (column 6), and the value of the low energy $U(1)$ coupling at the singularity (column 7). The first seven rows are scale invariant. The last two rows give infinite series of singularities which have a further dimensionful parameter Λ so are not scale invariant; they can be interpreted as IR free theories since $\tau_0 = i\infty$.

where τ is the $\mathfrak{su}(3)$ holomorphic gauge coupling. Taking the x discriminant of the r.h.s. of (2.23) we can then readily compute its quantum discriminant:

$$\left. \begin{array}{l} 1. \\ 2. \end{array} \right\} : \quad D_x^\Lambda \sim v^6(u^3 + \lambda_1 v^2)(u^3 + \lambda_2 v^2) \quad (2.24)$$

where $\lambda_i \in \mathbb{C}^*$ and we left out inessential numerical coefficients. As expected, the zeros of the quantum discriminant coincide with the discriminant locus, see (2.10a), but now we can extract extra information from the order of such zeros:

$$\ell_1 = 6, \quad \ell_{2,3} = 1. \quad (2.25)$$

Before interpreting these numbers it is useful to remind the physical interpretation of the I_n and I_n^* geometry in table 1 which are associated, respectively, to the CB of a $U(1)$ and a $\mathfrak{su}(2)$ $\mathcal{N} = 2$ gauge theory. The n in both cases refers to the coefficient of the gauge coupling beta function which is:

$$U(1) : \quad b_g = \sum_i q_i^2 \quad (2.26)$$

$$\mathfrak{su}(2) : \quad b_g = \sum_i \left(T(\mathbf{R}_{\text{hypers}}^i) - 4 \right) \quad (2.27)$$

where the q_i are the electric charges of the hypermultiplets and $T(\mathbf{R}_{\text{hypers}}^i)$ is the quadratic index of the $\mathfrak{su}(2)$ representation of the i -th hypermultiplet.⁷ Using (2.16) and (2.17) we

⁷We choose a normalization for which $T(\mathbf{3}) = 4$.

can straightforwardly apply these formula to conclude that for both theory 1. and 2. we expect the CB of $\mathcal{T}_1$ to be an I_6 while that of $\mathcal{T}_{2,3}$ to be an I_1 . We leave it as an exercise for the reader to check the matching of the values in (2.25) from the corresponding entry in table 1.

For theory 3. no SW curve is known in the form that we need for the computation of the quantum discriminant. Yet above we have determined the theory supported on the singular locus and could use this information to infer the expected quantum discriminant:

$$3. \quad D_x^\Lambda \sim (u^3 + \lambda v^2)^6 \quad (2.28)$$

where we used (2.18) and the fact that $T(\mathbf{3}) = 4$ and therefore the CB of the theory supported on the singular locus is a I_0^* .

As we saw from the examples, the quantum discriminant improves on the discriminant locus and provides considerable extra information about the low-energy limit of the superconformal theory. A big shortcoming of this nice story is that the value of the ℓ_i 's, which should be an invariant of the special Kahler geometry, is instead not invariant under coordinate reparametrization of the curve (2.20). In defining (2.21) we have secretly dealt with this ambiguity by assuming that the curve in (2.20) is in some sort of higher rank generalization of the Weierstrass form. This is obtained by extending the $\mathbb{C}^*$ action on $\mathcal{C}$ to $\Sigma_{\mathbf{u}} \cong \mathcal{C}_r$, fixing the $[x]$ and $[y]$ in terms of $[\mathbf{u}]$ via the definition of the SW differential:

$$\frac{\partial \lambda_{\text{SW}}}{\partial u_i} = \frac{x^{i-1} dx}{y^2} := \omega_i, \quad \mathbf{u} = (u_1, \dots, u_r) \quad (2.29)$$

where the ω_i , $i = 1, \dots, r$ is a basis of $H^0(\mathcal{C}_r, \Omega_{\mathcal{C}_r}^1)$, the space of globally defined holomorphic one-forms on $\Sigma_{\mathbf{u}}$. Once we appropriately extend the $\mathbb{C}^*$ action, we can harvest the power of scale invariance. It in fact follows that $\mathcal{P}(x, \mathbf{u})$ in (2.20) must be a weighted homogeneous polynomial of weight:

$$[\mathcal{P}(x, \mathbf{u})] = 2[y] \quad (2.30)$$

which immediately fixes the degree d of the highest term in x . We can fix the remaining reparametrization invariance setting to zero the term of degree $d - 1$ in x though the quantum discriminant does not depend on this last step.

The quantum discriminant is certainly a useful object to consider, yet our definition seems too convoluted for an object whose physical interpretation is instead sharp and appears to be intrinsic to the special Kahler geometry of $\mathcal{C}$. Furthermore hyper-elliptic curves are a set of measure zero in the space of complex curves of genus $g > 2$. Therefore the special Kahler geometry of a generic $\mathcal{N} = 2$ SCFT with rank ≥ 3 will not have a formulation in these terms and we don't know how to replace the definition of the quantum discriminant in a useful way. Finding an intrinsic algebraic definition of the quantum discriminant which does not make use of the explicit realization of special Kahler geometry in terms of a family of hyperelliptic curves will bring significant progress in the study of scale invariant CBs of arbitrary ranks. Most importantly, the quantum discriminant is not the natural quantity which enters the twisted partition function.

2.5 Physical discriminant

In this subsection we introduce an important definition for the discussion of the next section where we analyze the partition function of topologically twisted $\mathcal{N} = 2$ SCFTs at arbitrary rank. The partition function of a topologically twisted $\mathcal{N} = 2$ SCFT reduces to an integral over the CB parameter $\mathbf{u}$. The non trivial dependence of the integrand can be fixed by carefully analyzing its dependence near the singular locus $\bar{\mathcal{S}}$. The *physical discriminant* arises precisely in this context.

As we will review below, the $\mathcal{B}$ and $\mathcal{C}_a$ factors of the measure of the integrand of the partition can only vanish or diverge at the zeros of (2.2). In particular by the general consideration of [14–17], the $\mathcal{B}$ factor should be proportional to a close cousin of the quantum discriminant which was called *physical discriminant* in [10]:

$$\mathcal{B} = \beta \left(D_x^{\text{phys}} \right)^{1/8} \quad (2.31)$$

where β is a constant. D_x^{phys} is a close cousin of (2.21) in the sense that we expect D_x^{phys} to share the same zeros as the quantum discriminant and to also be non-reduced. We will parametrize our ignorance in the following way:

$$D_x^{\text{phys}} \sim \prod_{i \in I} \left[P_i(\mathbf{u}) \right]^{b_i}, \quad \bar{\mathcal{S}}_i =: \left\{ \mathbf{u} \in \mathcal{C} \mid P_i(\mathbf{u}) = 0 \right\} \quad (2.32)$$

which is indeed very resembling of (2.21)]. Determining the form of the b_i s will be one of the goals of the next section. It will be the determination of these exponents that leads to our central charge formulae (1.1a)–(1.1c).

3 Twisted partition function and central charge

$\mathcal{N} = 2$ supersymmetric gauge theories are related via a well-known twisting procedure to topological field theories [1], see [29, 30] for a pedagogical presentation of this subject. The starting point is the Euclidean version of $\mathcal{N} = 2$ supersymmetric theories with total symmetry group:

$$\mathcal{G} = \mathcal{K} \times \text{SU}(2)_R \times \text{U}(1)_R \quad (3.1)$$

where the $\mathcal{K} = \text{SU}(2)_+ \times \text{SU}(2)_-$ is the rotation group and $\text{SU}(2)_R \times \text{U}(1)_R$ the R-symmetry. Under this symmetry the supercharges transform as:

$$Q : \quad (\mathbf{1}, \mathbf{2}, \mathbf{2})^{-1} \quad (3.2a)$$

$$\tilde{Q} : \quad (\mathbf{2}, \mathbf{1}, \mathbf{2})^1 \quad (3.2b)$$

The topological twisting procedure consists in modifying the coupling to gravity by redefining the spin of the fields. To do this we introduce an external $\text{SU}(2)_R$ gauge potential setting it equal to the self-dual part of the spin connection. Or, in other words, re-define the rotation group as $\mathcal{K}' = \text{SU}(2)'_+ \times \text{SU}(2)_-$ where $\text{SU}(2)'_+ = [\text{SU}(2)_+ \times \text{SU}(2)_R]_{\text{diag}}$ is the

diagonal combination of the two remaining $SU(2)$'s. It is easy to see that under $\mathcal{K}' \times U(1)_R$ the supercharges transform:

$$Q : \quad (\mathbf{2}, \mathbf{2})^{-1} \quad (3.3)$$

$$\tilde{Q} : \quad (\mathbf{3} \oplus \mathbf{1}, \mathbf{1})^1 \quad (3.4)$$

therefore there is one supercharge which is a singlet under the rotation group. We call it Q_{BRST} and it is with respect to this supercharge that the supersymmetry preserved on a curve background (see below) is defined. The topological sector of the twisted theory is then obtained by projecting into the cohomology of the singlet supercharge. The resulting theory has many remarkable properties including the fact that its partition function, discussed in more detail momentarily, does not depend on the metric [1]. This is the sense in which the theory is topologically invariant.

Upon flowing to the IR on the CB, the partition function of the twisted theory is given by the path integral of the low-energy lagrangian after integrating out massive degrees of freedom. Since the twisted theory is topological, the contributions of the massive states can only result in terms which are proportional to topological densities [10, 14]:

$$\mathcal{L}_{\text{twisted}} \sim \mathcal{L}_{\text{IR}[V,H]} + (\log \mathcal{A}) \text{tr} R \wedge \tilde{R} + (\log \mathcal{B}) \text{tr} R \wedge R + \sum_a (\log \mathcal{C}_a) \text{tr} F_{\mathfrak{f}_a} \wedge F_{\mathfrak{f}_a} \quad (3.5)$$

R is the Riemann tensor of the four manifold and $F_{\mathfrak{f}_a}$ is an external gauge field coupled to the simple factor $\mathfrak{f}_a$ of the twisted theory flavor algebra $\mathfrak{f} = \oplus_a \mathfrak{f}_a$. $\mathcal{L}_{\text{IR}[V,H]}$ is the sum of a BRST exact term and a contribution which depends on the coupling of the low-energy Abelian theory on a generic point of the CB. The details of $\mathcal{L}_{\text{IR}[V,H]}$ will not play any role below so we will not write its dependence explicitly and refer to the literature for details [14, 16]. The factors $(\mathcal{A}, \mathcal{B}, \mathcal{C}_a)$ depend holomorphically on the globally defined CB coordinates $\mathbf{u}$ and we will spend a considerable part of this section to determine their detailed form. We will henceforth not discuss possible $U(1)$ flavor factor as in that case equation (3.5) becomes considerable more subtle and non-holomorphic terms might arise.⁸

Bringing all together we have the following expression for the twisted partition function [10, 14]

$$Z = \int [dV][dH] \mathcal{A}^\chi \mathcal{B}^\sigma \prod_a \mathcal{C}_a^{n_a} e^{S_{\text{IR}[V,H]}}. \quad (3.6)$$

where σ and χ are the signature and Euler characteristics of the four manifold and n_a is the instanton number for the $\mathfrak{f}_a$ factor. The path integral is over the n_V (IR free) massless neutral vector multiplet fields and n_H massless neutral hypermultiplet fields (if any) on the generic point of the CB $\mathcal{C}$. When there are free hypers on the generic point of $\mathcal{C}$, by giving a vev to them, it is possible to see $\mathcal{C}$ as a part of a larger branch of the moduli space usually called the extended CB (ECB). n_H is then naturally interpreted as the quaternionic dimension of the ECB h . n_V is instead naturally identified with the rank r of the theories. Therefore we have:

$$n_H = h, \quad n_V = r. \quad (3.7)$$

⁸We thank G. Moore for clarifying this point and sharing some unpublished results [31].

The integral (3.6) includes an ordinary integral over the 0-modes (constant modes) of the vector multiplet scalars, u .

The expression for the $\mathcal{A}$ factor was determined in full generality to be [15, 16, 18]:

$$\mathcal{A} = \alpha \left[\det \left(\frac{\partial u^i}{\partial a^j} \right) \right]^{1/2} \quad (3.8)$$

where α is a constant and for a theory of rank- r the indices i, j run up to r . The expression for the $\mathcal{B}$ and $\mathcal{C}_a$ factor are instead more involved. They can only vanish along the singular locus of the CB and therefore both $\mathcal{B}$ and $\mathcal{C}_a$ are somewhat related to the discriminant locus (2.2). In order to clarify this dependence we need to take a detour and remind the reader about the relationship between the $U(1)_R$ anomaly of the topologically twisted $\mathcal{N} = 2$ supersymmetric theory and the value of the $(a, c, k_{\mathfrak{f}})$ of the untwisted version. This relationship will also allow us to finally derive our central charge formulae.

3.1 $U(1)$ anomaly and central charges

The a and c central charges of the 4d conformal algebra are certain coefficients in OPEs of energy-momentum tensors, and the k central charges appear in the OPEs of flavor currents. We will assume throughout that the flavor algebra $\mathfrak{f} = \oplus_a \mathfrak{f}_a$ is a sum of simple factors and each factor will have a separate k_a central charge. The fact that these quantities appear in the ΔR anomaly follows from the fact that (a, c, k_a) appear in the scale anomaly in the presence of a background metric and background gauge fields for $\mathfrak{f}$. Then $\mathcal{N} = 2$ superconformal symmetry relates the scale anomaly to 't Hooft anomalies for the $U(1)_R \oplus SU(2)_R \oplus_a \mathfrak{f}_a$ global symmetry, with the result that in the presence of a background metric and background gauge fields for the global symmetries, the conservation of the $U(1)_R$ current is broken by terms proportional to the central charges times topological densities formed from the background fields. Notice that background metric and gauge fields describing an arbitrary smooth oriented 4-fold M with F -bundle (where F is the flavor symmetry group with Lie algebra $\mathfrak{f}$) generally break $\mathcal{N} = 2$ supersymmetry. However, as we reviewed above, the twisted topological version of it is still protected by a supersymmetry [1], despite a curve background, and therefore we can still talk about a $U(1)_R$. The result [10, 14] is that the partition function of the twisted theory on M with an F -bundle carries $U(1)_R$ charge⁹

$$\Delta R = (2a - c) \cdot \chi + \frac{3}{2} c \cdot \sigma - \frac{1}{2} \sum_a k_a \cdot n_a, \quad (3.9)$$

where χ and σ are the Euler characteristics and signature of M and n_a are the instanton numbers of the F -bundle. Here we have again assumed that the flavor symmetry of the SCFT is a semi-simple Lie algebra $\mathfrak{f} = \oplus_a \mathfrak{f}_a$ and k_a is the level of flavor currents of the simple factor $\mathfrak{f}_a$.

⁹Here we are using a normalization of the $U(1)_R$ charge such that $R(u) = \Delta(u)$. This differs from that used in [10] by a factor of two.

Eq. (3.9) corresponds to the standard normalizations of the central charges where for n_V free vector multiplets and n_H free hypermultiplets

$$24a = 5n_V + n_H, \quad 12c = 2n_V + n_H, \quad k_a = T_a(\mathbf{2h}). \quad (3.10)$$

Thus, in this case

$$\Delta R_{\text{free}} = \frac{1}{4}n_V \cdot \chi + \left(\frac{1}{4}n_V + \frac{1}{8}n_H \right) \cdot \sigma - \frac{1}{2} \sum_a T_a(\mathbf{2h}) \cdot n_a. \quad (3.11)$$

Here $\mathbf{2h}$ is the (reducible) representation of $\mathfrak{f}$ under which the $2n_H$ half-hypermultiplets transform. $T_a(\mathbf{2h})$ is the quadratic index of $\mathbf{2h}$ with respect to the $\mathfrak{f}_a$ factor.¹⁰ In case $n_H = 0$, there is no contribution from the last term in (3.11), so we adopt the convention that $T(\mathbf{0}) := 0$.

From the explicit expression of the partition function (3.6) its total $U(1)_R$ charge can be evaluated at a generic (i.e., non-singular) point on the CB to be

$$\Delta R = \left(R(\mathcal{A}) + \frac{1}{4}n_V \right) \cdot \chi + \left(R(\mathcal{B}) + \frac{1}{4}n_V + \frac{1}{8}n_H \right) \cdot \sigma + \sum_a \left(R(\mathcal{C}_a) - \frac{1}{2}T_a(\mathbf{2h}) \right) \cdot n_a, \quad (3.12)$$

where we have used (3.11) to evaluate the contribution from the $[dV][dH]$ measure and have left the R -charge of the $(\mathcal{A}, \mathcal{B}, \mathcal{C}_a)$ factors to be determined. Comparing this to (3.9) for arbitrary (χ, σ, n_a) and using (3.7) gives [10]:

$$24a = 5r + h + 12R(\mathcal{A}) + 8R(\mathcal{B}), \quad (3.13a)$$

$$12c = 2r + h + 8R(\mathcal{B}), \quad (3.13b)$$

$$k_a = T_a(\mathbf{2h}) - 2R(\mathcal{C}_a). \quad (3.13c)$$

Let's stress again that the twisting procedure is a purely calculational tool for us, the (a, c, k_a) above refer to the corresponding quantities of the untwisted, non-topological $\mathcal{N} = 2$ SCFT in flat space.

Computing $R(\mathcal{A})$, from (3.8), is a straightforward task. Indeed a^j , $j = 1, \dots, r$, are the special coordinates on the CB (see appendix A) which have scaling dimension one, and thus, in our conventions, $U(1)_R$ charge one. The prefactor α is u -independent. In the conformal case it can only depend on constants which are all dimensionless, so $R(\alpha) = 0$. We therefore have

$$R(\mathcal{A}) = \frac{\sum_{i=1}^r \Delta_{\mathbf{u}_i} - r}{2} \quad (3.14)$$

where $\Delta_{\mathbf{u}_i}$ is the scaling dimension of the i -th entry of the vector of r globally defined complex coordinates on the CB $\mathbf{u}$. We turn now to a detailed discussion of the $\mathcal{B}$ factor.

¹⁰If $\mathbf{2h}$ decomposes into irreps of $\oplus_{a=1}^L \mathfrak{f}_a$ according to $\mathbf{2h} = \oplus_{\alpha} (\mathbf{r}_{\alpha 1} \otimes \mathbf{r}_{\alpha 2} \otimes \dots \otimes \mathbf{r}_{\alpha L})$, then $T_a(\mathbf{2h}) = \sum_{\alpha} \left(\prod_{b \neq a} r_{\alpha b} \right) T(\mathbf{r}_{\alpha a})$. The quadratic index for a simple factor is proportional to the sum of the squared-lengths of weights in $\mathbf{2h}$, $T(\mathbf{2h}) := (1/\text{rank} \mathfrak{f}) \sum_{\lambda} (\lambda, \lambda)$, where the weights are normalized so that the long roots of $\mathfrak{f}$ have length-squared 2. This is the normalization for which $T(\mathbf{n}) = 1$ for $SU(n)$.

3.2 Final form of the partition function and central charge formulae

We already anticipated, as it has been argued for many years, that the $\mathcal{B}$ factors has to have the following form:

$$\mathcal{B} = \beta \left(D_x^{\text{phys}} \right)^{1/8} \quad (3.15)$$

and the form of D_x^{phys} was written explicitly in (2.32). We will turn now to the computation of the b_i .

When the twisted theory is put on a smooth spin 4-manifold, $\mathcal{B}^\sigma$ has to be single-valued functions on the CB.¹¹ Since for smooth spin 4-manifolds $\sigma \in 16\mathbb{Z}$ [32, 33], we see that $\mathcal{B}^{16}$ must be a single-valued holomorphic function of (u, v) . This immediately implies that

$$b_i \in \frac{\mathbb{Z}}{2}. \quad (3.16)$$

Recall that the ℓ_i in (2.21) are instead integers. To determine the quantities b_i explicitly, we will zoom in on the various complex co-dimension one strata of the singular locus $\mathcal{S}_i$. There, we can exploit the fact that the $U(1)_R$ of $\mathcal{T}_i$ (the rank-1 theory supported on the singular stratum $\mathcal{S}_i$) provides an accidental low-energy $U(1)_R$ symmetry which constraints the integrand of the partition function in (3.6) by its anomaly ΔR . Since this anomaly depends on the physical properties of the low-energy effective theory $\mathcal{T}_i$, we will be able to relate the b_i to properties of rank-1 theories. This strategy is very resembling of that followed in [14].

To do that, consider probing $\mathcal{S}_i$ away from the origin. We can choose a set of coordinates $(\mathbf{u}_\parallel, u_\perp)$ on $\mathcal{C}$ which are “adapted” to $\mathcal{S}_i$. $u_\perp$ is a coordinate transverse to $\mathcal{S}_i$ while $\mathbf{u}_\parallel$ are an $r - 1$ dimensional vector of coordinates *along* $\mathcal{S}_i$ in the following sense:¹²

$$P_i(\mathbf{u}_\parallel, u_\perp) = 0, \quad \text{for } \{u_\perp = 0, \mathbf{u}_\parallel \in \mathbb{C}^{r-1}\}. \quad (3.17)$$

The CB parameter of the rank-1 theory $\mathcal{T}_i$, which we indicated as u_i , is a natural choice, at least locally, for $u_\perp$ and therefore we will pick $u_\perp \equiv u_i$, this point is explained in more detail in appendix A.1. We can expand (2.32) around $u_i = 0$ finding:

$$D_x^{\text{phys}} \sim u_i^{b_i} \cdot P(u_i, \mathbf{u}_\parallel) \quad (3.18)$$

where $P(\mathbf{u}_\parallel, u_i)$, as long as $\mathbf{u}_\parallel \neq 0$, does not vanish as $u_i \rightarrow 0$.

As it is explained in appendix A.2, only the u_i is charged under the $U(1)_R$ symmetry arising near $\mathcal{S}_i$ and the $U(1)_R$ charge of the u_i is determined by its scaling dimension Δ_i defined in (2.14):

$$R(u_i) = \Delta_i, \quad R(\mathbf{u}_\parallel) = 0. \quad (3.19)$$

¹¹For non-spin 4-manifolds the $\mathcal{B}^\sigma$ measure factor may be multi-valued on the CB [14].

¹²Zeros of a polynomial form a closed set, therefore $P_i(\mathbf{u}) = 0$ defines $\overline{\mathcal{S}}_i$ not $\mathcal{S}_i$. Here we focus on $\mathcal{S}_i$ because we are interested in the set which supports a well-defined $\mathcal{T}_i$. Therefore, to be precise and a bit pedantic, we should restrict to those values of $\mathbf{u}_\parallel$ such that $P_i(\mathbf{u}_\parallel, u_\perp) = 0$ spans $\mathcal{S}_i$.

Bringing (2.31), (3.18) and (3.19) together we conclude that

$$\frac{b_i \Delta_i}{8} = R(\mathcal{B})|_i \quad (3.20)$$

where $R(\mathcal{B})|_i$ indicates the condition that the $\mathcal{B}$ R -charge has to satisfy near $\mathcal{S}_i$. From eq. (3.13b) we can solve for $R(\mathcal{B})|_i$ and get:

$$R(\mathcal{B})|_i = \frac{12c_i - 2 - h_i}{8} \quad (3.21)$$

where the i subscript indicates that the given quantity refers to the rank-1 theory $\mathcal{T}_i$, see for example the quantities calculated in (2.19).¹³ We then finally get an expression for b_i :

$$b_i = \frac{12c_i - 2 - h_i}{\Delta_i} \quad (3.22)$$

The quantity in (3.22) has already appeared in [34]¹⁴ and has a clear physical significance of being linked to the deformation pattern of the rank-1 theory $\mathcal{T}_i$ supported on the particular stratum of the singular locus. In other words the value of b_i is related to the way in which $\mathcal{T}_i$ behaves under relevant deformations. The most promising consequence of this observation is that the study of the mass deformations of higher rank scale invariant CBs might ultimately reduce to a clever application of techniques which were thoroughly analyzed and understood in the context of rank-1. What we are going to learn next studying $\mathcal{C}_a$, will further support this picture.

Before turning to $\mathcal{C}_a$, let us write explicitly $R(\mathcal{B})$. From (2.31) and apply the analysis above to all co-dimension one components we obtain:

$$8R(\mathcal{B}) = \sum_{i \in I} \Delta_i^{\text{sing}} b_i = \sum_{i \in I} \Delta_i^{\text{sing}} \frac{12c_i - 2 - h_i}{\Delta_i}. \quad (3.23)$$

where Δ_i^{sing} is defined in (2.3). Notice that the fact that the (3.22) must be semi-integers, can be converted in non-trivial constraints on the rank-1 theories $\mathcal{T}_i$ which have been discussed in detail in [34].

We have now determined $R(\mathcal{A})$ and $R(\mathcal{B})$ and we can finally use (3.13a)–(3.13b) to derive the first two central charge formulae which relate the (a, c) of a SCFT of arbitrary rank r to known quantities of the rank-1 theories supported on its CB singular locus:

$$24a = 5r + h + 6 \left(\sum_{\ell=1}^r \Delta_{\mathbf{u}_\ell} - 1 \right) + \sum_{i \in I} \Delta_i^{\text{sing}} \frac{12c_i - 2 - h_i}{\Delta_i}, \quad (3.24a)$$

$$12c = 2r + h + \sum_{i \in I} \Delta_i^{\text{sing}} \frac{12c_i - 2 - h_i}{\Delta_i}, \quad (3.24b)$$

where $i \in I$ labels the complex co-dimension one strata of $\mathcal{C}$. We will discuss the flavor level, and in general the flavor structure of $\mathcal{N} = 2$ SCFTs in the next section.

¹³Notice that to obtain the formula above we used $n_V = 1$ in plugging the value from (3.13b) since the theory $\mathcal{T}_i$ is a rank-1 theory.

¹⁴The b_i in (3.22) and those defined in [34] differ by an inessential factor of 2 due to a different definition of the physical discriminant.

Example: SCFTs with $\mathfrak{su}(3)$ gauge algebra. Let's apply the formulae that we just derived to our $\mathfrak{su}(3)$ examples. In particular we will see that while theories 1. and 2. shared the same quantum discriminant they will have a different physical discriminant. The calculation here is for pedagogical purposes only since we won't learn anything new on these very well studied lagrangian theories. We will leverage the power of our central charge formulae in [35] where we compute many new properties of $\mathcal{N} = 2$ SCFTs of rank-2 theories and their moduli space.

Let's start reminding the reader of the properties of these theories:

	$24a$	$12c$	$\mathfrak{f}_{k_f}$	
$N_f = 6$	58	34	$\mathfrak{su}(6)_6$	
$1(\mathbf{6}) \oplus 1(\mathbf{3})$	49	25	$\emptyset$	
$\mathcal{N} = 4$	48	24	$\mathfrak{sp}(1)_8$	(3.25)

These can be straightforwardly computed using (3.10) and counting the number of hypermultiplets and vector multiplets. We will reproduce these values from the structure of the CB, in particular using the quantities in (2.19). It will be a rewarding exercise. Here we will focus on the values for (a, c) while the flavor part will be discussed in the next section.

Following (2.32), we can parametrize the physical discriminants of the three theories as:

$$\left. \begin{array}{l} 1. \\ 2. \end{array} \right\} : D_x^{\text{phys}} \sim v^{b_1^{1,2}} (u^2 + \lambda_1 v^3)^{b_2^{1,2}} (u^2 + \lambda_2 v^3)^{b_3^{1,2}} \quad (3.26)$$

$$3. : D_x^{\text{phys}} \sim (u^2 + \lambda v^3)^{b^{\mathcal{N}=4}}. \quad (3.27)$$

We have already determined the low-energy theory along each component of the singularity, see (2.16), (2.17) and (2.18), therefore we can directly use (2.19) and equation (3.22) to find the following result:

$$\begin{aligned} b_1^1 &= 6, & b_1^2 &= 3, \\ b_2^{1,2} &= b_3^{1,2} = 1, \\ b^{\mathcal{N}=4} &= 3. \end{aligned} \quad (3.28)$$

From (2.32), this immediately gives the expression for the physical discriminant of three theories:

$$1. \quad D_x^{\text{phys}} \sim v^6 (u^3 + \lambda_1 v^2) (u^3 + \lambda_2 v^2) \quad (3.29)$$

$$2. \quad D_x^{\text{phys}} \sim v^3 (u^3 + \lambda_1 v^2) (u^3 + \lambda_2 v^2) \quad (3.30)$$

$$3. \quad D_x^{\text{phys}} \sim (u^3 + \lambda v^2)^3 \quad (3.31)$$

A quick comparison with (2.24) shows that the quantum and physical discriminants coincide only in one of the three cases.¹⁵

¹⁵There is a clear generalization of this fact, physical and quantum discriminant coincides iff the CB of all rank-1 theories supported on all $\mathcal{S}_i$'s deform, turning on all relevant deformations, into I_1 singularities [34, 36–38].

Now let's tackle the central charges of these theories. In this, $\mathfrak{su}(3)$, case, $\Delta_u=2$ and $\Delta_v=3$ and $r=2$ so (3.24a) and (3.24b) reduce to:

$$24a = 10 + h + 18 + \sum_{i \in I} \Delta_i^{\text{sing}} \frac{12c_i - 2 - h_i}{\Delta_i} = 28 + h + \sum_{i \in I} \Delta_i^{\text{sing}} b_i, \quad (3.32a)$$

$$12c = 4 + h + \sum_{i \in I} \Delta_i^{\text{sing}} \frac{12c_i - 2 - h_i}{\Delta_i} = 4 + h + \sum_{i \in I} \Delta_i^{\text{sing}} b_i, \quad (3.32b)$$

Starting from theory 3., the singular locus has only one component (see (2.10b)) and the sum will reduce to a single term. This is a rank-2 $\mathcal{N}=4$ theory, therefore there it has $h=2$:

$$24a_{\mathcal{N}=4} = 30 + \Delta^{\text{sing}} b^{\mathcal{N}=4} \quad (3.33)$$

$$12c_{\mathcal{N}=4} = 6 + \Delta^{\text{sing}} b^{\mathcal{N}=4} \quad (3.34)$$

From (2.10b), $\Delta^{\text{sing}} = 6$ and we beautifully reproduce the value for a and c in (3.25). A very similar calculation can be performed for theory 1. and 2., but now the sum in the central charge formulae includes three factors. We leave it up to the reader to check that things work neatly.

4 Flavor structure of general $\mathcal{N}=2$ SCFTs

Recall that in a SCFT with flavor symmetry $\mathfrak{f} = \oplus_a \mathfrak{f}_a$, where we restrict to $\mathfrak{f}_a$ simple, the term $\mathcal{C}_a$ arises from the contribution to the low-energy lagrangian (3.5) proportional to the topological density:

$$\text{Tr} F_{\mathfrak{f}_a} \wedge F_{\mathfrak{f}_a} \quad (4.1)$$

and $F_{\mathfrak{f}_a}$ is an external gauge field coupled to the global symmetry factor $\mathfrak{f}_a$. As discussed above, $\mathcal{C}_a$ is holomorphic in $\mathbf{u}$ by topological invariance but, to determine the form of $\mathcal{C}_a$, we have to understand how the flavor symmetry $\mathfrak{f}_a$ acts on the theory at a generic point of the CB. Notice that all CB operators are uncharged under the flavor symmetry therefore no spontaneous breaking takes place anywhere on $\mathcal{C}$. As a consequence, the full spectrum at any point of CB, including massive and massless states, should be organized in representations of the full flavor symmetry $\mathfrak{f}$. But each simple factor $\mathfrak{f}_a$ might or might not act on the light states. Determining this is particularly relevant as the zeros of $\mathcal{C}_a$ can only lie along those components of $\overline{\mathcal{S}}$ where the massless states are charged under $\mathfrak{f}_a$.

This is an important point so let's elaborate on it further. From the definition of the $\mathcal{C}_a$ factor (3.5), the zeros of $\mathcal{C}_a$ correspond to loci where the contribution to the low-energy Lagrangian, proportional to the topological density (4.1), diverges. This can only happen at the zeros of the discriminant locus (2.2). But, more specifically, a diverging contribution proportional to the topological quantity (4.1) can only be generated if the states becoming massless are charged under $\mathfrak{f}_a$, thus the previous statement. We label the special strata where this happens as $\mathcal{S}_i$ with $i \in I_{\mathfrak{f}_a}$. Notice that for $i \in I_{\mathfrak{f}_a}$, the rank-1 theory $\mathcal{T}_i$, describing the massless excitations on $\mathcal{S}_i$, is strongly constrained by the fact that its flavor symmetry should contain a $\mathfrak{f}_a$ factor.

From these considerations and the fact that all the components of the singular locus can be algebraically parametrized using the polynomial $P_i(\mathbf{u})$ in (2.2), it then follows that $\mathcal{C}_a$ should have the general form:

$$\mathcal{C}_a \sim \prod_{i \in I_{\mathfrak{f}_a}} [P_i(\mathbf{u})]^{e_i^a}, \quad (4.2)$$

where the e_i^a are for the moment arbitrary. One might wonder whether it is possible that only a subalgebra $\tilde{\mathfrak{f}}_a \subset \mathfrak{f}_a$ is realized on the massless spectrum. But $\mathfrak{f}_a$ is simple and since we argued that it is nowhere spontaneously broken, this possibility is forbidden. An IR flavor symmetry enhancement is instead possible.

Let's now study the zeros of $\mathcal{C}_a$. First let's indicate the cardinality of $I_{\mathfrak{f}_a}$ as $\mathbf{n}_a \geq 0$. $\mathcal{C}_a$ is a holomorphic function of $\mathbf{u}$ and $\mathcal{C}$ is a connected complex manifold ($\overline{\mathcal{S}}_{\text{cplx}} = \emptyset$). This implies that the zeros of $\mathcal{C}_a$ have to be in complex co-dimension one. $\mathfrak{f}_a$ acts on the massless spectrum at the superconformal vacuum by assumption, therefore $\mathcal{C}_a$ vanishes at $\mathbf{u} = 0$. This is enough to conclude that $\mathcal{C}_a$ has to have a non trivial dependence on at least one of the $P_i(\mathbf{u})$ and, for any simple factor $\mathfrak{f}_a \subset \mathfrak{f}$ of the flavor symmetry, $\mathbf{n}_a \neq 0$. In other words we have just proven the following remarkable fact:

Fact 1. *The Coulomb branch of any $\mathcal{N} = 2$ SCFT with a semi-simple global symmetry $\mathfrak{f} = \oplus_a \mathfrak{f}_a$ has, for each factor $\mathfrak{f}_a$, one or multiple connected co-dimension one strata $\mathcal{S}_i$ where the charged massless spectrum is organized in non-trivial irreducible representations of $\mathfrak{f}_a$.*

Since mass deformations of $\mathcal{N} = 2$ theories can be interpreted as background field configurations of the scalar component of a vector multiplet of a weakly gauged complexified flavor algebra $\mathfrak{f}$, the above facts proves the following general condition:

UV-IR simple flavor condition

All mass deformations of a rank- r $\mathcal{N} = 2$ SCFT deform the CB asymptotically and are realized, in the low-energy limit, as mass deformations of rank-1 theories supported on special complex co-dimension one loci.

These remarkable facts apply even beyond the superconformal case. In fact the gist of this result can be reframed as a statement that if there is any point P on the CB where a simple factor $\mathfrak{f}_a$ acts on the light spectrum, $\mathfrak{f}_a$ has to act on the light spectrum on a whole co-dimension one locus $\mathcal{V}_P$ going through P and no rank enhancement can appear along $\mathcal{V}_P$. This might sound counterintuitive as there are many examples of asymptotically free $\mathcal{N} = 2$ gauge theories which have loci on the CB with an enhanced simple flavor symmetry, e.g. [39, 40]. But in the asymptotically free case, the region near the origin is generally regular and the UV-IR simple flavor condition cannot be used to leverage the UV flavor data to constrain the flavor action on the CB. Conversely, in the SCFT case scale invariance implies that all complex co-dimension one singular components go through the superconformal vacuum. Therefore the CB action of the flavor symmetry on the light spectrum becomes extremely constrained.

We are now in a position to finally complete our discussion on the structure of the partition function. First, using (3.13c) and (4.2), we can immediately obtain an expression for k_a :

$$k_a = T_a(\mathbf{2h}) - 2 \sum_{i \in I_{f_a}} \Delta_i^{\text{sing}} e_i^a, \quad (4.3)$$

where Δ_i^{sing} is again defined in (2.3) and the sum is restricted to I_{f_a} . To determine the e_i^a we notice that for (3.6) to be well defined on the CB, $\mathcal{C}_a^{n_a}$ has to be a single valued function of $\mathbf{u}$. Since the instanton numbers $n_a \in \mathbb{Z}$ (at least for simply-connected flavor groups), $\mathcal{C}_a$ itself must be a single-valued holomorphic function of $\mathbf{u}$ and therefore $e_i^a \in \mathbb{Z}$.

To determine the value of individual e_i^a we can again zoom-in on the individual complex co-dimension one strata $\mathcal{S}_i$, as we did for the b_i , and apply (3.13c) to the rank-1 theory $\mathcal{T}_i$ supported there. In doing so, here we restrict to $i \in I_{f_a}$ since we are computing the level corresponding to the f_a factor. Following precisely the same steps which led us to the determination of the b_i s we obtain:

$$e_i^a = \frac{T_a(\mathbf{2h}_i) - k_a^i}{2d_i^a \Delta_i} \in \mathbb{Z}, \quad i \in I_{f_a}. \quad (4.4)$$

Here k_a^i is the level of the f_a flavor factor of the rank-1 theory $\mathcal{T}_i$, Δ_i is the scaling dimension of its CB parameter, and $T_a(\mathbf{2h}_i)$ is the quadratic index of its ECB. To account for a possible enhancement of the flavor symmetry in the IR, (4.4) also depends on d_i^a , the index of embedding of f_a into the appropriate simple factor of the flavor symmetry of $\mathcal{T}_i$.

Putting all together we obtain the last of our central charge formula which determines the level of the higher rank $\mathcal{N} = 2$ SCFT in terms of the levels of the rank-1 theories supported on the complex co-dimension one strata of its singular locus:

$$k_a = \sum_{i \in I_{f_a}} \frac{\Delta_i^{\text{sing}}}{d_i^a \Delta_i} (k_a^i - T_a(\mathbf{2h}_i)) + T_a(\mathbf{2h}). \quad (4.5)$$

Notice that the fact that the (4.4) must be integers, can be converted into non-trivial constraints which have been discussed in detail in [34].

Example: SCFTs with $\mathfrak{su}(3)$ gauge algebra. As we did throughout the manuscript, let's apply what we just learned to the $\mathfrak{su}(3)$ examples. In particular we will:

- i.* Check that the $\mathcal{N} = 2$ UV-IR simple flavor condition is indeed respected.
- ii.* Compute the flavor level in (3.25) directly from CB quantities.

To do the calculation of the flavor level we will analyze theory 1., that is $\mathfrak{su}(3)$ with $N_f = 6$ which has an $\mathfrak{su}(6)$ flavor symmetry at level 6, see (3.25). This result can be obtained using (3.10) and observing that the hypermultiplet fields transform in the fundamental of $\mathfrak{su}(3)$ and both the $\mathcal{N} = 1$ chiral and anti-chiral part contribute a $T(\mathbf{6}) = 1$ to (3.10). From our previous CB analysis, two of three strata support a $U(1)$ with a single hypermultiplet which carries no-simple flavor factor. The third stratum supports instead a

U(1) gauge theory with six massless hypermultiplets which carries a $\mathfrak{su}(6)$ flavor symmetry, see (2.19). This checks that the $\mathcal{N} = 2$ UV-IR simple flavor condition is indeed satisfied.

To compute the level of the $\mathfrak{su}(6)$ flavor symmetry of the $\mathfrak{su}(3)$ with $N_f = 6$, notice that (4.5) has an explicit dependence on Δ_i^{sing} . Therefore to reproduce the proper flavor level we need more information about the stratum where massless states carrying a $\mathfrak{su}(6)$ appear. From our previous analysis we know that the U(1) theory carrying an $\mathfrak{su}(6)$ is supported on the first entry of (2.11), implying $\Delta_1^{\text{sing}} = 3$. Finally, from (2.19), the level of the $\mathfrak{su}(6)$ of this U(1) theory is two, $k^1 = 2$ (the factor of three difference between the level of the $\mathfrak{su}(6)$ in the $\mathfrak{su}(3)$ theory and this case arises because now the hypers are no longer in a three dimensional representation but in a one-dimensional one). It is a matter of elementary algebra to obtain:

$$k_{\mathfrak{su}(6)} = \frac{3}{1}2 = 6. \quad (4.6)$$

Theory 2. carries no simple flavor symmetry, but a similar calculation can be carried out successfully for the $\mathcal{N} = 4$ $\mathfrak{su}(3)$.

5 Conclusions

In this paper we have thoroughly analyzed the dependence on the singularity structure of the Coulomb branch of the integrand of the partition function for an arbitrary topologically twisted $\mathcal{N} = 2$ superconformal field theory. This exercise allowed us to derive general formulae, (3.24a), (3.24b) and (4.5), relating the (a, c, k_f) conformal central charges of untwisted $\mathcal{N} = 2$ SCFTs to analogous quantities of rank-1 theories appearing in the low-energy description of the SCFTs at special loci of the their Coulomb branches. These formulae are valid for SCFTs of arbitrary rank and extend and generalize the work of Shapere and Tachikawa [10]. The main input which allowed this generalization is our improved understanding of the structure of the stratification of the singular locus of the Coulomb branch [35].

In performing our analysis we sharpen the definition of the discriminant of the Coulomb branch and its relevance for understanding the low-energy limit of $\mathcal{N} = 2$ SCFTs. We are also able to derive general lessons about the flavor structure of $\mathcal{N} = 2$ SCFTs and, in particular, on how the simple factors of their flavor symmetry are realized in the low-energy. Our main result in this regard is summarized in the $\mathcal{N} = 2$ UV-IR simple flavor condition.

The work described here as well as in a companion paper [35], is motivated by the classification program of four dimensional $\mathcal{N} = 2$ superconformal field theories [12] based on the systematic analysis of their Coulomb branch geometries. Particularly with the goal of extending the success story of rank-1 [34, 36–38] to arbitrary rank. At rank- r , this classification program involves two broad steps:

1. The characterization of all scale invariant CBs of complex dimension r .
2. The analysis of mass deformations of the entries in item 1.

The result described in this paper represent a, somewhat unexpected, progress on both of these fronts. The $\mathcal{N} = 2$ UV/IR simple flavor condition de-facto reduces the analysis of mass deformations of higher rank theories to a, perhaps involved, implementation of known rank-1 techniques. This therefore identifies a strategy to tackle item 2. for arbitrary ranks. Major progress in tackling item 1. is represented by the central charge formulae (3.24a), (3.24b) and (4.5). Even more so, our detailed understanding of the stratification of the CB singular locus presented in [35] lays the foundation for organizing a search of scale invariant special Kahler geometries of arbitrary complex dimensions. In [35] we will also present a more thorough report on the status of our classification program and the prospects to extend the work beyond rank-1.

Acknowledgments

The author would like to thank Fabio Apruzzi, Philip Argyres, Dnyanesh Kulkarni, Madalena Lemos, Greg Moore, Leonardo Rastelli, Andrea Rocchetto, Sakura Schafer-Nameki and Yuji Tachikawa for helpful conversations, useful suggestions and insightful comments on the draft. The work of the author is supported by NSF grants PHY-1151392 and PHY-1620610.

A Coulomb branch generalities

Because of the unbroken low energy $U(1)^r$ gauge invariance on $\mathcal{C}$, states in the low-energy theory are labeled by a set of $2r$ integral electric and magnetic charges, $Q \in \mathbb{Z}^{2r} = \Lambda$, the *charge lattice*. The Q s satisfy the standard Dirac-Zwanziger-Schwinger quantization condition

$$\langle Q, Q' \rangle \in \mathbb{Z} \quad (\text{A.1})$$

where the Dirac pairing $\langle \cdot, \cdot \rangle$ gives the charge lattice an integral symplectic structure.¹⁶ The matrix of low-energy holomorphic gauge couplings τ_{ij} of the r $U(1)$ factors at a generic point of $\mathcal{C}$ can be extracted by writing the Kahler metric on $\mathcal{C}_{\text{reg}}$ in a special basis $ds^2 = \text{Im}(da_j^D d\bar{a}^j) = \text{Im}(\tau_{ij}) da^i d\bar{a}^j$, where:

$$\tau_{ij} := \frac{\partial a_i^D}{\partial a^j} = \tau_{ji}, \quad \text{and} \quad \text{Im}(\tau_{ij}) \text{ is positive definite.} \quad (\text{A.3})$$

¹⁶The situation is slightly more subtle as in general $\langle Q, Q' \rangle := Q^T \mathbb{D} Q'$ where $\mathbb{D}$ is an integer non-degenerate skew-symmetric $2r \times 2r$ matrix which is called the *polarization* of the charge lattice. Here we will always assume it to have its canonical form:

$$\mathbb{D} = \begin{pmatrix} 0 & \mathbf{1}_r \\ -\mathbf{1}_r & 0 \end{pmatrix} \quad (\text{A.2})$$

The choice of $\mathbb{D}$ in (A.2) is also called *principal*. The physical meaning of alternative choices of $\mathbb{D}$ remains largely unclear.

$\mathbf{a}^D$ and $\mathbf{a}$ are separately good holomorphic coordinates on $\mathcal{C}_{\text{reg}}$ and are the celebrated *special coordinates*:

$$\sigma := \begin{pmatrix} \mathbf{a}^D \\ \mathbf{a} \end{pmatrix}, \quad \text{with} \quad \mathbf{a}^D := \begin{pmatrix} a_1^D \\ \vdots \\ a_r^D \end{pmatrix}, \quad \text{and} \quad \mathbf{a} := \begin{pmatrix} a^1 \\ \vdots \\ a^r \end{pmatrix}. \quad (\text{A.4})$$

A central fact about CB geometry is that there is no globally defined lagrangian description of the low energy $\mathcal{N} = 2$ $\text{U}(1)^r$ gauge theory, and non-trivial monodromies have to be considered to describe the physics on $\mathcal{C}_{\text{reg}}$. While the charge lattice remains the same over all points in $\mathcal{C}_{\text{reg}}$, upon dragging a given $Q \in \Lambda$ along a closed path $\gamma \subset \mathcal{C}_{\text{reg}}$, Q in general suffers a monodromy $Q \xrightarrow{\gamma} Q' = MQ$, where $M \in \text{sp}(2r, \mathbb{Z})$. σ also suffers monodromies around closed loops and it is therefore a holomorphic section of a rank- $2r$ complex $\text{sp}(2r, \mathbb{Z})$ vector bundle over $\mathcal{C}_{\text{reg}}$. This structure makes $\mathcal{C}$ a special Kahler variety. There are other formulation of special Kahler geometry, some of which will be discussed below, for a systematic account see [41].

Finally the complex central charge, Z_Q , of the low energy $\mathcal{N} = 2$ supersymmetry algebra of a vacuum in $\mathcal{C}_{\text{reg}}$ acting on the superselection sector of states with charge $Q \in \Lambda$:

$$Z_Q := Q^T \sigma, \quad (\text{A.5})$$

where T indicates transposition. It follows from the $\mathcal{N} = 2$ supersymmetry algebra that $|Z_Q|$ is a lower bound on the mass of any state with charge Q .

A.1 Special Kahler stratification

The singular locus $\bar{\mathcal{S}}$ plays a central role in our analysis, here we will provide a quick summary of its rich structure, for a more detailed discussion see [35]. It is well-known that the singular locus $\bar{\mathcal{S}}$ is a stratified space with closed subsets of increasingly higher complex co-dimensions. But a careful analysis shows that $\bar{\mathcal{S}}$ inherits an even more constraining structure from the ambient space $\mathcal{C}$, this is called *special Kahler stratification* in [35]. The main idea behind the special Kahler stratification, is that each component of the singular locus $\bar{\mathcal{S}}_i$, starting at complex co-dimension one, inherits a special Kahler structure from $\mathcal{C}$ and can be therefore seen as a scale invariant CB geometry in its own right with its own set of metric singularities. As it is the case for $\mathcal{C}_{\text{reg}}$, the set of regular points of $\bar{\mathcal{S}}_i$ will be an open set which we will indicate simply as $\mathcal{S}_i$ and which we will constitute the *strata* of the stratification. Iterating this procedure we can identify components and corresponding strata of higher complex co-dimensions.

As we discussed throughout the paper, each strata naturally supports a theory which in the text we called $\mathcal{T}_i$. The direction transverse to each $\mathcal{S}_i$ into $\mathcal{C}$ are naturally identified, at least locally, with the CB moduli of the low-energy theory $\mathcal{T}_i$ supported on the stratum. It therefore follows that the rank of $\mathcal{T}_i$ coincides with the complex co-dimension of the strata. Restricted in particular, as we did in the text, to complex co-dimension one strata, we only encounter rank-1 theories. Our notation in this section is a bit sloppy and maybe confusing at times since we are not keeping track of the increasing complex co-dimension

of the strata and the rank of the corresponding theories. We chose to avoid setting up the proper, lengthy, notation since here we don't really use the properties of higher complex co-dimension strata, for a proper discussion see [35].

The $\mathbb{C}^*$ action acts by restriction on all strata which are all separately scale invariant and therefore are all closed under it. Since the stratification works inductively, all properties that apply to complex co-dimension one strata extend to the higher complex co-dimension ones. This stratification is very reminiscent of the stratification of symplectic singularities [42, 43] which applies to Higgs branches of $\mathcal{N} = 2$ SCFTs in four dimensions. The special Kahler stratification is both more constrained and richer. It is more constrained because the complex dimension of the strata jump precisely by one at each step and a full list of allowed elementary slices is known, while an analogous list remains an open question for symplectic singularities. And it is richer because strata supporting $U(1)$ gauge theories with massless hypers and trivial Higgs branch are not necessarily special Kahler and a weaker condition applies [35].

A.2 Accidental $U(1)_R$ near strata

The $U(1)_R$ component of the R -symmetry of the theory at the superconformal vacuum, acts non-trivially on $\mathcal{C}$ and for a generic CB vacuum $\mathbf{u}$ it will be spontaneously broken. An accidental $U(1)_R$ symmetry arises instead along any strata $\mathcal{S}_i$ (and along strata of higher co-dimensions), where this low-energy $U(1)_R$ is identified with the appropriate component of the global symmetry of $\mathcal{T}_i$. Because of that, near each stratum, we can define a scaling action which is different from the globally defined one and which does not act globally on $\mathcal{C}$. This new scaling action scales *into* the stratum and it only really acts on the transverse slice. The global scaling action scales instead into *the origin* of the moduli space.

The accidental $U(1)_R$ symmetry combines with the $\mathbb{R}^+$ of $\mathcal{T}_i$ giving rise to a $\mathbb{C}_i^*$ action which scales away from the stratum $\mathcal{S}_i$. Notice that we did not label this new scaling action as $\mathbb{C}^*|_i$ because $\mathbb{C}_i^*$ is *not* the restriction of $\mathbb{C}^*$. For example $\mathbb{C}_i^*$ is not globally defined and the weight of the coordinates $\mathbf{u}$ under the $\mathbb{C}^*$ and $\mathbb{C}_i^*$ are in general radically different. The scaling dimension Δ_i^{sing} in (2.3) is defined with respect to the global $\mathbb{C}^*$ action while the scaling dimension of the CB of the low-energy theories $\mathcal{T}_i$, Δ_i see (2.14), is defined with respect to $\mathbb{C}_i^*$.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] E. Witten, *Topological Quantum Field Theory*, *Commun. Math. Phys.* **117** (1988) 353 [[INSPIRE](#)].
- [2] E. Witten, *Monopoles and four manifolds*, *Math. Res. Lett.* **1** (1994) 769 [[hep-th/9411102](#)] [[INSPIRE](#)].
- [3] E. Witten, *Supersymmetric Yang-Mills theory on a four manifold*, *J. Math. Phys.* **35** (1994) 5101 [[hep-th/9403195](#)] [[INSPIRE](#)].

- [4] S.K. Donaldson, *The Seiberg-Witten equations and 4-manifold topology*, *Bull. Am. Math. Soc.* **33** (1996) 45.
- [5] M. Dedushenko, S. Gukov and P. Putrov, *Vertex algebras and 4-manifold invariants*, in *Nigel Hitchin's 70th Birthday Conference. Vol. 1*, Oxford U.K. (2017), pg. 249 [[arXiv:1705.01645](#)] [[INSPIRE](#)].
- [6] S. Gukov, *Trisecting non-Lagrangian theories*, *JHEP* **11** (2017) 178 [[arXiv:1707.01515](#)] [[INSPIRE](#)].
- [7] G.W. Moore and I. Nidaiev, *The Partition Function Of Argyres-Douglas Theory On A Four-Manifold*, [arXiv:1711.09257](#) [[INSPIRE](#)].
- [8] N.A. Nekrasov, *Seiberg-Witten prepotential from instanton counting*, *Adv. Theor. Math. Phys.* **7** (2003) 831 [[hep-th/0206161](#)] [[INSPIRE](#)].
- [9] N. Nekrasov and A. Okounkov, *Seiberg-Witten theory and random partitions*, *Prog. Math.* **244** (2006) 525 [[hep-th/0306238](#)] [[INSPIRE](#)].
- [10] A.D. Shapere and Y. Tachikawa, *Central charges of $N = 2$ superconformal field theories in four dimensions*, *JHEP* **09** (2008) 109 [[arXiv:0804.1957](#)] [[INSPIRE](#)].
- [11] V. Pestun, *Localization of gauge theory on a four-sphere and supersymmetric Wilson loops*, *Commun. Math. Phys.* **313** (2012) 71 [[arXiv:0712.2824](#)] [[INSPIRE](#)].
- [12] P. Argyres and M. Martone, *Construction and classification of Coulomb branch geometries*, [arXiv:2003.04954](#) [[INSPIRE](#)].
- [13] P.C. Argyres and M. Martone, *4d $\mathcal{N} = 2$ theories with disconnected gauge groups*, *JHEP* **03** (2017) 145 [[arXiv:1611.08602](#)] [[INSPIRE](#)].
- [14] E. Witten, *On S duality in Abelian gauge theory*, *Selecta Math.* **1** (1995) 383 [[hep-th/9505186](#)] [[INSPIRE](#)].
- [15] A. Losev, N. Nekrasov and S.L. Shatashvili, *Issues in topological gauge theory*, *Nucl. Phys. B* **534** (1998) 549 [[hep-th/9711108](#)] [[INSPIRE](#)].
- [16] G.W. Moore and E. Witten, *Integration over the u plane in Donaldson theory*, *Adv. Theor. Math. Phys.* **1** (1997) 298 [[hep-th/9709193](#)] [[INSPIRE](#)].
- [17] M. Mariño and G.W. Moore, *Integrating over the Coulomb branch in $N = 2$ gauge theory*, *Nucl. Phys. B Proc. Suppl.* **68** (1998) 336 [[hep-th/9712062](#)] [[INSPIRE](#)].
- [18] M. Mariño and G.W. Moore, *The Donaldson-Witten function for gauge groups of rank larger than one*, *Commun. Math. Phys.* **199** (1998) 25 [[hep-th/9802185](#)] [[INSPIRE](#)].
- [19] M. Mariño, G.W. Moore and G. Peradze, *Superconformal invariance and the geography of four manifolds*, *Commun. Math. Phys.* **205** (1999) 691 [[hep-th/9812055](#)] [[INSPIRE](#)].
- [20] P.C. Argyres and M. Martone, *Coulomb branches with complex singularities*, *JHEP* **06** (2018) 045 [[arXiv:1804.03152](#)] [[INSPIRE](#)].
- [21] O. Aharony, N. Seiberg and Y. Tachikawa, *Reading between the lines of four-dimensional gauge theories*, *JHEP* **08** (2013) 115 [[arXiv:1305.0318](#)] [[INSPIRE](#)].
- [22] N. Seiberg and E. Witten, *Electric-magnetic duality, monopole condensation, and confinement in $N = 2$ supersymmetric Yang-Mills theory*, *Nucl. Phys. B* **426** (1994) 19 [*Erratum ibid.* **430** (1994) 485] [[hep-th/9407087](#)] [[INSPIRE](#)].
- [23] N. Seiberg and E. Witten, *Monopoles, duality and chiral symmetry breaking in $N = 2$ supersymmetric QCD*, *Nucl. Phys. B* **431** (1994) 484 [[hep-th/9408099](#)] [[INSPIRE](#)].
- [24] L. Álvarez-Gaumé and S.F. Hassan, *Introduction to S duality in $N = 2$ supersymmetric gauge theories: A Pedagogical review of the work of Seiberg and Witten*, *Fortsch. Phys.* **45** (1997) 159 [[hep-th/9701069](#)] [[INSPIRE](#)].

- [25] W. Lerche, *Introduction to Seiberg-Witten theory and its stringy origin*, *Nucl. Phys. (Proc. Suppl.)* **55** (1997) 83 [[hep-th/9611190](#)] [[INSPIRE](#)].
- [26] M. Martone, *The constraining power of Coulomb Branch Geometry: lectures on Seiberg-Witten theory*, in *Young Researchers Integrability School and Workshop 2020: A modern primer for superconformal field theories*, Hamburg Germany (2020) [[arXiv:2006.14038](#)] [[INSPIRE](#)].
- [27] P.C. Argyres, M. Plesser and A.D. Shapere, *The Coulomb phase of $N = 2$ supersymmetric QCD*, *Phys. Rev. Lett.* **75** (1995) 1699 [[hep-th/9505100](#)] [[INSPIRE](#)].
- [28] K. Landsteiner, E. Lopez and D.A. Lowe, *$N = 2$ supersymmetric gauge theories, branes and orientifolds*, *Nucl. Phys. B* **507** (1997) 197 [[hep-th/9705199](#)] [[INSPIRE](#)].
- [29] J. Labastida and M. Marino, *Mathematical Physics Studies. Vol. 25: Topological quantum field theory and four manifolds*, Springer, Dordrecht The Netherlands (2005).
- [30] G.W. Moore, *Lectures On The Physical Approach To Donaldson And Seiberg-Witten Invariants Of Four-Manifolds*, <http://www.physics.rutgers.edu/~gmoore/SCGP-FourManifoldsNotes-2017.pdf>.
- [31] G.W. Moore and J. Manschot, to appear.
- [32] V. Rohlin, *New results in the theory of 4-dimensional manifolds*, *Dokl. Akad. Nauk. SSSR* **84** (1952) 221.
- [33] P. Teichner, *On the signature of four-manifolds with universal covering spin*, *Math. Ann.* **295** (1993) 745 [[INSPIRE](#)].
- [34] P. Argyres, M. Lotito, Y. Lü and M. Martone, *Geometric constraints on the space of $\mathcal{N} = 2$ SCFTs. Part III. Enhanced Coulomb branches and central charges*, *JHEP* **02** (2018) 003 [[arXiv:1609.04404](#)] [[INSPIRE](#)].
- [35] P.C. Argyres and M. Martone, *Towards a classification of rank r $\mathcal{N} = 2$ SCFTs Part II. Special Kähler stratification of the Coulomb branch*, *JHEP* **12** (2020) 022 [[arXiv:2007.00012](#)] [[INSPIRE](#)].
- [36] P. Argyres, M. Lotito, Y. Lü and M. Martone, *Geometric constraints on the space of $\mathcal{N} = 2$ SCFTs. Part I. Physical constraints on relevant deformations*, *JHEP* **02** (2018) 001 [[arXiv:1505.04814](#)] [[INSPIRE](#)].
- [37] P.C. Argyres, M. Lotito, Y. Lü and M. Martone, *Geometric constraints on the space of $\mathcal{N} = 2$ SCFTs. Part II. Construction of special Kähler geometries and RG flows*, *JHEP* **02** (2018) 002 [[arXiv:1601.00011](#)] [[INSPIRE](#)].
- [38] P.C. Argyres, M. Lotito, Y. Lü and M. Martone, *Expanding the landscape of $\mathcal{N} = 2$ rank 1 SCFTs*, *JHEP* **05** (2016) 088 [[arXiv:1602.02764](#)] [[INSPIRE](#)].
- [39] T. Eguchi, K. Hori, K. Ito and S.-K. Yang, *Study of $N = 2$ superconformal field theories in four-dimensions*, *Nucl. Phys. B* **471** (1996) 430 [[hep-th/9603002](#)] [[INSPIRE](#)].
- [40] T. Eguchi and K. Hori, *$N = 2$ superconformal field theories in four-dimensions and A-D-E classification*, in *Conference on the Mathematical Beauty of Physics (In Memory of C. Itzykson)*, Saclay France (1996), pg. 67 [[hep-th/9607125](#)] [[INSPIRE](#)].
- [41] D.S. Freed, *Special Kähler manifolds*, *Commun. Math. Phys.* **203** (1999) 31 [[hep-th/9712042](#)] [[INSPIRE](#)].
- [42] A. Beauville, *Symplectic singularities*, [[math/9903070](#)].
- [43] D. Kaledin, *Symplectic singularities from the Poisson point of view*, *J. Reine Angew. Math.* **2006** (2006) 135 [[math/0310186](#)].