

RECEIVED: June 26, 2023

REVISED: October 25, 2023

ACCEPTED: November 22, 2023

PUBLISHED: December 4, 2023

D3-branes and gaugino condensation

Iosif Bena,^a Emilian Dudaș,^b Mariana Graña,^a Gabriele Lo Monaco^c and
Dimitrios Toulikas^a

^a*Institut de Physique Théorique, Université Paris Saclay, CEA, CNRS,
Orme des Merisiers, 91191 Gif-sur-Yvette CEDEX, France*

^b*CPHT, CNRS, Ecole Polytechnique, IP Paris,
F-91128 Palaiseau, France*

^c*Università degli Studi di Palermo, Dipartimento di Fisica e Chimica — Emilio Segrè,
via Archirafi 36, I-90123 Palermo, Italy*

E-mail: iosif.bena@ipht.fr, mariana.grana@ipht.fr,
gabriele.lomonaco@ipht.fr, emilian.dudas@polytechnique.edu,
dimitrios.toulikas@ipht.fr

ABSTRACT: Anti-D3 branes at the bottom of warped throats, commonly used to uplift the cosmological constant in String-Theory de Sitter proposals, source a plethora of supersymmetry-breaking fluxes, that can interact nontrivially with other ingredients of the flux compactification. In this paper we perform a complex-structure decomposition of these fluxes, and compute the effect of the (0,3) flux component on the stabilization of Kähler moduli via D7-branes gaugino condensation. This allows us to obtain a new constraint on the validity of this stabilization mechanism. This effect does not appear hard to satisfy in de Sitter construction proposals that use long warped throats, but may be problematic in proposals where the warping is small.

KEYWORDS: Flux Compactifications, String and Brane Phenomenology, Confinement, Supersymmetry Breaking

ARXIV EPRINT: [2211.14381](https://arxiv.org/abs/2211.14381)

Contents

1	Introduction	1
2	$\overline{\text{D3}}$formed KS geometry	4
2.1	Review of the KS solution and its non-supersymmetric deformations	4
2.2	Adding $\overline{\text{D3}}$ -branes	6
3	The flux zoo	7
4	Fermion masses	9
4.1	Gaugino mass and gaugino condensation	11

1 Introduction

Many proposals to construct de Sitter vacua in String Theory involve uplifting the negative cosmological constant that one typically obtains in flux compactifications after fixing the moduli. The original and most popular uplifting ingredient are anti-D3 branes placed in long warped throats within the compactification manifold [1]. They are argued to give a tunably small uplift energy, which would make the cosmological constant positive without perturbing the stabilization of the other moduli.

The prototypical example of such a warped throat is the Klebanov-Strassler (KS) geometry, obtained by adding fluxes to the deformed conifold [2]. There is by now an extensive body of work investigating the physics of anti-D3 branes in the KS geometry, in several regimes of parameters. Many of the results of these calculation point towards the existence of pathologies and instabilities when multiple anti-D3 branes are placed at the bottom of the KS geometry [3–7]. Furthermore, if the flux on the three-cycle of the deformed conifold is not large, even a single anti-D3 brane appears to have the power to cause a runaway behavior in the conifold deformation modulus [7–10], collapsing the whole KS geometry into the singular Klebanov-Tseytlin one [11] and annihilating against the singularity.¹ A similar behavior is found when considering black holes in the KS geometry [13, 14]. Hence, the only corner of parameter space where KS antibranes have any chance of being metastable is when there is a single antibrane and the flux on the three-cycle of the deformed conifold is large.

However, even a single anti-brane has a significant back-reaction, and is capable of significantly affecting the other ingredients of the de Sitter construction. It is the purpose of this paper to calculate the strength the interactions between this single antibrane at the bottom of a KS solution and other ingredients that are used in the KKLT proposal [1] to construct de Sitter vacua with stabilized moduli in String Theory.

¹For a recent argument against this scenario see [12].

If we start from a supersymmetric solution in which a KS throat is glued to a CY geometry, the complex structure of the KS throat will match the complex structure of the bulk compact CY. Hence, the fluxes sourced by the anti-D3 branes can be decomposed according to the complex-structure of the KS throat and, as expected, have all possible components. The values of the field strength fluxes at the top of the KS geometry, where it is glued to the compactification manifold, are expected to be of the same order as the values of these fluxes in the rest of this manifold, away from the throat, and they determine the magnitude of the effect of antibranes on the other ingredients needed in de Sitter construction proposals.

As we have shown in a previous paper [15], the $(0, 3)$ component of the fluxes sourced by the anti-D3 branes gives rise to a nontrivial constant term in the superpotential. This avoids the need of one of the ingredients of the KKLT proposal, namely a finite but very small constant term in the superpotential, introduced ad hoc by turning on $(0, 3)$ fluxes. Hence, taking the interactions between antibranes and the rest of the flux-compactification ingredients fully into account can result in simplified “bare-bones” de Sitter proposals [15], which use one less ingredient than KKLT.

Another possible consequence of the anti-brane fluxes is to affect gaugino condensation on D7 branes wrapping certain divisors of the CY geometry. In the absence of D7 branes, the volumes of these divisors (which correspond to Kähler moduli of the compactification) are flat directions. The low-energy physics of the D7 branes wrapping a holomorphic divisor is an $\mathcal{N} = 1$ Super-Yang-Mills theory, which confines in the infrared. The non-perturbative Affleck-Dine-Seiberg superpotential [16] of this confining theory depends nontrivially on the Kähler moduli, and gives rise to a term in the potential that is responsible for stabilizing the size of the divisors [1].

The fields sourced by the anti-D3 brane can in principle interfere with this sensitive mechanism of Kähler-moduli stabilization. Indeed, as shown in [17–19], both the $(1, 2)$ and the $(0, 3)$ components of the complex three-form field strength give rise to mass terms for the fermions on the D7 brane. These mass terms break the supersymmetry of the $\mathcal{N} = 1$ theory on the D7 branes, and can modify the RG flow and the gaugino-condensation scale. More precisely, they can affect the Affleck-Dine-Seiberg superpotential and can introduce extra terms in the potential for the Kähler moduli, potentially ruining their stabilization.

In this paper we use the expression of the supersymmetry-breaking fluxes sourced by the anti-D3 branes [3, 20] to calculate precisely their effect on the stabilization of Kähler moduli via D7-brane gaugino condensation. We obtain a new bound relating various parameters of compactification with long warped throats, which must be satisfied in order to be able to stabilize Kähler moduli via D7-brane gaugino condensation. We combine this bound with other constraints appearing in vanilla-type KKLT compactifications, and find that it is generically satisfied. We leave the exploration of the importance of this bound on de Sitter proposals that do involve large warping² to future exploration.

²Such as the LVS proposals in [21, 22]. Note though that for large warping both KKLT and LVS were argued to suffer from control issues, the former by the so-called singular bulk problem [23], and the latter by large α' corrections [24]. Here we are assuming a constant warp factor in the bulk, such that the singular bulk problem does not arise.

Beside their effect on the gaugino condensation on D7 branes, the extra fluxes and fields sourced by the antibranes can also affect the number of fermion zero modes on the D3 instantons that also give rise to non-perturbative terms in the superpotential [25] that can stabilize other Kähler moduli, possibly switching off or on these terms. This is a subtle effect [26–34], which potentially can be more drastic than the effect on D7 gaugino condensation: even the tiniest supersymmetry-breaking fluxes sourced by the antibranes are enough to uplift fermion zero modes and change completely the Kähler moduli stabilisation process.³ We leave its investigation for future work.

As a byproduct of our calculation, we decode some aspects of the holographic dictionary corresponding to anti-D3 branes in the Klebanov-Strassler geometry. If a single anti-D3 brane in the KS geometry with large three-form flux on the S^3 of the deformed conifold is indeed metastable, the resulting solution would be holographically dual to a metastable vacuum of the quiver gauge theory dual to the KS geometry. The structure of vacua of this theory is quite rich [35], and the existence of a metastable vacuum at strong coupling would be a nontrivial prediction of holography. It would be very interesting to try construct this putative vacuum directly in field theory, using for example ISS methods [36] (see [37] for the dual of a similar supersymmetry-breaking vacuum). The fact that this vacuum can only exist in a very restricted region of parameter space may also explain why earlier attempts at finding it have not been successful [38].

Our analysis gives several clear holographic indications as to what the physics of this vacuum is. In particular, the fall-offs of the three-form fluxes with the radius can be used to show that certain dimension-three operators corresponding to fermion bilinears and certain dimension-seven operators corresponding to fermion bilinears multiplied by $F_{\mu\nu}F^{\mu\nu}$ acquire non-trivial vacuum expectation values. Furthermore, the $(1, 2)$ fluxes sourced by the antibranes fall off as $1/r^4$, and hence give rise to a nontrivial vacuum expectation value of a dimension-4 operator corresponding to a marginal deformation of the superpotential [39]. Since the dictionary between the bulk three-form fields and the fermion bilinears is well understood [40, 41], we believe this information will be useful in searching for the holographic dual of the putative KS metastable vacuum.

The paper is organized as follows: in section 2 we review the Klebanov-Strassler geometry, as well as the most general deformation preserving its $SU(2) \times SU(2) \times \mathbb{Z}_2$ symmetry and describe how the solution corresponding to anti-D3 branes smeared at the tip of the throat is obtained. In section 3 we write the analytic expression, as well as the UV expansion, of all G_3 flux components that are generated by the addition of the anti-D3 branes and comment on their holographic duals. In section 4 we compute the D7-brane gaugino mass induced by the $G_{(0,3)}$ flux sourced by the anti-D3 branes and compare it with a four-dimensional supergravity description of supersymmetry-breaking gaugino masses finding parametric agreement. Moreover, we derive a bound that all KKLT-like constructions should satisfy in order for the Kähler moduli stabilization via gaugino condensation to work, and argue that it is easily satisfied by the existing constructions.

³Fluxes generically uplift some of the zero modes, changing the arithmetic genus of the divisor wrapped by the instanton, generating (turning off) a superpotential on a cycle that would otherwise forbid (allow) it.

2 $\overline{\text{D3}}$ formed KS geometry

2.1 Review of the KS solution and its non-supersymmetric deformations

A long warped Klebanov-Strassler-like throat, at the bottom of which anti-D3 branes (denoted as $\overline{\text{D3}}$ -branes in the following) can sit, is a key element in the KKLT proposal for constructing de Sitter vacua in String Theory [1]. In this section we review the supersymmetric Klebanov-Strassler (KS) geometry [2], as well as the ansatz that describes the most general deformation (with vanishing RR axion C_0) that preserves its $\text{SU}(2) \times \text{SU}(2) \times \mathbb{Z}_2$ symmetry [42]. The ten-dimensional spacetime consists of a warped product of four-dimensional Minkowski space and the deformed conifold:

$$ds_{10}^2 = e^{2A+2W-X} ds_{1,3}^2 + e^{-6W-X} d\tau^2 + e^{X+Y} (g_1^2 + g_2^2) + e^{X-Y} (g_3^2 + g_4^2) + e^{-6W-X} g_5^2, \quad (2.1)$$

where $\{A, W, X, Y\}$ are functions of the radial coordinate, τ , and the one-forms g_i are:

$$\begin{aligned} g_1 &= -\frac{1}{\sqrt{2}} \text{Im}(w_1 + w_2), & g_2 &= \frac{1}{\sqrt{2}} \text{Re}(w_1 - w_2), \\ g_3 &= -\frac{1}{\sqrt{2}} \text{Im}(w_1 - w_2), & g_4 &= \frac{1}{\sqrt{2}} \text{Re}(w_1 + w_2), \\ g_5 &= d\psi + \sum_{i=1}^2 \cos \theta_i d\phi_i, & g_6 &= d\tau, \end{aligned} \quad (2.2)$$

with $0 \leq \psi \leq 4\pi$, $0 \leq \theta_i \leq \pi$, $0 \leq \phi_i \leq 2\pi$. Here we introduced the forms $w_1 \equiv d\theta_1 + i \sin \theta_1 d\phi_1$ and $e^{i\psi} w_2 \equiv d\theta_2 + i \sin \theta_2 d\phi_2$, with \mathbb{Z}_2 exchanging the two S^2 's defined by w_i [42].

The NSNS and RR forms, H_3 , F_3 and F_5 , are all non-vanishing and their form is fixed by Bianchi identities and isometries:

$$\begin{aligned} H_3 &= \frac{1}{2}(k-f)g_5 \wedge (g_1 \wedge g_3 + g_2 \wedge g_4) + d\tau \wedge (f' g_1 \wedge g_2 + k' g_3 \wedge g_4), \\ F_3 &= F g_1 \wedge g_2 \wedge g_5 + (2P-F) g_3 \wedge g_4 \wedge g_5 + F' d\tau \wedge (g_1 \wedge g_3 + g_2 \wedge g_4), \\ F_5 &= \left[\frac{\pi Q}{4} + (k-f)F + 2P f \right] (1 + \star_{10}) g_1 \wedge g_2 \wedge g_3 \wedge g_4 \wedge g_5, \\ \Phi &= \Phi(\tau), \quad C_0 = 0 \end{aligned} \quad (2.3)$$

with Q a constant and $\{k, f, F\}$ functions of τ .

The radial dependence of the functions appearing in the KS metric is:

$$\begin{aligned} e^{X_{\text{KS}}} &= \frac{1}{4} h(\tau)^{1/2} \left(\frac{1}{2} \sinh(2\tau) - \tau \right)^{1/3}, & e^{6A_{\text{KS}}} &= S^2 \frac{\sinh(\tau)^2}{3 \cdot 2^5} e^{2X_{\text{KS}}}, \\ e^{6W_{\text{KS}}} &= \frac{24}{h(\tau) \sinh(\tau)^2} \left(\frac{1}{2} \sinh(2\tau) - \tau \right)^{1/3}, & e^{Y_{\text{KS}}} &= \tanh(\tau/2), \end{aligned} \quad (2.4)$$

where S is a complex-structure modulus and $h(\tau)$ is the solution that vanishes at infinity to the following differential equation:

$$\frac{dh}{d\tau} = 32 P^2 g_s \frac{\tau \coth \tau - 1}{\sinh(\tau)^2} \left(\frac{1}{2} \sinh(2\tau) - \tau \right)^{1/3}, \quad (2.5)$$

with $P = \frac{1}{4}\alpha' M$ and M the number of units of F_3 flux on the large S^3 of the warped deformed conifold (also known as the compact A -cycle). To simplify the expressions, it is useful to define the constants $h_0 \equiv h(\tau=0)$ and $\mathcal{I}_0 \equiv \frac{h_0}{32P^2 g_s}$.

Moreover, the functions $\{f, k, F\}$ appearing in the fluxes are given by:

$$\begin{aligned} f_{\text{KS}} &= -g_s P \frac{(\tau \coth(\tau) - 1)(\cosh(\tau) - 1)}{\sinh(\tau)}, \\ k_{\text{KS}} &= -g_s P \frac{(\tau \coth(\tau) - 1)(\cosh(\tau) + 1)}{\sinh(\tau)}, \\ F_{\text{KS}} &= P \frac{\sinh(\tau) - \tau}{\sinh(\tau)}, \end{aligned} \tag{2.6}$$

and, when there are not mobile D3 branes, Q is zero.

It is worth noting that in the KS solution the complexified three-form, G_3 , satisfies the ISD condition, $(i + \star_6)G_3 = 0$. Furthermore, G_3 has only $(2, 1)$ components with respect to the choice of holomorphic vielbeins picked by supersymmetry:

$$\begin{aligned} h_1 &= E_1 + i(\cos \omega E_2 + \sin \omega E_4), \\ h_2 &= E_3 + i(\sin \omega E_2 - \cos \omega E_4), \\ h_3 &= E_5 + i E_6, \end{aligned} \tag{2.7}$$

where $\sin \omega = -\tanh Y$ and:

$$\begin{aligned} E_1 &= \frac{e^{X/2}}{\sqrt{2 \cosh(Y)}}(g_2 + g_4), & E_2 &= \frac{e^{X/2}}{\sqrt{2 \cosh(Y)}}(g_1 + g_3), \\ E_3 &= \frac{e^{X/2} \sqrt{\cosh(Y)}}{\sqrt{2}}(g_4 - g_2 - (g_2 + g_4) \tanh Y), \\ E_4 &= \frac{e^{X/2} \sqrt{\cosh(Y)}}{\sqrt{2}}(g_3 - g_1 - (g_1 + g_3) \tanh Y), \\ E_5 &= e^{-X/2 - 3W} d\tau, & E_6 &= e^{-X/2 - 3W} g_5. \end{aligned} \tag{2.8}$$

We will denote the holomorphic 3-form on the *warped* geometry as $\Omega \equiv n h_1 \wedge h_2 \wedge h_3$, where the normalization constant, n , is such that Ω^{KS} , the unwarped Klebanov-Strassler 3-form, satisfies:

$$\int_A \Omega^{\text{KS}} = \int_A H^{-3/4} \Omega = S \quad \Rightarrow \quad n = -\frac{\sqrt{6}}{4\pi^2}, \tag{2.9}$$

where $H = e^{-4A - 4W + 2x}$ is the warp factor.⁴ The explicit form of the normalized Ω^{KS} in terms of the set of $\{g_i\}$ is the following:

$$\Omega^{\text{KS}} = -\frac{S}{16\pi^2} \sinh \tau \left(\tanh\left(\frac{\tau}{2}\right) g_1 \wedge g_2 - \coth\left(\frac{\tau}{2}\right) g_3 \wedge g_4 + i g_1 \wedge g_3 + i g_2 \wedge g_4 \right) \wedge (g_5 - i g_6). \tag{2.10}$$

Note that with this normalization constant, $\Omega \wedge \bar{\Omega} = \frac{3i}{\pi^4} \text{vol}_6$, instead of $\Omega \wedge \bar{\Omega} = 8i \text{vol}_6$, which is the usual convention in the literature.⁵

⁴The KS metric (2.1) can be written in terms of H as $ds^2 = H^{-1/2} ds_4^2 + H^{1/2} ds_6^2$.

⁵Remember that vol_6 is the volume of the six-dimensional internal space including the warp factor.

2.2 Adding $\overline{\text{D3}}$ -branes

Anti-D3 branes have a charge that is opposite to that of the Klebanov-Strassler geometry and, in the probe approximation, fall to the S^3 tip of the throat. When the antibranes are localized on the S^3 , the fields they source have a complicated dependence on τ and the S^3 coordinates, and are hard to compute analytically. However, if we are interested in the solution away from the tip, one can assume the antibranes to be smeared, and then the solution will have $\text{SU}(2) \times \text{SU}(2) \times \mathbb{Z}_2$ symmetry. Its metric and fluxes will then be described by the eight functions of τ appearing in the Papadopoulos-Tseytlin ansatz (2.1)–(2.3):

$$\{\Phi^i\} = \{X - 2W - 5A, Y, X + 3W, X - 2W - 2A, f, k, F, \Phi\}, \quad (2.11)$$

where we performed the above redefinition for convenience.

When the number of the anti-D3 branes, \overline{N} , is small one can describe their solution as a small perturbation around the KS geometry [3, 20, 43, 44]:

$$\Phi^i = \Phi_{\text{KS}}^i + \varepsilon \varphi^i + O(\varepsilon^2) \quad (2.12)$$

where ε is an expansion parameter that can be taken to be:

$$\varepsilon = \frac{\overline{N}}{g_s M^2}. \quad (2.13)$$

We then require the equations of motion of type-IIB supergravity to be satisfied at leading order in ε . Plugging the particular ansatz (2.1)–(2.3) in the type IIB supergravity action, one gets an action for the fields $\{\Phi^i\}$ that can be cast in the following form [42]:

$$\mathcal{L} = -\frac{1}{2} G_{ij} \left(\frac{d\Phi^i}{d\tau} - \frac{1}{2} G^{ik} \frac{\partial W}{\partial \Phi^k} \right) \left(\frac{d\Phi^j}{d\tau} - \frac{1}{2} G^{jl} \frac{\partial W}{\partial \Phi^l} \right) - \frac{1}{2} \frac{\partial W}{\partial \tau}, \quad (2.14)$$

where G and W are respectively a Φ -dependent metric and superpotential whose exact functional dependence is not relevant in the following. To study perturbations around a supersymmetric solution governed by this action [45] it is useful to introduce the set of functions, $\{\xi^i\}$, conjugate to the perturbations, $\{\varphi^i\}$. They are defined as:

$$\xi_i \equiv G_{ij}(\Phi_{\text{KS}}) \left(\frac{d\varphi^j}{d\tau} - M^j_k(\Phi_{\text{KS}}) \varphi^k \right), \quad M^j_k \equiv \frac{1}{2} \frac{\partial}{\partial \Phi^k} \left(G^{jl} \frac{\partial W}{\partial \Phi^l} \right). \quad (2.15)$$

The supersymmetric KS background corresponds to $\xi_i = 0$ for all i . Our goal is to find other solutions to the equations of motion:⁶

$$\begin{aligned} \frac{d\xi_i}{d\tau} + \xi_j M^j_i(\Phi_{\text{KS}}) &= 0, \\ \frac{d\varphi^i}{d\tau} - M^i_j(\Phi_{\text{KS}}) \varphi^j &= G^{ij} \xi_j. \end{aligned} \quad (2.16)$$

An analytical form for the most general perturbation can be found and it involves various nested integrals [20]. Nevertheless, such integrals can be evaluated both as series

⁶These equations are actually supplemented by the zero-energy condition $\xi_i \frac{d\Phi_{\text{KS}}^i}{d\tau} = 0$.

expansions in the UV or IR limits, or numerically throughout the whole solution. The numerical evaluation allows one to match the parameters of the UV and IR expansions. The simplest example of this matching is the evaluation of the “momentum” ξ_1 , which controls the force on a probe D3-brane:

$$F_{\text{D3}} = \frac{2\varepsilon}{3} e^{-2X_{\text{KS}}} \xi_1, \quad \xi_1 = X_1 h(\tau), \quad (2.17)$$

where $X_1 = \frac{3\pi}{8\mathcal{L}_0} \frac{S^{4/3}}{h_0}$ is an integration constant. As we have already observed, the function $h(\tau)$ can be expressed as a definite integral, and its value at an arbitrary τ can only be evaluated numerically. However, its asymptotic IR and UV expansions can be evaluated straightforwardly:

$$h_{\text{IR}} = h_0 - \frac{16}{3} \left(\frac{2}{3}\right)^{\frac{1}{3}} g_s P^2 \tau^2 + O(\tau^4), \quad h_{\text{UV}} = 12 \times 2^{\frac{1}{3}} g_s P^2 (4\tau - 1) e^{-4\tau/3} + O(e^{-10\tau/3}). \quad (2.18)$$

The expressions for the other perturbations and momenta are considerably more involved and can be found in [3].

3 The flux zoo

When the KS throat is deformed because of the presence of anti-D3 branes, the G_3 flux takes the most general form:

$$G_3 = G_{(3,0)} + G_{(0,3)} + G_{(1,2)} + G_{(2,1)}. \quad (3.1)$$

The first three components, which break supersymmetry, are sourced by the anti-D3 branes and appear at $O(\varepsilon)$. This happens because the fluxes H_3 and F_3 , defined in (2.3), do not combine anymore into a (2, 1) form (with respect to the complex structure (2.7) of the zeroth order KS solution) for a generic choice of the functions Φ^i .

Using the complex structure (2.7), we can extract the various components of G_3 . For example, the (0, 3) component of G_3 for an arbitrary set of Φ^i is:

$$G_{(0,3)} = -\frac{1}{8} \gamma_{(0,3)} \bar{h}_1 \wedge \bar{h}_2 \wedge \bar{h}_3, \quad (3.2)$$

$$\gamma_{(0,3)} = e^{3W - \phi - X/2} \left(2e^\phi \left(e^Y P - \cosh Y F - F' \right) + e^Y k' - e^{-Y} f' + k - f \right).$$

This expression holds at any order in ε . The function $\gamma_{(0,3)}$ vanishes in the KS limit, while its first-order term in the ε expansion is:

$$\gamma_{(0,3)} = -\frac{8\sqrt{6} g_s^{-1}}{\frac{3}{h^4} \sinh(\tau)^2} \partial_\tau \left(g_s \sinh \tau \varphi_7 + \cosh\left(\frac{\tau}{2}\right)^2 \varphi_5 - \sinh\left(\frac{\tau}{2}\right)^2 \varphi_6 \right) + O(\varepsilon^2). \quad (3.3)$$

Let us stress that in (3.2) the vielbeins, \bar{h}_i , can be taken to be the Klebanov-Strassler ones, since any correction to them would come at order $O(\varepsilon^2)$. Thus, the first correction to $G_{(0,3)}$ is:

$$G_{(0,3)} = \frac{\pi^2}{2\sqrt{6}} \gamma_{(0,3)} \bar{\Omega}^{\text{KS}} + O(\varepsilon^2), \quad (3.4)$$

with $\gamma_{(0,3)}$ given by (3.3) and Ω^{KS} normalized as in (2.9). Following the same procedure, we obtain the expressions at order $O(\varepsilon)$ for the other components of G_3 . The (3,0) component is given by:

$$G_{(3,0)} = -\frac{8\pi^2 h^{1/4}}{S^{4/3} \sinh(\tau)^2} \partial_\tau (\sinh \tau \xi_6 - \tau \xi_5) \Omega^{\text{KS}} + O(\varepsilon^2), \quad (3.5)$$

and the (1,2) component by:

$$\begin{aligned} G_{(1,2)} = & -\frac{2\sqrt{6}h^{1/4}}{\sinh(\tau)^2} \left[\left(\xi_5 - \text{sech}(\tau)\xi_6 \right) \left(h_1 \wedge \bar{h}_2 \wedge \bar{h}_3 + h_2 \wedge \bar{h}_1 \wedge \bar{h}_3 \right. \right. \\ & \left. \left. + \sinh(\tau) \left(h_1 \wedge \bar{h}_1 \wedge \bar{h}_3 - h_2 \wedge \bar{h}_2 \wedge \bar{h}_3 \right) \right) + (-\xi_5 + \tau \partial_\tau \xi_5 \right. \\ & \left. + \cosh(\tau)\xi_6 - \sinh(\tau)\partial_\tau \xi_6 \right) h_3 \wedge \bar{h}_1 \wedge \bar{h}_2 \left. \right] + O(\varepsilon^2). \end{aligned} \quad (3.6)$$

Finally, as we have already noted, the (2,1) component is the only one that has a non-vanishing term at zeroth order in ε , given by:

$$\begin{aligned} G_{(2,1)}^0 = & \frac{2\sqrt{6}}{h^{3/4}} P \text{csch}(\tau) \text{sech}(\tau) \left(2 \coth(\tau) (-1 + \tau \coth(\tau)) h_1 \wedge h_2 \wedge \bar{h}_3 + (\cosh(\tau) \right. \\ & \left. - \tau \text{csch}(\tau)) (h_1 \wedge h_3 \wedge \bar{h}_1 + \text{csch}(\tau) (h_1 \wedge h_3 \wedge \bar{h}_2 + h_2 \wedge h_3 \wedge \bar{h}_1) - h_2 \wedge h_3 \wedge \bar{h}_2) \right), \end{aligned} \quad (3.7)$$

whereas, at first-order in ε , the non-vanishing $G_{(2,1)}$ components, $G_{ijk}^1 h_i \wedge h_j \wedge \bar{h}_k$, are:

$$\begin{aligned} G_{12\bar{3}}^1 = & \frac{\sqrt{3} \text{csch}(\tau)}{5\sqrt{2}g_s h^{3/4}} \left(5\varphi_5 + \text{csch}(\tau) (4g_s P (-1 + \tau \coth(\tau)) (6\varphi_1 + 4\varphi_3 - 5(3\varphi_4 + \varphi_8)) \right. \\ & \left. - 5(\varphi'_5 + \varphi'_6)) - 5(\varphi_6 + \coth(\tau) (2g_s \varphi_7 + \varphi'_5 - \varphi'_6) - 2g_s \varphi'_7) \right), \end{aligned} \quad (3.8)$$

$$\begin{aligned} G_{13\bar{1}}^1 = & -G_{23\bar{2}} = \frac{\sqrt{6} \text{csch}(\tau)}{5h^{3/4} S^{4/3}} \left(10h(\xi_5 - \xi_6 \text{sech}(\tau)) \right. \\ & \left. + S^{4/3} (P (6(1 - \tau \text{csch}(\tau)) \text{sech}(\tau)) \varphi_1 \right. \\ & \left. + 10\tau \text{sech}(\tau) \tanh(\tau) \varphi_2 - (-1 + \tau \text{csch}(\tau)) \text{sech}(\tau) (4\varphi_3 - 15\varphi_4) + 10\text{sech}(\tau) \varphi_7) \right), \end{aligned} \quad (3.9)$$

$$\begin{aligned} G_{13\bar{2}}^1 = & G_{23\bar{1}} = \frac{2\sqrt{6} \text{csch}(\tau) \text{csch}(2\tau)}{5h^{3/4} S^{4/3}} \left(10h(\xi_6 - \xi_5 \cosh(\tau)) \right. \\ & \left. + S^{4/3} (-6P(\cosh(\tau) - \tau \text{csch}(\tau)) \varphi_1(\tau) + 5P(1 + \cosh(2\tau) - 4\tau \coth(2\tau)) \varphi_2(\tau) \right. \\ & \left. - P(\cosh(\tau) - \tau \text{csch}(\tau)) (4\varphi_3(\tau) - 15\varphi_4(\tau)) - 10\varphi_7(\tau) \right). \end{aligned} \quad (3.10)$$

Note that the (2,1) flux remains primitive at first order: $G_{(2,1)}^1 \wedge J = 0$, where the Kähler form is:

$$J = \frac{1}{2} i \left(h_1 \wedge \bar{h}_1 + h_2 \wedge \bar{h}_2 + h_3 \wedge \bar{h}_3 \right). \quad (3.11)$$

The behavior in the UV of all components of G_3 can be obtained using the UV expansion of the functions φ_i and ξ_i of [4]. The (0, 3) component is given by:

$$G_{(0,3)}^{\text{UV}} = \left(\frac{2}{3}\right)^{3/4} \frac{81\pi\varepsilon}{5\mathcal{I}_0} \frac{1}{\sqrt{\alpha'M}g_s^{3/4}} \log(r_{\text{UV}}^3/|S|)^{5/4} \left(\frac{|S|}{r_{\text{UV}}^3}\right)^{7/3} \bar{h}_1 \wedge \bar{h}_2 \wedge \bar{h}_3 + O(\varepsilon^2), \quad (3.12)$$

where we introduced a new radial coordinate, r :

$$r \equiv \frac{3^{1/2}}{2^{5/6}} |S|^{1/3} e^{\tau/3}, \quad (3.13)$$

such that for large values of τ , the six-dimensional metric of the deformed conifold approaches the conifold metric: $dr^2 + r^2 ds_{T^{11}}^2$.

Note that $G_{(0,3)}$ falls down as r^{-7} in the UV and is dual to an operator of dimension $\Delta = 7$. Such operator is a fermion bilinear of the (schematic) form $F_{\mu\nu} F^{\mu\nu} \lambda\lambda$ where λ is the gaugino and $F_{\mu\nu}$ the field strength.

In the same way, we can compute the asymptotic behavior of the (3, 0) component of G_3 :

$$G_{(3,0)}^{\text{UV}} = \left(\frac{3}{2}\right)^{29/4} \frac{2\pi\varepsilon}{5\mathcal{I}_0^2} \frac{1}{\sqrt{\alpha'M}g_s^{3/4}} \log(r_{\text{UV}}^3/|S|)^{5/4} \left(\frac{|S|}{r_{\text{UV}}^3}\right)^{11/3} h_1 \wedge h_2 \wedge h_3 + O(\varepsilon^2). \quad (3.14)$$

The asymptotic decay of $G_{(3,0)}$ indicates that it is dual to an operator of dimension $\Delta = 11$ that is again a combination of a fermion bilinear and the field strength of the gauge field of the (schematic) form $(F_{\mu\nu} F^{\mu\nu})^2 \lambda\lambda$. In a similar way, using the asymptotic expansion of the more involved $G_{(1,2)}$ component, we find that the leading term is primitive (as one could also see from (3.6)) and is given by:

$$G_{(1,2)}^{\text{UV}} = \left(\frac{3}{2}\right)^{7/4} \frac{2\pi\varepsilon}{\mathcal{I}_0} \frac{1}{\sqrt{\alpha'M}g_s^{3/4}} \log\left(\frac{r_{\text{UV}}^3}{|S|}\right)^{1/4} \left(\frac{|S|}{r_{\text{UV}}^3}\right)^{4/3} (h_1 \wedge \bar{h}_1 \wedge \bar{h}_3 - h_2 \wedge \bar{h}_2 \wedge \bar{h}_3) + O(\varepsilon^2), \quad (3.15)$$

which means that this must be holographically dual to the expectation value of an operator of dimension $\Delta = 4$, which corresponds to a marginal deformation of the superpotential [39].

Finally, the leading asymptotic terms in the (2,1) component at zeroth order in ε is given by:

$$G_{(2,1)}^0 = \left(\frac{2}{3}\right)^{1/4} \frac{1}{\sqrt{\alpha'M}g_s^{3/4}} \log\left(\frac{r_{\text{UV}}^3}{|S|}\right)^{-3/4} (h_1 \wedge h_3 \wedge \bar{h}_1 - h_2 \wedge h_3 \wedge \bar{h}_2), \quad (3.16)$$

whereas the leading asymptotic behavior of the (2,1) component at first order in ε is

$$G_{(2,1)}^1 = \frac{24\varepsilon}{\sqrt{\alpha'M}g_s^{3/4}} \log\left(\frac{r_{\text{UV}}^3}{|S|}\right)^{-7/4} (h_1 \wedge h_3 \wedge \bar{h}_1 - h_2 \wedge h_3 \wedge \bar{h}_2). \quad (3.17)$$

4 Fermion masses

The presence of new components of the fluxes can strongly affect the worldvolume dynamics of D7-branes in the UV. In fact, any component of G_3 that is not (2, 1) generates fermion

masses, possibly breaking supersymmetry on the brane. For instance, a non-trivial $G_{(0,3)}$ component is responsible for a non-vanishing gaugino mass $m_\lambda \neq 0$ [17–19]. A gaugino mass that is larger than the gaugino condensation scale, Λ_{YM} , signals a breakdown of the $\mathcal{N} = 1$ supersymmetric dynamics at this scale, and affects the expression of the ADS superpotential. Hence, in order for the KKLTT “*moduli stabilization via gaugino condensation*” scenario to apply we need $\frac{m_\lambda}{\Lambda_{\text{YM}}} \ll 1$. Our purpose is to calculate this ratio.

The mass of the canonically normalized gaugino is given by [19]:

$$m_\lambda = \left(-\frac{4\pi^2}{\sqrt{6}} \right) \frac{1}{4} \frac{\int \delta_\Sigma^{(0)} e^{\phi/2} G_3 \wedge \Omega}{\int_\Sigma \frac{1}{2} J \wedge J}, \tag{4.1}$$

where we work in Einstein frame and $\delta_\Sigma^{(0)}$ localizes the integral on the cycle Σ wrapped by the seven-branes. The three-form, Ω , is again normalized such that $\Omega \wedge \bar{\Omega} = \frac{3i}{\pi^4} \text{vol}_6$ and the two-form J such that $\frac{1}{2} J \wedge J = \text{vol}_4$. Here vol_6 is the volume form of the warped Calabi-Yau threefold and vol_4 is the volume of the 4-cycle wrapped by the D7-branes. The $(0, 3)$ fluxes (3.12) then generate the D7 worldvolume gaugino mass:⁷

$$m_\lambda = (-i) \left(\frac{2}{3} \right)^{3/4} \frac{162 \pi \varepsilon g_s^{-1/4}}{5 \mathcal{I}_0 \sqrt{\alpha' M}} \log(r_{\text{UV}}^3 / |S|)^{5/4} \left(\frac{|S|}{r_{\text{UV}}^3} \right)^{7/3}, \tag{4.2}$$

where r_{UV} is the radial cut-off, whose value can be determined if we require the warping to be of order one in the UV.⁸

We now compare this with a four-dimensional supergravity computation of the supersymmetry breaking gaugino masses. In order to compare correctly, we should use the Gukov-Vafa-Witten superpotential, without including the non-perturbative superpotential coming from gaugino condensation on the D7-branes, since the computation of the $\overline{\text{D3}}$ brane induced fluxes is performed without it. The general form of Majorana gaugino masses in four-dimensional supergravity is given, up to an irrelevant phase, by

$$m_\lambda = \frac{1}{2 \text{Re} f} \mathcal{K}^{i\bar{j}} e^{\frac{\mathcal{K}}{2}} \overline{D_{\bar{j}} W} \frac{\partial f}{\partial z^i}, \tag{4.3}$$

where \mathcal{K} is the Kähler potential, $D_i W = \partial_i W + (\partial_i \mathcal{K}) W$ is the Kähler covariant derivative of the superpotential W , $f(z^i)$ is the gauge kinetic function and z^i denote the complex scalars in the chiral multiplets.

The relevant quantities in the 4d SUGRA Lagrangian for a single Kähler modulus are⁹

$$\mathcal{K} = -3 \log r - \log(2/g_s) - \log \left(\frac{|\Omega|^2 V_w^2}{\kappa_4^{12}} \right), \quad W = \frac{1}{\kappa_4^8} \int_M G_3 \wedge \Omega, \quad f_{D7} = T, \tag{4.4}$$

where

$$r = T + \bar{T} - \frac{3c' g_s (\alpha' M)^2}{\pi |\Omega|^2 V_w} |S|^{2/3}. \tag{4.5}$$

⁷Note that this is the fall-off corresponding to an operator of dimension 7 in the holographic theory dual to the KS solution, and not to an operator of dimension 3, which would correspond to a gaugino mass in this theory.

⁸This is the natural assumption if the Calabi-Yau manifold is weakly warped.

⁹For more details of this particular form of the Kähler potential and superpotential see [4].

Here V_w is a fiducial volume that we take to be equal to one, $\kappa_4 = M_{pl}^{-2}$, $c' = 1.18$ is a numerical factor coming from taking into account warping effects in the effective field theory [46] and Ω has the usual normalization: $|\Omega|^2 = 8$. From (4.4) one can easily see that

$$D_T W = -\frac{3}{\kappa_4^8 r} \int G_3 \wedge \Omega. \tag{4.6}$$

Combining all the above relations, the gaugino mass (4.3) is

$$m_\lambda = -\frac{\sqrt{g_s}}{4\kappa_4^8 (T + \bar{T}) r^{1/2}} \int G_3 \wedge \Omega. \tag{4.7}$$

In order to compare this expression with (4.2) we should note that in Einstein frame, $T = \frac{\text{vol}_4}{2\pi\alpha'^2}$ and that for a single Kähler modulus $\text{vol}_6 = \frac{\sqrt{2}}{3} T^{3/2}$. Taking into account the $\delta_\Sigma^{(0)}$ localization factor in (4.1), it is not hard to see that, in the regime where we can neglect the conifold contribution in (4.5), the two results agree parametrically.¹⁰

4.1 Gaugino mass and gaugino condensation

As already mentioned, the component $G_{(0,3)}$ is non-vanishing and gives rise to a mass for the world-volume gaugino of the D7-branes. Since this mass is generated by the addition of $\overline{\text{D3}}$ branes in the IR throat, we can think of it as the manifestation of a $\overline{\text{D3}}$ -D7 interaction. In order to see whether and how this mass can affect the dynamics of the gaugino condensation responsible for Kähler-moduli stabilization in the KKLТ scenario, we need to compute the various energy scales of our system. In the following, we will mostly follow the same conventions as in [8, 14]. First let us recall that:

$$M = \frac{1}{4\pi^2\alpha'} \int_A F_3, \quad K = \frac{1}{4\pi^2\alpha'} \int_B H_3, \tag{4.8}$$

where the A -cycle is the S^3 at the bottom of the throat. The B -cycle is a bit more subtle, extending to the brim of the KS throat and into the Calabi-Yau compactification manifold. Here we approximate K with the integral of H_3 over the part of this cycle inside the throat, ignoring the contribution from the rest of the compactification manifold.

If we call τ_{UV} the distance from the tip of the KS throat to the region where the throat merges with the CY,¹¹ then

$$K = \frac{1}{4\pi^2\alpha'} \int_0^{\tau_{UV}} d\tau \int_{S^2} H_3, \tag{4.9}$$

where the two-sphere is taken at $\psi = 0$, $\theta_1 = -\theta_2$, $\phi_1 = -\phi_2$. Using the form (2.3) for the fluxes and working at zeroth order in ε , we can compute the units of NSNS-flux as:

$$K = -\frac{2}{\pi\alpha'} f(\tau) \Big|_0^{\tau_{UV}} \approx \frac{g_s M}{2\pi} \tau_{UV}, \tag{4.10}$$

¹⁰An independent check of the agreement between the 10d versus the 4d description would be a ten-dimensional computation of the gravitino mass, along the lines of [47], taking into account the backreaction effects from the antibrane.

¹¹In KS holography this corresponds to the UV cutoff, Λ_{UV} .

where in the last expression we evaluated $f(\tau)$ assuming τ_{UV} to be large. This gives the radial cutoff of the throat and its associated energy scale:

$$r_{UV}^3 = \Lambda_{UV}^3 = \frac{3^{3/2}}{2^{5/2}} |S| e^{\frac{2\pi K}{g_s M}}, \quad (4.11)$$

where we used the relation between τ and r given in (3.13). Using holographic KS terminology, one can also define an infrared energy scale, which can be taken to coincide with the value of r at the bottom of the throat ($\tau = 0$). We can thus introduce a parameter, η , measuring the hierarchy between the ultraviolet and infrared scales:

$$\eta = 3 \ln \frac{\Lambda_{UV}}{\Lambda_{IR}} = \frac{2\pi K}{g_s M}. \quad (4.12)$$

Even if we do not know the full metric and fluxes of the Calabi-Yau compactification, it is reasonable to assume that, when the size of the compactification manifold is not very large, the fields at the location of the N_{D7} seven-branes wrapping a four-cycle Σ are of the same order as the fields at the brim of the KS throat, at $r_{UV} = \Lambda_{UV}$.

In the absence of a gaugino mass, the low-energy world-volume theory of D7 branes at a non-singular locus is 8-dimensional $SU(N_{D7})$ SYM theory, which one further compactifies on Σ .¹² The coupling constant of the resulting $\mathcal{N} = 1$ four-dimensional gauge theory runs logarithmically with the energy and the theory confines in the infrared at a scale

$$\Lambda_{YM} \approx \mu_0 e^{-\frac{2\pi \text{Re } T}{3N_{D7}}}, \quad (4.13)$$

where μ_0 is the ‘‘UV scale’’ that depends on (stabilized) complex-structure moduli and will be assumed to be of order one (in Planck units). If the world-volume theory has a more general gauge group, G , the exponent in (4.13) must be replaced by $-2\pi \text{Re } T / (3 \#_C(G))$, with $\#_C(G)$ the dual Coxeter number of G . Furthermore, in the confined phase the gaugino condenses, giving rise to a nontrivial contribution to the superpotential that depends on the coupling constant of the four-dimensional $\mathcal{N} = 1$ theory, and hence on the volume of the four-cycle wrapped by the D7 branes.

In the presence of gaugino mass, m_λ , induced by the three-form fields sourced by the antibranes, this scenario can change. In particular, when m_λ is larger than Λ_{YM} , the theory will confine at an energy scale proportional to m_λ , and the resulting potential will be independent of the four-cycle volume, thus ruining the Kähler moduli stabilization by gaugino condensation. Hence, gaugino condensation can only stabilize the Kähler moduli when:

$$\frac{m_\lambda}{\Lambda_{YM}} \ll 1. \quad (4.14)$$

To estimate this ratio one can consider a KKLT-like scenario, in which the value of the potential at the AdS minimum is proportional to the square of the non-perturbative superpotential, while the contribution to the scalar potential from the anti-D3 branes (which sit at the tip of the throat, where their energy is minimized) is proportional to H^{-1} . In a realistic de Sitter compactification, where we want the cosmological constant to be as small

¹²When the D7 branes are on top of an O7-plane, the gauge group can be orthogonal or symplectic.

as possible, these two quantities should have approximately the same magnitude, which means that the following relation should hold¹³

$$V_{D3} \approx V_{\text{AdS}} \rightarrow \frac{|S|^{4/3}}{h_0} \approx e^{-\frac{4\pi}{N_c} \text{Re}(T)}. \quad (4.16)$$

In our conventions the hierarchy is given by $e^{-\eta}$ and we note that $\tau_{UV} \approx \eta$. We also require that the warping is $O(1)$ in the UV . All these give a relation between the complex-structure modulus, S , and the flux-induced D3 charge of the throat, $Q_{D3}^{\text{Throat}} = KM$:

$$H = \frac{h(\tau)}{|S|^{4/3}} \approx 1 \quad \rightarrow \quad |S|^{4/3} \approx h(\tau_{UV}) \approx 6\pi \cdot 2^{1/3} (\alpha')^2 e^{-\frac{4\eta}{3}} Q_{D3}^{\text{Throat}}. \quad (4.17)$$

Combining (4.16) (more precisely (4.15)) and (4.17), we can estimate the value of the stabilized Kähler modulus

$$\text{Re } T \sim \frac{2N_c K}{3g_s M} = \frac{N_c}{3\pi} \eta, \quad (4.18)$$

where we have ignored logarithmic corrections coming from the non-exponential terms in (4.15).

Using this approximation, we can evaluate the ratio between the gaugino mass and the condensation scale in the KKLT scenario:

$$\frac{m_\lambda}{\Lambda_{\text{YM}}} \approx 10^2 \bar{N} M^{-5/2} g_s^{-5/4} \eta^{5/4} e^{-\frac{19}{9}\eta}. \quad (4.19)$$

Our calculation indicates that this ratio must be smaller than one in all flux compactifications where Kähler moduli are stabilized via D7 gaugino condensation and where the cosmological constant is uplifted using anti-D3 branes in warped Klebanov-Strassler-like throats. It is not hard to see that for the range of parameters one uses in “vanilla” KKLT scenarios this bound will be satisfied. For example for the de Sitter minimum of [7], where $M = K = 70$ and $g_s = 1/2$ we find this ratio to be:

$$\frac{m_\lambda}{\Lambda_{\text{YM}}} \sim 10^{-13}. \quad (4.20)$$

Hence, in these scenarios, the stabilization of the Kähler moduli via D7 gaugino condensation is not affected by the fluxes produced by the anti-D3 branes.

In [48] it was also argued, by evaluating the D7-brane gauge coupling function in the perturbed solution, that the anti-D3 branes do not affect the gaugino condensate for a sufficiently large throat. It would be interesting to try to span larger families of possible flux compactifications and other de Sitter scenarios which use a smaller hierarchy (such as those of [21, 22]) to see whether this bound may become harder to satisfy and may result in nontrivial constraints on the parameters of the compactification.

¹³If one takes into account the explicit expression for both V_{D3} [10] and V_{AdS} , one finds using (4.17) a more precise expression

$$e^{-\frac{2\pi \text{Re } T}{3N_c}} \approx \left(\frac{N_c}{A}\right)^{1/3} \left(\frac{\eta}{\text{Re } T}\right)^{1/6} e^{-2\eta/9}, \quad (4.15)$$

where A is the Pfaffian factor that appears in the non-perturbative superpotential. Since we assume both A and N_c to be of order one, the difference between the results obtained with (4.15) and those obtained with (4.16) is negligible.

Acknowledgments

We would like to thank Veronica Collazuol, Nicolás Kovensky, Severin Lüst and Salvatore Raucci for useful discussions. This work was supported in part by the ERC Grants 772408 “Stringlandscape” and 787320 “QBH Structure”, and by the John Templeton Foundation grant 61149.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] S. Kachru, R. Kallosh, A.D. Linde and S.P. Trivedi, *De Sitter vacua in string theory*, *Phys. Rev. D* **68** (2003) 046005 [[hep-th/0301240](https://arxiv.org/abs/hep-th/0301240)] [[INSPIRE](https://inspirehep.net/literature/9891)].
- [2] I.R. Klebanov and M.J. Strassler, *Supergravity and a confining gauge theory: Duality cascades and χ SB resolution of naked singularities*, *JHEP* **08** (2000) 052 [[hep-th/0007191](https://arxiv.org/abs/hep-th/0007191)] [[INSPIRE](https://inspirehep.net/literature/5421)].
- [3] I. Bena et al., *The backreaction of anti-D3 branes on the Klebanov-Strassler geometry*, *JHEP* **06** (2013) 060 [[arXiv:1106.6165](https://arxiv.org/abs/1106.6165)] [[INSPIRE](https://inspirehep.net/literature/1000000)].
- [4] I. Bena, M. Graña and N. Halmagyi, *On the Existence of Meta-stable Vacua in Klebanov-Strassler*, *JHEP* **09** (2010) 087 [[arXiv:0912.3519](https://arxiv.org/abs/0912.3519)] [[INSPIRE](https://inspirehep.net/literature/8430)].
- [5] I. Bena, M. Graña, S. Kuperstein and S. Massai, *Giant Tachyons in the Landscape*, *JHEP* **02** (2015) 146 [[arXiv:1410.7776](https://arxiv.org/abs/1410.7776)] [[INSPIRE](https://inspirehep.net/literature/1270000)].
- [6] U.H. Danielsson and T. Van Riet, *Fatal attraction: more on decaying anti-branes*, *JHEP* **03** (2015) 087 [[arXiv:1410.8476](https://arxiv.org/abs/1410.8476)] [[INSPIRE](https://inspirehep.net/literature/1270000)].
- [7] R. Blumenhagen, D. Kläwer and L. Schlechter, *Swampland Variations on a Theme by KKLT*, *JHEP* **05** (2019) 152 [[arXiv:1902.07724](https://arxiv.org/abs/1902.07724)] [[INSPIRE](https://inspirehep.net/literature/1660000)].
- [8] I. Bena, E. Dudas, M. Graña and S. Lüst, *Uplifting Runaways*, *Fortsch. Phys.* **67** (2019) 1800100 [[arXiv:1809.06861](https://arxiv.org/abs/1809.06861)] [[INSPIRE](https://inspirehep.net/literature/1660000)].
- [9] L. Randall, *The Boundaries of KKLT*, *Fortsch. Phys.* **68** (2020) 1900105 [[arXiv:1912.06693](https://arxiv.org/abs/1912.06693)] [[INSPIRE](https://inspirehep.net/literature/1700000)].
- [10] E. Dudas and S. Lüst, *An update on moduli stabilization with antibrane uplift*, *JHEP* **03** (2021) 107 [[arXiv:1912.09948](https://arxiv.org/abs/1912.09948)] [[INSPIRE](https://inspirehep.net/literature/1700000)].
- [11] I.R. Klebanov and A.A. Tseytlin, *Gravity duals of supersymmetric $SU(N) \times SU(N+M)$ gauge theories*, *Nucl. Phys. B* **578** (2000) 123 [[hep-th/0002159](https://arxiv.org/abs/hep-th/0002159)] [[INSPIRE](https://inspirehep.net/literature/5421)].
- [12] S. Lüst and L. Randall, *Effective Theory of Warped Compactifications and the Implications for KKLT*, *Fortsch. Phys.* **70** (2022) 2200103 [[arXiv:2206.04708](https://arxiv.org/abs/2206.04708)] [[INSPIRE](https://inspirehep.net/literature/2000000)].
- [13] A. Buchel, *Klebanov-Strassler black hole*, *JHEP* **01** (2019) 207 [[arXiv:1809.08484](https://arxiv.org/abs/1809.08484)] [[INSPIRE](https://inspirehep.net/literature/1660000)].
- [14] I. Bena, A. Buchel and S. Lüst, *Throat destabilization (for profit and for fun)*, [arXiv:1910.08094](https://arxiv.org/abs/1910.08094) [[INSPIRE](https://inspirehep.net/literature/1700000)].
- [15] I. Bena et al., *Bare-bones de Sitter vacua*, *Phys. Rev. D* **108** (2023) L021901 [[arXiv:2202.02327](https://arxiv.org/abs/2202.02327)] [[INSPIRE](https://inspirehep.net/literature/2000000)].

- [16] I. Affleck, M. Dine and N. Seiberg, *Dynamical Supersymmetry Breaking in Supersymmetric QCD*, *Nucl. Phys. B* **241** (1984) 493 [INSPIRE].
- [17] P.G. Camara, L.E. Ibanez and A.M. Uranga, *Flux-induced SUSY-breaking soft terms on D7-D3 brane systems*, *Nucl. Phys. B* **708** (2005) 268 [hep-th/0408036] [INSPIRE].
- [18] H. Jockers and J. Louis, *The effective action of D7-branes in $N = 1$ Calabi-Yau orientifolds*, *Nucl. Phys. B* **705** (2005) 167 [hep-th/0409098] [INSPIRE].
- [19] M. Graña, N. Kovensky and A. Retolaza, *Gaugino mass term for D-branes and Generalized Complex Geometry*, *JHEP* **06** (2020) 047 [arXiv:2002.01481] [INSPIRE].
- [20] I. Bena et al., *On Metastable Vacua and the Warped Deformed Conifold: Analytic Results*, *Class. Quant. Grav.* **30** (2013) 015003 [arXiv:1102.2403] [INSPIRE].
- [21] C. Crinò, F. Quevedo and R. Valandro, *On de Sitter String Vacua from Anti-D3-Branes in the Large Volume Scenario*, *JHEP* **03** (2021) 258 [arXiv:2010.15903] [INSPIRE].
- [22] B.V. Bento, D. Chakraborty, S.L. Parameswaran and I. Zavala, *A new de Sitter solution with a weakly warped deformed conifold*, *JHEP* **12** (2021) 124 [arXiv:2105.03370] [INSPIRE].
- [23] X. Gao, A. Hebecker and D. Junghans, *Control issues of KKLT*, *Fortsch. Phys.* **68** (2020) 2000089 [arXiv:2009.03914] [INSPIRE].
- [24] D. Junghans, *LVS de Sitter vacua are probably in the swampland*, *Nucl. Phys. B* **990** (2023) 116179 [arXiv:2201.03572] [INSPIRE].
- [25] E. Witten, *Nonperturbative superpotentials in string theory*, *Nucl. Phys. B* **474** (1996) 343 [hep-th/9604030] [INSPIRE].
- [26] E. Bergshoeff et al., *An index for the Dirac operator on D3 branes with background fluxes*, *JHEP* **10** (2005) 102 [hep-th/0507069] [INSPIRE].
- [27] D. Lüüst, S. Reffert, W. Schulgin and P.K. Tripathy, *Fermion zero modes in the presence of fluxes and a non-perturbative superpotential*, *JHEP* **08** (2006) 071 [hep-th/0509082] [INSPIRE].
- [28] D. Robbins and S. Sethi, *A Barren landscape?*, *Phys. Rev. D* **71** (2005) 046008 [hep-th/0405011] [INSPIRE].
- [29] P.K. Tripathy and S.P. Trivedi, *D3 brane action and fermion zero modes in presence of background flux*, *JHEP* **06** (2005) 066 [hep-th/0503072] [INSPIRE].
- [30] R. Kallosh, A.-K. Kashani-Poor and A. Tomasiello, *Counting fermionic zero modes on M5 with fluxes*, *JHEP* **06** (2005) 069 [hep-th/0503138] [INSPIRE].
- [31] N. Saulina, *Topological constraints on stabilized flux vacua*, *Nucl. Phys. B* **720** (2005) 203 [hep-th/0503125] [INSPIRE].
- [32] L. Martucci, J. Rosseel, D. Van den Bleeken and A. Van Proeyen, *Dirac actions for D-branes on backgrounds with fluxes*, *Class. Quant. Grav.* **22** (2005) 2745 [hep-th/0504041] [INSPIRE].
- [33] P. Berglund and P. Mayr, *Non-perturbative superpotentials in F-theory and string duality*, *JHEP* **01** (2013) 114 [hep-th/0504058] [INSPIRE].
- [34] M. Bianchi, G. Inverso and L. Martucci, *Brane instantons and fluxes in F-theory*, *JHEP* **07** (2013) 037 [arXiv:1212.0024] [INSPIRE].
- [35] A. Dymarsky, I.R. Klebanov and N. Seiberg, *On the moduli space of the cascading $SU(M+p) \times SU(p)$ gauge theory*, *JHEP* **01** (2006) 155 [hep-th/0511254] [INSPIRE].

- [36] K.A. Intriligator, N. Seiberg and D. Shih, *Dynamical SUSY breaking in meta-stable vacua*, *JHEP* **04** (2006) 021 [[hep-th/0602239](#)] [[INSPIRE](#)].
- [37] R. Argurio, M. Bertolini, S. Franco and S. Kachru, *Meta-stable vacua and D-branes at the conifold*, *JHEP* **06** (2007) 017 [[hep-th/0703236](#)] [[INSPIRE](#)].
- [38] N. Seiberg, private communication.
- [39] M. Graña and J. Polchinski, *Gauge / gravity duals with holomorphic dilaton*, *Phys. Rev. D* **65** (2002) 126005 [[hep-th/0106014](#)] [[INSPIRE](#)].
- [40] S. Kuperstein and J. Sonnenschein, *Analytic nonsupersymmetric background dual of a confining gauge theory and the corresponding plane wave theory of hadrons*, *JHEP* **02** (2004) 015 [[hep-th/0309011](#)] [[INSPIRE](#)].
- [41] O. Aharony, A. Buchel and A. Yarom, *Holographic renormalization of cascading gauge theories*, *Phys. Rev. D* **72** (2005) 066003 [[hep-th/0506002](#)] [[INSPIRE](#)].
- [42] G. Papadopoulos and A.A. Tseytlin, *Complex geometry of conifolds and five-brane wrapped on two sphere*, *Class. Quant. Grav.* **18** (2001) 1333 [[hep-th/0012034](#)] [[INSPIRE](#)].
- [43] A. Dymarsky, *On gravity dual of a metastable vacuum in Klebanov-Strassler theory*, *JHEP* **05** (2011) 053 [[arXiv:1102.1734](#)] [[INSPIRE](#)].
- [44] A. Dymarsky and S. Massai, *Uplifting the baryonic branch: a test for backreacting anti-D3-branes*, *JHEP* **11** (2014) 034 [[arXiv:1310.0015](#)] [[INSPIRE](#)].
- [45] V. Borokhov and S.S. Gubser, *Nonsupersymmetric deformations of the dual of a confining gauge theory*, *JHEP* **05** (2003) 034 [[hep-th/0206098](#)] [[INSPIRE](#)].
- [46] M.R. Douglas, J. Shelton and G. Torroba, *Warping and supersymmetry breaking*, [arXiv:0704.4001](#) [[INSPIRE](#)].
- [47] O. DeWolfe and S.B. Giddings, *Scales and hierarchies in warped compactifications and brane worlds*, *Phys. Rev. D* **67** (2003) 066008 [[hep-th/0208123](#)] [[INSPIRE](#)].
- [48] S. Kachru, M. Kim, L. Mcallister and M. Zimet, *de Sitter vacua from ten dimensions*, *JHEP* **12** (2021) 111 [[arXiv:1908.04788](#)] [[INSPIRE](#)].