

Regulation of Atherogenesis by Chemokines and Chemokine Receptors

Wuzhou Wan · Philip M. Murphy

Received: 4 June 2012 / Accepted: 18 November 2012 / Published online: 7 December 2012
© L. Hirszfeld Institute of Immunology and Experimental Therapy, Wrocław, Poland 2012

Abstract Atherosclerosis is a chronic inflammatory and metabolic disorder affecting large- and medium-sized arteries, and the leading cause of mortality worldwide. The pathogenesis of atherosclerosis involves accumulation of lipids and leukocytes in the intima of blood vessel walls creating plaque. How leukocytes accumulate in plaque remains poorly understood; however, chemokines acting at specific G protein-coupled receptors appear to be important. Studies using knockout mice suggest that chemokine receptor signaling may either promote or inhibit atherogenesis, depending on the receptor. These proof of concept studies have spurred efforts to develop drugs targeting the chemokine system in atherosclerosis, and several have shown beneficial effects in animal models. This study will review key discoveries in basic and translational research in this area.

Keywords Atherosclerosis · Cardiology · Immunology · Inflammation · Antagonist

Introduction

Atherosclerosis is the pathologic process underlying most strokes and heart attacks, which together are now the leading cause of death worldwide (Roger et al. 2012). Risk factors for atherosclerosis include age, gender, a high ratio of low-density lipoprotein (LDL) to high-density lipoprotein (HDL)

in the blood, hypertension, diabetes, obesity, smoking and inheritance (Berger et al. 2010). It is generally accepted that atherosclerosis is a chronic metabolic and inflammatory disease (Hansson 2005). The pathologic hallmark is the atherosclerotic plaque, composed of lipids, collagen, platelets, fibroblasts, smooth muscle cells (SMCs) and leukocytes. Rupture of unstable plaques may result in thrombosis and ischemia in surrounding tissues (Weber et al. 2008; Weber and Noels 2011).

Both innate and adaptive immunity appear to be involved in the development of plaque (Packard et al. 2009). Innate immune cells, including macrophages, neutrophils, mast cells and platelets, accumulate early and express reactive oxygen species (ROS), proteinases, lipid mediators and various cytokines, leading to SMC proliferation, angiogenesis and additional inflammatory cell activation. Adaptive immune cells, including T and B lymphocytes, accumulate later and may accelerate disease progression through complex and poorly understood mechanisms (Galkina and Ley 2009; Weber et al. 2008). In particular, in mouse models, the B1 subset of B cells is atheroprotective, whereas the B2 subset is pro-atherogenic; Th1 cells are pro-atherogenic, and regulatory T cells (Tregs) are atheroprotective, whereas the roles of Th2 and Th17 cells are still controversial (Butcher and Galkina 2011; Kyaw et al. 2011; Robertson and Hansson 2006).

The classic mechanism of immune cell recruitment from the blood into tissues, including the blood vessel wall, is thought to involve sequential interactions between leukocytes and endothelial cells (ECs). These include selectin-mediated leukocyte rolling on endothelium, followed by leukocyte activation by chemokines, then integrin-mediated firm arrest of leukocytes to endothelium, culminating in leukocyte transendothelial migration (Ley et al. 2007). Chemokines are a large protein family of small

W. Wan · P. M. Murphy (✉)
Laboratory of Molecular Immunology,
National Institute of Allergy and Infectious Diseases,
National Institutes of Health, 9000 Rockville Pike,
Bldg 10, Rm 11N113, Bethesda, MD 20892, USA
e-mail: pmm@nih.gov

(8–10 kDa) leukocyte chemoattractants that can be divided into four classes: C, CC, CXC and CX3C according to the number and spacing of conserved cysteines in the N-terminal domain of the molecule (Murphy et al. 2000). In human, approximately 46 chemokines and 20 chemokine receptors have been identified (Murphy 2002). Since the discovery in 1998 that the chemokine Ccl2 and its receptor Ccr2 are positive regulators of both the apolipoprotein E-deficient (*ApoE*^{-/-}) and LDL receptor-deficient (*Ldlr*^{-/-}) mouse models of atherosclerosis (Boring et al. 1998; Dawson et al. 1999; Gu et al. 1998), roles for many other chemokines and chemokine receptors have been identified using genetic and, in some cases, pharmacological criteria (Koenen and Weber 2011). In this review, we will discuss current concepts of the chemokine system as an immunoregulator and potential therapeutic target in atherosclerotic cardiovascular disease.

The Chemokine System as an Immunoregulator in Atherogenesis

Chemokines and chemokine receptors are able to regulate leukocyte trafficking in both homeostasis and inflammation (Murphy et al. 2000). Chemokine receptors identified in atherogenesis include inflammatory receptors (e.g., CCR2, CCR5, CXCR2 and CX3CR1), classical homeostatic receptors (e.g., CCR7) as well as receptors that have both inflammatory and homeostatic functions (e.g., CCR6) (Koenen and Weber 2011). We will discuss the chemokines and their receptors in three different categories: atherogenic, atheroprotective and controversial (Table 1).

Atherogenic Chemokines/Chemokine Receptors

CCL2–CCR2

CCL2–CCR2 was the first chemokine/chemokine receptor pair to be implicated in the pathogenesis of atherosclerosis, and the one studied in the greatest detail. *CCL2* (also known as monocyte chemoattractant protein-1) is mainly produced by monocytes, macrophages, ECs and SMCs, and has been identified in both mouse and human atherosclerotic lesions (Nelken et al. 1991; Rayner et al. 2000). In human, single nucleotide polymorphisms (SNPs) in the promoter of *CCL2*, *CCL2-2518G*, and in the open reading frame of *CCR2*, *CCR2-V64I*, have been associated with increased risk of myocardial infarction (McDermott et al. 2005; Ortlepp et al. 2003; Szalai et al. 2001). *CCL2-2518G* appears to be functional since individuals with this allele have significantly higher serum *CCL2* levels (McDermott et al. 2005). The biochemical effect of *CCR2-V64I* has not been clearly defined. In mice, genetic deletion of either *Ccl2* or *Ccr2* in both the *ApoE*^{-/-} and *Ldlr*^{-/-} models of atherosclerosis significantly reduced the size of lesions in the aorta (Boring et al. 1998; Dawson et al. 1999; Gu et al. 1998). In a third model, *Ccl2* deficiency decreased atherosclerotic plaque burden in mice that overexpress human apolipoprotein B (Gosling et al. 1999). A pro-atherogenic role for *Ccl2/Ccr2* was also demonstrated by blocking *Ccr2* with 7ND, an N-terminally truncated mutant form of *Ccl2*, which attenuated the initiation and progression of atherosclerosis in *ApoE*^{-/-} mice (Inoue et al. 2002; Ni et al. 2001). A study of radiation chimeric mice has attributed the pro-atherogenic effect of *Ccr2* to expression on hematopoietic cells. In particular, *Ccr2*^{-/-} but not *Ccr2*^{+/+} bone marrow

Table 1 Different effects of chemokines/chemokine receptors in atherogenesis

Effect in atherosclerosis	Chemokines/chemokine receptors	References
Pathogenic	<i>CCL2–CCR2</i>	Boring et al. (1998), Dawson et al. (1999), Gosling et al. (1999), Gu et al. (1998) and Guo et al. (2003)
	<i>CCL5–CCR5</i>	Braunersreuther et al. (2007), Potteaux et al. (2006) and Quinones et al. (2007)
	<i>CCR6</i>	Doran et al. (2012) and Wan et al. (2011)
	<i>CCL17</i>	Weber and Noels (2011)
	<i>CX3CL1–CX3CR1</i>	Combadiere et al. (2003), Lesnik et al. (2003), Liu et al. (2008) and Teupser et al. (2004)
	<i>CXCL1–CXCR2</i>	Boisvert et al. (1998, 2006) and Huo et al. (2001)
	<i>CXCL10–CXCR3</i>	Heller et al. (2006) and Veillard et al. (2005)
	<i>CXCR6</i>	Galkina et al. (2007)
	Protective	<i>CCR1</i>
<i>CXCR4</i>		Zernecke et al. (2008)
Controversial	<i>CXCL16</i>	Aslanian and Charo (2006) and Yi et al. (2011)
	<i>CCR7</i>	Feig et al. (2011a, b), Luchtefeld et al. (2010), Potteaux et al. (2011) and Trogan et al. (2006)

transplantation into ApoE3-leiden mice reduced the size of atherosclerotic lesions; moreover, overexpression of Ccl2 on hematopoietic cells in *ApoE*^{-/-} mice increased atherosclerotic lesion size (Aiello et al. 1999; Guo et al. 2003). However, blockade of Ccr2 by either 7ND treatment or *Ccr2*^{-/-} bone marrow transplantation did not affect the progression of established atherosclerotic lesion development (de Waard et al. 2010; Guo et al. 2005). Further studies in mice showed that *Ccr2* deficiency abolished the egress of monocytes from the bone marrow and markedly reduced Ccl2-induced recruitment of monocytes from the blood into inflammatory sites (Boring et al. 1997; Tacke et al. 2007; Tsou et al. 2007). Considering that monocytes from the bone marrow and the spleen are critical for both the initiation and progression of atherosclerosis (Fig. 1)

(Robbins et al. 2012; Swirski et al. 2009; Weber et al. 2008), this may partially explain why Ccl2 and Ccr2 deficiencies inhibit the development of atherosclerosis.

CCL5–CCR5

The inflammatory chemokine CCL5 [also known as regulated upon activation, normal T cell expressed and secreted (RANTES)] is expressed by monocytes, macrophages, T cells and SMCs in both mouse and human atherosclerotic lesions, and acts at the chemokine receptors CCR1, CCR3 and CCR5 (Krohn et al. 2007; Pattison et al. 1996). In human, the gain-of-function polymorphism *CCL5-G403A* affects basal CCL5 expression levels and is associated with a higher risk of coronary artery disease (CAD) (Boger et al.




Fig. 1 Schematic representation of chemokines/chemokine receptors and their target cells involved in the progress of atherosclerosis. **CCL2–CCR2** and **CCL20–CCR6** induce the egress of monocytes from the bone marrow and the spleen into the circulating blood (arrows), and then together with **CCL5–CCR5** and **CX3CL1–CX3CR1**, they mediate the arrest of monocytes on the endothelium and their migration into the atherosclerotic plaque. In the plaque, monocytes differentiate into macrophages (foam cells) after digestion of oxLDL through **CXCL16** and other scavenger receptors. **CX3CL1–CX3CR1** is also important for the survival/retention of macrophages in the plaque and the formation of monocyte–platelet complex during atherogenesis. The egress of neutrophils from the bone marrow and their recruitment into the vessel wall are mediated by **CXCL1/8–CXCR2** and **CXCL12–CXCR4**. The adhesion and

migration of Th1 cells are controlled by **CXCL10–CXCR3** and **CCL5–CCR1/5**, while the recruitment of Th2 and Th17 cells relies on **CCR4** and **CCL20–CCR6**, respectively. The proliferation of Tregs in the plaque is controlled by **CCR4** and its ligand **CCL17**, which is secreted by DCs. The recruitment of HSCs, B cells and T cells may depend on their specific chemokine receptors (dashed arrow). Macrophages, DCs and Tregs all express **CCR7** and during atherosclerosis regression these cells may egress into the lymph node by signaling through **CCL19** and **CCL21**. DCs Dendritic cells, EC endothelial cell, HSCs hematopoietic stem cells, Mono monocytes, oxLDL oxidized low-density lipoprotein, PMNs polymorphonuclear leukocytes (neutrophils), SMC smooth muscle cell, Tregs regulatory T cells

2005; Simeoni et al. 2004), and loss-of-function polymorphism *CCR5Δ32* may be associated with a lower risk of myocardial infarction (Gonzalez et al. 2001; Simeoni et al. 2004). Although genetic deletion of *Ccr5* in *ApoE*^{-/-} mice was not shown to reduce early spontaneous atherosclerosis in the initial study (Kuziel et al. 2003), later investigations suggested that it reduced both early- and late-stage atherosclerosis development in *ApoE*^{-/-} mice fed either a chow or high-fat diet (Braunersreuther et al. 2007; Quinones et al. 2007). Also, *Ldlr*^{-/-} mice reconstituted with *Ccr5*^{-/-} bone marrow showed reduced macrophage accumulation in atherosclerotic lesions and improved plaque stability (Potteaux et al. 2006). The mechanism of CCL5/CCR5 modulation of atherosclerosis has not been clearly defined. Possibilities include regulation of leukocyte trafficking to lesions and modulation of adaptive immunity. As indirect supportive evidence, CCR5 is important for the spreading and *trans*-endothelial migration of monocytes, neutrophils and Th1 cells, and the atheroprotective effects seen in *Ccr5*^{-/-} mice may be associated with up-regulation of interleukin (IL)-10 and reduction of Th1-type immune responses (Fig. 1) (Braunersreuther et al. 2007; Drechsler et al. 2010; Potteaux et al. 2006; Weber et al. 2001).

CCR6

Both CCR6 and its sole chemokine ligand CCL20 (also known as macrophage inflammatory protein 3 α) are expressed at increased levels in human atherosclerotic plaques, and the circulating level of CCL20 is significantly increased in hypercholesterolemic subjects (Calvayrac et al. 2011; Yilmaz et al. 2007). In mice, *Ccr6* and *Ccl20* are both present constitutively in healthy aortas and in atherosclerotic plaques, and *Ccr6* deletion in *ApoE*^{-/-} mice fed a Western diet significantly reduced the atherosclerotic lesion size in both the whole aorta and the aortic root, accompanied by a reduction of macrophage content in the plaques (Wan et al. 2011). Bone marrow transplantation suggested that the lesion reduction seen in *Ccr6*^{-/-}*ApoE*^{-/-} mice is caused by *Ccr6* expression on hematopoietic cells. CCL20 is produced by epithelial cells, ECs and SMCs, whereas CCR6 is mainly expressed by immature dendritic cells (DCs), B cells, T cells, neutrophils and monocytes (Schutyser et al. 2003; Wan et al. 2011). Interestingly, *Ccr6*-deficient mice on the *ApoE* knockout background were relatively monocytopenic, whereas bone marrow monocyte content was slightly increased. Thus, modulation of atherosclerosis could be due to *Ccr6* function on monocytes, mediating their egress from the bone marrow and their recruitment from the blood into the vessel wall (Fig. 1) (Koenen and Weber 2011; Wan et al. 2011). However, the effect of *Ccr6* on atherogenesis may be cell-type specific since a second study suggested that *Ccr6* might be involved

in B-cell-mediated atheroprotection (Doran et al. 2012). Because mice globally deficient in *Ccr6* have reduced atherosclerosis, the effect of *Ccr6* deficiency on non-B cells appears to be dominant over the effect of B-cell *Ccr6* deficiency (Wan and Murphy 2011). Additional work will be necessary to further clarify the underlying mechanism.

CCL17

CCL17 (also known as thymus and activation-regulated chemokine) is a DC-derived chemokine that induces chemotaxis of T cells through its receptor CCR4 (Imai et al. 1997a). CCL17 has been detected in both advanced mouse and human atherosclerotic lesions, and its expression is upregulated in human atherosclerotic plaques compared with healthy arteries (Greaves et al. 2004; Weber et al. 2011). Genetic targeting of *Ccl17* in *ApoE*^{-/-} mice resulted in reduced atherosclerotic lesion size, along with a decrease of macrophages and T cells in the plaques, which was confirmed by *Ccl17*^{-/-} bone marrow transplantation and *Ccl17*-specific antibody treatment (Weber et al. 2011). The beneficial effect may be dependent on Tregs that accumulate in the lymph nodes and aortas in the double knockout mice (Fig. 1). Moreover, co-culture of CD4⁺ T cells with *Ccl17*^{-/-} DCs resulted in less apoptosis and enhanced expansion of Tregs, suggesting that *Ccl17* is a central regulator of Treg homeostasis (Weber et al. 2011). Taken together, this suggests that CCL17 expressed by DCs may drive atherosclerosis development through inhibiting expansion of Tregs.

CX3CL1–CX3CR1

CX3CL1 (Fractalkine), the sole ligand for CX3CR1, is a unique chemokine that exists in both membrane-tethered and soluble shed forms, thus mediating both cell adhesion and chemotaxis (Bazan et al. 1997). CX3CL1 is primarily produced by ECs and CX3CR1 is expressed on monocytes, lymphocytes, platelets, and DCs (Barlic and Murphy 2007a; Imai et al. 1997b). In addition, both molecules have been identified on foam cells and SMCs in mouse and human atherosclerotic lesions, and CX3CR1 is upregulated on monocytes from coronary artery disease (CAD) patients (Apostolakis et al. 2007; Lucas et al. 2003). In several retrospective cohort studies and the population-based prospective Framingham Heart Study Offspring Cohort, two SNPs of CX3CR1, *V249I* and *T280M*, were associated with a markedly reduced risk of CAD (Ghilardi et al. 2004; McDermott et al. 2001, 2003; Moatti et al. 2001; Norata et al. 2006). However, two other studies showed that these polymorphisms were not associated with peripheral arterial disease (522 patients) but were associated with increased risk of restenosis after coronary stenting (365 patients)

(Gugl et al. 2003; Niessner et al. 2005). The biochemical effects of these polymorphisms on CX3CR1 function are also in dispute with two groups reporting that they affect function but one finding a gain of function and the other a loss of function (Daoudi et al. 2004; McDermott et al. 2003). In atherosclerosis mouse models (*ApoE*^{-/-} mice and *Ldlr*^{-/-} mice), genetic deletion of *Cx3cl1* and *Cx3cr1* significantly reduced the size of atherosclerotic lesions and inhibited the recruitment of macrophages/DCs into the vessel wall (Combadiere et al. 2003; Lesnik et al. 2003; Liu et al. 2008; Teupser et al. 2004). The role of *Cx3cl1* and *Cx3cr1* in atherogenesis appears to be independent of *Ccl2*, *Ccr2* and *Ccr5* since additive protection was observed in the *ApoE* knockout model in both *Cx3cl1*^{-/-}*Ccr2*^{-/-} mice and *Ccl2*^{-/-}*Cx3cr1*^{-/-} mice treated with Met-RANTES, a pharmacological inhibitor of *Ccr5* (Combadiere et al. 2008; Saederup et al. 2008). The proposed atherogenic mechanisms include *Cx3cr1*-dependent migration/retention of monocyte/macrophage in the vessel wall (Barlic and Murphy 2007b; Tacke et al. 2007); *Cx3cr1*-conferred monocyte/macrophage survival in atherosclerotic plaques (Landsman et al. 2009) and *Cx3cr1* supported the formation of platelet-monocyte complexes in hyperlipidemic mice (Fig. 1) (Postea et al. 2012).

CXCL1–CXCR2

Both CXCR2 and its major ligands, CXCL1 (also known as growth-related oncogene α) and CXCL8 (also known as IL-8), have been identified in human atheromata, and mouse *Cxcr2* and *Cxcl1* have been found in plaque in mouse models of atherosclerosis (Apostolopoulos et al. 1996; Boisvert et al. 2000; Wang et al. 1996). Consistent with this, the plasma level of CXCL1 is significantly increased in patients with CAD (Breland et al. 2008), and expression of CXCL8 and CXCR2 are strongly upregulated on ECs and monocytes after oxidized LDL (oxLDL) stimulation (Lei et al. 2002; Yeh et al. 2001). A pro-atherogenic role for *Cxcr2* was suggested by studies in which *Cxcr2*^{-/-} bone marrow was transplanted into irradiated *Ldlr*^{-/-} mice, resulting in much smaller lesions with fewer lesional macrophages compared to controls (Boisvert et al. 1998). Also, genetic deletion of *Cxcl1* significantly reduced atherosclerotic lesion size in both the whole aorta and the aortic root of *Ldlr*^{-/-} mice (Boisvert et al. 2006). CXCL1 is mainly produced by macrophages, neutrophils and epithelial cells, whereas CXCR2 is found most prominently on neutrophils, with some expression on monocytes and mast cells (Murphy et al. 2000). The pro-atherogenic effect of the CXCL1–CXCR2 axis may be due to CXCL1-triggered monocyte arrest on early atherosclerotic endothelium and CXCR2-mediated macrophage accumulation in established lesions (Boisvert et al. 2006; Huo et al. 2001;

Papadopoulou et al. 2008). In humans, CXCL8 levels were significantly increased in the setting of acute myocardial infarction (Neumann et al. 1995). Recent studies have suggested that neutrophils are also pro-atherogenic, and thus CXCR2 might be involved in the recruitment of neutrophils to the vessel wall during the initiation of atherosclerotic plaque formation (Fig. 1) (Drechsler et al. 2011).

CXCL10–CXCR3

CXCR3 and its ligand CXCL10 [also known as interferon (IFN)- γ -induced protein of 10 kDa or IP-10] are highly expressed in lesional T cells (Th1 cells) as well as ECs, SMCs and macrophages in human atheromata (Mach et al. 1999). The percentage of CXCR3⁺ lymphocytes has been reported to be significantly increased in the blood of CAD patients (Fernandes et al. 2004) and the plasma concentration of CXCL10 was also much higher in CAD patients compared to controls (Kawamura et al. 2003). In *ApoE*^{-/-} mice, both *Cxcr3* and *Cxcl10* deficiencies resulted in a reduction of atherosclerotic lesion formation, accompanied by a decrease of CD4⁺ T-cell accumulation in the plaques (Heller et al. 2006; Veillard et al. 2005). This was associated with an elevated number of Treg cells and increased expression of the anti-inflammatory cytokine IL-10 within lesions, suggesting that the CXCL10–CXCR3 axis may promote atherogenesis by regulating the recruitment and balance of effector T cells and Tregs (Fig. 1) (Heller et al. 2006). The role of *Cxcr3* in atherogenesis is not redundant since *Cxcr3*^{-/-}*Ccr2*^{-/-}*ApoE*^{-/-} triple knockout mice show a further decrease of lesion formation compared with deletion of either *Cxcr3* or *Ccr2* alone in *ApoE*^{-/-} mice (Veillard et al. 2005). *Cxcl10* deficiency may also be harmful to the cardiovascular system in the *ApoE*^{-/-} mouse model, since in addition to reduced atherosclerosis, increased aneurysm formation was identified in these mice (King et al. 2009).

CXCR6

CXCR6 is expressed on monocytes, macrophages, T cells and SMCs in both mouse and human atherosclerotic lesions (Wuttge et al. 2004). Genetic deletion of *Cxcr6* in *ApoE*^{-/-} mice significantly decreased the size of atherosclerotic plaques. This was accompanied by a reduced percentage of macrophages and *Cxcr6*⁺ T cells within the aortas (Fig. 1) (Galkina et al. 2007). The production of IFN- γ within the aortas of *Cxcr6*^{-/-}*ApoE*^{-/-} mice was also diminished, consistent with the finding that *Cxcr6*⁺ T cells express high amounts of IFN- γ upon activation (Calabresi et al. 2002). These results suggested that *Cxcr6* might accelerate the progress of atherosclerosis by regulating the migration of T

cells into the aortic wall thus influencing the accumulation of macrophages indirectly. The ligand for CXCR6 is CXCL16 and its effect in atherogenesis will be discussed in the following section since an opposite phenotype has been suggested (Aslanian and Charo 2006).

Atheroprotective Chemokines and Chemokine Receptors

CCR1

CCR1 is a chemokine receptor for many pro-inflammatory CC chemokines, including CCL3 and CCL5, both of which have been identified in atherosclerotic plaques; it is mainly expressed on monocytes, macrophages and T cells in humans, but important functional roles have been identified for Ccr1 on mouse neutrophils (Murphy et al. 2000; Wilcox et al. 1994). In contrast to the pro-atherogenic role of Ccl5, *Ccr1*^{-/-} bone marrow reconstitution of *Ldlr*^{-/-} mice resulted in markedly increased atherosclerotic lesion size in both the thoracic aorta and the aortic root (Potteaux et al. 2005). Similarly, *Ccr1*^{-/-}*ApoE*^{-/-} mice showed increased atherosclerotic lesion formation compared with control mice and there is a significant increase of CD3⁺ T cells and IFN- γ production in the plaques, but the lesional macrophage content was not affected (Braunersreuther et al. 2007). Also, it has been reported that although both CCR1 and CCR5 support *trans*-endothelial chemotaxis toward CCL5, only CCR1 mediates CCL5-induced arrest of monocytes and Th1 cells on activated endothelium (Fig. 1) (Weber et al. 2001). These results suggest that CCR1 plays a protective role in atherogenesis, possibly by regulating T-cell activation and the trafficking of monocytes and T cells into the vessel wall.

CXCR4

CXCR4 is a receptor only for the chemokine CXCL12 (also known as stromal cell-derived factor-1 α), which has been identified in ECs and SMCs of human atherosclerotic plaques (Abi-Younes et al. 2000). It is well known that the CXCL12–CXCR4 axis plays an important role in hematopoietic stem cell mobilization, organ development and angiogenesis, and recent data suggest that it may also be involved in atherosclerosis (Zernecke and Weber 2010). In particular, plasma levels of CXCL12 in CAD patients were significantly reduced compared to those in healthy controls (Damas et al. 2002) and aged *ApoE*^{-/-} mice had much lower serum and bone marrow levels of Cxcl12 (Xu et al. 2011), indicating that the CXCL12–CXCR4 axis may exert a protective effect in atherogenesis. Consistent with this, chimeric *ApoE*^{-/-} mice transplanted with *Cxcr4*^{-/-} bone marrow showed a marked increase of atherosclerotic

lesions in both the whole aorta and the aortic root (Zernecke et al. 2008). This effect was confirmed by either blocking *Cxcr4* through long-term administration of a specific pharmacologic antagonist, AMD3465, in *ApoE*^{-/-} mice or by repopulating *Ldlr*^{-/-} mice with bone marrow that had been transduced by a recombinant lentivirus encoding a *Cxcr4*-specific “degrakine”, which traps *Cxcr4* in the endoplasmic reticulum (Zernecke et al. 2008). Blocking *Cxcr4* with AMD3465 in *ApoE*^{-/-} mice caused a pronounced leukocytosis and an expansion of circulating neutrophils, and correspondingly the recruitment of neutrophils into the plaques was significantly increased. There was also a significant reduction of SMCs and CD3⁺ T cells in aortic root plaques, implying that the plaques become more vulnerable (Zernecke et al. 2008). Taken together, these results suggest that the CXCL12–CXCR4 axis may play a protective role in atherogenesis possibly by controlling homeostasis of neutrophils and their recruitment to atherosclerotic lesions (Fig. 1).

Controversial Chemokines/Chemokine Receptors in Atherogenesis

CXCL16

CXCL16 (originally named scavenger receptor for phosphatidylserine and oxidized lipoprotein [SR-PSOX]) is the only other chemokine besides CX3CL1 that exists in both membrane-tethered and soluble shed forms, and it is expressed by macrophages, SMCs and T cells in mouse, rabbit and human atherosclerotic plaques (Hofnagel et al. 2011; Minami et al. 2001a, b; Wuttge et al. 2004). The role of CXCL16 in human atherosclerosis is controversial. One study showed that patients with stable angina pectoris and myocardial infarction have significantly lower plasma levels of CXCL16 (Sheikine et al. 2006), whereas several other groups found that patients with angina, stroke or acute coronary syndromes have elevated plasma CXCL16 levels compared with healthy controls (Lehrke et al. 2007; Smith et al. 2008; Sun et al. 2008; Ueland et al. 2012; Wang et al. 2010; Yi and Zeng 2008). In candidate gene analysis, two SNPs within the *CXCL16* gene, rs3744700 and *CXCL16-A181V*, were reported to be independently associated with CAD development and the severity of coronary stenosis (Huang et al. 2010; Lundberg et al. 2005). In *Ldlr*^{-/-} mice, genetic deletion of *Cxcl16* significantly aggravated atherosclerosis and enhanced macrophage recruitment into the plaques (Aslanian and Charo 2006). However, this is in contrast to the deletion of its receptor *Cxcr6* in *ApoE*^{-/-} mice, which resulted in reduced atherosclerosis (Galkina et al. 2007). Conceptually, an atheroprotective role is plausible since CXCL16 is known to function not just as a chemokine but also as a

scavenger receptor for the pro-atherogenic factors, phosphatidylserine and oxLDL (Fig. 1) (Fukumoto et al. 2004; Minami et al. 2001a). Consistent with this, CXCL16 promoted the internalization of oxLDL in human macrophages and *Cxcl16*^{-/-} mouse macrophages have a significant reduction in the capacity to bind and internalize oxLDL (Aslanian and Charo 2006; Barlic et al. 2009). However, a recent study showed that the plasma level of Cxcl16 was much higher in *ApoE*^{-/-} mice receiving a high-fat diet (Yi et al. 2011). Moreover, overexpression of Cxcl16 in *ApoE*^{-/-} mice did not affect the size of existing atherosclerotic lesions but instead promoted their evolution to vulnerable plaques, suggesting that CXCL16 may be an atherogenic marker in plasma (Yi et al. 2011). Considering the limited studies in atherosclerosis mouse models and the contradictory findings in human CAD patients, more work will be needed to clarify the role of CXCL16 in atherogenesis.

CCR7

CCR7 and its ligands, CCL19 and CCL21, have been identified in both mouse and human atherosclerotic lesions, especially in macrophages and T-cell rich areas (Damas et al. 2007). In humans, the plasma levels of CCL19 and CCL21 were found to be increased in patients with stable and unstable angina (Damas et al. 2007). In *Ldlr*^{-/-} mice, genetic deletion of *Ccr7* attenuated the size of atherosclerotic plaques and reduced the macrophage accumulation in those plaques (Luchtefeld et al. 2010). In contrast, *Ccr7* deficiency did not affect the size of atherosclerotic plaques or the plaque macrophage content in *ApoE*^{-/-} mice (Feig et al. 2010; Potteaux et al. 2011). In an atherosclerosis regression mouse model in which a segment of atherosclerotic *ApoE*^{-/-} mouse aortic arch was transplanted into wild-type mice, it was found that *Ccr7* was upregulated in foam cells during the regression of atherosclerotic lesions, accompanied by a significant reduction of macrophage content in the lesions (Trojan et al. 2006). Plaque regression and foam cell content reduction were both greatly inhibited by the blockade of Ccl19 and Ccl21 with specific antibodies, suggesting that *Ccr7* may drive the egress of macrophages from the lesions and have a protective role in atherosclerosis development (Fig. 1) (Trojan et al. 2006). Consistent with this, atorvastatin and rosuvastatin treatment significantly increased the expression of *Ccr7* in lesional macrophages and enhanced the emigration of macrophages from the plaques in *ApoE*^{-/-} mice, while macrophages from *Ccr7*^{-/-}*ApoE*^{-/-} mice failed to emigrate upon statin treatment (Feig et al. 2011a). Also, it was found that liver-X-receptor agonist- and HDL-induced atherosclerotic lesion regression in different atherogenic mouse models are associated with increased expression of

Ccr7 on plaque macrophages (Feig et al. 2010, 2011b; Verschuren et al. 2009). However, in a non-surgical model, it was found that *Ccr7* had no effect on atherosclerotic plaque regression in *ApoE*^{-/-} mice, and the loss of plaque macrophages during disease regression was attributed to suppressed monocyte recruitment (Potteaux et al. 2011). In fact, although expression of CCR7 is increased in atherosclerotic plaques, its expression on circulating T cells was significantly decreased in angina patients, indicating that CCR7 may also be involved in infiltration and egress of T cells from the atherosclerotic vessel wall (Fig. 1) (Damas et al. 2007). Clearly, more studies are necessary to dissect the exact role of CCR7 in the development of atherosclerosis.

Chemokine/Chemokine Receptor Antagonists in Atherogenesis

Considering the above evidence from animal models that chemokines and chemokine receptors may modulate atherogenesis (Fig. 1), it is reasonable to consider them as potential drug targets for the treatment of cardiovascular disease. In the USA, more than 30 % of drugs on the market target G protein-coupled receptors and numerous antagonists of chemokine receptors have been developed (Horuk 2009). We will focus on the antagonists that have been tested in atherosclerosis (Table 2). The difficulty in demonstrating efficacy for these agents is the endpoint that must be considered: clinical events or inflammation of the blood vessel wall. The former is difficult to assess in a short trial, and the latter is difficult to quantitate with precision.

Drugs targeting CCR2 and its ligand CCL2 have been the most extensively evaluated in preclinical and clinical studies. In a phase II clinical trial of patients at risk for atherosclerotic cardiovascular disease, **MLN1202** (Millennium), a specific humanized monoclonal antibody directed against CCR2, was found to significantly reduce the serum levels of C-reactive protein, a surrogate marker of inflammation in cardiovascular disease (Gilbert et al. 2011). **CCX140** (Chemocentryx) is a small molecule inhibitor of CCR2 that has successfully completed a phase II clinical trial in type 2 diabetes, but has not yet been tested in cardiovascular disease (Koenen and Weber 2011). The CCR2 small molecule antagonists, **INCB-3344** and **GSK1344386B** (Glaxo-Smith-Kline), did not affect the size of atherosclerotic lesions in *ApoE*^{-/-} mice, even though they markedly reduced the number of circulating inflammatory monocytes and macrophage content in plaques (Aiello et al. 2010; Olzinski et al. 2010). As for CCL2 antagonists, **11K2** (an inhibitory antibody against CCL2 and CCL12; Biogen Idec Inc.) treatment of *ApoE*^{-/-} mice significantly reduced both atherosclerotic plaque formation

Table 2 Antagonists to inhibit chemokines/chemokine receptors in atherogenesis

Chemokines/chemokine receptors	Antagonists	Atherosclerotic lesion change after treatment	References
CCR2	GSK1344386B	No reduction in mice	Olzinski et al. (2010)
	INCB-3344	No reduction in mice	Aiello et al. (2010)
	MLN1202	Reduced plasma CRP levels in patients	Gilbert et al. (2011)
CCL2	11K2	Reduced in mice	Lutgens et al. (2005)
	PA508	Reduced in mice	Liehn et al. (2010)
CCR5/CXCR3	TAK-779	Reduced in mice	van Wanrooij et al. (2005)
CCL5	Met-RANTES	Reduced in mice	Combadiere et al. (2008) and Veillard et al. (2004)
	[⁴⁴ AANA ⁴⁷]-CCL5	Reduced in mice	Braunersreuther et al. (2008)
CCL5/CXCL4	MKEY	Reduced in mice	Koenen et al. (2009)
CXCR2	SB-517785-M	Reduced arteriolar leukocyte recruitment in rats	Nabah et al. (2007)
CXCR3	NBI-74330	Reduced in mice	van Wanrooij et al. (2008)
CXCR4	AMD-3465	Increased in mice	Zernecke et al. (2008)
Chemokines	Evasin-3	Reduced post-ischemic infarct size in mice	Montecucco et al. (2010)
	M-T7	Reduced intimal hyperplasia and aortic allograft vasculopathy in rats and mice	Dai et al. (2010) and Liu et al. (2000, 2004)
	NR58-3.14.3	No reduction in mice	Reckless et al. (2005)
	35 K	Reduced in mice	Ali et al. (2005) and Bursill et al. (2004)

and lesional macrophage content (Lutgens et al. 2005). Treatment of *ApoE*^{-/-} mice with **PA508**, a modified CCL2 form with increased glycosaminoglycan binding activity but reduced affinity for CCR2, markedly reduced neointimal plaque formation after arterial injury, due to reduced inflammatory monocyte recruitment to the lesions (Liehn et al. 2010). A humanized monoclonal antibody against CCL2, **ABN-912** (Novartis), has been tested in patients with rheumatoid arthritis, but the treatment did not show any benefit and whether it may affect atherosclerosis still remains unknown (Haringman et al. 2006).

CCR5 and its ligand CCL5 is another chemokine/chemokine receptor pair that has been targeted in atherogenesis. **TAK-779** is a small molecule antagonist for both CCR5 and CXCR3 that was originally developed as an HIV entry inhibitor (Baba et al. 1999; Gao et al. 2003). TAK-779 treatment of *Ldlr*^{-/-} mice significantly reduced atherosclerotic plaque formation and Th1 cell infiltration into the lesions (van Wanrooij et al. 2005). **Maraviroc** (Pfizer) is a small molecule CCR5 antagonist approved by the FDA for HIV treatment and **HGS004/HGS101** (antibodies raised against CCR5; Human Genome Sciences) has also been used to treat HIV in clinical trials (Latinovic et al. 2011), but whether they can affect atherosclerosis is

still unclear. Treatment of *Ldlr*^{-/-} mice with Met-RANTES, an *N*-methionylated variant of CCL5 with potent CCR5 antagonist activity (Proudfoot et al. 1996), reduced the size of atherosclerotic lesions in both the whole aorta and the aortic root, accompanied by increased plaque stability (Veillard et al. 2004). Met-RANTES treatment of *ApoE*^{-/-}*Ccl2*^{-/-}*Cx3cr1*^{-/-} triple knockout mice induced a further decrease of atherosclerotic plaque size, suggesting that it might also be considered alone or in combination with other chemokine receptor antagonists to treat atherosclerosis (Combadiere et al. 2008). Another antagonist for CCL5 is [⁴⁴AANA⁴⁷]-CCL5, a CCL5 variant with specific mutations in the principal RANTES/GAG binding site, that was shown to prevent the progression of established atherosclerotic lesions in *Ldlr*^{-/-} mice (Braunersreuther et al. 2008). **MKEY** (Carolus Therapeutics Inc.) is a synthetic peptide designed to disrupt the heteromerization of CCL5 and CXCL4, and administration of MKEY into *ApoE*^{-/-} mice markedly reduced the formation of atherosclerotic lesions as well as macrophage accumulation in plaque (Koenen et al. 2009).

Antagonists against CXCR2, CXCR3 and CXCR4 have also been developed and tested in atherosclerosis models. **SB-517785-M** (Glaxo-Smith-Kline) is an antagonist that

selectively inhibits CXCR2. This agent significantly reduced angiotensin II-induced infiltration of mononuclear cells and neutrophils into rat arterioles, suggesting that it may inhibit atherogenesis by controlling recruitment of neutrophils into the vessel wall (Drechsler et al. 2011; Nabah et al. 2007). Treatment of *Ldlr*^{-/-} mice with **NBI-74330**, a CXCR3 antagonist, markedly increased the content of Tregs in atherosclerotic plaques but reduced effector T cells in lymph nodes draining the aortic arch, thus attenuating atherosclerotic lesion formation in these mice (van Wanrooij et al. 2008). As discussed in the previous section, CXCR4 blockade by its antagonist **AMD3465** led to aggravated atherosclerotic lesion formation in *ApoE*^{-/-} mice, probably due to increased neutrophil expansion in the blood and their recruitment into plaque (Zernecke et al. 2008). **AMD3100** (Plerixafor; Genzyme) is a CXCR4 antagonist that has been approved by the FDA for stem cell mobilization for autologous transplantation in the setting of ablative chemotherapy for multiple myeloma and non-Hodgkin's lymphoma, but its role in atherogenesis is still unknown (Keating 2011).

Human CX3CL1 has been modified at the N-terminus to produce a CX3CR1 antagonist named **F1** (Dorgham et al. 2009). F1 did not induce signaling through CX3CR1 but potently inhibited CX3CL1-induced chemotaxis, calcium flux and cell adhesion. In a peritonitis mouse model, F1 was found to strongly inhibit macrophage accumulation, indicating that it has anti-inflammatory activity (Dorgham et al. 2009). However, it is still not clear whether this antagonist may affect the progress of atherosclerosis.

There are also some naturally occurring chemokine-blocking agents that target more than one chemokine. For example, Evasins are a group of multipotential chemokine binding proteins found in tick saliva. Evasin-1 binds to CCL3, CCL4, and CCL18; Evasin-3 binds to CXCL1 and CXCL8; and Evasin-4 binds to CCL5 and CCL11 (Deruaz et al. 2008). A single administration of **Evasin-3** significantly reduced post-ischemic infarct size during myocardial ischemia in C57BL/6 mice, associated with reduced infiltration of neutrophils into the site of injury and decreased production of ROS (Montecucco et al. 2010). These results suggest that Evasins have anti-inflammatory functions that may be useful for the treatment of atherosclerosis. **M-T7** is a 37-kDa glycoprotein that also binds to a broad range of C, CC and CXC chemokines through the conserved C-terminal GAG binding domain of the chemokines (Lalani et al. 1997). Administration of purified M-T7 into rats caused a significant reduction of intimal hyperplasia after angioplasty injury (Liu et al. 2000) and inhibited aortic allograft vasculopathy associated with reduced inflammatory cell invasion (Dai et al. 2010; Liu et al. 2004), indicating that it may prevent recurrent atherosclerotic plaque growth. **NR58-3.14.3** is a broad-

spectrum chemokine-blocking peptide that effectively inhibits the activities of CCL2, CCL3, CXCL8 and CXCL12, and treatment of *ApoE*^{-/-} mice with this peptide significantly reduced macrophage accumulation in vascular lesions and increased the content of collagen and SMCs, although the vascular lipid lesion area was not changed (Reckless et al. 2005). Also, tail-vein injection of a recombinant adenovirus encoding soluble protein **35K**, a broad-spectrum CC-chemokine blocking agent encoded by Vaccinia virus, into *ApoE*^{-/-} mice markedly reduced atherosclerotic plaque formation and atherosclerosis in carotid-caval vein grafts, accompanied by decreased macrophage recruitment into the lesions (Ali et al. 2005; Bursill et al. 2004). These results have suggested that broad-spectrum chemokine-blocking peptides might be useful to inhibit the progress of atherosclerosis and promote the stabilization of atherosclerotic plaques.

Recent studies suggest that statins (HMG-CoA reductase inhibitors), a class of drugs used for the primary and secondary prevention of CAD, may also affect the expression of chemokines and chemokine receptors during atherosclerosis (Feig et al. 2011a; Waehre et al. 2003). Statins are known to lower cholesterol levels in hypercholesterolemic patients and it has been shown that normalization of plasma lipid levels may cause atherosclerosis regression in both humans and atherosclerosis-prone mouse models (Nissen et al. 2006; Trogan et al. 2006). The regression of atherosclerosis is accompanied by reduced plaque inflammatory cell content (e.g., macrophages), which may be due to increased efflux of leukocytes from plaques (Trogan et al. 2006) and decreased influx of inflammatory cells into plaques (Potteaux et al. 2011). The increased efflux and reduced influx of leukocytes are associated with increased expression of Ccr7 and reduced expression of Ccl2, respectively (Lieu et al. 2003; Trogan et al. 2006). In support of this, statin treatment significantly increased the expression of Ccr7 in lesional macrophages and enhanced their emigration from the plaques of *ApoE*^{-/-} mice (Feig et al. 2011a). In vitro statin treatment significantly inhibited the expression of CCL2 in peripheral blood mononuclear cell (PBMC), ECs and monocytes/macrophages (Jougasaki et al. 2010; Morikawa et al. 2002; Romano et al. 2000; Veillard et al. 2006). In vivo simvastatin treatment reduced expression of CCL2 and CCR2 on circulating monocytes in both hypercholesterolemic patients and healthy people (Han et al. 2005; Rezaie-Majd et al. 2002); both atorvastatin and simvastatin significantly inhibited the expression of CCR2 and CX3CR1 on PBMCs in CAD patients (Damas et al. 2005; Waehre et al. 2003). In addition, statin treatment has been reported to downregulate the expression of CCR1, CCL3, CCL4, CCL19, CCL21, CXCL8, CXCL16 and CX3CL1 in different studies (Damas et al. 2005, 2007; Smith et al. 2008; Waehre et al. 2003),

suggesting that statins may have a broad effect on the expression of atherogenic chemokines during the reversal of hypercholesterolemia in cardiovascular patients. However, not all patients can benefit from using statins and as many as 20 % of the patients who use statins develop adverse effects such as muscle fatigue and weakness and impaired cognition (Maningat and Breslow 2011). New drugs are needed to treat those statin-intolerant patients and chemokine and chemokine receptor antagonists are worth considering in this setting, either as prevention or in combination with statins in treatment of atherosclerosis.

Conclusions

Atherosclerosis is the leading cause of mortality worldwide. Although many drugs in several classes are approved for the treatment of cardiovascular disease, so far only statins target atherosclerosis by lowering the cholesterol level. However, not all patients benefit from statins, and new therapeutics are needed. In the past decade, numerous studies in mice and humans have suggested that chemokines and their corresponding G protein-coupled chemokine receptors play important roles in the pathophysiology of atherosclerosis development. For example, CCL2–CCR2, CCL5–CCR5 and CX3CL1–CX3CR1 are found to be pathogenic whereas CCR1 and CXCR4 are protective, indicating that antagonists for CCR2, CCR5, CX3CR1 and agonists for CCR1 and CXCR4 may be beneficial for therapeutic intervention. However, although more than 30 % of the drugs on the USA market are targeted against G protein-coupled receptors and many chemokine receptor antagonists have been tested in different mouse models and clinical trials, only two small molecule antagonists of chemokine receptors have been approved by the FDA: the CCR5 antagonist Maraviroc for HIV treatment and the CXCR4 antagonist AMD3100 (Plerixafor) for multiple myeloma and non-Hodgkin's lymphoma. Despite substantial progress in identifying chemokine system targets and developing lead compounds, much more research will be necessary to define the cellular mechanisms by which they operate as well as to evaluate current drug candidates and identify new ones. Also, it would be necessary to evaluate the effect of these new drugs on the immune system and antimicrobial host defense during long-term use.

Acknowledgments This work was supported by the Intramural Research Program of the Division of Intramural Research, National Institute of Allergy and Infectious Diseases, National Institutes of Health (Bethesda, MD, USA).

Conflict of interest None.

Note added in proof We have recently reported that Ccr7-deficient ApoE knockout mice fed a Western diet have markedly increased atherosclerotic plaque. This effect was due to Ccr7 expression on a bone marrow-derived cell and was associated with increased lesional and blood T cells. (Reference: Wan W, Lionakis MS, Liu Q et al (2012) Genetic deletion of chemokine receptor Ccr7 exacerbates atherogenesis in ApoE-deficient mice. *Cardiovasc Res* PMID: 23180724).

References

- Abi-Younes S, Sauty A, Mach F et al (2000) The stromal cell-derived factor-1 chemokine is a potent platelet agonist highly expressed in atherosclerotic plaques. *Circ Res* 86:131–138
- Aiello RJ, Bourassa PA, Lindsey S et al (1999) Monocyte chemoattractant protein-1 accelerates atherosclerosis in apolipoprotein E-deficient mice. *Arterioscler Thromb Vasc Biol* 19:1518–1525
- Aiello RJ, Perry BD, Bourassa PA et al (2010) CCR2 receptor blockade alters blood monocyte subpopulations but does not affect atherosclerotic lesions in apoE(−/−) mice. *Atherosclerosis* 208:370–375
- Ali ZA, Bursill CA, Hu Y et al (2005) Gene transfer of a broad spectrum CC-chemokine inhibitor reduces vein graft atherosclerosis in apolipoprotein E-knockout mice. *Circulation* 112:1235–1241
- Apostolakis S, Krambovitis E, Vlata Z et al (2007) CX3CR1 receptor is up-regulated in monocytes of coronary artery diseased patients: impact of pre-inflammatory stimuli and renin-angiotensin system modulators. *Thromb Res* 121:387–395
- Apostolopoulos J, Davenport P, Tipping PG (1996) Interleukin-8 production by macrophages from atheromatous plaques. *Arterioscler Thromb Vasc Biol* 16:1007–1012
- Aslanian AM, Charo IF (2006) Targeted disruption of the scavenger receptor and chemokine CXCL16 accelerates atherosclerosis. *Circulation* 114:583–590
- Baba M, Nishimura O, Kanzaki N et al (1999) A small-molecule, nonpeptide CCR5 antagonist with highly potent and selective anti-HIV-1 activity. *Proc Natl Acad Sci USA* 96:5698–5703
- Barlic J, Murphy PM (2007a) Chemokine regulation of atherosclerosis. *J Leukoc Biol* 82:226–236
- Barlic J, Murphy PM (2007b) An oxidized lipid-peroxisome proliferator-activated receptor gamma-chemokine pathway in the regulation of macrophage-vascular smooth muscle cell adhesion. *Trends Cardiovasc Med* 17:269–274
- Barlic J, Zhu W, Murphy PM (2009) Atherogenic lipids induce high-density lipoprotein uptake and cholesterol efflux in human macrophages by up-regulating transmembrane chemokine CXCL16 without engaging CXCL16-dependent cell adhesion. *J Immunol* 182:7928–7936
- Bazan JF, Bacon KB, Hardiman G et al (1997) A new class of membrane-bound chemokine with a CX3C motif. *Nature* 385:640–644
- Berger JS, Jordan CO, Lloyd-Jones D et al (2010) Screening for cardiovascular risk in asymptomatic patients. *J Am Coll Cardiol* 55:1169–1177
- Boger CA, Fischereder M, Deinzer M et al (2005) RANTES gene polymorphisms predict all-cause and cardiac mortality in type 2 diabetes mellitus hemodialysis patients. *Atherosclerosis* 183: 121–129
- Boisvert WA, Santiago R, Curtiss LK et al (1998) A leukocyte homologue of the IL-8 receptor CXCR-2 mediates the accumulation of macrophages in atherosclerotic lesions of LDL receptor-deficient mice. *J Clin Invest* 101:353–363

- Boisvert WA, Curtiss LK, Terkeltaub RA (2000) Interleukin-8 and its receptor CXCR2 in atherosclerosis. *Immunol Res* 21:129–137
- Boisvert WA, Rose DM, Johnson KA et al (2006) Up-regulated expression of the CXCR2 ligand KC/GRO- α in atherosclerotic lesions plays a central role in macrophage accumulation and lesion progression. *Am J Pathol* 168:1385–1395
- Boring L, Gosling J, Chensue SW et al (1997) Impaired monocyte migration and reduced type 1 (Th1) cytokine responses in C–C chemokine receptor 2 knockout mice. *J Clin Invest* 100:2552–2561
- Boring L, Gosling J, Cleary M et al (1998) Decreased lesion formation in CCR2^{-/-} mice reveals a role for chemokines in the initiation of atherosclerosis. *Nature* 394:894–897
- Braunersreuther V, Zernecke A, Steffens S et al (2007) Ccr5 but not Ccr1 deficiency reduces development of diet-induced atherosclerosis in mice. *Arterioscler Thromb Vasc Biol* 27:373–379
- Braunersreuther V, Steffens S, Arnaud C et al (2008) A novel RANTES antagonist prevents progression of established atherosclerotic lesions in mice. *Arterioscler Thromb Vasc Biol* 28:1090–1096
- Breland UM, Halvorsen B, Hol J et al (2008) A potential role of the CXC chemokine GRO α in atherosclerosis and plaque destabilization: downregulatory effects of statins. *Arterioscler Thromb Vasc Biol* 28:1005–1011
- Bursill CA, Choudhury RP, Ali Z et al (2004) Broad-spectrum CC-chemokine blockade by gene transfer inhibits macrophage recruitment and atherosclerotic plaque formation in apolipoprotein E-knockout mice. *Circulation* 110:2460–2466
- Butcher M, Galkina E (2011) Current views on the functions of interleukin-17A-producing cells in atherosclerosis. *Thromb Haemost* 106:787–795
- Calabresi PA, Yun SH, Allie R et al (2002) Chemokine receptor expression on MBP reactive T cells: CXCR6 is a marker of IFN γ -producing effector cells. *J Neuroimmunol* 127:96–105
- Calvayrac O, Rodríguez-Calvo R, Alonso J et al (2011) CCL20 is increased in hypercholesterolemic subjects and is upregulated by LDL in vascular smooth muscle cells: role of NF- κ B. *Arterioscler Thromb Vasc Biol* 31:2733–2741
- Combadiere C, Potteaux S, Gao JL et al (2003) Decreased atherosclerotic lesion formation in CX3CR1/apolipoprotein E double knockout mice. *Circulation* 107:1009–1016
- Combadiere C, Potteaux S, Rodero M et al (2008) Combined inhibition of CCL2, CX3CR1, and CCR5 abrogates Ly6C(hi) and Ly6C(lo) monocytosis and almost abolishes atherosclerosis in hypercholesterolemic mice. *Circulation* 117:1649–1657
- Dai E, Liu LY, Wang H et al (2010) Inhibition of chemokine-glycosaminoglycan interactions in donor tissue reduces mouse allograft vasculopathy and transplant rejection. *PLoS One* 5:e10510
- Damas JK, Waehre T, Yndestad A et al (2002) SDF-1 α in unstable angina: potential antiinflammatory and matrix-stabilizing effects. *Circulation* 106:36–42
- Damas JK, Boullier A, Waehre T et al (2005) Expression of fractalkine (CX3CL1) and its receptor, CX3CR1, is elevated in coronary artery disease and is reduced during statin therapy. *Arterioscler Thromb Vasc Biol* 25:2567–2572
- Damas JK, Smith C, Oie E et al (2007) Enhanced expression of the homeostatic chemokines CCL19 and CCL21 in clinical and experimental atherosclerosis: possible pathogenic role in plaque destabilization. *Arterioscler Thromb Vasc Biol* 27:614–620
- Daoudi M, Lavergne E, Garin A et al (2004) Enhanced adhesive capacities of the naturally occurring Ile249-Met280 variant of the chemokine receptor CX3CR1. *J Biol Chem* 279:19649–19657
- Dawson TC, Kuziel WA, Osahar TA et al (1999) Absence of CC chemokine receptor-2 reduces atherosclerosis in apolipoprotein E-deficient mice. *Atherosclerosis* 143:205–211
- de Waard V, Bot I, de Jager SC et al (2010) Systemic MCP1/CCR2 blockade and leukocyte specific MCP1/CCR2 inhibition affect aortic aneurysm formation differently. *Atherosclerosis* 211:84–89
- Deruaz M, Frauenschuh A, Alessandri AL et al (2008) Ticks produce highly selective chemokine binding proteins with antiinflammatory activity. *J Exp Med* 205:2019–2031
- Doran AC, Lipinski MJ, Oldham SN et al (2012) B-cell aortic homing and atheroprotection depend on Id3. *Circ Res* 110:e1–e12
- Dorgham K, Ghadiri A, Hermand P et al (2009) An engineered CX3CR1 antagonist endowed with anti-inflammatory activity. *J Leukoc Biol* 86:903–911
- Drechsler M, Megens RT, van Zandvoort M et al (2010) Hyperlipidemia-triggered neutrophilia promotes early atherosclerosis. *Circulation* 122:1837–1845
- Drechsler M, Döring Y, Megens RT et al (2011) Neutrophilic granulocytes—promiscuous accelerators of atherosclerosis. *Thromb Haemost* 106:839–848
- Feig JE, Pineda-Torra I, Sanson M et al (2010) LXR promotes the maximal egress of monocyte-derived cells from mouse aortic plaques during atherosclerosis regression. *J Clin Invest* 120:4415–4424
- Feig JE, Shang Y, Rotllan N et al (2011a) Statins promote the regression of atherosclerosis via activation of the CCR7-dependent emigration pathway in macrophages. *PLoS One* 6:e28534
- Feig JE, Rong JX, Shamir R et al (2011b) HDL promotes rapid atherosclerosis regression in mice and alters inflammatory properties of plaque monocyte-derived cells. *Proc Natl Acad Sci USA* 108:7166–7171
- Fernandes JL, Mamoni RL, Orford JL et al (2004) Increased Th1 activity in patients with coronary artery disease. *Cytokine* 26:131–137
- Fukumoto N, Shimaoka T, Fujimura H et al (2004) Critical roles of CXC chemokine ligand 16/scavenger receptor that binds phosphatidylserine and oxidized lipoprotein in the pathogenesis of both acute and adoptive transfer experimental autoimmune encephalomyelitis. *J Immunol* 173:1620–1627
- Galkina E, Ley K (2009) Immune and inflammatory mechanisms of atherosclerosis. *Annu Rev Immunol* 27:165–197
- Galkina E, Harry BL, Ludwig A et al (2007) CXCR6 promotes atherosclerosis by supporting T-cell homing, interferon- γ production, and macrophage accumulation in the aortic wall. *Circulation* 116:1801–1811
- Gao P, Zhou XY, Yashiro-Ohtani Y et al (2003) The unique target specificity of a nonpeptide chemokine receptor antagonist: selective blockade of two Th1 chemokine receptors CCR5 and CXCR3. *J Leukoc Biol* 73:273–280
- Ghilardi G, Biondi ML, Turri O et al (2004) Internal carotid artery occlusive disease and polymorphisms of fractalkine receptor CX3CR1: a genetic risk factor. *Stroke* 35:1276–1279
- Gilbert J, Lekstrom-Himes J, Donaldson D et al (2011) Effect of CC chemokine receptor 2 CCR2 blockade on serum C-reactive protein in individuals at atherosclerotic risk and with a single nucleotide polymorphism of the monocyte chemoattractant protein-1 promoter region. *Am J Cardiol* 107:906–911
- Gonzalez P, Alvarez R, Batalla A et al (2001) Genetic variation at the chemokine receptors CCR5/CCR2 in myocardial infarction. *Genes Immun* 2:191–195
- Gosling J, Slaymaker S, Gu L et al (1999) MCP-1 deficiency reduces susceptibility to atherosclerosis in mice that overexpress human apolipoprotein B. *J Clin Invest* 103:773–778
- Greaves DR, Hakkinen T, Lucas AD et al (2004) Linked chromosome 16q13 chemokines, macrophage-derived chemokine, fractalkine, and thymus- and activation-regulated chemokine, are expressed in human atherosclerotic lesions. *Arterioscler Thromb Vasc Biol* 21:923–929

- Gu L, Okada Y, Clinton SK et al (1998) Absence of monocyte chemoattractant protein-1 reduces atherosclerosis in low density lipoprotein receptor-deficient mice. *Mol Cell* 2:275–281
- Gugl A, Renner W, Seino G et al (2003) Two polymorphisms in the fractalkine receptor CX3CR1 are not associated with peripheral arterial disease. *Atherosclerosis* 166:339–343
- Guo J, Van Eck M, Twisk J et al (2003) Transplantation of monocyte CC-chemokine receptor 2-deficient bone marrow into ApoE3-Leiden mice inhibits atherogenesis. *Arterioscler Thromb Vasc Biol* 23:447–453
- Guo J, de Waard V, Van Eck M et al (2005) Repopulation of apolipoprotein E knockout mice with CCR2-deficient bone marrow progenitor cells does not inhibit ongoing atherosclerotic lesion development. *Arterioscler Thromb Vasc Biol* 25:1014–1019
- Han KH, Ryu J, Hong KH et al (2005) HMG-CoA reductase inhibition reduces monocyte CC chemokine receptor 2 expression and monocyte chemoattractant protein-1-mediated monocyte recruitment in vivo. *Circulation* 111:1439–1447
- Hansson GK (2005) Inflammation, atherosclerosis, and coronary artery disease. *N Engl J Med* 352:1685–1695
- Haringman JJ, Gerlag DM, Smeets TJ et al (2006) A randomized controlled trial with an anti-CCL2 (anti-monocyte chemotactic protein 1) monoclonal antibody in patients with rheumatoid arthritis. *Arthritis Rheum* 54:2387–2392
- Heller EA, Liu E, Tager AM et al (2006) Chemokine CXCL10 promotes atherogenesis by modulating the local balance of effector and regulatory T cells. *Circulation* 113:2301–2312
- Hofnagel O, Engel T, Severs NJ et al (2011) SR-PSOX at sites predisposed to atherosclerotic lesion formation mediates monocyte–endothelial cell adhesion. *Atherosclerosis* 217:371–378
- Horuk R (2009) Chemokine receptor antagonists: overcoming developmental hurdles. *Nat Rev Drug Discov* 8:23–33
- Huang M, Han Y, Zhang X et al (2010) An intron polymorphism in the CXCL16 gene is associated with increased risk of coronary artery disease in Chinese Han population: a large angiography-based study. *Atherosclerosis* 210:160–165
- Huo Y, Weber C, Forlow SB et al (2001) The chemokine KC, but not monocyte chemoattractant protein-1, triggers monocyte arrest on early atherosclerotic endothelium. *J Clin Invest* 108:1307–1314
- Imai T, Baba M, Nishimura M et al (1997a) The T cell-directed CC chemokine TARC is a highly specific biological ligand for CC chemokine receptor 4. *J Biol Chem* 272:15036–15042
- Imai T, Hieshima K, Haskell C et al (1997b) Identification and molecular characterization of fractalkine receptor CX3CR1, which mediates both leukocyte migration and adhesion. *Cell* 91:521–530
- Inoue S, Egashira K, Ni W et al (2002) Anti-monocyte chemoattractant protein-1 gene therapy limits progression and destabilization of established atherosclerosis in apolipoprotein E-knockout mice. *Circulation* 106:2700–2706
- Jougasaki M, Ichiki T, Takenoshita Y et al (2010) Statins suppress interleukin-6-induced monocyte chemo-attractant protein-1 by inhibiting Janus kinase/signal transducers and activators of transcription pathways in human vascular endothelial cells. *Br J Pharmacol* 159:1294–1303
- Kawamura A, Miura S, Fujino M et al (2003) CXCR3 chemokine receptor–plasma IP10 interaction in patients with coronary artery disease. *Circ J* 67:851–854
- Keating GM (2011) Plerixafor: a review of its use in stem-cell mobilization in patients with lymphoma or multiple myeloma. *Drugs* 71:1623–1647
- King VL, Lin AY, Kristo F et al (2009) Interferon-gamma and the interferon-inducible chemokine CXCL10 protect against aneurysm formation and rupture. *Circulation* 119:426–435
- Koenen RR, Weber C (2011) Chemokines: established and novel targets in atherosclerosis. *EMBO Mol Med* 3:713–725
- Koenen RR, von Hundelshausen P, Nesmelova IV et al (2009) Disrupting functional interactions between platelet chemokines inhibits atherosclerosis in hyperlipidemic mice. *Nat Med* 15:97–103
- Krohn R, Raffetseder U, Bot I et al (2007) Y-box binding protein-1 controls CC chemokine ligand-5 (CCL5) expression in smooth muscle cells and contributes to neointima formation in atherosclerosis-prone mice. *Circulation* 116:1812–1820
- Kuziel WA, Dawson TC, Quinones M et al (2003) CCR5 deficiency is not protective in the early stages of atherogenesis in apoE knockout mice. *Atherosclerosis* 167:25–32
- Kyaw T, Tipping P, Toh BH et al (2011) Current understanding of the role of B cell subsets and intimal and adventitial B cells in atherosclerosis. *Curr Opin Lipidol* 22:373–379
- Lalani AS, Graham K, Mossman K et al (1997) The purified myxoma virus gamma interferon receptor homolog M-T7 interacts with the heparin-binding domains of chemokines. *J Virol* 71:4356–4363
- Landsman L, Bar-On L, Zernecke A et al (2009) CX3CR1 is required for monocyte homeostasis and atherogenesis by promoting cell survival. *Blood* 113:963–972
- Latinovic O, Reitz M, Le NM et al (2011) CCR5 antibodies HGS004 and HGS101 preferentially inhibit drug-bound CCR5 infection and restore drug sensitivity of Maraviroc-resistant HIV-1 in primary cells. *Virology* 411:32–40
- Lehrke M, Millington SC, Lefterova M et al (2007) CXCL16 is a marker of inflammation, atherosclerosis, and acute coronary syndromes in humans. *J Am Coll Cardiol* 49:442–449
- Lei ZB, Zhang Z, Jing Q et al (2002) OxLDL upregulates CXCR2 expression in monocytes via scavenger receptors and activation of p38 mitogen-activated protein kinase. *Cardiovasc Res* 53:524–532
- Lesnik P, Haskell CA, Charo IF (2003) Decreased atherosclerosis in CX3CR1^{-/-} mice reveals a role for fractalkine in atherogenesis. *J Clin Invest* 111:333–340
- Ley K, Laudanna C, Cybulsky MI et al (2007) Getting to the site of inflammation: the leukocyte adhesion cascade updated. *Nat Rev Immunol* 7:678–689
- Liehn EA, Piccinini AM, Koenen RR et al (2010) A new monocyte chemotactic protein-1/chemokine CC motif ligand-2 competitor limiting neointima formation and myocardial ischemia/reperfusion injury in mice. *J Am Coll Cardiol* 56:1847–1857
- Lieu HD, Withycombe SK, Walker Q et al (2003) Eliminating atherogenesis in mice by switching off hepatic lipoprotein secretion. *Circulation* 107:1315–1321
- Liu L, Lalani A, Dai E et al (2000) The viral anti-inflammatory chemokine-binding protein M-T7 reduces intimal hyperplasia after vascular injury. *J Clin Invest* 105:1613–1621
- Liu L, Dai E, Miller L et al (2004) Viral chemokine-binding proteins inhibit inflammatory responses and aortic allograft transplant vasculopathy in rat models. *Transplantation* 77:1652–1660
- Liu P, Yu YR, Spencer JA et al (2008) CX3CR1 deficiency impairs dendritic cell accumulation in arterial intima and reduces atherosclerotic burden. *Arterioscler Thromb Vasc Biol* 28:243–250
- Lucas AD, Bursill C, Guzik TJ et al (2003) Smooth muscle cells in human atherosclerotic plaques express the fractalkine receptor CX3CR1 and undergo chemotaxis to the CX3C chemokine fractalkine (CX3CL1). *Circulation* 108:2498–2504
- Luchtefeld M, Grothusen C, Gagalick A et al (2010) Chemokine receptor 7 knockout attenuates atherosclerotic plaque development. *Circulation* 122:1621–1628
- Lundberg GA, Kellin A, Samnegard A et al (2005) Severity of coronary artery stenosis is associated with a polymorphism in the CXCL16/SR-PSOX gene. *J Intern Med* 257:415–422

- Lutgens E, Faber B, Schapira K et al (2005) Gene profiling in atherosclerosis reveals a key role for small inducible cytokines: validation using a novel monocyte chemoattractant protein monoclonal antibody. *Circulation* 111:3443–3452
- Mach F, Sauty A, Iarossi AS et al (1999) Differential expression of three T lymphocyte-activating CXC chemokines by human atheroma-associated cells. *J Clin Invest* 104:1041–1050
- Maningat P, Breslow JL (2011) Needed: pragmatic clinical trials for statin-intolerant patients. *N Engl J Med* 365:2250–2251
- McDermott DH, Halcox JPI, Schenke WH et al (2001) Association between polymorphism in the chemokine receptor CX3CR1 and coronary vascular endothelial dysfunction and atherosclerosis. *Circ Res* 89:401–407
- McDermott DH, Fong AM, Yang Q et al (2003) Chemokine receptor mutant CX3CR1-M280 has impaired adhesive function and correlates with protection from cardiovascular disease in humans. *J Clin Invest* 111:1241–1250
- McDermott DH, Yang Q, Kathiresan S et al (2005) CCL2 polymorphisms are associated with serum monocyte chemoattractant protein-1 levels and myocardial infarction in the Framingham Heart Study. *Circulation* 112:1113–1120
- Minami M, Kume N, Shimaoka T et al (2001a) Expression of SR-PSOX, a novel cell-surface scavenger receptor for phosphatidylserine and oxidized LDL in human atherosclerotic lesions. *Arterioscler Thromb Vasc Biol* 21:1796–1800
- Minami M, Kume N, Shimaoka T et al (2001b) Expression of scavenger receptor for phosphatidylserine and oxidized lipoprotein (SR-PSOX) in human atheroma. *Ann N Y Acad Sci* 947:373–376
- Moatti D, Faure S, Fumeron F et al (2001) Polymorphism in the fractalkine receptor CX3CR1 as a genetic risk factor for coronary artery disease. *Blood* 97:1925–1928
- Montecucco F, Lenglet S, Braunerreuther V et al (2010) Single administration of the CXC chemokine-binding protein Evasin-3 during ischemia prevents myocardial reperfusion injury in mice. *Arterioscler Thromb Vasc Biol* 30:1371–1377
- Morikawa S, Takabe W, Mataka C et al (2002) The effect of statins on mRNA levels of genes related to inflammation, coagulation, and vascular constriction in HUVEC. *J Atheroscler Thromb* 9:178–183
- Murphy PM (2002) International Union of Pharmacology. XXX. Update on chemokine receptor nomenclature. *Pharmacol Rev* 54:227–229
- Murphy PM, Baggiolini M, Charo IF et al (2000) International union of pharmacology. XXII. Nomenclature for chemokine receptors. *Pharmacol Rev* 52:145–176
- Nabah YNA, Losada M, Estelles R et al (2007) CXCR2 blockade impairs angiotensin II induced CC chemokine synthesis and mononuclear leukocyte infiltration. *Arterioscler Thromb Vasc Biol* 27:2370–2376
- Nelken NA, Coughlin SR, Gordon D et al (1991) Monocyte chemoattractant protein-1 in human atheromatous plaques. *J Clin Invest* 88:1121–1127
- Neumann FJ, Ott I, Gawaz M et al (1995) Cardiac release of cytokines and inflammatory responses in acute myocardial infarction. *Circulation* 92:748–755
- Ni W, Egashira K, Kitamoto S et al (2001) New anti-monocyte chemoattractant protein-1 gene therapy attenuates atherosclerosis in apolipoprotein E-knockout mice. *Circulation* 103:2096–2101
- Niessner A, Marculescu R, Kvakan H et al (2005) Fractalkine receptor polymorphisms V2491 and T280M as genetic risk factors for restenosis. *Thromb Haemost* 94:1251–1256
- Nissen SE, Nicholls SJ, Sipahi I et al (2006) Effect of very high-intensity statin therapy on regression of coronary atherosclerosis: the ASTEROID trial. *JAMA* 295:1556–1565
- Norata GD, Garlaschelli K, Ongari M et al (2006) Effects of fractalkine receptor variants on common carotid artery intima-media thickness. *Stroke* 37:1558–1561
- Olzinski AR, Turner GH, Bernard RE et al (2010) Pharmacological inhibition of C–C chemokine receptor 2 decreases macrophage infiltration in the aortic root of the human C–C chemokine receptor 2/apolipoprotein E–/– mouse: magnetic resonance imaging assessment. *Arterioscler Thromb Vasc Biol* 30:253–259
- Ortlepp JR, Vesper K, Mevissen V et al (2003) Chemokine receptor (CCR2) genotype is associated with myocardial infarction and heart failure in patients under 65 years of age. *J Mol Med* 81:363–367
- Packard RR, Lichtman AH, Libby P (2009) Innate and adaptive immunity in atherosclerosis. *Semin Immunopathol* 31:5–22
- Papadopoulou C, Corrigan V, Taylor PR et al (2008) The role of the chemokines MCP-1, GRO- α , IL-8 and their receptors in the adhesion of monocytic cells to human atherosclerotic plaques. *Cytokine* 43:181–186
- Pattison JM, Nelson PJ, Huie P et al (1996) RANTES chemokine expression in transplant-associated accelerated atherosclerosis. *J Heart Lung Transplant* 15:1194–1199
- Postea O, Vasina EM, Cauwenberghs S et al (2012) Contribution of platelet CX3CR1 to platelet–monocyte complex formation and vascular recruitment during hyperlipidemia. *Arterioscler Thromb Vasc Biol* 32:1186–1193
- Potteaux S, Combadiere C, Esposito B et al (2005) Chemokine receptor CCR1 disruption in bone marrow cells enhances atherosclerotic lesion development and inflammation in mice. *Mol Med* 11:16–20
- Potteaux S, Combadiere C, Esposito B et al (2006) Role of bone marrow-derived CC-chemokine receptor 5 in the development of atherosclerosis of low-density lipoprotein receptor knockout mice. *Arterioscler Thromb Vasc Biol* 26:1858–1863
- Potteaux S, Gautier EL, Hutchison SB et al (2011) Suppressed monocyte recruitment drives macrophage removal from atherosclerotic plaques of ApoE^{–/–} mice during disease regression. *J Clin Invest* 121:2025–2036
- Proudfoot AE, Power CA, Hoogewerf AJ et al (1996) Extension of recombinant human RANTES by the retention of the initiating methionine produces a potent antagonist. *J Biol Chem* 271:2599–2603
- Quinones MP, Martinez HG, Jimenez F et al (2007) CC chemokine receptor 5 influences late-stage atherosclerosis. *Atherosclerosis* 195:e92–e103
- Rayner K, Van ES, Groot PH et al (2000) Localization of mRNA for JE/MCP-1 and its receptor CCR2 in atherosclerotic lesions of the ApoE knockout mouse. *J Vasc Res* 37:93–102
- Reckless J, Tatalick L, Wilbert S et al (2005) Broad-spectrum chemokine inhibition reduces vascular macrophage accumulation and collagenolysis consistent with plaque stabilization in mice. *J Vasc Res* 42:492–502
- Rezaie-Majid A, Maca T, Bucek RA et al (2002) Simvastatin reduces expression of cytokines interleukin-6, interleukin-8, and monocyte chemoattractant protein-1 in circulating monocytes from hypercholesterolemic patients. *Arterioscler Thromb Vasc Biol* 22:1194–1199
- Robbins CS, Chudnovskiy A, Rauch PJ et al (2012) Extramedullary hematopoiesis generates Ly-6C(high) monocytes that infiltrate atherosclerotic lesions. *Circulation* 125:364–374
- Robertson AK, Hansson GK (2006) T cells in atherogenesis: for better or for worse? *Arterioscler Thromb Vasc Biol* 26:2421–2432
- Roger VL, Go AS, Lloyd-Jones DM et al (2012) Heart disease and stroke statistics—2012 update: a report from the American Heart Association. *Circulation* 125:e2–e220
- Romano M, Diomedea L, Sironi M et al (2000) Inhibition of monocyte chemotactic protein-1 synthesis by statins. *Lab Invest* 80:1095–1100

- Saederup N, Chan L, Lira SA et al (2008) Fractalkine deficiency markedly reduces macrophage accumulation and atherosclerotic lesion formation in CCR2^{-/-} mice: evidence for independent chemokine functions in atherogenesis. *Circulation* 117:1642–1648
- Schutysse E, Struyf S, Van Damme J (2003) The CC chemokine CCL20 and its receptor CCR6. *Cytokine Growth Factor Rev* 14:409–426
- Sheikine Y, Bang CS, Nilsson L et al (2006) Decreased plasma CXCL16/SR-PSOX concentration is associated with coronary artery disease. *Atherosclerosis* 188:462–466
- Simeoni E, Winkelmann BR, Hoffmann MM et al (2004) Association of RANTES G-403A gene polymorphism with increased risk of coronary arteriosclerosis. *Eur Heart J* 25:1438–1446
- Smith C, Halvorsen B, Otterdal K et al (2008) High levels and inflammatory effects of soluble CXC ligand 16 (CXCL16) in coronary artery disease: down-regulatory effects of statins. *Cardiovasc Res* 79:195–203
- Sun Y, Chang Z, Zhang S (2008) Increased serum CXCL16 level is a marker for acute coronary syndromes. *Arch Med Res* 39:332–337
- Swirski FK, Nahrendorf M, Etzrodt M et al (2009) Identification of splenic reservoir monocytes and their deployment to inflammatory sites. *Science* 325:612–616
- Szalai C, Duba J, Prohaszka Z et al (2001) Involvement of polymorphisms in the chemokine system in the susceptibility for coronary artery disease (CAD). Coincidence of elevated Lp(a) and MCP-1 -2518 G/G genotype in CAD patients. *Atherosclerosis* 158:233–239
- Tacke F, Alvarez D, Kaplan TJ et al (2007) Monocyte subsets differentially employ CCR2, CCR5, and CX3CR1 to accumulate within atherosclerotic plaques. *J Clin Invest* 117:185–194
- Teupser D, Pavlides S, Tan M et al (2004) Major reduction of atherosclerosis in fractalkine (CX3CL1)-deficient mice is at the brachiocephalic artery, not the aortic root. *Proc Natl Acad Sci USA* 101:17795–17800
- Trogan E, Feig JE, Dogan S et al (2006) Gene expression changes in foam cells and the role of chemokine receptor CCR7 during atherosclerosis regression in ApoE-deficient mice. *Proc Natl Acad Sci USA* 103:3781–3786
- Tsou CL, Peters W, Si Y et al (2007) Critical roles for CCR2 and MCP-3 in monocyte mobilization from bone marrow and recruitment to inflammatory sites. *J Clin Invest* 117:902–909
- Ueland T, Smedbakken LM, Hallén J et al (2012) Soluble CXCL16 and long-term outcome in acute ischemic stroke. *Atherosclerosis* 220:244–249
- van Wanrooij EJ, Happe H, Hauer AD et al (2005) HIV entry inhibitor TAK-779 attenuates atherogenesis in low-density lipoprotein receptor-deficient mice. *Arterioscler Thromb Vasc Biol* 25:2642–2647
- van Wanrooij EJ, de Jager SC, van Es T et al (2008) CXCR3 antagonist NBI-74330 attenuates atherosclerotic plaque formation in LDL receptor deficient mice. *Arterioscler Thromb Vasc Biol* 28:251–257
- Veillard NR, Kwak B, Pelli G et al (2004) Antagonism of RANTES receptors reduces atherosclerotic plaque formation in mice. *Circ Res* 94:253–261
- Veillard NR, Steffens S, Pelli G et al (2005) Differential influence of chemokine receptors CCR2 and CXCR3 in development of atherosclerosis in vivo. *Circulation* 112:870–878
- Veillard NR, Braunersreuther V, Arnaud C et al (2006) Simvastatin modulates chemokine and chemokine receptor expression by geranylgeranyl isoprenoid pathway in human endothelial cells and macrophages. *Atherosclerosis* 188:51–58
- Verschuren L, de Vries-van der Weij J, Zadelaar S et al (2009) LXR agonist suppresses atherosclerotic lesion growth and promotes lesion regression in apoE*3Leiden mice: time course and mechanisms. *J Lipid Res* 50:301–311
- Waehre T, Damås JK, Gullestad L et al (2003) Hydroxymethylglutaryl coenzyme a reductase inhibitors down-regulate chemokines and chemokine receptors in patients with coronary artery disease. *J Am Coll Cardiol* 41:1460–1467
- Wan W, Murphy PM (2011) Regulation of atherogenesis by chemokine receptor CCR6. *Trends Cardiovasc Med* 21:140–144
- Wan W, Lim JK, Lionakis MS et al (2011) Genetic deletion of chemokine receptor Ccr6 decreases atherogenesis in ApoE-deficient mice. *Circ Res* 109:374–381
- Wang N, Tabas I, Winchester R et al (1996) Interleukin 8 is induced by cholesterol loading of macrophages and expressed by macrophage foam cells in human atheroma. *J Biol Chem* 271:8837–8842
- Wang KD, Liu ZZ, Wang RM et al (2010) Chemokine CXC Ligand 16 serum concentration but not A181V genotype is associated with atherosclerotic stroke. *Clin Chim Acta* 411:1447–1451
- Weber C, Noels H (2011) Atherosclerosis: current pathogenesis and therapeutic options. *Nat Med* 17:1410–1422
- Weber C, Weber KS, Klier C et al (2001) Specialized roles of the chemokine receptors CCR1 and CCR5 in the recruitment of monocytes and T(H)1-like/CD45RO(+) T cells. *Blood* 97:1144–1146
- Weber C, Zernecke A, Libby P (2008) The multifaceted contributions of leukocyte subsets to atherosclerosis: lessons from mouse models. *Nat Rev Immunol* 8:802–815
- Weber C, Meiler S, Döring Y et al (2011) CCL17-expressing dendritic cells drive atherosclerosis by restraining regulatory T cell homeostasis in mice. *J Clin Invest* 121:2898–2910
- Wilcox JN, Nelken NA, Coughlin SR et al (1994) Local expression of inflammatory cytokines in human atherosclerotic plaques. *J Atheroscler Thromb* 1:S10–S13
- Wuttge DM, Zhou X, Sheikine Y et al (2004) CXCL16/SR-PSOX is an interferon-gamma-regulated chemokine and scavenger receptor expressed in atherosclerotic lesions. *Arterioscler Thromb Vasc Biol* 24:750–755
- Xu Q, Wang J, He J et al (2011) Impaired CXCR4 expression and cell engraftment of bone marrow-derived cells from aged atherogenic mice. *Atherosclerosis* 219:92–99
- Yeh M, Leitinger N, de Martin R et al (2001) Increased transcription of IL-8 in endothelial cells is differentially regulated by TNF-alpha and oxidized phospholipids. *Arterioscler Thromb Vasc Biol* 21:1585–1591
- Yi GW, Zeng QT (2008) Circulating CXCL16 is related to the severity of coronary artery stenosis. *Arch Med Res* 39:531–535
- Yi GW, Zeng QT, Mao XB et al (2011) Overexpression of CXCL16 promotes a vulnerable plaque phenotype in apolipoprotein E-knockout mice. *Cytokine* 53:320–326
- Yilmaz A, Lipfert B, Cicha I et al (2007) Accumulation of immune cells and high expression of chemokines/chemokine receptors in the upstream shoulder of atherosclerotic carotid plaques. *Exp Mol Pathol* 82:245–255
- Zernecke A, Weber C (2010) Chemokines in the vascular inflammatory response of atherosclerosis. *Cardiovasc Res* 86:192–201
- Zernecke A, Bot I, Djalali-Talab Y et al (2008) Protective role of CXC receptor 4/CXC ligand 12 unveils the importance of neutrophils in atherosclerosis. *Circ Res* 102:209–217