



Brill–Noether Theory of Stable Vector Bundles on Ruled Surfaces

L. Costa and Irene Macías Tarrío

Abstract. Let X be a ruled surface over a nonsingular curve C of genus $g \geq 0$. Let $M_H := M_{X,H}(2; c_1, c_2)$ be the moduli space of H -stable rank 2 vector bundles E on X with fixed Chern classes $c_i := c_i(E)$ for $i = 1, 2$. The main goal of this paper is to contribute to a better understanding of the geometry of the moduli space M_H in terms of its Brill–Noether locus $W_H^k(2; c_1, c_2)$, whose points correspond to stable vector bundles in M_H having at least k independent sections. We deal with the non-emptiness of this Brill–Noether locus, getting in most of the cases sharp bounds for the values of k such that $W_H^k(2; c_1, c_2)$ is non-empty.

Mathematics Subject Classification. 14J60, 14J26.

Contents

1. Introduction	
2. Preliminaries	
3. Walls and Chambers Structure in Brill–Noether Loci	
4. Emptiness and Non-emptiness of Brill–Noether Loci	
References	

1. Introduction

Let X be a smooth projective variety of dimension n over an algebraically closed field K of characteristic 0 and let $M_H = M_{X,H}(r; c_1, \dots, c_s)$ be the moduli space of stable rank r vector bundles E with respect to an ample divisor H on X and with fixed Chern classes $c_i := c_i(E)$ for $i = 1, \dots, s := \min\{r, n\}$.

Moduli spaces of stable vector bundles were defined in 1970 by Murayama. Since then, many authors have done a systematic study of these

L. Costa was partially supported by PID2020-113674GB-I00. I. M. Tarrío was partially supported by PID2020-113674GB-I00.

moduli spaces but in general, in spite of great contributions, a lot of questions remain open.

One way of studying the geometry of these moduli spaces is by studying their subvarieties. A Brill–Noether locus $W_H^k = W_H^k(r; c_1, \dots, c_s)$ is a subvariety of $M_{X,H}(r; c_1, \dots, c_s)$ whose points correspond to stable vector bundles having at least k independent sections and, roughly speaking, the Brill–Noether Theory describes the geometry of these subvarieties.

Classical Brill–Noether Theory deals with line bundles on smooth curves (see Ref. [1]). In the case of line bundles on projective curves C of genus g , Brill–Noether theory is concerned with the subvarieties of $\text{Pic}^d(C)$ whose line bundles have at least k independent global sections. Basic questions related to non-emptiness, connectedness, irreducibility, dimension, singularities, cohomological classes, etc. have been answered when the curve C is generic on the moduli space of curves of genus g .

There are several natural ways to generalize the classical theory of Brill–Noether. The first natural generalization concerns higher rank vector bundles on algebraic curves. During the last 2 decades, a great amount of job has been made around the Brill–Noether stratification of the moduli space of stable rank r vector bundles of degree d on algebraic curves, giving rise to nice and interesting descriptions of these subvarieties (see for instance Refs. [2,3]). Nevertheless, it should be mentioned that in spite of big efforts, many questions concerning their geometry still remain open. The other way of generalizing this theory is considering line bundles on varieties of arbitrary dimension. Finally, we can go in both directions simultaneously. The Brill–Noether theory for rank r bundles on a variety of arbitrary dimension was established in Ref. [4]. In fact, if X is a smooth projective variety of dimension n and H an ample divisor on X , the Brill–Noether locus $W_H^k(r; c_1, \dots, c_s)$ in $M_{X,H}(r; c_1, \dots, c_s)$ is defined as the set of stable vector bundles in $M_{X,H}(r; c_1, \dots, c_s)$ having at least k independent sections. It is proved in Ref. [4] that $W_H^k(r; c_1, \dots, c_s)$ has a natural structure of k -determinantal variety, that each of its irreducible components has dimension greater than the expected dimension $\rho_H^k(r; c_1, \dots, c_s)$ (see Definition 2.9) and that these subvarieties give a filtration of the moduli space. The general case was studied in Refs. [5,6] for K3 surfaces, but also for other varieties in Refs. [7,8], for example.

In the context of the Brill–Noether Theory, one can immediately raise the following natural problems:

- Problem 1.1.** (a) Determine when the Brill–Noether locus W_H^k is non-empty.
 (b) Study the smoothness of the Brill–Noether locus.
 (c) See if it is true that $\rho_H^k > 0$ implies that $W_H^k \neq \emptyset$ and $\rho_H^k < 0$ implies that $W_H^k = \emptyset$.
 (d) Determine the irreducibility of W_H^k .

In this paper, we will focus the attention on moduli spaces of rank 2 stable vector bundles on ruled surfaces. This case was also studied for Hirzebruch surfaces in Ref. [9] and for ruled surfaces in general in Ref. [10]. In Ref. [11], Qin developed a theory of walls and chambers and proved that

moduli spaces $M_{X,H}(2; c_1, c_2)$ only depend on the chamber of H . In this paper, we will analyze how this theory has implications on the Brill–Noether loci, and in addition, using this structure, we will obtain our first contribution to Problem 1.1(a) (see Theorem 3.6).

Concerning Problem 1.1(a), we will also prove in Theorem 4.1:

Theorem 1.2. *Let X be a ruled surface over a nonsingular curve C of genus $g \geq 0$, $m \in \{0, 1\}$, $c_2 \gg 0$ an integer and $H \equiv \alpha C_0 + \beta f$ an ample divisor on X with*

$$\alpha(e + m) < \beta.$$

Then, for any k in the range

$$\max\{1, g\} \leq k < \frac{1}{2\alpha}[\beta - \alpha(e - m + 2g - 2)],$$

the Brill–Noether locus $W_H^k(2; C_0 + mf, c_2) \neq \emptyset$ with $m = \deg(\mathfrak{m})$.

The fact that $W_H^k(2; C_0 + mf, c_2) \neq \emptyset$ for all $\max\{1, g\} \leq k < \frac{1}{2\alpha}[\beta - \alpha(e - m + 2g - 2)]$ implies that $W_H^k(2; C_0 + mf, c_2) \neq \emptyset$ for all $1 \leq k < \frac{1}{2\alpha}[\beta - \alpha(e - m + 2g - 2)]$. Moreover, we will see that this bound is sharp (see Theorem 4.5).

On the other hand, we will prove

Theorem 1.3. *Let X be a ruled surface over a nonsingular curve C of genus $g \geq 0$, $c_2 \gg 0$ an integer and $H \equiv C_0 + \beta f$ an ample divisor on X . Then, $W_H^1(2; f, c_2) \neq \emptyset$ and $W_H^k(2; f, c_2) = \emptyset$ for all $k \geq 3$.*

Finally, we will study in detail the locus $W_H^1(2; f, c_2) \setminus W_H^2(2; f, c_2)$ and we will contribute to Problem 1.1(b) showing that

Proposition 1.4. *Let X be a ruled surface over a nonsingular curve C of genus $g \geq 0$, $c_2 \gg 0$ an integer and $H \equiv C_0 + \beta f$ an ample divisor on X . Then, $W_H^1(2; f, c_2) \setminus W_H^2(2; f, c_2)$ is smooth and has the expected dimension, namely*

$$\rho_H^1 = 3c_2 + g - 1.$$

Now, we will outline the structure of the paper. In Sect. 2, we will fix some notation and we will prove some basic facts on cohomology of line bundles on ruled surfaces and we will recall the notion of stability of a vector bundle. We will also give the general definition of the Brill–Noether locus $W_H^k(r, c_1, \dots, c_s)$ and we will recall a result about its existence and its structure as a k -determinantal variety. Finally, we will introduce some technical lemmas that we will use in the next sections. In Sect. 3, we will apply the theory of walls and chambers due to Qin to describe how Brill–Noether loci change when the ample divisor crosses a wall between two chambers and we will obtain results concerning non-emptiness of $W_H^k(2; c_1, c_2)$. In Sect. 4, we will push forward the study of the non-emptiness of the Brill–Noether locus $W_H^k(2; c_1, c_2)$ (see Theorems 4.1, 4.5, 4.8 and 4.10). Finally, we will prove the smoothness and irreducibility of $W_H^1(2; f, c_2) \setminus W_H^2(2; f, c_2)$ in Proposition 4.12.

2. Preliminaries

The goal of this section is to fix some notation and prove some results concerning the cohomology of line bundles on ruled surfaces. In addition, we will recall some definitions needed in the sequel.

A ruled surface is a surface X , together with a surjective map $\pi : X \rightarrow C$ to a nonsingular curve C of genus g such that, for every point $y \in C$, the fiber X_y is isomorphic to \mathbb{P}^1 , and such that π admits a section. It is also defined as the projectivization $X = \mathbb{P}(\mathcal{E})$ of a normalized rank 2 bundle \mathcal{E} on C . Let ϵ be the divisor on C corresponding to the invertible sheaf $\wedge^2 \mathcal{E}$ and let us define $e := -\deg(\epsilon)$.

Let $C_0 \subseteq X$ be a section and f be a fiber. We have that $\text{Pic}(X) \cong \mathbb{Z} \oplus \pi^* \text{Pic}(C)$, where \mathbb{Z} is generated by C_0 . In addition, $\text{Num}(X) \cong \mathbb{Z} \oplus \mathbb{Z}$, generated by C_0 and f satisfying $C_0^2 = -e$, $C_0 \cdot f = 1$ and $f^2 = 0$. If \mathfrak{b} is a divisor on C , we will write $\mathfrak{b}f$ instead of $\pi^*\mathfrak{b}$. Thus, a divisor D on X can be written uniquely as $D = aC_0 + \mathfrak{b}f$, being $\mathfrak{b} \in \text{Pic}(C)$, and any element of $\text{Num}(X)$ can be written as $D \equiv aC_0 + \mathfrak{b}f$. The canonical divisor K_X on X is in the class $-2C_0 + (\mathfrak{k} + \epsilon)f$, where \mathfrak{k} is the canonical divisor on C of degree $2g - 2$.

If the genus of the curve C is zero then, for any integer $e \geq 0$, the ruled surface is known as a Hirzebruch surface X_e . In this case, we have $X_e \cong \mathbb{P}(\mathcal{E}) = \mathbb{P}(\mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(-e))$. Since $\text{Pic}(X_e) \cong \mathbb{Z}^2$, dealing with divisors on Hirzebruch surfaces we will not make any distinction between $D = aC_0 + \mathfrak{b}f$ and its numerical class $D \equiv aC_0 + \mathfrak{b}f$.

Going back to the general case, for any divisor $D = aC_0 + \mathfrak{b}f$ on X with $a \geq 0$, it follows from Lemma V.2.4, Exercises III.8.3 and III.8.4 of [12], that

$$h^i(X, \mathcal{O}_X(D)) = h^i(C, (S^a \mathcal{E})(\mathfrak{b})) \tag{1}$$

where $S^a \mathcal{E}$ stands for the a -th symmetric power of \mathcal{E} . Moreover,

$$h^0(C, \mathcal{O}_C(\mathfrak{b})) \leq h^0(C, (S^a \mathcal{E})(\mathfrak{b})) \leq \sum_{i=0}^a h^0(C, \mathcal{O}_C(\mathfrak{b} + i\epsilon)), \tag{2}$$

for each divisor \mathfrak{b} on C (see for instance [13, Section 2]).

Notation 2.1. Usually, we will write $h^0 \mathcal{O}_X(D)$ to refer to $h^0(X, \mathcal{O}_X(D))$.

From now on, we will assume that $e \geq 0$. In this case, a divisor $D = aC_0 + \mathfrak{b}f$ is ample if and only if $a > 0$ and $\deg(\mathfrak{b}) > ae$ (see [12, Chapter V, Corollary 2.8]).

For effective divisors on X , we have the following:

Lemma 2.2. *Let X be a ruled surface over a nonsingular curve C of genus $g \geq 0$ and $D = aC_0 + \mathfrak{b}f$ be a divisor on X with $b := \deg(\mathfrak{b})$. If D is effective then $a, b \geq 0$.*

Proof. Let us first assume that D is effective. If $D = 0$ there is nothing to prove. Therefore, let us assume that $D = aC_0 + \mathfrak{b}f \neq 0$.

First, we will see that $a \geq 0$. Since D is an effective divisor, $D \cdot H > 0$ for any ample divisor H on X . In particular, taking the ample divisor on X , $H \equiv C_0 + \beta f$ with $\beta \gg 0$, we have

$$0 < D \cdot H = -ae + a\beta + b = a(\beta - e) + b$$

what implies that $a \geq 0$.

Now, we will see that $b \geq 0$. Since D is effective and $a \geq 0$, it follows from (1) and (2) that

$$0 < h^0 \mathcal{O}_X(D) = h^0 S^a \mathcal{E}(\mathbf{b}) \leq \sum_{i=0}^a h^0 \mathcal{O}_C(\mathbf{b} + i\epsilon).$$

In particular, since $\deg(\epsilon) = -e \leq 0$, $h^0 \mathcal{O}_C(\mathbf{b}) \neq 0$ and hence $b \geq 0$. □

Lemma 2.3. *Let X be a ruled surface over a nonsingular curve C of genus $g \geq 0$ and $D = aC_0 + \mathbf{b}f$ on X such that $a \geq 0$. Then, if $b := \deg(\mathbf{b})$, we have*

$$h^0 \mathcal{O}_X(aC_0 + \mathbf{b}f) \leq \begin{cases} 0 & \text{for } b < 0 \\ (a + 1)(b + 1) & \text{for } 0 \leq b \leq g - 1, g \geq 1 \\ (a + 1)g & \text{for } g \leq b \leq 2g - 1, g \geq 1 \\ (a + 1)(b + 1 - g) & \text{for } b \geq 2g. \end{cases}$$

Proof. Let $D = aC_0 + \mathbf{b}f$ be a divisor on X with $a \geq 0$.

If $b < 0$, it follows from Lemma 2.2 that $h^0 \mathcal{O}_X(aC_0 + \mathbf{b}f) = 0$. Let us now consider the case $0 \leq b \leq g - 1$. Since $a \geq 0$, by (1) and (2) we have

$$h^0 \mathcal{O}_X(aC_0 + \mathbf{b}f) = h^0 S^a \mathcal{E}(\mathbf{b}) \leq \sum_{i=0}^a h^0 \mathcal{O}_C(\mathbf{b} + i\epsilon).$$

By [12, Chapter II, Exercise 1.15], for any divisor \mathfrak{d} on C

$$h^0 \mathcal{O}_C(\mathfrak{d}) \leq \deg(\mathfrak{d}) + 1. \tag{3}$$

In particular, we have

$$h^0 \mathcal{O}_C(\mathbf{b} + i\epsilon) \leq b - i e + 1 \leq b + 1,$$

and thus

$$h^0 \mathcal{O}_X(aC_0 + \mathbf{b}f) \leq (a + 1)(b + 1).$$

Let us now assume that $g \leq b \leq 2g - 1$. Since $a \geq 0$, again by (1) and (2),

$$h^0 \mathcal{O}_X(aC_0 + \mathbf{b}f) = h^0 S^a \mathcal{E}(\mathbf{b}) \leq \sum_{i=0}^a h^0 \mathcal{O}_C(\mathbf{b} + i\epsilon) \leq (a + 1) h^0 \mathcal{O}_C(\mathbf{b}) \leq (a + 1)g,$$

where the last inequality follows from the fact that, since $b \leq 2g - 1$, if \mathfrak{d} is a divisor of degree $2g - 1$, we get

$$h^0 \mathcal{O}_C(\mathbf{b}) \leq h^0 \mathcal{O}_C(\mathfrak{d}) = g.$$

Finally, let us prove the case $b \geq 2g$. Since $a \geq 0$, by (1) and (2),

$$\begin{aligned} h^0 \mathcal{O}_X(aC_0 + \mathbf{b}f) &= h^0 S^a \mathcal{E}(\mathbf{b}) \leq \sum_{i=0}^a h^0 \mathcal{O}_C(\mathbf{b} + i\epsilon) \\ &\leq (a + 1) h^0 \mathcal{O}_C(\mathbf{b}) = (a + 1)(b + 1 - g), \end{aligned}$$

where the last equality follows from the fact that, since $b \geq 2g > 2g - 1$, we have $h^0 \mathcal{O}_C(\mathbf{b}) = b + 1 - g$. \square

The following is a well-known result.

Proposition 2.4. *Let X be a smooth projective surface, D a divisor on X and $Z \subset X$ a general 0-dimensional subscheme. If $|Z| \geq h^0 \mathcal{O}_X(D)$ then $h^0 I_Z(D) = 0$.*

Let E be a rank 2 vector bundle on a smooth projective surface X . We denote by c_i the Chern Classes of E , $c_i = c_i(E) \in H^{2i}(X, \mathbb{Z})$, for $i = 1, 2$.

Our aim is to study the moduli space of stable vector bundles on X by means of its Brill–Noether locus. To do so, we end the section by recalling basic facts concerning stable vector bundles and Brill–Noether loci.

Definition 2.5. Let X be a smooth projective surface, H an ample divisor on X and E a vector bundle on X with fixed Chern classes c_i for $i = 1, 2$. Then, E is stable with respect to H (or H -stable) if, for any subbundle F of E , we have

$$\mu_H(F) := \frac{c_1(F) \cdot H}{\text{rank}(F)} < \frac{c_1(E) \cdot H}{\text{rank}(E)} := \mu_H(E).$$

Notice that the notion of stability depends on the ample divisor. Nevertheless, when there is no confusion, we will simply say that E is stable.

Given H an ample divisor on X , $c_1 \in \text{Num}(X)$ and $c_2 \in H^4(X, \mathbb{Z}) \cong \mathbb{Z}$, we denote by $M_{X,H}(r; c_1, c_2)$ the moduli space of H -stable rank r vector bundles E on X with fixed Chern classes c_i . When there is no confusion, we will denote it by $M_H(r; c_1, c_2)$ or by M_H .

For big values of c_2 , we have a general result concerning irreducibility, smoothness and dimension of these moduli spaces. In the particular case of smooth ruled surfaces, it can be stated as follows (see for instance [14]):

Theorem 2.6. *Let X be a ruled surface over a nonsingular curve C of genus $g \geq 0$, H an ample divisor on X and $c_1, c_2 \in H^{2i}(X, \mathbb{Z})$ Chern classes. For all $c_2 \gg 0$, the moduli space $M_{X,H}(r; c_1, c_2)$ is a smooth, irreducible, quasiprojective variety of dimension $2rc_2 - (r - 1)c_1^2 - (r^2 - 1)(1 - g)$.*

One possible way to study the moduli space of stable vector bundles is by describing the Brill–Noether locus $W_H^k := W_H^k(r; c_1, c_2)$ in M_H which parametrizes stable rank r vector bundles having at least k independent sections, that is

$$\text{Supp}(W_H^k(r; c_1, c_2)) = \{E \in M_{X,H}(r; c_1, c_2) \mid h^0 E \geq k\}.$$

The Brill–Noether locus has the structure of a k -determinantal variety $W_H^k(r; c_1, c_2)$, and so we can talk about its expected dimension and singular locus. This is given by the following result (see [4, Theorem 2.3]).

Theorem 2.7. *Let X be a smooth projective surface and consider the moduli space $M_{X,H}(r; c_1, c_2)$ of rank r , H -stable vector bundles E on X with fixed Chern classes c_i . Assume that for any $E \in M_H$, $h^2 E = 0$. Then, for any $k \geq 0$, there exists a determinantal variety $W_H^k(r; c_1, c_2)$ such that*

$$\text{Supp}(W_H^k(r; c_1, c_2)) = \{E \in M_H \mid h^0 E \geq k\}.$$

Moreover, each non-empty irreducible component of $W_H^k(r; c_1, c_2)$ has dimension at least

$$\dim(M_H) - k(k - \chi(r; c_1, c_2)),$$

and

$$W_H^{k+1}(r; c_1, c_2) \subset \text{Sing}(W_H^k(r; c_1, c_2))$$

whenever $W_H^{k+1}(r; c_1, c_2) \neq M_H$.

Remark 2.8. By definition, the different Brill–Noether loci form a filtration of the moduli space $M_{X,H}(r; c_1, c_2)$,

$$M_H \supseteq W_H^1(r; c_1, c_2) \supseteq W_H^2(r; c_1, c_2) \supseteq \cdots \supseteq W_H^k(r; c_1, c_2) \supseteq W_H^{k+1}(r; c_1, c_2) \supseteq \cdots$$

Definition 2.9. The expected dimension of the Brill–Noether locus is defined as

$$\rho_H^k := \rho_H^k(r; c_1, c_2) = \dim M_{X,H}(r; c_1, c_2) - k(k - \chi(r; c_1, c_2)).$$

By the Riemann–Roch theorem, $\chi(r; c_1, c_2)$ can be expressed as

$$\chi(r; c_1, c_2) = r(1 + p_a(X)) - \frac{c_1 K_X}{2} + \frac{c_1^2}{2} - c_2.$$

In particular, if X is a ruled surface over a smooth curve C of genus g , then

$$\chi(r; c_1, c_2) = r(1 - g) - \frac{c_1 \cdot [-2C_0 + (\mathfrak{k} + \mathfrak{e})f]}{2} + \frac{c_1^2}{2} - c_2.$$

Remark 2.10. Under the assumption $c_2 \gg 0$, the moduli space has the dimension given by Theorem 2.6 and hence, if X is a smooth ruled surface over a nonsingular curve of genus g ,

$$\rho_H^k := \rho_H^k(r; c_1, c_2) = 2rc_2 - (r - 1)c_1^2 - (r^2 - 1)(1 - g) - k(k - \chi(r; c_1, c_2)).$$

Lemma 2.11. *Let E be a rank r , H -stable vector bundle on a smooth projective surface X with $(c_1(E) - rK_X) \cdot H \geq 0$. Then, $h^2 E = 0$.*

Proof. Let us assume $h^2 E > 0$. This means, by Serre duality, that $h^0 E^*(K_X) > 0$, what implies $\mathcal{O}_X(-K_X) \hookrightarrow E^*$. Since E is H -stable, the same is true for E^* , and thus we have

$$\mu_H(\mathcal{O}_X(-K_X)) < \mu_H(E^*),$$

which is equivalent to

$$(-rK_X - c_1(E^*)) \cdot H = (-rK_X + c_1(E)) \cdot H < 0,$$

what is a contradiction. Hence, $h^2 E = 0$. □

From now on, we will assume that $(c_1(E) - rK_X) \cdot H \geq 0$ for all $E \in M_{X,H}(r; c_1, c_2)$, and therefore, by Theorem 2.7 and applying Lemma 2.11, the existence of the Brill–Noether locus $W_H^k(r; c_1, c_2)$ is guaranteed.

While dealing with the non-emptiness of the Brill–Noether locus $W_H^k(r; c_1, c_2)$, notice that if $c_1 \cdot H \leq 0$, then $W_H^k(r; c_1, c_2) = \emptyset$ for all $k \geq 1$. In fact, if we have $E \in W_H^k(r; c_1, c_2)$, then $h^0 E \neq 0$ and thus, $\mathcal{O}_X \hookrightarrow E$. By stability of E and because $c_1 \cdot H \leq 0$, we have $0 = \mu_H(\mathcal{O}_X) < \mu_H(E) \leq 0$, what is a contradiction. Hence, $W_H^1(r; c_1, c_2) = \emptyset$ whenever $c_1 \cdot H \leq 0$.

In particular, $W_H^1(r; 0, c_2) = \emptyset$. Therefore, we will assume that c_1 is effective and non-zero, and thus $c_1 \cdot H > 0$ for any ample divisor H . Furthermore, in the case of a rank 2 vector bundle E , we have that E is H -stable if and only if $E \otimes L$ is H -stable, for any line bundle L on X , so it is natural to focus the attention on the cases $c_1 \in \{C_0, C_0 + f, f\}$.

3. Walls and Chambers Structure in Brill–Noether Loci

We start the section recalling the theory of walls and chambers introduced by Qin, in the context of smooth surfaces (see for instance [11]). The main idea is that if X is a smooth surface, the ample cone of X , C_X , has a structure of walls and chambers such that the moduli space $M_H(2; c_1, c_2)$ only depends on the chamber of H .

To introduce the main results of this theory, let us remember some basic definitions.

Definition 3.1. (i) Let $\zeta \in \text{Num}(X) \otimes \mathbb{R}$. We define

$$W^\zeta = C_X \cap \{x \in \text{Num}(X) \mid x \cdot \zeta = 0\}.$$

- (ii) Define $\mathcal{W}(c_1, c_2)$ as the set whose elements consist of W^ζ , where ζ is the numerical equivalence class of a divisor D on X such that $D + c_1$ is divisible by 2 in $\text{Pic}(X)$, $-(4c_2 - c_1^2) \leq D^2 < 0$ and $|Z| = c_2 + \frac{D^2 - c_1^2}{4}$ for some locally complete intersection codimension-two cycle Z in X .
- (iii) A wall of type (c_1, c_2) is an element in $\mathcal{W}(c_1, c_2)$. A chamber of type (c_1, c_2) is a connected component of $C_X \setminus \mathcal{W}(c_1, c_2)$. A Z -chamber of type (c_1, c_2) is the intersection of $\text{Num}(X)$ with some chamber of type (c_1, c_2) .
- (iv) A face of type (c_1, c_2) is $\mathcal{F} = W^\zeta \cap \mathcal{C}$, where W^ζ is a wall of type (c_1, c_2) and \mathcal{C} is a chamber of type (c_1, c_2) .

Definition 3.2. Let ζ be a numerical equivalence class defining a wall of type (c_1, c_2) . We define $E_\zeta(c_1, c_2)$ as the set of rank 2 vector bundles E on X given by a non-trivial extension of type

$$0 \rightarrow \mathcal{O}_X(D) \rightarrow E \rightarrow I_Z(c_1 - D) \rightarrow 0$$

where D is a divisor with $2D - c_1 \equiv \zeta$ and Z is a locally complete intersection 0-cycle of length $c_2 + \frac{\zeta^2 - c_1^2}{4}$.

Qin proved in Refs. [11, Proposition 2.2.5] and [11, Remark 2.2.6] that the moduli space $M_H(2; c_1, c_2)$ only depends on the chamber of H , so he set $M_{\mathcal{C}}(c_1, c_2)$ to be $M_H(2; c_1, c_2)$ for some $H \in \mathcal{C}$.

Once we know that $M_H(2; c_1, c_2)$ only depends on the chamber of H , the natural question that arises is how the moduli space changes when the polarization H crosses a wall W^ζ between two chambers. The phenomena that occurs is summarized in the following result.

Theorem 3.3. [11, Proposition 1.3.1] *Let \mathcal{C} be a chamber and \mathcal{F} be one of its faces. Then, as sets,*

$$M_{\mathcal{C}}(c_1, c_2) = M_{\mathcal{F}}(c_1, c_2) \amalg (\amalg_{\zeta} E_{\zeta}^k(c_1, c_2))$$

where ζ satisfies $\zeta \cdot H < 0$ for some $H \in \mathcal{C}$, and runs over all numerical equivalence classes which define the wall containing \mathcal{F} .

As a consequence

Theorem 3.4. [11, Theorem 1.3.3] *Let \mathcal{C}_1 and \mathcal{C}_2 be two adjacent chambers of type (c_1, c_2) sharing a common wall W^ζ such that \mathcal{C}_1 lies above \mathcal{C}_2 . Then, as sets,*

$$M_{\mathcal{C}_2}(c_1, c_2) = (M_{\mathcal{C}_1}(c_1, c_2) \setminus \amalg_{\zeta} E_{-\zeta}(c_1, c_2)) \amalg (\amalg_{\zeta} E_{\zeta}(c_1, c_2))$$

where ζ runs over all normalized numerical equivalence classes (that is $\zeta \cdot f > 0$) which represents the wall W^ζ .

It is natural to ask ourselves what happens at the level of Brill–Noether loci. To this end, let us define

$$W_{\mathcal{C}}^k(c_1, c_2) := \{E \in \mathcal{M}_{\mathcal{C}}(c_1, c_2) \mid h^0 E \geq k\}$$

and

$$E_{\zeta}^k(c_1, c_2) := \{E \in E_{\zeta} \mid h^0 E \geq k\}.$$

As a consequence of Theorems 3.3 and 3.4, we can deduce

Proposition 3.5. (a) *Let \mathcal{C} be a chamber and \mathcal{F} one of its faces. Then, as sets,*

$$W_{\mathcal{C}}^k(c_1, c_2) = W_{\mathcal{F}}^k(c_1, c_2) \amalg (\amalg_{\zeta} E_{\zeta}^k(c_1, c_2))$$

where ζ satisfies $\zeta \cdot H < 0$ for some $H \in \mathcal{C}$, and runs over all numerical equivalence classes which define the wall containing \mathcal{F} .

(b) *Let \mathcal{C}_1 and \mathcal{C}_2 be two adjacent chambers of type (c_1, c_2) sharing a common wall W^ζ such that \mathcal{C}_1 lies above \mathcal{C}_2 . Then, as sets,*

$$W_{\mathcal{C}_2}^k(c_1, c_2) = (W_{\mathcal{C}_1}^k(c_1, c_2) \setminus \amalg_{\zeta} E_{-\zeta}^k(c_1, c_2)) \amalg (\amalg_{\zeta} E_{\zeta}^k(c_1, c_2)).$$

We will use this wall crossing to prove the following result.

Theorem 3.6. *Let X be a ruled surface over a nonsingular curve C of genus $g \geq 0$, $m \in \{0, 1\}$ and $c_2 \gg 0$ an integer. Let us consider the family of numerical equivalence classes*

$$\zeta_b \equiv (2b - m)f - C_0$$

with $0 < b = k - 1 + g < c_2$ and let \mathcal{C}_b be the chamber such that $W^{\zeta_b} \cap \text{Closure}(\mathcal{C}_b) \neq \emptyset$ and $\zeta_b \cdot H < 0$ for all $H \in \mathcal{C}_b$. Then, for any ample divisor $H \in \mathcal{C}_b$, $W_H^k(2; C_0 + \mathbf{m}f, c_2) \neq \emptyset$ whenever $M_H \neq \emptyset$.

Proof. First, since we have the filtration $W_H^{l+1} \subseteq W_H^l \subseteq \dots \subseteq W_H^1 \subseteq M_H$, we can assume $k \geq \max\{2, g\}$ and thus $b \geq \max\{g + 1, 2g - 1\}$.

Notice that $\zeta_b \equiv (2b - m)f - C_0$ defines a non-empty wall of type $(C_0 + \mathbf{m}f, c_2)$. In fact,

$$(\zeta_b)^2 = ((2b - m)f - C_0)^2 = -4b - e + 2m.$$

Since $m \in \{0, 1\}$ and by the lower and upper bounds of b , we have

$$-4c_2 - e \leq -4b - e + 2m < 0.$$

On the other hand, $\zeta_b + C_0 + \mathbf{m}f = 2\mathbf{b}f$ is divisible by 2 in $\text{Pic}(X)$ and $H \equiv C_0 + (2b + e - m)f$ is an ample divisor on X such that $\zeta_b \cdot H = 0$. Hence, ζ_b defines a non-empty wall of type $(C_0 + \mathbf{m}f, c_2)$.

Let us consider $E_{\zeta_b}(C_0 + \mathbf{m}f, c_2)$, which is by definition the family of rank 2 vector bundles E given by non-trivial extensions of type

$$0 \rightarrow \mathcal{O}_X(\mathbf{b}f) \rightarrow E \rightarrow I_Z(C_0 + (\mathbf{m} - \mathbf{b})f) \rightarrow 0, \tag{4}$$

where $\mathbf{b}f$ is a divisor such that $\zeta_b \equiv (2b - m)f - C_0$ with $b = \text{deg}(\mathbf{b})$ and Z a 0-dimensional subscheme of length $|Z| = c_2 - b$.

Claim: $E_{\zeta_b}(C_0 + \mathbf{m}f, c_2)$ is non-empty.

Proof of the Claim: It is enough to see that $\text{Ext}^1(I_Z(C_0 + (\mathbf{m} - \mathbf{b})f), \mathcal{O}_X(\mathbf{b}f)) \neq 0$. We have that $\text{ext}^1(I_Z(C_0 + (\mathbf{m} - \mathbf{b})f), \mathcal{O}_X(\mathbf{b}f)) = h^1 I_Z(C_0 + (\mathbf{m} - 2\mathbf{b})f + K_X)$. If we consider the short exact sequence

$$0 \rightarrow I_Z \rightarrow \mathcal{O}_X \rightarrow \mathcal{O}_Z \rightarrow 0, \tag{5}$$

we tensor it by $\mathcal{O}_X(C_0 + (\mathbf{m} - 2\mathbf{b})f + K_X)$ and we take cohomology, we get

$$h^0 I_Z(C_0 + (\mathbf{m} - 2\mathbf{b})f + K_X) \leq h^0 \mathcal{O}_X(C_0 + (\mathbf{m} - 2\mathbf{b})f + K_X) = 0,$$

where the last equality follows from Lemma 2.2.

On the other hand, if we consider again the exact sequence (5) and we take cohomology, we get

$$h^2 I_Z(C_0 + (\mathbf{m} - 2\mathbf{b})f + K_X) = h^2 \mathcal{O}_X(C_0 + (\mathbf{m} - 2\mathbf{b})f + K_X)$$

and, by duality and Lemma 2.2,

$$h^2 \mathcal{O}_X(C_0 + (\mathbf{m} - 2\mathbf{b})f + K_X) = h^0 \mathcal{O}_X(-C_0 - (\mathbf{m} - 2\mathbf{b})f) = 0.$$

Therefore,

$$h^1 I_Z(C_0 + (\mathbf{m} - 2\mathbf{b})f + K_X) = -\chi I_Z(C_0 + (\mathbf{m} - 2\mathbf{b})f + K_X).$$

Considering the exact sequence (5) and the Riemann–Roch Theorem,

$$-\chi I_Z(C_0 + (\mathbf{m} - 2\mathbf{b})f + K_X) = |Z| - \chi \mathcal{O}_X(C_0 + (\mathbf{m} - 2\mathbf{b})f + K_X) = |Z|.$$

Putting altogether, we have

$$\text{ext}^1(I_Z(C_0 + (\mathbf{m} - \mathbf{b})f), \mathcal{O}_X(\mathbf{b}f)) = |Z| = c_2 - b > 0,$$

where the last inequality follows from the upper bound of b .

Therefore, we can take a rank 2 vector bundle $E \in E_{\zeta_b}(C_0 + \mathbf{m}f, c_2)$. It follows from Theorem 3.3 that $E_{\zeta_b}(C_0 + \mathbf{m}f) \subseteq M_{C_b}(C_0 + \mathbf{m}f, c_2)$ and thus E is H -stable for any $H \in \mathcal{C}_b$.

Finally, since E is given by the exact sequence (4), taking cohomology, we get

$$h^0 E \geq h^0 \mathcal{O}_X(\mathbf{b}f) = h^0 \mathcal{O}_C(\mathbf{b}) = \chi \mathcal{O}_C(\mathbf{b}) = b + 1 - g = k,$$

where the second equality follows from the fact that, since $k \geq \max\{1, g\}$, $h^1 \mathcal{O}_C(\mathbf{b}) = 0$.

Thus, we have seen that $\emptyset \neq E_{\zeta_b}^k(C_0 + \mathbf{m}f, c_2) \subseteq W_{C_b}^k(2; C_0 + \mathbf{m}f, c_2)$ and thus $W_H^k(2; C_0 + \mathbf{m}f, c_2) \neq \emptyset$ for all $H \in \mathcal{C}_b$. □

Notice that with the above notations, if $H \equiv \alpha C_0 + \beta f$ the condition $\zeta_b \cdot H < 0$ is equivalent to

$$k < \frac{1}{2\alpha}[\beta - \alpha(e - m + 2g - 2)].$$

Hence,

Corollary 3.7. *Let X be a ruled surface over a nonsingular curve C of genus $g \geq 0$, $m \in \{0, 1\}$ and $c_2 \gg 0$ an integer. Let us consider the family of numerical equivalence classes*

$$\zeta_b \equiv (2b - m)f - C_0$$

with $0 < b = k - 1 + g < c_2$. Assume $M_H \neq \emptyset$. Then, $W_H^k(2; C_0 + \mathbf{m}f, c_2) \neq \emptyset$ for

$$1 \leq k < \frac{1}{2\alpha}[\beta + \alpha(e - m + 2g - 2)]$$

and $H \in \mathcal{C}$ being \mathcal{C} the chamber such that $W^{\zeta_b} \cap \text{Closure}(\mathcal{C}) \neq \emptyset$.

4. Emptiness and Non-emptiness of Brill–Noether Loci

The goal of this section is to determine bounds of k , depending on an ample divisor H , that guarantees that the Brill–Noether locus $W_H^k(2; c_1, c_2)$ is non-empty. Moreover, we will see that these bounds are sharp. In contrast with the above section, we will obtain non-emptiness results by explicitly constructing families of H -stable rank 2 vector bundles for, in many cases, almost all ample divisors H and with values of the first Chern class c_1 in $\{C_0, C_0 + f, f\}$.

We will start with the case $c_1 = C_0 + \mathbf{m}f$ with $\deg(\mathbf{m}) = m \in \{0, 1\}$.

Theorem 4.1. *Let X be a ruled surface over a nonsingular curve C of genus $g \geq 0$, $m \in \{0, 1\}$, $c_2 \gg 0$ an integer and $H \equiv \alpha C_0 + \beta f$ an ample divisor on X with*

$$\alpha(e + m) < \beta.$$

Then, for any k in the range

$$\max\{1, g\} \leq k < \frac{1}{2\alpha}[\beta - \alpha(e - m + 2g - 2)],$$

the Brill–Noether locus $W_H^k(2; C_0 + \mathbf{m}f, c_2) \neq \emptyset$ whenever $M_H \neq \emptyset$.

Proof. First, notice that one can extend [15, Lemma 1.10] to any ample divisor $H \equiv \alpha C_0 + \beta f$ and prove that if $M_H(2; C_0 + \mathbf{m}f, c_2) \neq \emptyset$ then $\beta < \alpha(2c_2 + e - m)$. Hence, we can assume $\beta < \alpha(2c_2 + e - m)$. Let us consider the family \mathcal{G} of rank two vector bundles E on X given by a non-trivial extension

$$0 \rightarrow \mathcal{O}_X(\mathbf{b}f) \rightarrow E \rightarrow I_Z(C_0 + (\mathbf{m} - \mathbf{b})f) \rightarrow 0, \tag{6}$$

where Z is a generic 0-dimensional subscheme of length $|Z| = c_2 - b$ with

$$b = \deg(\mathbf{b}) = k - 1 + g$$

and

$$\max\{1, g\} \leq k < \frac{1}{2\alpha}[\beta - \alpha(e - m + 2g - 2)].$$

Claim: \mathcal{G} is non-empty.

Proof of the Claim: Notice that since $h^0 I_Z(C_0 + (\mathbf{m} - 2\mathbf{b})f + K_X) = 0 = h^2 I_Z(C_0 + (\mathbf{m} - 2\mathbf{b})f + K_X)$, using Riemann–Roch Theorem and the upper bound of b we deduce that

$$\text{ext}^1(I_Z(C_0 + (\mathbf{m} - \mathbf{b})f), \mathcal{O}_X(\mathbf{b}f)) = |Z| = c_2 - b > 0,$$

what implies $\text{Ext}^1(I_Z(C_0 + (\mathbf{m} - \mathbf{b})f), \mathcal{O}_X(\mathbf{b}f)) \neq 0$.

Hence, any non-zero element in $\text{Ext}^1(I_Z(C_0 + (\mathbf{m} - \mathbf{b})f), \mathcal{O}_X(\mathbf{b}f))$ defines a non-trivial extension of type (6). Notice that by the Cayley–Bacharach property [16, Theorem 12], E is indeed a rank two vector bundle, with $c_1(E) = C_0 + \mathbf{m}f$ and $c_2(E) = c_2 \gg 0$. Hence, \mathcal{G} is non-empty.

On the other hand, if we consider the exact sequence (6), we get

$$h^0 E \geq h^0 \mathcal{O}_X(\mathbf{b}f) = h^0 \mathcal{O}_C(\mathbf{b})$$

and, since $k \geq \max\{1, g\}$, we get $h^1 \mathcal{O}_C(\mathbf{b}) = 0$ and thus $h^0 \mathcal{O}_C(\mathbf{b}) = \chi \mathcal{O}_C(\mathbf{b}) = b + 1 - g = k$.

Now, let us prove that any rank two vector bundle E in \mathcal{G} is stable with respect to the polarization $H \equiv \alpha C_0 + \beta f$ with $\alpha(e + m) < \beta < \alpha(2c_2 + e - m)$.

To this end, since E is a rank two vector bundle on X , we have to check that for any line subbundle $\mathcal{O}_X(G) \hookrightarrow E$, we have

$$\mu_H(\mathcal{O}_X(G)) = G \cdot H < \mu_H(E) = \frac{c_1(E) \cdot H}{2}.$$

By construction, E is given by a non-trivial extension of type (6). Therefore, we have two possibilities: $\mathcal{O}_X(G) \hookrightarrow \mathcal{O}_X(\mathbf{b}f)$ or $\mathcal{O}_X(G) \hookrightarrow I_Z(C_0 + (\mathbf{m} - \mathbf{b})f)$.

Case 1 Assume that $\mathcal{O}_X(G) \hookrightarrow \mathcal{O}_X(\mathbf{b}f)$. In this case, since

$$k < \frac{1}{2\alpha}[\beta - \alpha(e - m + 2g - 2)],$$

we get

$$\begin{aligned} \mu_H(\mathcal{O}_X(G)) &\leq (\mathbf{b}f) \cdot H = \alpha b = \alpha k + \alpha(-1 + g) \\ &< \frac{\alpha}{2\alpha}[\beta - \alpha(e - m + 2g - 2)] + \alpha(-1 + g) = \frac{1}{2}(\beta - \alpha e + \alpha m) = \mu_H(E). \end{aligned}$$

Case 2 Assume that $\mathcal{O}_X(G) \hookrightarrow I_Z(C_0 + (\mathbf{m} - \mathbf{b})f)$, where $G = sC_0 + \mathbf{t}f$ with $t = \deg(\mathbf{t})$. In this case,

$$C_0 + (\mathbf{m} - \mathbf{b})f - G = (1 - s)C_0 + (\mathbf{m} - \mathbf{b} - \mathbf{t})f$$

is an effective divisor, which by Lemma 2.2 implies that $s \leq 1$ and $t \leq m - b$.

Assume that $s \leq 0$. Since $t \leq m - b$, we have

$$\mu_H(\mathcal{O}_X(G)) = s(\beta - \alpha e) + t\alpha \leq \alpha(m - b) < \frac{1}{2}(\beta - \alpha e + \alpha m) = \mu_H(E),$$

where the last inequality follows from the fact that $b \geq 0$ and $\beta > \alpha(e + m)$.

Finally assume that $s = 1$. In this case $G = C_0 + \mathbf{t}f$ and since

$$\mathcal{O}_X(G) \hookrightarrow I_Z(C_0 + (\mathbf{m} - \mathbf{b})f),$$

we have $h^0 I_Z(C_0 + (\mathbf{m} - \mathbf{b})f - G) = h^0 I_Z((\mathbf{m} - \mathbf{b} - \mathbf{t})f) \neq 0$. Since Z is a generic 0-dimensional subscheme, we must have

$$|Z| < h^0 \mathcal{O}_X((\mathbf{m} - \mathbf{b} - \mathbf{t})f) = h^0 \mathcal{O}_C(\mathbf{m} - \mathbf{b} - \mathbf{t}) \leq m - b - t + 1,$$

where the last inequality is given by (3).

Thus, since $|Z| = c_2 - b$,

$$t \leq m - c_2. \tag{7}$$

Now, let us check that $\mu_H(\mathcal{O}_X(C_0 + \mathbf{t}f)) < \mu_H(E)$. Notice that $\mu_H(\mathcal{O}_X(C_0 + \mathbf{t}f)) < \mu_H(E)$ if and only if

$$-\alpha e + \beta + t\alpha < \frac{1}{2}(\beta - \alpha e + \alpha m)$$

what is equivalent to

$$-\alpha e + \beta - \alpha m < -2t\alpha.$$

This inequality holds since, by the upper bound of β , and according to (7),

$$-2t\alpha \geq -2\alpha(m - c_2) > -\alpha e + \beta - \alpha m.$$

Therefore, $\mu_H(\mathcal{O}_X(C_0 + \mathbf{t}f)) < \mu_H(E)$ and putting altogether, we get that E is H -stable.

We have proved that any E in \mathcal{G} is a rank two H -stable vector bundle with $h^0 E \geq k$, which implies that $\mathcal{G} \hookrightarrow W_H^k(2; C_0 + \mathbf{m}f, c_2)$ and hence $W_H^k(2; C_0 + \mathbf{m}f, c_2) \neq \emptyset$. □

Remark 4.2. Notice that the condition $\beta > \alpha(e + m)$ in Theorem 4.1 is not restrictive. In fact, it is verified for almost all ample divisors, since $m \in \{0, 1\}$ and $\beta > \alpha e$ for any ample divisor $H \equiv \alpha C_0 + \beta f$.

Corollary 4.3. *Let X be a ruled surface over a nonsingular curve C of genus $g \geq 0$, $m \in \{0, 1\}$, $c_2 \gg 0$ an integer and $H \equiv \alpha C_0 + \beta f$ an ample divisor on X with*

$$\alpha(e + m) < \beta.$$

Then, for any

$$1 \leq k < \frac{1}{2\alpha}[\beta - \alpha(e - m + 2g - 2)],$$

$W_H^k(2; C_0 + \mathbf{m}f, c_2) \neq \emptyset$ whenever $M_H \neq \emptyset$.

Proof. If $\max\{1, g\} \leq k < \frac{1}{2}\alpha[\beta - \alpha(e + m + 2g - 2)]$, the result follows from Theorem 4.1. Finally, since we have the filtration $W_H^1 \supseteq W_H^2 \supseteq \dots \supseteq W_H^q$ with $q = \max\{1, g\}$, the fact that $W_H^q \neq \emptyset$ implies that $W_H^l \neq \emptyset$ for all $1 \leq l \leq q$. \square

Remark 4.4. Notice that the expected dimension of the Brill–Noether locus $W_H^k(2; C_0 + \mathbf{m}f, c_2)$ is

$$\rho_H^k(2; C_0 + \mathbf{m}f, c_2) = (4 - k)c_2 + 3k - 3 + 3(1 - k)g + (1 - k)e + 2(k - 1)m - k^2.$$

Hence, for $k \geq 4$, we have $\rho_H^k < 0$, but for

$$1 \leq g \leq k < \frac{1}{2\alpha}[\beta - \alpha(e - m + 2g - 2)]$$

we have seen that $W_H^k \neq \emptyset$. Therefore, we have an example of a non-empty Brill–Noether locus with negative expected dimension, and therefore, it is clear that, in general, $\rho_H^k < 0$ does not imply that $W_H^k = \emptyset$.

Our next goal is to see that the above bound is sharp, in the sense that if

$$k \geq \frac{1}{2\alpha}[\beta - \alpha(e - m + 2g - 2)],$$

then $W_H^k(2; C_0 + \mathbf{m}f, c_2) = \emptyset$.

Theorem 4.5. *Let X be a ruled surface over a nonsingular curve C of genus $g \geq 0$, $m \in \{0, 1\}$, $c_2 \gg 0$ an integer and $H \equiv \alpha C_0 + \beta f$ be an ample divisor on X such that*

$$\max\{\alpha(8 + e - m + 2g), \alpha(6g - 4 + e - m)\} < \beta.$$

Then, $W_H^k(2; C_0 + \mathbf{m}f, c_2) = \emptyset$ for all $k \geq \frac{1}{2\alpha}[\beta - \alpha(e - m + 2g - 2)]$.

Proof. Let us assume that $W_H^k(2; C_0 + \mathbf{m}f, c_2) \neq \emptyset$. Since $h^0 E \neq 0$, we can take a non-zero section s of E . We denote by Y its scheme of zeros and by $D = aC_0 + \mathbf{b}f$ the maximal effective divisor contained in Y . Then, s can be regarded as a section of $E(-D)$ and its scheme of zeros has codimension greater or equal than two. Thus, we have a short exact sequence

$$0 \rightarrow \mathcal{O}_X(D) \rightarrow E \rightarrow I_Z(C_0 + \mathbf{m}f - D) \rightarrow 0 \tag{8}$$

where Z is a locally complete intersection 0-cycle of length $|Z| = c_2 - D(C_0 + \mathbf{m}f - D)$.

Since D is effective, by Lemma 2.2, $a \geq 0$ and $b := \deg(\mathbf{b}) \geq 0$.

If $D = 0$, considering the exact sequence (8) and taking cohomology, we have that

$$k \leq h^0 E \leq h^0 \mathcal{O}_X + h^0 I_Z(C_0 + \mathbf{m}f) = 1 + h^0 I_Z(C_0 + \mathbf{m}f).$$

By the short exact sequence

$$0 \rightarrow I_Z \rightarrow \mathcal{O}_X \rightarrow \mathcal{O}_Z \rightarrow 0 \tag{9}$$

and by Lemma 2.3, we have

$$h^0 I_Z(C_0 + \mathbf{m}f) \leq h^0 \mathcal{O}_X(C_0 + \mathbf{m}f) \leq 2(m + 1) - e \leq 4.$$

This implies $k \leq 5$, but this contradicts the fact that $k \geq \frac{1}{2\alpha}[\beta - \alpha(e - m + 2g - 2)] > 5$ since $\beta > \alpha(8 + e - m + 2g)$. Hence, $D \neq 0$.

Now, let us see that if $D \neq 0$, then $D = \mathfrak{b}f$ with $b > 0$.

Since E is a rank two H -stable vector bundle, $\mu_H(\mathcal{O}_X(D)) < \mu_H(E)$ and this is equivalent to

$$\alpha(2b - m) < (1 - 2a)(\beta - \alpha e). \tag{10}$$

Let us see that $a = 0$.

If $a > 0$, since $\beta - \alpha e > \alpha(8 + 2g - m)$ and considering the inequality in (10), we get $\alpha(2b + m) < -\alpha(8 + 2g - m)$ which implies $b < 0$, a contradiction.

Thus, $a = 0$ and hence $D = \mathfrak{b}f$. In addition, since D is a non-zero effective divisor, we get $b > 0$. Therefore, by (8), E sits in the exact sequence

$$0 \rightarrow \mathcal{O}_X(\mathfrak{b}f) \rightarrow E \rightarrow I_Z(C_0 + (\mathfrak{m} - \mathfrak{b})f) \rightarrow 0. \tag{11}$$

with $b \geq 1$ and $|Z| = c_2 - b$.

If we consider the exact sequence (11) and we take cohomology, we get

$$k \leq h^0 E \leq h^0 \mathcal{O}_X(\mathfrak{b}f) + h^0 I_Z(C_0 + (\mathfrak{m} - \mathfrak{b})f).$$

Notice that since $m \in \{0, 1\}$ and $b \geq 1$, we get $m - b \leq 0$.

Let us first assume $m - b = 0$. Then, since $b \geq 1$, $m = b = 1$ and we get

$$k \leq h^0 \mathcal{O}_X(f) + h^0 I_Z(C_0).$$

In this case, since

$$h^0 I_Z(C_0) \leq h^0 \mathcal{O}_X(C_0) \leq h^0 \mathcal{O}_C + h^0 \mathcal{O}_C(-e) \leq 2 - e \leq 2,$$

and

$$h^0 \mathcal{O}_X(f) = h^0 \mathcal{O}_C(1) \leq 2,$$

we get that $k \leq 4$, which is a contradiction.

Let us now study the case $m - b < 0$. This implies $h^0 I_Z(C_0 + (\mathfrak{m} - \mathfrak{b})f) = 0$ and then $k \leq h^0 E = h^0 \mathcal{O}_X(\mathfrak{b}f)$.

First, let us assume $b \geq 2g - 1$. In this case, we get $k \leq h^0 \mathcal{O}_C(\mathfrak{b}) = b + 1 - g$. On the other hand, since $\mathcal{O}_X(\mathfrak{b}f) \hookrightarrow E$, by stability $b < \frac{1}{2\alpha}[\beta - \alpha(e - m)]$.

Putting altogether, we get $k \leq b + 1 - g < \frac{1}{2\alpha}[\beta - \alpha(e - m + 2g - 2)]$, which is a contradiction.

Finally, let us assume $b < 2g - 1$. This implies that $k \leq h^0 \mathcal{O}_C(\mathfrak{b}) \leq b + 1 < 2g$, which again is a contradiction since $k \geq \frac{1}{2\alpha}[\beta - \alpha(e - m + 2g - 2)] > 2g$ by the fact that $\beta > \alpha(6g - 4 + e - m)$.

Hence, $W_H^k(2; C_0 + \mathfrak{m}f, c_2) = \emptyset$ for all $k \geq \frac{1}{2\alpha}[\beta - \alpha(e - m + 2g - 2)]$. □

Remark 4.6. Notice that for $g \geq 3$ we have that

$$\max\{\alpha(8 + e - m + 2g), \alpha(6g - 4 + e - m)\} = \alpha(6g - 4 + e - m).$$

Thus, for these values of g , in Theorem 4.5, we only need to assume that $\beta > \alpha(6g - 4 + e - m)$. Analogously, for $g \leq 2$, it is sufficient to assume $\beta > \alpha(8 + e - m + 2g)$.

Furthermore, if we consider the particular case $g = 1$, the result is also true if we expand the family of ample line bundles on X to the one of $H \equiv \alpha C_0 + \beta f$ with

$$\alpha(6 + e - m) < \beta.$$

As a consequence of the above results, we have the following equivalence.

Corollary 4.7. *Let X be a ruled surface over a nonsingular curve C of genus $g \geq 0$, $m \in \{0, 1\}$, $c_2 \gg 0$ an integer and $H \equiv \alpha C_0 + \beta f$ an ample divisor on X such that*

$$\max\{\alpha(8 + e - m + 2g), \alpha(6g - 4 + e - m)\} < \beta.$$

Assume that $M_H \neq \emptyset$. Then, $W_H^k(2; C_0 + mf, c_2) \neq \emptyset$ if and only if

$$1 \leq k < \frac{1}{2\alpha}[\beta - \alpha(e - m + 2g - 2)].$$

Proof. It follows from Corollary 4.3 and Theorem 4.5. □

Let us now turn our attention to the case $c_1 = f$ and see under which conditions $W_H^k(2; f, c_2)$ is non-empty. The first result is about non-emptiness.

Theorem 4.8. *Let X be a ruled surface over a nonsingular curve C of genus $g \geq 0$, $c_2 \gg 0$ an integer and $H \equiv C_0 + \beta f$ an ample divisor on X . Then, $W_H^1(2; f, c_2) \neq \emptyset$ whenever $M_H \neq \emptyset$.*

Proof. Let us consider the family \mathcal{G} of rank two vector bundles E on X given by a non-trivial extension

$$0 \rightarrow \mathcal{O}_X \rightarrow E \rightarrow I_Z(f) \rightarrow 0, \tag{12}$$

where Z is a 0-dimensional subscheme of length $|Z| = c_2$.

Claim: \mathcal{G} is non-empty.

Proof of the Claim: first, let us see that $\text{Ext}^1(I_Z(f), \mathcal{O}_X) \neq 0$. By Serre’s duality, $\text{ext}^1(I_Z(f), \mathcal{O}_X) = h^1 I_Z(f + K_X)$.

From the short exact sequence (9), by Lemma 2.2 and Serre’s duality, we have

$$h^0 I_Z(f + K_X) \leq h^0 \mathcal{O}_X(f + K_X) = 0$$

and also

$$h^2 I_Z(f + K_X) = h^2 \mathcal{O}_X(f + K_X) = h^0 \mathcal{O}_X(-f) = 0.$$

Therefore,

$$\begin{aligned} h^1 I_Z(f + K_X) &= -\chi I_Z(C_0 + (m - 2b)f + K_X) = |Z| - \chi \mathcal{O}_X(f + K_X) \\ &= |Z| - 1 + g. \end{aligned}$$

Hence, $\text{ext}^1(I_Z(f), \mathcal{O}_X) = |Z| - 1 + g = c_2 - 1 + g \neq 0$.

Any non-zero element in $\text{Ext}^1(I_Z(f), \mathcal{O}_X)$ defines a non-trivial extension of type (12). Notice that by the Cayley–Bacharach property [16, Theorem 12], E is indeed a rank two vector bundle, with $c_1(E) = f$ and $c_2(E) = c_2 \gg 0$. Hence, \mathcal{G} is non-empty and by construction, for any $E \in \mathcal{G}$, $h^0 E \geq 1$.

Now, let us prove that any rank two vector bundle E in \mathcal{G} is stable with respect to $H \equiv C_0 + \beta f$. To this end, since E is a rank two vector bundle on X , we have to check that for any line subbundle $\mathcal{O}_X(G) \hookrightarrow E$ we have $\mu_H(\mathcal{O}_X(G)) = G \cdot H < \mu_H(E) = \frac{c_1(E) \cdot H}{2}$.

By construction, E is given by a non-trivial extension of type (12). Therefore, we have two possibilities: $\mathcal{O}_X(G) \hookrightarrow \mathcal{O}_X$ or $\mathcal{O}_X(G) \hookrightarrow I_Z(f)$.

Case 1 Assume that $\mathcal{O}_X(G) \hookrightarrow \mathcal{O}_X$. Then,

$$\mu_H(\mathcal{O}_X(G)) \leq 0 < \frac{1}{2} = \frac{c_1(E) \cdot H}{2} = \mu_H(E).$$

Case 2 Assume that $\mathcal{O}_X(G) \hookrightarrow I_Z(f)$, where $G = sC_0 + tf$ with $t = \deg(t)$. In this case,

$$f - G = f - sC_0 - tf$$

is an effective divisor, which by Lemma 2.2 implies that $s \leq 0$ and $t \leq 1$.

Notice that since $h^0 I_Z = 0$, $f \neq G$. Since $f - G$ is effective, we have $(f - G) \cdot H > 0$, which is equivalent to $t \leq s(e - \beta)$, and hence

$$\mu_H(\mathcal{O}_X(G)) = s(\beta - e) + t \leq 0 < \mu_H(E).$$

Thus, putting altogether, we have already seen that E is H -stable.

We have proved that any E in \mathcal{G} is a rank two H -stable vector bundle with $h^0 E \geq 1$, what implies that $\mathcal{G} \hookrightarrow W_H^1(2; f, c_2)$. Thus, $W_H^1(2; f, c_2) \neq \emptyset$. □

Let us see now what happens for values of $k \geq 2$. It follows from [8, Proposition 7.7] that $W_H^2(2; f, c_2) \neq \emptyset$. We will prove the case $k \geq 3$. To this end, we will use the following technical Lemma.

Lemma 4.9. *Let X be a ruled surface over a nonsingular curve C of genus $g \geq 0$ and E a rank 2 vector bundle with $c_1(E) = f$. If E is stable respect to the ample divisor $H \equiv C_0 + \beta f$, then $h^0 E \leq h^0 E|_f$.*

Proof. First, let us see that $h^0 E(-f) = 0$. If $h^0 E(-f) \neq 0$, then $\mathcal{O}_X(f) \hookrightarrow E$. Since E is H -stable, we get

$$\mu_H(\mathcal{O}_X(f)) = 1 < \frac{1}{2} = \mu_H(E),$$

which is a contradiction. Hence, $h^0 E(-f) = 0$.

If we twist the short exact sequence

$$0 \rightarrow \mathcal{O}_X(-f) \rightarrow \mathcal{O}_X \rightarrow \mathcal{O}_f \rightarrow 0$$

by E and we take cohomology, using the fact that $h^0 E(-f) = 0$, we get $h^0 E \leq h^0 E|_f$. □

Now, we are ready to study the Brill–Noether locus $W_H^k(2; f, c_2)$ for values of $k \geq 3$.

Theorem 4.10. *Let X be a ruled surface over a nonsingular curve C of genus $g \geq 0$, $c_2 \gg 0$ an integer and $H \equiv C_0 + \beta f$ an ample divisor on X . Then, $W_H^k(2; f, c_2) = \emptyset$ for all $k \geq 3$.*

Proof. Let us assume that $W_H^k(2; f, c_2) \neq \emptyset$ and consider a vector bundle $E \in W_H^k(2; f, c_2)$. Since $h^0 E \geq k > 0$, we can take a non-zero section s of E . We denote by Y its scheme of zeros and by $D = aC_0 + bf$ the maximal effective divisor contained in Y . Then, s can be regarded as a section of $E(-D)$ and its scheme of zeros has codimension greater or equal than two. Thus, we have a short exact sequence

$$0 \rightarrow \mathcal{O}_X(D) \rightarrow E \rightarrow I_Z(f - D) \rightarrow 0 \tag{13}$$

where Z is a locally complete intersection 0-cycle of length $|Z| = c_2 - D(f - D)$.

Since D is effective, by Lemma 2.2, $a \geq 0$ and $b := \deg(\mathfrak{b}) \geq 0$.

Let us see that $D = 0$. Assume that $D \neq 0$. Since D is effective, $D \cdot L > 0$ for any ample divisor L on X . On the other hand, since E is stable respect to $H \equiv C_0 + \beta f$, we have that $\mu_H(\mathcal{O}_X(D)) < \mu_H(E)$.

Putting altogether, we get

$$0 < \mu_H(\mathcal{O}_X(D)) = D \cdot H < \frac{1}{2} = \mu_H(E),$$

which is a contradiction. Hence, $D = 0$.

Therefore, by (13), any vector bundle $E \in W_H^k(2; f, c_2)$ sits in a short exact sequence

$$0 \rightarrow \mathcal{O}_X \rightarrow E \rightarrow I_Z(f) \rightarrow 0. \tag{14}$$

If $h^0 I_Z(f) = 0$, then $h^0 E = h^0 \mathcal{O}_X = 1$. On the other hand, if $h^0 I_Z(f) \neq 0$ this would mean that the points of Z are contained in a line.

Let l be a line such that $Z \cap l = \emptyset$. Restricting the exact sequence (14) to l , we get the short exact sequence

$$0 \rightarrow \mathcal{O}_l \rightarrow E|_l \rightarrow \mathcal{O}_l \rightarrow 0,$$

and taking cohomology, we obtain $h^0 E|_l \leq 2$.

By Lemma 4.9, this implies that $h^0 E \leq 2$ and thus $W_H^k(2; f, c_2) = \emptyset$ for all $k \geq 3$. □

Putting altogether,

Corollary 4.11. *Let X be a ruled surface over a nonsingular curve C of genus $g \geq 0$, $c_2 \gg 0$ an integer and $H \equiv C_0 + \beta f$ an ample divisor on X . Then, $W_H^k(2; f, c_2) \neq \emptyset$ if and only if $1 \leq k \leq 2$.*

Proof. It follows from Theorem 4.8, [8, Proposition 7.7] and Theorem 4.10. □

See [8, Proposition 7.7] for an alternative proof of Theorems 4.8 and 4.10.

Now, we will analyze in more detail the locus $W_H^1(2; f, c_2) \setminus W_H^2(2; f, c_2)$.

Proposition 4.12. *Let X be a ruled surface over a nonsingular curve C of genus $g \geq 0$, $c_2 \gg 0$ an integer and $H \equiv C_0 + \beta f$ an ample divisor on X . Then, $W_H^1(2; f, c_2) \setminus W_H^2(2; f, c_2)$ is smooth and irreducible of the expected dimension, namely*

$$\rho_H^1 = 3c_2 + g - 1.$$

Proof. Let us take a vector bundle $E \in W_H^1(2; f, c_2) \setminus W_H^2(2; f, c_2)$. Since $h^0 E = 1$, we can take a non-zero section s of E . We denote by Y its scheme of zeros and by $D = aC_0 + \mathfrak{b}f$ the maximal effective divisor contained in Y . Then, s can be regarded as a section of $E(-D)$ and its scheme of zeros has codimension greater or equal than two. Thus, we have a short exact sequence

$$0 \rightarrow \mathcal{O}_X(D) \rightarrow E \rightarrow I_Z(f - D) \rightarrow 0 \tag{15}$$

where Z is a locally complete intersection 0-cycle of length $|Z| = c_2 - D(f - D)$.

Following the same arguments as in proof of Theorem 4.10, we see that $D = 0$. Hence, any vector bundle E in $W_H^1(2; f, c_2) \setminus W_H^2(2; f, c_2)$ sits in a short exact sequence of the following type:

$$0 \rightarrow \mathcal{O}_X \rightarrow E \rightarrow I_Z(f) \rightarrow 0. \tag{16}$$

Let us now check that any vector bundle E in $W_H^1(2; f, c_2) \setminus W_H^2(2; f, c_2)$ is a smooth point of $M_H(2; f, c_2)$. To this end, it is enough to see that $h^2(E^* \otimes E) = 0$.

First, let us see that $h^2(E^*) = 0$. If we tensor the short exact sequence (16) by $\mathcal{O}_X(K_X)$ and we take cohomology, we get the long exact sequence

$$0 \rightarrow H^0 \mathcal{O}_X(K_X) \rightarrow H^0 E(K_X) \rightarrow H^0 I_Z(f + K_X) \rightarrow \dots$$

From the fact that $H^0 \mathcal{O}_X(K_X) = H^0 I_Z(f + K_X) = 0$, we deduce that

$$0 = h^0 E(K_X) = h^2 E^*,$$

where the last equality follows from Serre duality. Now, we will see that

$$h^2(E^* \otimes \mathcal{O}_X(f) \otimes I_Z) = 0.$$

Tensoring the short exact sequence (16) by $\mathcal{O}_X(K_X - f)$ and taking cohomology, we get the long exact sequence

$$0 \rightarrow H^0 \mathcal{O}_X(K_X - f) \rightarrow H^0 E(K_X - f) \rightarrow H^0 I_Z(K_X) \rightarrow \dots$$

Since $H^0 \mathcal{O}_X(K_X - f) = H^0 I_Z(K_X) = 0$, we get $H^0 E(K_X - f) = 0$ and by Serre’s duality $H^2 E^*(f) = 0$. Hence, $h^2(E^* \otimes \mathcal{O}_X(f) \otimes I_Z) = h^2(E^* \otimes \mathcal{O}_X(f)) = 0$.

Finally, if we tensor the short exact sequence (16) by E^* and we take cohomology, we get

$$\dots \rightarrow H^2(E^*) \rightarrow H^2(E^* \otimes E) \rightarrow H^2(E^* \otimes \mathcal{O}_X(f) \otimes I_Z) \rightarrow 0$$

and since $H^2(E^*) = H^2(E^* \otimes \mathcal{O}_X(f) \otimes I_Z) = 0$ we deduce that $H^2(E^* \otimes E) = 0$. Hence, E is a smooth point of $M_H(2; f, c_2)$.

To end the proof, let us see that the map

$$\mu_E : H^0(E) \otimes H^1(E^* \otimes \mathcal{O}_X(K_X)) \rightarrow H^1(E \otimes E^* \otimes \mathcal{O}_X(K_X))$$

is injective. Since $E \in W_H^1(2; f, c_2) \setminus W_H^2(2; f, c_2)$, $H^0(E) \cong K$ and hence this is equivalent to prove that the map

$$\tilde{\mu}_E : H^1(E^* \otimes \mathcal{O}_X(K_X)) \rightarrow H^1(E \otimes E^* \otimes \mathcal{O}_X(K_X))$$

is injective.

If we tensor the exact sequence (16) by $\mathcal{O}_X(-f)$ and we take cohomology, we get the long exact sequence

$$\cdots \rightarrow H^2 \mathcal{O}_X(-f) \rightarrow H^2 E(-f) \rightarrow H^2 I_Z \rightarrow 0.$$

Since by duality $h^2 \mathcal{O}_X(-f) = h^0 \mathcal{O}_X(f + K_X) = 0$ and $h^2 I_Z = h^2 \mathcal{O}_X = 0$, we deduce that $h^2 E(-f) = 0$ and this implies, again by Serre’s duality, that $h^0 E^*(K_X + f) = 0$. Moreover, if $h^0 E^*(K_X + f) = 0$, we get $h^0(E^* \otimes_{I_Z}(K_X + f)) = 0$.

Therefore, since $h^0(E^* \otimes_{I_Z}(K_X + f)) = 0$, if we tensor the exact sequence (16) by $E^* \otimes \mathcal{O}_X(K_X)$ and we take cohomology, we get

$$0 \rightarrow H^1(E^* \otimes \mathcal{O}_X(K_X)) \xrightarrow{\tilde{\mu}_E} H^1(E \otimes E^* \otimes \mathcal{O}_X(K_X)) \rightarrow \cdots$$

which implies that $\tilde{\mu}_E$ is injective. Hence, μ_E is injective.

We have proved that, for any vector bundle $E \in W_H^1(2; f, c_2) \setminus W_H^2(2; f, c_2)$, E is a smooth point of $M_{X,H}(2; f, c_2)$ and μ_E is injective.

Therefore, by [17, Corollary 2.6], $W_H^1(2; f, c_2) \setminus W_H^2(2; f, c_2)$ is smooth and of the expected dimension at E , for all $E \in W_H^1(2; f, c_2) \setminus W_H^2(2; f, c_2)$. Hence, $W_H^1(2; f, c_2) \setminus W_H^2(2; f, c_2)$ is smooth and of the expected dimension.

It only remains to prove the irreducibility of $W_H^1(2; f, c_2) \setminus W_H^2(2; f, c_2)$. We have already seen that any $E \in W_H^1(2; f, c_2) \setminus W_H^2(2; f, c_2)$ sits in an extension of type (16). Denote by \mathcal{S} the family of rank two vector bundles E on X given by an extension of type (16). Notice that, since $\text{ext}^1(I_Z(f), \mathcal{O}_X) = c_2 + g$, \mathcal{S} is a $(c_2 + g - 1)$ -projective bundle over de Hilbert scheme $\mathcal{H}ilb^{c_2}(X)$, which parametrizes 0-dimensional subschemes Z of length $|Z| = c_2$. Thus, \mathcal{S} is an irreducible family of bundles of dimension $3c_2 + g - 1 = \rho_H^1(2; f, c_2)$ and there is a modular map π from an open dense subset of \mathcal{S} to $W_H^1(2; f, c_2) \setminus W_H^2(2; f, c_2)$. Denote by W the irreducible component of $W_H^1(2; f, c_2) \setminus W_H^2(2; f, c_2)$ containing $\text{im}(\pi)$. By the above arguments, the general point of $\text{im}(\pi)$ is a smooth point of W . Hence, W has the expected dimension $3c_2 + g - 1$. Since $\dim W = \dim \mathcal{S}$, π must be generically finite onto an open subset of W , and thus \mathcal{S} finitely dominates W . Finally, since we have seen that any isomorphism class $[E]$ in $W_H^1(2; f, c_2) \setminus W_H^2(2; f, c_2)$ must arise as a non-trivial extension of type (16) for some $Z \in \mathcal{H}ilb^{c_2}(X)$, \mathcal{S} finitely dominates the whole $W_H^1(2; f, c_2) \setminus W_H^2(2; f, c_2)$, and then $W = W_H^1(2; f, c_2) \setminus W_H^2(2; f, c_2)$. Therefore, $W_H^1(2; f, c_2) \setminus W_H^2(2; f, c_2)$ is irreducible. \square

Acknowledgements

I. M. Tarrío thanks professor Marian Aprodu for his advice and comments and for his hospitality during her stay in Bucharest. Both authors want to thank the anonymous referee that in particular improved the statement and proof of Proposition 4.12.

Author contributions All authors wrote the main parts of the manuscript.

Funding Open Access funding provided thanks to the CRUE-CSIC agreement with Springer Nature.

Data Availability No datasets were generated or analysed during the current study.

Declarations

Conflict of Interest The authors declare no competing interests.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

References

- [1] Arbarello, E., Cornalba, M., Griffiths, P.A., Harris, J.: Geometry of Algebraic Curves, vol. I. In: Grundlehren der Mathematischen Wissenschaften (Fundamental Principles of Mathematical Sciences), vol. 267. Springer, New York (1985)
- [2] Brambila-Paz, L., Grzegorzczak, I., Newstead, P.E.: Geography of Brill–Noether loci for small slopes. *J. Algebr. Geom.* **6**(4), 645–669 (1997)
- [3] Grzegorzczak, I., Teixidor i Bigas, M.: Brill–Noether theory for stable vector bundles. *Lond. Math. Soc. Lect. Note Ser.* **359**, 29–50 (2009)
- [4] Costa, L., Miró-Roig, R.M.: Brill–Noether theory for moduli spaces of sheaves on algebraic varieties. *Forum Math.* **22**(3), 411–432 (2010)
- [5] Leyenson, M.: On the Brill–Noether theory for K3 surfaces. *Cent. Eur. J. Math.* **10**(4), 1486–1540 (2012)
- [6] Leyenson, M.: On the Brill–Noether theory for K3 II surfaces. Preprint (2018). <https://arXiv.org/pdf/math/0602358>
- [7] Nakashima, T.: Brill–Noether problems in higher dimensions. *Forum Math.* **20**, 145–161 (2008)
- [8] Coskun, I., Huizenga, J., Nuer, H.: Brill–Noether theory of moduli spaces of sheaves on surfaces. Preprint (2023). <https://arXiv.org/pdf/2306.11033.pdf>
- [9] Costa, L., Miró-Roig, R.M.: Brill–Noether theory on Hirzebruch surfaces. *J. Pure Appl. Algebra* **214**(9), 1612–1622 (2010)
- [10] Reyes-Ahumada, G., Roa-Leguizamón, L., Torres-López, H.: A note on rank two stable bundles over surfaces. *Glasg. Math. J.* **64**(2), 397–410 (2022)
- [11] Qin, Z.: Equivalence classes of polarizations and moduli spaces of sheaves. *J. Differ. Geom.* **37**, 397–415 (1993)

- [12] Hartshorne, R.: Algebraic Geometry. G.T.M., vol. 52. Springer, Berlin (1977)
- [13] Aprodu, M., Casnati, G., Costa, L., Miró-Roig, R.M., Teixidor i Bigas, M.: Theta divisors and Ulrich bundles on geometrically ruled surfaces. *Ann. Mat. Pura Appl. (4)* **199**(1), 199–216 (2020)
- [14] Huybrechts, D., Lehn, M.: The Geometry of Moduli Spaces of Sheaves, 2nd edn. Cambridge University Press, Cambridge (2010)
- [15] Qin, Z.: Moduli spaces of stable rank-2 bundles on ruled surfaces. *Invent. Math.* **110**, 615–626 (1992)
- [16] Friedman, R.: Algebraic Surfaces and Holomorphic Vector Bundles. Springer, New York (1998)
- [17] Filimon, L.: On translated rank-2 Brill–Noether loci on regular surfaces. *Arch. Math.* **118**(3), 271–281 (2022)

L. Costa and Irene Macías Tarrío
Departament de Matemàtiques i Informàtica
Facultat de Matemàtiques i Informàtica
Gran Via de les Corts Catalanes 585
08007 Barcelona
Spain
e-mail: costa@ub.edu

Irene Macías Tarrío
e-mail: irene.macias@ub.edu

Received: January 27, 2024.

Revised: April 15, 2024.

Accepted: April 17, 2024.