



Mathematical Theory of Compressible Magnetohydrodynamics Driven by Non-conservative Boundary Conditions

Eduard Feireisl, Piotr Gwiazda, Young-Sam Kwon and Agnieszka Świerczewska-Gwiazda

Communicated by G. P. Galdi

Abstract. We propose a new concept of weak solution to the equations of compressible magnetohydrodynamics driven by inhomogeneous boundary data. The system of the underlying field equations is solvable globally in time in the out of equilibrium regime characteristic for turbulence. The weak solutions comply with the weak–strong uniqueness principle; they coincide with the classical solution of the problem as long as the latter exists. The choice of constitutive relations is motivated by applications in stellar magnetoconvection.

Keywords. Compressible MHD system, Weak solution, Stellar magnetoconvection.

Contents

| | |
|---|----|
| 1. Introduction | 2 |
| 2. Weak Formulation | 5 |
| 2.1. Equation of State | 5 |
| 2.2. Transport Coefficients | 6 |
| 2.3. Boundary Data | 6 |
| 2.4. Weak Solutions | 6 |
| 3. Relative Energy | 9 |
| 3.1. Relative Energy Inequality | 10 |
| 3.1.1. Momentum Perturbation | 10 |
| 3.1.2. Density Perturbation | 11 |
| 3.1.3. Magnetic Field Perturbation and the Final form of Relative Energy Inequality | 11 |
| 4. Weak–Strong Uniqueness Principle | 12 |
| 4.1. Magnetic Field | 13 |
| 4.2. Pressure and Entropy | 16 |
| 4.3. Diffusive Terms | 18 |
| 4.4. Refined Estimates | 19 |
| 4.4.1. Temperature Gradient | 19 |
| 4.4.2. Viscous Stress and Magnetic Diffusion | 21 |
| 4.5. Completing the Proof of Weak–Strong Uniqueness | 22 |
| 5. Existence of Weak Solutions | 22 |
| 5.1. Approximation Scheme | 23 |

The work of E.F. was partially supported by the Czech Sciences Foundation (GAČR), Grant Agreement 21-02411S. The Institute of Mathematics of the Academy of Sciences of the Czech Republic is supported by RVO:67985840. This work was partially supported by the Thematic Research Programme, University of Warsaw, Excellence Initiative Research University. The work of P.G. and A. Ś-G. was partially supported by National Science Centre (Poland), Agreement No. 2021/43/B/ST1/02851. The work of Y.-S. Kwon was partially supported by the National Research Foundation of Korea (NRF2022R1F1A1073801).

| | |
|----------------------------------|----|
| 5.2. Existence of Weak Solutions | 24 |
| 6. Concluding Remarks | 25 |
| References | 26 |

1. Introduction

The large scale dynamics arising in stellar magnetoconvection is driven in an essential way by the boundary conditions, see Gough [12]. The flux separation and pattern formation arise in the turbulent regime when the fluid flow is far from equilibrium. As the underlying field equations are non-linear, the existence of global in time smooth solutions in such a regime is not known. Indeed the vast majority of *global* existence results in the class of classical solutions is restricted to the initial state close to a stable equilibrium, see e.g. Matsumura and Nishida [17], Valli [24], among others. Moreover, the recent results of Buckmaster et al. [4], and Merle et al. [18, 19] indicate that classical solutions may develop singularities in a finite time.

In view of the above arguments, the concept of *weak solution* represents a suitable alternative to restore global-in-time existence even for problems with large data and solutions remaining out of equilibrium in the long run. We consider a mathematical model of fully compressible three dimensional fluid convection driven by an externally imposed magnetic field. The fluid occupies a bounded domain $\Omega \subset \mathbb{R}^3$ with regular boundary. The time evolution of the density $\varrho = \varrho(t, x)$, the (absolute) temperature $\vartheta = \vartheta(t, x)$, the velocity field $\mathbf{u} = \mathbf{u}(t, x)$, and the magnetic field $\mathbf{B} = \mathbf{B}(t, x)$ is governed by the *compressible MHD system*.

Equation of continuity

$$\partial_t \varrho + \operatorname{div}_x(\varrho \mathbf{u}) = 0. \quad (1.1)$$

Momentum equation

$$\partial_t(\varrho \mathbf{u}) + \operatorname{div}_x(\varrho \mathbf{u} \otimes \mathbf{u}) + \nabla_x p(\varrho, \vartheta) = \operatorname{div}_x \mathbb{S}(\vartheta, \nabla_x \mathbf{u}) + \operatorname{curl}_x \mathbf{B} \times \mathbf{B} + \varrho \mathbf{g}. \quad (1.2)$$

Induction equation

$$\partial_t \mathbf{B} + \operatorname{curl}_x(\mathbf{B} \times \mathbf{u}) + \operatorname{curl}_x(\zeta(\vartheta) \operatorname{curl}_x \mathbf{B}) = 0, \quad \operatorname{div}_x \mathbf{B} = 0. \quad (1.3)$$

Internal energy balance

$$\begin{aligned} \partial_t(\varrho e(\varrho, \vartheta)) + \operatorname{div}_x(\varrho e(\varrho, \vartheta) \mathbf{u}) + \operatorname{div}_x \mathbf{q}(\vartheta, \nabla_x \vartheta) \\ = \mathbb{S}(\vartheta, \nabla_x \mathbf{u}) : \nabla_x \mathbf{u} + \zeta(\vartheta) |\operatorname{curl}_x \mathbf{B}|^2 - p(\varrho, \vartheta) \operatorname{div}_x \mathbf{u}. \end{aligned} \quad (1.4)$$

The pressure $p = p(\varrho, \vartheta)$ and the internal energy $e = e(\varrho, \vartheta)$ are interrelated through *Gibbs' equation*

$$\vartheta Ds = De + pD\left(\frac{1}{\varrho}\right), \quad (1.5)$$

where $s = s(\varrho, \vartheta)$ is the entropy. Consequently, the internal energy balance (1.4) may be reformulated in the form of *entropy balance equation*

$$\begin{aligned} \partial_t(\varrho s(\varrho, \vartheta)) + \operatorname{div}_x(\varrho s(\varrho, \vartheta) \mathbf{u}) + \operatorname{div}_x \left(\frac{\mathbf{q}(\vartheta, \nabla_x \vartheta)}{\vartheta} \right) \\ = \frac{1}{\vartheta} \left(\mathbb{S}(\vartheta, \nabla_x \mathbf{u}) : \nabla_x \mathbf{u} - \frac{\mathbf{q}(\vartheta, \nabla_x \vartheta) \cdot \nabla_x \vartheta}{\vartheta} + \zeta(\vartheta) |\operatorname{curl}_x \mathbf{B}|^2 \right), \end{aligned} \quad (1.6)$$

see e.g. Gough [12], Weiss and Proctor [26], Tao et al. [22].

The right-hand side of equation (1.4) represents the entropy production rate, which, in accordance with the Second law of thermodynamics, must be non-negative. Accordingly, we consider *Newtonian fluid*, with the viscous stress tensor

$$\mathbb{S}(\vartheta, \nabla_x \mathbf{u}) = \mu(\vartheta) \left(\nabla_x \mathbf{u} + \nabla_x^t \mathbf{u} - \frac{2}{3} \operatorname{div}_x \mathbf{u} \mathbb{I} \right) + \eta(\vartheta) \operatorname{div}_x \mathbf{u} \mathbb{I}, \tag{1.7}$$

where the viscosity coefficients $\mu > 0$ and $\eta \geq 0$ are continuously differentiable functions of the temperature. Similarly, the heat flux obeys *Fourier’s law*,

$$\mathbf{q}(\vartheta, \nabla_x \vartheta) = -\kappa(\vartheta) \nabla_x \vartheta, \tag{1.8}$$

where the heat conductivity coefficient $\kappa > 0$ is a continuously differentiable function of the temperature.

The existence of global-in-time weak solutions for the compressible MHD system (1.1)–(1.8) was shown in [7] under certain physically relevant restrictions imposed on constitutive equations and transport coefficients. The boundary conditions considered in [7] are conservative,

$$\mathbf{u}|_{\partial\Omega} = 0, \quad \nabla_x \vartheta \cdot \mathbf{n}|_{\partial\Omega} = 0, \quad \mathbf{B} \cdot \mathbf{n}|_{\partial\Omega} = 0, \quad [\mathbf{B} \times \mathbf{u} + \zeta \operatorname{curl}_x \mathbf{B}] \times \mathbf{n}|_{\partial\Omega} = 0 \tag{1.9}$$

characteristic for closed systems. If the driving force \mathbf{g} in the momentum equation is conservative, meaning $\mathbf{g} = \nabla_x G$, $G = G(x)$, the total energy of the system is conserved and the dynamics obeys the rather “boring” scenario formulated by the celebrated statement of Clausius:

The energy of the world is constant; its entropy tends to a maximum.

A rigorous mathematical proof of this statement for the Navier–Stokes–Fourier system was given in [11].

As pointed out by Gough [12], a rich fluid behaviour is conditioned by a proper choice of boundary conditions. Motivated by models in astrophysics, we suppose the boundary $\partial\Omega$ is impermeable and the tangential component of the normal viscous stress vanishes on it,

$$\mathbf{u} \cdot \mathbf{n}|_{\partial\Omega} = 0, \quad (\mathbb{S} \cdot \mathbf{n}) \times \mathbf{n}|_{\partial\Omega} = 0. \tag{1.10}$$

However, the theory presented in this paper can accommodate more general boundary conditions for the velocity field discussed in Sect. 6.

Similarly to the well known Rayleigh–Bénard problem, see e.g. Davidson [6], we impose the inhomogeneous Dirichlet boundary conditions for the temperature,

$$\vartheta|_{\partial\Omega} = \vartheta_B, \quad \vartheta_B > 0. \tag{1.11}$$

To incorporate the effect of an exterior magnetic field, we suppose that \mathbf{B} is (not necessarily small) perturbation of a background magnetic field \mathbf{B}_B , $\operatorname{div}_x \mathbf{B}_B = 0$,

$$\mathbf{B} = \mathbf{B}_B + \mathbf{b}.$$

In the context of astrophysics, we may think of \mathbf{B}_B as being the magnetic field imposed by a massive star. As for \mathbf{b} , we suppose either

$$\mathbf{b} \times \mathbf{n}|_{\partial\Omega} = 0 \quad \Rightarrow \quad \mathbf{B} \times \mathbf{n}|_{\partial\Omega} = \mathbf{B}_B \times \mathbf{n}|_{\partial\Omega}, \tag{1.12}$$

or

$$\mathbf{b} \cdot \mathbf{n}|_{\partial\Omega} = 0 \quad \Rightarrow \quad \mathbf{B} \cdot \mathbf{n}|_{\partial\Omega} = \mathbf{B}_B \cdot \mathbf{n}|_{\partial\Omega}. \tag{1.13}$$

The boundary condition (1.12) is of Dirichlet type, compatible with the requirement of solenoidality of \mathbf{B} . The condition (1.13) is of flux type, similar to the first condition (1.10) and must be accompanied by another flux condition related to the electric field, namely

$$[\mathbf{B} \times \mathbf{u} + \zeta \operatorname{curl}_x \mathbf{B}] \times \mathbf{n}|_{\partial\Omega} = 0. \tag{1.14}$$

The flux in (1.14) may be inhomogeneous as well. The borderline case $\mathbf{B}_B = 0$ in (1.12) and (1.13) corresponds to a perfectly isolating and perfectly conducting boundary, respectively (see e.g. Alekseev [1]). There are other types of boundary conditions discussed in the literature, see e.g. Ladyzhenskaya and Solonnikov [14], Lassner [15], Ströhmer [21], among others.

Due to possibly only technical but so far unsurmountable difficulties, a mathematically tractable *weak formulation* of compressible MHD system cannot be based on merely rewriting the system (1.1)–(1.4) in the sense of distributions. The available *a priori* bounds are not strong enough to render certain terms, notably $p(\varrho, \vartheta)\operatorname{div}_x \mathbf{u}$ in the internal energy balance (1.4), integrable. A remedy proposed in [7] is replacing (1.4) by the entropy equation (1.6). Unfortunately, the entropy production rate is *a priori* bounded only in the non-reflexive space L^1 of integrable functions. As a result, any approximation scheme provides merely an *inequality*

$$\begin{aligned} & \partial_t(\varrho s(\varrho, \vartheta)) + \operatorname{div}_x(\varrho s(\varrho, \vartheta)\mathbf{u}) + \operatorname{div}_x\left(\frac{\mathbf{q}(\vartheta, \nabla_x \vartheta)}{\vartheta}\right) \\ & \geq \frac{1}{\vartheta}\left(\mathbb{S}(\vartheta, \nabla_x \mathbf{u}) : \nabla_x \mathbf{u} - \frac{\mathbf{q}(\vartheta, \nabla_x \vartheta) \cdot \nabla_x \vartheta}{\vartheta} + \zeta(\vartheta)|\operatorname{curl}_x \mathbf{B}|^2\right) \end{aligned} \tag{1.15}$$

satisfied in the sense of distributions.

Replacing entropy equation by inequality definitely enlarges the class of possible solutions and gives rise to an underdetermined problem. To save well posedness, at least formally, the integrated *total energy balance/inequality* is appended to the system (1.1)–(1.3), (1.15) in [7]. The total energy balance reads

$$\begin{aligned} & \partial_t\left(\frac{1}{2}\varrho|\mathbf{u}|^2 + \varrho e(\varrho, \vartheta) + \frac{1}{2}|\mathbf{B}|^2\right) + \operatorname{div}_x\left[\left(\frac{1}{2}\varrho|\mathbf{u}|^2 + \varrho e(\varrho, \vartheta) + \frac{1}{2}|\mathbf{B}|^2 + p(\varrho, \vartheta)\right)\mathbf{u}\right] \\ & + \operatorname{div}_x\left[(\mathbf{B} \times \mathbf{u}) \times \mathbf{B} + \zeta(\vartheta)\operatorname{curl}_x \mathbf{B} \times \mathbf{B}\right] - \operatorname{div}_x(\mathbb{S}(\vartheta, \nabla_x \mathbf{u}) \cdot \mathbf{u}) + \operatorname{div}_x \mathbf{q}(\vartheta, \nabla_x \vartheta) = \varrho \mathbf{g} \cdot \mathbf{u}. \end{aligned} \tag{1.16}$$

If the system is *conservative*, the total energy flux vanishes on the boundary and (1.16) integrated over Ω yields

$$\frac{d}{dt} \int_{\Omega} \left(\frac{1}{2}\varrho|\mathbf{u}|^2 + \varrho e(\varrho, \vartheta) + \frac{1}{2}|\mathbf{B}|^2\right) dx = \int_{\Omega} \varrho \mathbf{g} \cdot \mathbf{u} dx. \tag{1.17}$$

It turns out that the equations (1.1)–(1.3), the inequality (1.15), together with the integral identity (1.17) represent a suitable weak formulation for the *conservative* system. Indeed the existence of global-in-time weak solutions for any finite energy (initial) data was proved in [7]. The weak solutions comply with a natural *compatibility* principle. Any sufficiently smooth weak solution is a classical solution of the problem. In addition, the weak solutions satisfy the weak–strong uniqueness principle: A weak solution coincides with the strong solution on the life span of the latter, see [8].

The situation becomes more delicate for open (non-conservative) systems, with inhomogeneous Dirichlet boundary conditions. The heat flux $\mathbf{q} \cdot \mathbf{n}$ as well as the induction flux (the normal component of the electric field) $[\mathbf{B} \times \mathbf{u} \times \mathbf{B} + \zeta \operatorname{curl}_x \mathbf{B} \times \mathbf{B}] \cdot \mathbf{n}$ do not necessarily vanish on $\partial\Omega$ and give rise to additional uncontrollable sources of energy in (1.17). In [5] (see also the monograph [10]), a new approach has been developed to handle the inhomogeneous Dirichlet boundary conditions for the temperature. The total energy balance (1.17) is replaced by a similar inequality for the ballistic energy

$$\left[\frac{1}{2}\varrho|\mathbf{u}|^2 + \varrho e(\varrho, \vartheta) + \frac{1}{2}|\mathbf{B}|^2 - \tilde{\vartheta}\varrho s(\varrho, \vartheta)\right],$$

where $\tilde{\vartheta}$ is an arbitrary extension of the boundary temperature ϑ_B inside Ω . Pursuing the same strategy, we introduce a “magnetic” variant of the ballistic energy

$$E_{BM} = \left[\frac{1}{2}\varrho|\mathbf{u}|^2 + \varrho e + \frac{1}{2}|\mathbf{B}|^2 - \tilde{\vartheta}\varrho s - \mathbf{B}_B \cdot \mathbf{B}\right], \tag{1.18}$$

After a straightforward manipulation, we deduce equality

$$\begin{aligned} & \frac{d}{dt} \int_{\Omega} \left(\frac{1}{2}\varrho|\mathbf{u}|^2 + \varrho e + \frac{1}{2}|\mathbf{B}|^2 - \tilde{\vartheta}\varrho s - \mathbf{B}_B \cdot \mathbf{B}\right) dx \\ & + \int_{\Omega} \frac{\tilde{\vartheta}}{\vartheta} \left(\mathbb{S} : \nabla_x \mathbf{u} - \frac{\mathbf{q} \cdot \nabla_x \vartheta}{\vartheta} + \zeta|\operatorname{curl}_x \mathbf{B}|^2\right) dx \end{aligned}$$

$$\begin{aligned}
 &= - \int_{\Omega} \left(\varrho s \partial_t \tilde{\vartheta} + \varrho s \mathbf{u} \cdot \nabla_x \tilde{\vartheta} + \frac{\mathbf{q}}{\vartheta} \cdot \nabla_x \tilde{\vartheta} \right) dx \\
 &\quad - \int_{\Omega} \left(\mathbf{B} \cdot \partial_t \mathbf{B}_B - (\mathbf{B} \times \mathbf{u}) \cdot \mathbf{curl}_x \mathbf{B}_B - \zeta \mathbf{curl}_x \mathbf{B} \cdot \mathbf{curl}_x \mathbf{B}_B \right) dx \\
 &\quad + \int_{\Omega} \varrho \mathbf{g} \cdot \mathbf{u} dx
 \end{aligned} \tag{1.19}$$

as soon as $\tilde{\vartheta}|_{\partial\Omega} = \vartheta_B$, $\tilde{\vartheta} > 0$, $\mathbf{B} \times \mathbf{n}|_{\partial\Omega} = \mathbf{B}_B \times \mathbf{n}|_{\partial\Omega}$. The same is true if $\mathbf{B} \cdot \mathbf{n}|_{\partial\Omega} = \mathbf{B}_B \cdot \mathbf{n}|_{\partial\Omega}$ and the vanishing tangential flux condition (1.14) holds. Note carefully that (1.19) does not contain the boundary flux integrals; whence it is suitable for a weak formulation of the problem.

The mathematical theory we propose is based on imposing the weak form of the equations (1.1)–(1.3), together with the entropy inequality (1.15), and the ballistic energy balance (1.19) as a weak formulation of the compressible MHD system. The basic hypotheses concerning the state equation and the transport coefficients as well as the exact definition of weak solution are given in Sect. 2. In Sect. 3 we introduce the basic tool to investigate stability properties in the class of weak solutions—the *relative energy inequality*. In Sect. 4, we show that the weak solutions enjoy the weak–strong uniqueness property. In Sect. 5, the existence of global-in-time weak solutions is established by means of a multilevel approximation scheme. The paper is concluded in Sect. 6 by a brief discussion on possible extensions of the theory to a larger class of boundary conditions.

2. Weak Formulation

We start by a list of structural hypotheses imposed on the equation of state and the transport coefficients.

2.1. Equation of State

Our choice of the equation of state is motivated by [10, Chapter 4]. In particular, we consider the radiation contribution to the pressure/internal energy relevant to problems in astrophysics, cf. Battaner [3]. In accordance with Gibbs’ relation (1.5) we suppose that

$$\begin{aligned}
 p(\varrho, \vartheta) &= p_M(\varrho, \vartheta) + p_R(\vartheta), \quad p_M(\varrho, \vartheta) = \vartheta^{\frac{5}{2}} P\left(\frac{\varrho}{\vartheta^{\frac{3}{2}}}\right), \quad p_R(\vartheta) = \frac{a}{3} \vartheta^4, \\
 e(\varrho, \vartheta) &= e_M(\varrho, \vartheta) + e_R(\varrho, \vartheta), \quad e_M(\varrho, \vartheta) = \frac{3}{2} \frac{\vartheta^{\frac{5}{2}}}{\varrho} P\left(\frac{\varrho}{\vartheta^{\frac{3}{2}}}\right), \quad e_R(\varrho, \vartheta) = \frac{a}{\varrho} \vartheta^4, \quad a > 0,
 \end{aligned} \tag{2.1}$$

where the function $P \in C^1[0, \infty)$ satisfies

$$P(0) = 0, \quad P'(Z) > 0 \text{ for } Z \geq 0, \quad 0 < \frac{\frac{5}{3}P(Z) - P'(Z)Z}{Z} \leq c \text{ for } Z > 0. \tag{2.2}$$

This implies, in particular, that $Z \mapsto P(Z)/Z^{\frac{5}{3}}$ is decreasing, and we suppose

$$\lim_{Z \rightarrow \infty} \frac{P(Z)}{Z^{\frac{5}{3}}} = p_{\infty} > 0. \tag{2.3}$$

Note that p_M, e_M satisfy the relation characteristic for monoatomic gas,

$$p_M = \frac{2}{3} \varrho e_M,$$

p_R is the radiation pressure with the associated internal energy e_R , and (2.3) corresponds to the presence of electron pressure under the degenerate gas regime. It follows from (2.2) that p and e satisfy the *hypothesis of thermodynamic stability*:

$$\frac{\partial p(\varrho, \vartheta)}{\partial \varrho} > 0, \quad \frac{\partial e(\varrho, \vartheta)}{\partial \vartheta} > 0. \tag{2.4}$$

In accordance with (1.5), the entropy takes the form

$$s(\varrho, \vartheta) = s_M(\varrho, \vartheta) + s_R(\varrho, \vartheta), \quad s_M(\varrho, \vartheta) = \mathcal{S}\left(\frac{\varrho}{\vartheta^{\frac{3}{2}}}\right), \quad s_R(\varrho, \vartheta) = \frac{4a}{3} \frac{\vartheta^3}{\varrho}, \tag{2.5}$$

where

$$\mathcal{S}'(Z) = -\frac{3}{2} \frac{\frac{5}{3}P(Z) - P'(Z)Z}{Z^2}. \tag{2.6}$$

2.2. Transport Coefficients

We suppose the viscosity coefficients in the Newtonian stress $\mathbb{S}(\vartheta, \nabla_x \mathbf{u})$ are continuously differentiable functions of the temperature satisfying

$$\begin{aligned} 0 < \underline{\mu}(1 + \vartheta^\alpha) \leq \mu(\vartheta) \leq \bar{\mu}(1 + \vartheta^\alpha), \quad |\mu'(\vartheta)| \leq c \text{ for all } \vartheta \geq 0, \\ 0 \leq \underline{\eta}(1 + \vartheta^\alpha) \leq \eta(\vartheta) \leq \bar{\eta}(1 + \vartheta^\alpha) \text{ for some } \frac{1}{2} \leq \alpha \leq 1. \end{aligned} \tag{2.7}$$

Note that $\alpha = \frac{1}{2}$ corresponds to Sutherland’s law relevant in astrophysics, see Yang et al. [27]. We also allow $\underline{\eta} = 0$.

Similarly, the heat conductivity coefficient in the Fourier heat flux $\mathbf{q}(\vartheta, \nabla_x \vartheta)$ is a continuously differentiable function of the temperature satisfying

$$0 < \underline{\kappa}(1 + \vartheta^\beta) \leq \kappa(\vartheta) \leq \bar{\kappa}(1 + \vartheta^\beta) \text{ for some } \beta \geq 3. \tag{2.8}$$

Here, the case $\beta = 3$ reflects the effect of radiation.

Finally, we suppose the magnetic diffusivity coefficient $\zeta = \zeta(\vartheta)$ is a continuously differentiable function of the temperature,

$$0 < \underline{\zeta}(1 + \vartheta) \leq \zeta(\vartheta) \leq \bar{\zeta}(1 + \vartheta), \quad |\zeta'(\vartheta)| \leq c \text{ for all } \vartheta \geq 0. \tag{2.9}$$

2.3. Boundary Data

We suppose the background magnetic field $\mathbf{B}_B = \mathbf{B}_B(t, x)$ is sufficiently smooth at least of class $C^{1,2}([0, T] \times \bar{\Omega}; R^3)$,

$$\operatorname{div}_x \mathbf{B}_B = 0 \text{ in } (0, T) \times \Omega. \tag{2.10}$$

Similarly, we suppose the boundary temperature ϑ_B can be extended inside Ω , $\vartheta_B \in C^{1,2}([0, T] \times \bar{\Omega})$,

$$\vartheta_B > 0 \text{ in } [0, T] \times \bar{\Omega}. \tag{2.11}$$

2.4. Weak Solutions

Before discussing the concept of weak solution, it is useful to introduce the function spaces

$$\begin{aligned} H_{0,\tau}(\Omega; R^3) &= \left\{ \mathbf{b} \in L^2(\Omega; R^3) \mid \operatorname{curl}_x \mathbf{b} \in L^2(\Omega; R^d), \operatorname{div}_x \mathbf{b} = 0, \mathbf{b} \times \mathbf{n}|_{\partial\Omega} = 0 \right\}, \\ H_{0,n}(\Omega; R^3) &= \left\{ \mathbf{b} \in L^2(\Omega; R^3) \mid \operatorname{curl}_x \mathbf{b} \in L^2(\Omega; R^d), \operatorname{div}_x \mathbf{b} = 0, \mathbf{b} \cdot \mathbf{n}|_{\partial\Omega} = 0 \right\}. \end{aligned}$$

Both $H_{0,\tau}$ and $H_{0,n}$ are endowed with the Hilbert norm

$$\|\mathbf{b}\|_{H_0}^2 = \|\operatorname{curl}_x \mathbf{b}\|_{L^2(\Omega; R^3)}^2 + \|\mathbf{b}\|_{L^2(\Omega; R^3)}^2,$$

see e.g. Alexander and Auchmuty [2].

Definition 2.1. (*Weak solution to the compressible MHD system*) A quantity $(\varrho, \vartheta, \mathbf{u}, \mathbf{B})$ is termed *weak solution* of the compressible MHD system (1.1)–(1.4), with the boundary conditions (1.10), (1.11), (1.12) (respectively (1.13)), and the initial data

$$\varrho(0, \cdot) = \varrho_0, \quad \vartheta(0, \cdot) = \vartheta_0, \quad \mathbf{u}(0, \cdot) = \mathbf{u}_0, \quad \mathbf{B}(0, \cdot) = \mathbf{B}_0, \tag{2.12}$$

if the following holds:

- **Equation of continuity.** $\varrho \in C_{\text{weak}}([0, T]; L^{\frac{5}{3}}(\Omega))$, $\varrho \geq 0$ in $(0, T) \times \Omega$, and the integral identity

$$\int_0^\tau \int_\Omega \left(\varrho \partial_t \varphi + \varrho \mathbf{u} \cdot \nabla_x \varphi \right) dx dt = - \left[\int_\Omega \varrho \varphi dx \right]_{t=0}^{t=\tau}, \quad \varrho(0, \cdot) = \varrho_0, \tag{2.13}$$

holds for any $\varphi \in C^1([0, T] \times \overline{\Omega})$. In addition, the renormalized version of (2.13)(2.13)

$$\int_0^\tau \int_\Omega \left(b(\varrho) \partial_t \varphi + b(\varrho) \mathbf{u} \cdot \nabla_x \varphi + (b(\varrho) - b'(\varrho)\varrho) \operatorname{div}_x \mathbf{u} \right) dx dt = \left[\int_\Omega b(\varrho) \varphi dx \right]_{t=0}^{t=\tau} \tag{2.14}$$

holds for any $\varphi \in C^1([0, T] \times \overline{\Omega})$ and any $b \in C^1(R)$, $b' \in C_c(R)$.

- **Momentum equation.** $\varrho \mathbf{u} \in C_{\text{weak}}([0, T]; L^{\frac{5}{4}}(\Omega; R^3))$, $\mathbf{u} \in L^2(0, T; W^{1,q}(\Omega; R^3))$, $q = \frac{8}{5-\alpha}$, $\mathbf{u} \cdot \mathbf{n}|_{\partial\Omega} = 0$, and the integral identity

$$\begin{aligned} & \int_0^\tau \int_\Omega \left(\varrho \mathbf{u} \cdot \partial_t \varphi + \varrho \mathbf{u} \otimes \mathbf{u} : \nabla_x \varphi + p(\varrho, \vartheta) \operatorname{div}_x \varphi \right) dx \\ &= \int_0^\tau \int_\Omega \mathbb{S}(\vartheta, \nabla_x \mathbf{u}) : \nabla_x \varphi dx dt - \int_0^\tau \int_\Omega \left(\mathbf{B} \otimes \mathbf{B} - \frac{1}{2} |\mathbf{B}|^2 \mathbb{I} \right) : \nabla_x \varphi dx dt \\ & \quad - \int_0^\tau \int_\Omega \varrho \mathbf{g} \cdot \varphi dx dt + \left[\int_\Omega \varrho \mathbf{u} \cdot \varphi dx \right]_{t=0}^{t=\tau}, \quad (\varrho \mathbf{u})(0, \cdot) = \varrho_0 \mathbf{u}_0 \end{aligned} \tag{2.15}$$

holds for any $\varphi \in C^1([0, T] \times \overline{\Omega}; R^3)$, $\varphi \cdot \mathbf{n}|_{\partial\Omega} = 0$.

- **Induction equation.** $\mathbf{B} \in C_{\text{weak}}([0, T]; L^2(\Omega; R^3))$,

$$\operatorname{div}_x \mathbf{B}(\tau, \cdot) = 0 \text{ for any } \tau \in [0, T]. \tag{2.16}$$

In addition

$$(\mathbf{B} - \mathbf{B}_B) \in L^2(0, T; H_{0,\tau}(\Omega; R^3)) \tag{2.17}$$

if the boundary condition (1.12) is imposed or

$$(\mathbf{B} - \mathbf{B}_B) \in L^2(0, T; H_{0,n}(\Omega; R^3)) \tag{2.18}$$

for the boundary conditions (1.13), (1.14). The integral identity

$$\begin{aligned} & \int_0^\tau \int_\Omega \left(\mathbf{B} \cdot \partial_t \varphi - (\mathbf{B} \times \mathbf{u}) \cdot \operatorname{curl}_x \varphi - \zeta(\vartheta) \operatorname{curl}_x \mathbf{B} \cdot \operatorname{curl}_x \varphi \right) dx dt \\ &= \left[\int_\Omega \mathbf{B} \cdot \varphi dx \right]_{t=0}^{t=\tau}, \quad \mathbf{B}(0, \cdot) = \mathbf{B}_0, \end{aligned} \tag{2.19}$$

holds for any $\varphi \in C^1([0, T] \times \overline{\Omega}; R^3)$, $\operatorname{div}_x \varphi = 0$,

$$\varphi \times \mathbf{n}|_{\partial\Omega} = 0 \tag{2.20}$$

in the case of boundary conditions (1.12),

$$\varphi \cdot \mathbf{n}|_{\partial\Omega} = 0 \tag{2.21}$$

for the boundary conditions (1.13), (1.14).

- **Entropy inequality.** $\vartheta \in L^\infty(0, T; L^4(\Omega))$, $\vartheta > 0$ a.a. in $(0, T) \times \Omega$, $\vartheta - \vartheta_B \in L^2(0, T; W_0^{1,2}(\Omega))$, $\log(\vartheta) \in L^2(0, T; W^{1,2}(\Omega))$, and the integral inequality

$$\begin{aligned} & \int_{\tau_1}^{\tau_2} \int_{\Omega} \left(\varrho s(\varrho, \vartheta) \partial_t \varphi + \varrho s(\varrho, \vartheta) \mathbf{u} \cdot \nabla_x \varphi + \frac{\mathbf{q}(\vartheta, \nabla_x \vartheta)}{\vartheta} \cdot \nabla_x \varphi \right) dx dt \\ & \leq - \int_{\tau_1}^{\tau_2} \int_{\Omega} \frac{1}{\vartheta} \left(\mathbb{S}(\vartheta, \nabla_x \mathbf{u}) : \nabla_x \mathbf{u} - \frac{\mathbf{q}(\vartheta, \nabla_x \vartheta) \cdot \nabla_x \vartheta}{\vartheta} + \zeta(\vartheta) |\mathbf{curl}_x \mathbf{B}|^2 \right) \varphi dx dt \\ & \quad + \left[\int_{\Omega} \varrho s(\varrho, \vartheta) \varphi dx \right]_{t=\tau_1-}^{t=\tau_2+}, \quad 0 \leq \tau_1 \leq \tau_2 \leq T, \quad \varrho s(\varrho, \vartheta)(0-, \cdot) = \varrho_0 s(\varrho_0, \vartheta_0), \end{aligned} \tag{2.22}$$

holds for any $\varphi \in C_c^1([0, T] \times \Omega)$, $\varphi \geq 0$.

- **Ballistic energy inequality.** The inequality

$$\begin{aligned} & \int_{\Omega} \left(\frac{1}{2} \varrho |\mathbf{u}|^2 + \varrho e(\varrho, \vartheta) + \frac{1}{2} |\mathbf{B}|^2 - \tilde{\vartheta} \varrho s(\varrho, \vartheta) - \mathbf{B}_B \cdot \mathbf{B} \right) (\tau, \cdot) dx \\ & \quad + \int_0^\tau \int_{\Omega} \frac{\tilde{\vartheta}}{\vartheta} \left(\mathbb{S}(\vartheta, \nabla_x \mathbf{u}) : \nabla_x \mathbf{u} - \frac{\mathbf{q}(\vartheta, \nabla_x \vartheta) \cdot \nabla_x \vartheta}{\vartheta} + \zeta(\vartheta) |\mathbf{curl}_x \mathbf{B}|^2 \right) dx dt \\ & \leq - \int_0^\tau \int_{\Omega} \left(\varrho s(\varrho, \vartheta) \partial_t \tilde{\vartheta} + \varrho s(\varrho, \vartheta) \mathbf{u} \cdot \nabla_x \tilde{\vartheta} + \frac{\mathbf{q}(\vartheta, \nabla_x \vartheta)}{\vartheta} \cdot \nabla_x \tilde{\vartheta} \right) dx \\ & \quad - \int_0^\tau \int_{\Omega} \left(\mathbf{B} \cdot \partial_t \mathbf{B}_B - (\mathbf{B} \times \mathbf{u}) \cdot \mathbf{curl}_x \mathbf{B}_B - \zeta(\vartheta) \mathbf{curl}_x \mathbf{B} \cdot \mathbf{curl}_x \mathbf{B}_B \right) dx dt \\ & \quad + \int_0^\tau \int_{\Omega} \varrho \mathbf{g} \cdot \mathbf{u} dx dt \\ & \quad + \int_{\Omega} \left(\frac{1}{2} \varrho_0 |\mathbf{u}_0|^2 + \varrho_0 e(\varrho_0, \vartheta_0) + \frac{1}{2} |\mathbf{B}_0|^2 - \tilde{\vartheta}(0, \cdot) \varrho_0 s(\varrho_0, \vartheta_0) - \mathbf{B}_B(0, \cdot) \cdot \mathbf{B}_0 \right) dx \end{aligned} \tag{2.23}$$

holds for any $\tilde{\vartheta} \in C^1([0, T] \times \bar{\Omega})$, $\tilde{\vartheta} > 0$, $\tilde{\vartheta}|_{\partial\Omega} = \vartheta_B$.

Remark 2.2. As the magnetic field \mathbf{B} satisfies (2.16), validity of (2.19) can be extended to any (not necessarily solenoidal) test function φ satisfying (2.20), (2.21), respectively. Indeed, if φ satisfies the boundary condition (2.20), we consider the potential Ψ solving

$$\Delta_x \Psi = \operatorname{div}_x \varphi \text{ in } \Omega, \quad \Psi|_{\partial\Omega} = 0,$$

and write

$$\varphi = (\varphi - \nabla_x \Psi) + \nabla_x \Psi.$$

On the one hand, as Ψ vanishes on $\partial\Omega$ and \mathbf{B} is solenoidal, we have

$$\int_{\Omega} \mathbf{B} \cdot \nabla_x \Psi dx = \int_{\Omega} \mathbf{B} \cdot \partial_t \nabla_x \Psi dx = 0;$$

whence the integral identity (2.19) holds for $\nabla_x \Psi$. On the other hand, the function $(\varphi - \nabla_x \Psi)$ is solenoidal, and, as $\nabla_x \Psi \times \mathbf{n}|_{\partial\Omega} = 0$, a legal test function in (2.19).

If φ satisfies (2.21), we simply consider

$$\Delta_x \Psi = \operatorname{div}_x \varphi, \quad \nabla_x \Psi \cdot \mathbf{n}|_{\partial\Omega} = \varphi \cdot \mathbf{n}|_{\partial\Omega} = 0$$

- the standard potential in the Helmholtz decomposition.

Remark 2.3. Unlike ϱ , $\varrho \mathbf{u}$, and \mathbf{B} , the total entropy $\varrho s(\varrho, \vartheta)$ is not weakly continuous in time. However, it can be deduced from the entropy inequality that the one sided limits

$$\lim_{t \rightarrow \tau-} \int_{\Omega} \varrho s(\varrho, \vartheta)(t, \cdot) \phi dx, \quad \lim_{t \rightarrow \tau+} \int_{\Omega} \varrho s(\varrho, \vartheta)(t, \cdot) \phi dx$$

exist for any $\phi \in C(\bar{\Omega})$, and

$$\lim_{t \rightarrow \tau^-} \int_{\Omega} \varrho s(\varrho, \vartheta)(t, \cdot) \phi \, dx \leq \lim_{t \rightarrow \tau^+} \int_{\Omega} \varrho s(\varrho, \vartheta)(t, \cdot) \phi \, dx$$

whenever $\phi \geq 0$, see [10, Chapter 3, Section 3.3.3] for details.

Remark 2.4. As we shall see below, the “magnetic” ballistic energy E_{BM} introduced in (1.18) is in fact a strictly convex l.s.c function of the variables $(\varrho, S = \varrho s(\varrho, \vartheta), \mathbf{m} = \varrho \mathbf{u}, \mathbf{B})$. In particular, its integral is a weakly lower semi-continuous function; whence (2.23) holds for *any* time τ with the convention for entropy discussed Remark 2.3. Alternatively, we may impose a stronger integrated version of (2.23) in the form

$$\begin{aligned} & \int_0^T \partial_t \psi \int_{\Omega} \left(\frac{1}{2} \varrho |\mathbf{u}|^2 + \varrho e(\varrho, \vartheta) + \frac{1}{2} |\mathbf{B}|^2 - \tilde{\vartheta} \varrho s(\varrho, \vartheta) - \mathbf{B}_B \cdot \mathbf{B} \right) \, dx \, dt \\ & - \int_0^T \psi \int_{\Omega} \frac{\tilde{\vartheta}}{\vartheta} \left(\mathbb{S}(\vartheta, \nabla_x \mathbf{u}) : \nabla_x \mathbf{u} - \frac{\mathbf{q}(\vartheta, \nabla_x \vartheta) \cdot \nabla_x \vartheta}{\vartheta} + \zeta(\vartheta) |\mathbf{curl}_x \mathbf{B}|^2 \right) \, dx \, dt \\ & \geq \int_0^T \psi \int_{\Omega} \left(\varrho s(\varrho, \vartheta) \partial_t \tilde{\vartheta} + \varrho s(\varrho, \vartheta) \mathbf{u} \cdot \nabla_x \tilde{\vartheta} + \frac{\mathbf{q}(\varrho, \vartheta)}{\vartheta} \cdot \nabla_x \tilde{\vartheta} \right) \, dx \, dt \\ & + \int_0^T \psi \int_{\Omega} \left(\mathbf{B} \cdot \partial_t \mathbf{B}_B - (\mathbf{B} \times \mathbf{u}) \cdot \mathbf{curl}_x \mathbf{B}_B - \zeta \mathbf{curl}_x \mathbf{B} \cdot \mathbf{curl}_x \mathbf{B}_B \right) \, dx \, dt \\ & - \int_0^T \psi \int_{\Omega} \varrho \mathbf{g} \cdot \mathbf{u} \, dx \, dt \\ & - \psi(0) \int_{\Omega} \left(\frac{1}{2} \varrho_0 |\mathbf{u}_0|^2 + \varrho_0 e(\varrho_0, \vartheta_0) + \frac{1}{2} |\mathbf{B}_0|^2 - \tilde{\vartheta}(0, \cdot) \varrho_0 s(\varrho_0, \vartheta_0) - \mathbf{B}_B(0, \cdot) \cdot \mathbf{B}_0 \right) \, dx \end{aligned} \tag{2.24}$$

for any $\psi \in C_c^1[0, T]$, $\psi \geq 0$.

Remark 2.5. Note carefully that the ballistic energy inequality (2.23) remains valid if we replace \mathbf{B}_B by any other extension $\tilde{\mathbf{B}}$ such that

$$\operatorname{div}_x \tilde{\mathbf{B}} = 0,$$

and

$$\tilde{\mathbf{B}} \times \mathbf{n}|_{\partial\Omega} = \mathbf{B}_B \times \mathbf{n}|_{\partial\Omega}, \text{ or } \tilde{\mathbf{B}} \cdot \mathbf{n}|_{\partial\Omega} = \mathbf{B}_B \cdot \mathbf{n}|_{\partial\Omega},$$

respectively. Indeed the difference $\mathbf{B}_B - \tilde{\mathbf{B}}$ becomes an eligible test function for for the weak form of the induction equation (2.19), endowed with the boundary conditions (2.20), (2.21), respectively.

The existence of global-in-time weak solutions in the sense of Definition 2.1 will be shown in Sect. 5 below.

3. Relative Energy

A suitable form of the *relative energy* for the compressible MHD system reads

$$\begin{aligned} E \left(\varrho, \vartheta, \mathbf{u}, \mathbf{B} \mid r, \Theta, \mathbf{U}, \mathbf{H} \right) &= \frac{1}{2} \varrho |\mathbf{u} - \mathbf{U}|^2 + \frac{1}{2} |\mathbf{B} - \mathbf{H}|^2 \\ &+ \varrho e(\varrho, \vartheta) - \Theta \left(\varrho s(\varrho, \vartheta) - r s(r, \Theta) \right) - \left(e(r, \Theta) - \Theta s(r, \Theta) + \frac{p(r, \Theta)}{r} \right) (\varrho - r) - r e(r, \Theta). \end{aligned} \tag{3.1}$$

In applications, the quantity $(\varrho, \vartheta, \mathbf{u}, \mathbf{B})$ stands for a weak solution of the compressible MHD system while $(r, \Theta, \mathbf{U}, \mathbf{H})$ are arbitrary sufficiently smooth functions satisfying the compatibility conditions

$$r > 0, \Theta > 0 \text{ in } [0, T] \times \bar{\Omega}, \Theta|_{\partial\Omega} = \vartheta_B, \mathbf{U} \cdot \mathbf{n}|_{\partial\Omega} = 0,$$

$$\operatorname{div}_x \mathbf{H} = 0 \text{ in } (0, T) \times \Omega, \quad \mathbf{H} \times \mathbf{n}|_{\partial\Omega} = \mathbf{B}_B \times \mathbf{n}|_{\partial\Omega}, \text{ or } \mathbf{H} \cdot \mathbf{n}|_{\partial\Omega} = \mathbf{B}_B \cdot \mathbf{n}|_{\partial\Omega}, \text{ respectively.} \quad (3.2)$$

As a consequence of hypothesis of thermodynamic stability stated in (2.4), the energy

$$E(\varrho, \vartheta, \mathbf{u}, \mathbf{B}) = \frac{1}{2} \varrho |\mathbf{u}|^2 + \varrho e(\varrho, \vartheta) + \frac{1}{2} |\mathbf{B}|^2$$

is a strictly convex l.s.c. function if expressed in the variables $(\varrho, S = \varrho s(\varrho, \vartheta), \mathbf{m} = \varrho \mathbf{u}, \mathbf{B})$, see [10, Chapter 3, Section 3.1] for details. Moreover, we have

$$\frac{\partial E(\varrho, S, \mathbf{m}, \mathbf{B})}{\partial \varrho} = \left(e(\varrho, \vartheta) - \vartheta s(\varrho, \vartheta) + \frac{p(\varrho, \vartheta)}{\varrho} \right), \quad \frac{\partial E(\varrho, S, \mathbf{m}, \mathbf{B})}{\partial S} = \vartheta,$$

where $\left(e(\varrho, \vartheta) - \vartheta s(\varrho, \vartheta) + \frac{p(\varrho, \vartheta)}{\varrho} \right)$ is Gibbs' free energy. Consequently, the relative energy can be seen as Bregman divergence associate to the convex function E . In particular,

$$E(\varrho, \vartheta, \mathbf{u}, \mathbf{B} \mid r, \Theta, \mathbf{U}, \mathbf{H}) \geq 0, \text{ and } E(\varrho, \vartheta, \mathbf{u}, \mathbf{B} \mid r, \Theta, \mathbf{U}, \mathbf{H}) = 0 \Leftrightarrow (\varrho, \vartheta, \mathbf{u}, \mathbf{B}) = (r, \Theta, \mathbf{U}, \mathbf{H})$$

as long as $r > 0$.

3.1. Relative Energy Inequality

In the remaining part of this section, we derive a relative energy inequality provided $(\varrho, \vartheta, \mathbf{u}, \mathbf{B})$ is a weak solution of the compressible MHD system specified in Definition 2.1 and $(r, \Theta, \mathbf{U}, \mathbf{H})$ arbitrary smooth functions satisfying (3.2). Our starting point is the ballistic energy inequality (2.24). As noted in Remark 2.5, we may flip \mathbf{B}_B for \mathbf{H} , and, of course, $\tilde{\vartheta}$ for Θ obtaining

$$\begin{aligned} & \int_{\Omega} \left(\frac{1}{2} \varrho |\mathbf{u}|^2 + \varrho e(\varrho, \vartheta) + \frac{1}{2} |\mathbf{B}|^2 - \Theta \varrho s(\varrho, \vartheta) - \mathbf{H} \cdot \mathbf{B} \right) (\tau, \cdot) \, dx \\ & + \int_0^\tau \int_{\Omega} \frac{\Theta}{\vartheta} \left(\mathbb{S}(\vartheta, \nabla_x \mathbf{u}) : \nabla_x \mathbf{u} - \frac{\mathbf{q}(\vartheta, \nabla_x \vartheta) \cdot \nabla_x \vartheta}{\vartheta} + \zeta(\vartheta) |\operatorname{curl}_x \mathbf{B}|^2 \right) \, dx \, dt \\ & \leq - \int_0^\tau \int_{\Omega} \left(\varrho s(\varrho, \vartheta) \partial_t \Theta + \varrho s(\varrho, \vartheta) \mathbf{u} \cdot \nabla_x \Theta + \frac{\mathbf{q}(\vartheta, \nabla_x \vartheta)}{\vartheta} \cdot \nabla_x \Theta \right) \, dx \\ & - \int_0^\tau \int_{\Omega} \left(\mathbf{B} \cdot \partial_t \mathbf{H} - (\mathbf{B} \times \mathbf{u}) \cdot \operatorname{curl}_x \mathbf{H} - \zeta(\vartheta) \operatorname{curl}_x \mathbf{B} \cdot \operatorname{curl}_x \mathbf{H} \right) \, dx \, dt \\ & + \int_0^\tau \int_{\Omega} \varrho \mathbf{g} \cdot \mathbf{u} \, dx \, dt \\ & + \int_{\Omega} \left(\frac{1}{2} \varrho_0 |\mathbf{u}_0|^2 + \varrho_0 e(\varrho_0, \vartheta_0) + \frac{1}{2} |\mathbf{B}_0|^2 - \Theta(0, \cdot) \varrho_0 s(\varrho_0, \vartheta_0) - \mathbf{H}(0, \cdot) \cdot \mathbf{B}_0 \right) \, dx \end{aligned} \quad (3.3)$$

3.1.1. Momentum Perturbation. Our first goal is to replace $|\mathbf{u}|^2$ by $|\mathbf{u} - \mathbf{U}|^2$. This can be achieved by considering \mathbf{U} as a test function in the momentum balance equation (2.15). After a straightforward manipulation explained in detail in [10, Chapter 3, Section 3.2.1], we obtain

$$\begin{aligned} & \int_{\Omega} \left(\frac{1}{2} \varrho |\mathbf{u} - \mathbf{U}|^2 + \varrho e(\varrho, \vartheta) + \frac{1}{2} |\mathbf{B}|^2 - \Theta \varrho s(\varrho, \vartheta) - \mathbf{H} \cdot \mathbf{B} \right) (\tau, \cdot) \, dx \\ & + \int_0^\tau \int_{\Omega} \frac{\Theta}{\vartheta} \left(\mathbb{S}(\vartheta, \nabla_x \mathbf{u}) : \nabla_x \mathbf{u} - \frac{\mathbf{q}(\vartheta, \nabla_x \vartheta) \cdot \nabla_x \vartheta}{\vartheta} + \zeta(\vartheta) |\operatorname{curl}_x \mathbf{B}|^2 \right) \, dx \, dt \\ & \leq - \int_0^\tau \int_{\Omega} \left(\varrho (\mathbf{u} - \mathbf{U}) \otimes (\mathbf{u} - \mathbf{U}) + p(\varrho, \vartheta) \mathbb{I} - \mathbb{S}(\vartheta, \nabla_x \mathbf{u}) \right) : \nabla_x \mathbf{U} \, dx \, dt \\ & - \int_0^\tau \int_{\Omega} (\operatorname{curl}_x \mathbf{B} \times \mathbf{B}) \cdot \mathbf{U} \, dx \, dt \end{aligned}$$

$$\begin{aligned}
& - \int_0^\tau \int_\Omega \varrho \left(\partial_t \mathbf{U} + \mathbf{U} \cdot \nabla_x \mathbf{U} \right) \cdot (\mathbf{u} - \mathbf{U}) \, dx \, dt \\
& - \int_0^\tau \int_\Omega \left(\varrho s(\varrho, \vartheta) \partial_t \Theta + \varrho s(\varrho, \vartheta) \mathbf{u} \cdot \nabla_x \Theta + \frac{\mathbf{q}(\vartheta, \nabla_x \vartheta)}{\vartheta} \cdot \nabla_x \Theta \right) \, dx + \int_0^\tau \int_\Omega \varrho \mathbf{g} \cdot (\mathbf{u} - \mathbf{U}) \, dx \, dt \\
& - \int_0^\tau \int_\Omega \left(\mathbf{B} \cdot \partial_t \mathbf{H} - (\mathbf{B} \times \mathbf{u}) \cdot \operatorname{curl}_x \mathbf{H} - \zeta(\vartheta) \operatorname{curl}_x \mathbf{B} \cdot \operatorname{curl}_x \mathbf{H} \right) \, dx \, dt \\
& + \int_\Omega \left(\frac{1}{2} \varrho_0 |\mathbf{u}_0 - \mathbf{U}(0, \cdot)|^2 + \varrho_0 e(\varrho_0, \vartheta_0) + \frac{1}{2} |\mathbf{B}_0|^2 - \Theta(0, \cdot) \varrho_0 s(\varrho_0, \vartheta_0) - \mathbf{H}(0, \cdot) \cdot \mathbf{B}_0 \right) \, dx \quad (3.4)
\end{aligned}$$

for any $(\Theta, \mathbf{U}, \mathbf{H})$ as in (3.2).

3.1.2. Density Perturbation. Next, exactly as in [10, Chapter 3, Section 3.4], we use the quantity

$$\left(e(r, \Theta) - \Theta s(r, \Theta) + \frac{p(r, \Theta)}{r} \right)$$

as a test function in the equation of continuity (2.13):

$$\begin{aligned}
& \int_\Omega \left(\frac{1}{2} \varrho |\mathbf{u} - \mathbf{U}|^2 + \varrho e(\varrho, \vartheta) + \frac{1}{2} |\mathbf{B}|^2 - \Theta \varrho s(\varrho, \vartheta) - \mathbf{H} \cdot \mathbf{B} \right) (\tau, \cdot) \, dx \\
& + \int_\Omega \left(\Theta r s(r, \Theta) - \left(e(r, \Theta) - \Theta s(r, \Theta) + \frac{p(r, \Theta)}{r} \right) (\varrho - r) - r e(r, \Theta) \right) (\tau, \cdot) \, dx \\
& + \int_0^\tau \int_\Omega \frac{\Theta}{\vartheta} \left(\mathbb{S}(\vartheta, \mathbf{u}) : \nabla_x \mathbf{u} - \frac{\mathbf{q}(\vartheta, \nabla_x \vartheta) \cdot \nabla_x \vartheta}{\vartheta} + \zeta(\vartheta) |\operatorname{curl}_x \mathbf{B}|^2 \right) \, dx \, dt \\
& \leq - \int_0^\tau \int_\Omega \left(\varrho (\mathbf{u} - \mathbf{U}) \otimes (\mathbf{u} - \mathbf{U}) + p(\varrho, \vartheta) \mathbb{I} - \mathbb{S}(\vartheta, \nabla_x \mathbf{u}) \right) : \nabla_x \mathbf{U} \, dx \, dt \\
& + \int_0^\tau \int_\Omega \frac{\varrho}{r} (\mathbf{u} - \mathbf{U}) \cdot \nabla_x p(r, \Theta) \, dx \, dt \\
& - \int_0^\tau \int_\Omega (\operatorname{curl}_x \mathbf{B} \times \mathbf{B}) \cdot \mathbf{U} \, dx \, dt \\
& - \int_0^\tau \int_\Omega \varrho \left(\partial_t \mathbf{U} + \mathbf{U} \cdot \nabla_x \mathbf{U} + \frac{1}{r} \nabla_x p(r, \Theta) \right) \cdot (\mathbf{u} - \mathbf{U}) \, dx \, dt \\
& - \int_0^\tau \int_\Omega \left(\varrho (s(\varrho, \vartheta) - s(r, \Theta)) \partial_t \Theta + \varrho (s(\varrho, \vartheta) - s(r, \Theta)) \mathbf{u} \cdot \nabla_x \Theta + \frac{\mathbf{q}(\vartheta, \nabla_x \vartheta)}{\vartheta} \cdot \nabla_x \Theta \right) \, dx \\
& + \int_0^\tau \int_\Omega \varrho \mathbf{g} \cdot (\mathbf{u} - \mathbf{U}) \, dx \, dt \\
& + \int_0^\tau \int_\Omega \left(\left(1 - \frac{\varrho}{r} \right) \partial_t p(r, \Theta) - \frac{\varrho}{r} \mathbf{u} \cdot \nabla_x p(r, \Theta) \right) \, dx \, dt \\
& - \int_0^\tau \int_\Omega \left(\mathbf{B} \cdot \partial_t \mathbf{H} - (\mathbf{B} \times \mathbf{u}) \cdot \operatorname{curl}_x \mathbf{H} - \zeta(\vartheta) \operatorname{curl}_x \mathbf{B} \cdot \operatorname{curl}_x \mathbf{H} \right) \, dx \, dt \\
& + \int_\Omega \left(\frac{1}{2} \varrho_0 |\mathbf{u}_0 - \mathbf{U}(0, \cdot)|^2 + \varrho_0 e(\varrho_0, \vartheta_0) + \frac{1}{2} |\mathbf{B}_0|^2 - \Theta(0, \cdot) \varrho_0 s(\varrho_0, \vartheta_0) - \mathbf{H}(0, \cdot) \cdot \mathbf{B}_0 \right) \, dx \\
& + \int_\Omega \left(\Theta r s(r, \Theta) - \left(e(r, \Theta) - \Theta s(r, \Theta) + \frac{p(r, \Theta)}{r} \right) (\varrho_0 - r) - r e(r, \Theta) \right) (0, \cdot) \, dx, \quad (3.5)
\end{aligned}$$

whenever $r \in C^1([0, T] \times \bar{\Omega})$, $r > 0$.

3.1.3. Magnetic Field Perturbation and the Final form of Relative Energy Inequality. Since

$$\int_\Omega \frac{1}{2} |\mathbf{H}|^2(\tau, \cdot) \, dx - \int_\Omega \frac{1}{2} |\mathbf{H}|^2(0, \cdot) \, dx = \int_0^\tau \int_\Omega \mathbf{H} \cdot \partial_t \mathbf{H} \, dx \, dt. \quad (3.6)$$

we may rewrite (3.5) in the final form

$$\begin{aligned}
 & \left[\int_{\Omega} E(\varrho, \vartheta, \mathbf{u}, \mathbf{B} \mid r, \Theta, \mathbf{U}, \mathbf{H}) \, dx \right]_{t=0}^{t=\tau} \\
 & + \int_0^\tau \int_{\Omega} \frac{\Theta}{\vartheta} \left(\mathbb{S}(\vartheta, \nabla_x \mathbf{u}) : \nabla_x \mathbf{u} - \frac{\mathbf{q}(\vartheta, \nabla_x \vartheta) \cdot \nabla_x \vartheta}{\vartheta} + \zeta(\vartheta) |\mathbf{curl}_x \mathbf{B}|^2 \right) \, dx \, dt \\
 & \leq - \int_0^\tau \int_{\Omega} \left(\varrho(\mathbf{u} - \mathbf{U}) \otimes (\mathbf{u} - \mathbf{U}) + p(\varrho, \vartheta) \mathbb{I} - \mathbb{S}(\vartheta, \nabla_x \mathbf{u}) \right) : \nabla_x \mathbf{U} \, dx \, dt \\
 & + \int_0^\tau \int_{\Omega} \frac{\varrho}{r} (\mathbf{u} - \mathbf{U}) \cdot \nabla_x p(r, \Theta) \, dx \, dt \\
 & - \int_0^\tau \int_{\Omega} (\mathbf{curl}_x \mathbf{B} \times \mathbf{B}) \cdot \mathbf{U} \, dx \, dt \\
 & - \int_0^\tau \int_{\Omega} \varrho \left(\partial_t \mathbf{U} + \mathbf{U} \cdot \nabla_x \mathbf{U} + \frac{1}{r} \nabla_x p(r, \Theta) - \mathbf{g} \right) \cdot (\mathbf{u} - \mathbf{U}) \, dx \, dt \\
 & - \int_0^\tau \int_{\Omega} \left(\varrho (s(\varrho, \vartheta) - s(r, \Theta)) \partial_t \Theta + \varrho (s(\varrho, \vartheta) - s(r, \Theta)) \mathbf{u} \cdot \nabla_x \Theta + \frac{\mathbf{q}(\vartheta, \nabla_x \vartheta) \cdot \nabla_x \Theta}{\vartheta} \right) \, dx \\
 & + \int_0^\tau \int_{\Omega} \left(\left(1 - \frac{\varrho}{r} \right) \partial_t p(r, \Theta) - \frac{\varrho}{r} \mathbf{u} \cdot \nabla_x p(r, \Theta) \right) \, dx \, dt \\
 & - \int_0^\tau \int_{\Omega} \left(\mathbf{B} \cdot \partial_t \mathbf{H} - (\mathbf{B} \times \mathbf{u}) \cdot \mathbf{curl}_x \mathbf{H} - \zeta(\vartheta) \mathbf{curl}_x \mathbf{B} \cdot \mathbf{curl}_x \mathbf{H} \right) \, dx \, dt \\
 & + \int_0^\tau \int_{\Omega} \mathbf{H} \cdot \partial_t \mathbf{H} \, dx \, dt
 \end{aligned} \tag{3.7}$$

for any “test” functions $(r, \Theta, \mathbf{U}, \mathbf{H})$ specified in (3.2).

Remark 3.1. In view of future applications, in particular the weak–strong uniqueness principle discussed in the forthcoming section, the class of functions $(r, \Theta, \mathbf{U}, \mathbf{H})$ in (3.2) can be extended to the “maximal regularity” framework

$$r \in W^{1,q}((0, T) \times \Omega), \quad (\Theta, \mathbf{U}, \mathbf{B}) \in L^q(0, T; W^{2,q}(\Omega; R^7)) \cap W^{1,q}(0, T; L^q(\Omega; R^7)), \tag{3.8}$$

by a density argument. The exponent q must be taken large enough for all terms in (3.7) to be well defined.

4. Weak–Strong Uniqueness Principle

The first important property of the weak solutions introduced in Definition 2.1 is the *weak–strong uniqueness principle*. We suppose that $(r, \Theta, \mathbf{U}, \mathbf{H})$ is a smooth solution of the problem, specifically,

$$\begin{aligned}
 \partial_t r + \operatorname{div}_x(r\mathbf{U}) &= 0, \quad \partial_t \mathbf{U} + \mathbf{U} \cdot \nabla_x \mathbf{U} + \frac{1}{r} \nabla_x p(r, \Theta) = \frac{1}{r} \operatorname{div}_x \mathbb{S}(\Theta, \nabla_x \mathbf{U}) + \mathbf{g} + \frac{1}{r} \mathbf{curl}_x \mathbf{H} \times \mathbf{H}, \\
 \partial_t \mathbf{H} + \mathbf{curl}_x(\mathbf{H} \times \mathbf{U}) + \mathbf{curl}_x(\zeta(\Theta) \mathbf{curl}_x \mathbf{H}) &= 0, \quad \operatorname{div}_x \mathbf{H} = 0, \\
 r \partial_t s(r, \Theta) + r \mathbf{U} \cdot \nabla_x s(r, \Theta) + \operatorname{div}_x \left(\frac{\mathbf{q}(\Theta, \nabla_x \Theta)}{\Theta} \right) \\
 &= \frac{1}{\Theta} \left(\mathbb{S}(\Theta, \nabla_x \mathbf{U}) : \nabla_x \mathbf{U} - \frac{\mathbf{q}(\Theta, \nabla_x \Theta) \cdot \nabla_x \Theta}{\Theta} + \zeta(\Theta) |\mathbf{curl}_x \mathbf{H}|^2 \right),
 \end{aligned} \tag{4.1}$$

together with the relevant initial and boundary conditions.

Theorem 4.1. (Weak–strong uniqueness) *Let $\Omega \subset R^3$ be a bounded Lipschitz domain. Suppose the thermodynamic functions p, e, s satisfy the hypotheses (2.1)–(2.6), and the transport coefficients μ, η, κ, ζ comply with (2.7)–(2.9), with*

$$\frac{1}{2} \leq \alpha \leq 1, \quad \beta \geq 3. \quad (4.2)$$

Let $(\varrho, \vartheta, \mathbf{u}, \mathbf{B})$ be a weak solution of the compressible MHD system (1.1)–(1.4) in $(0, T) \times \Omega$ specified in Definition 2.1. Suppose that the same problem, with the same initial and boundary data, admits a classical solution $(r, \Theta, \mathbf{U}, \mathbf{H})$ in the class

$$r \in C^1([0, T] \times \bar{\Omega}), \quad \Theta \in C^{1,2}([0, T] \times \bar{\Omega}), \quad \mathbf{U}, \mathbf{H} \in C^{1,2}([0, T] \times \bar{\Omega}; R^3). \quad (4.3)$$

Then

$$\varrho = r, \quad \mathbf{u} = \mathbf{U}, \quad \vartheta = \Theta, \quad \mathbf{B} = \mathbf{H} \quad \text{a.e. in } [0, T] \times \bar{\Omega}.$$

Remark 4.2. As we assume existence of a classical solution in Theorem 4.1, the initial data are also regular, in particular,

$$\inf_{\Omega} \varrho_0 > 0, \quad \inf_{\Omega} \vartheta_0 > 0.$$

Remark 4.3. Remarkably, the result holds in the full range of exponents α, β specified in (4.2). In particular, we remove the gap between the weak–strong uniqueness principle stated in [5, 10] for $\beta = 3$ and the existence result requiring $\beta > 6$. The main novelty is formulated in Lemma 4.4 below.

The rest of this section is devoted to the proof of Theorem 4.1. Note that existence of local in time strong solutions to the compressible MHD system can be established by the nowadays well understood technique proposed by Valli [24], Valli and Zajaczkowski [25], or by a more recent approach via maximal regularity in the spirit of Kotschote [13]. As noted in Remark 3.1, the conclusion of Theorem 4.1 remains valid for the strong solutions in the maximal regularity class used in [13].

The idea of the proof of Theorem 4.1 is simple: We consider the strong solution $(r, \Theta, \mathbf{U}, \mathbf{H})$ as test functions in the relative energy inequality (3.7) and use Gronwall’s type argument. We proceed in several steps specified below.

4.1. Magnetic Field

As \mathbf{U} solves the momentum equation, we get

$$\begin{aligned} & - \int_0^\tau \int_{\Omega} \varrho \left(\partial_t \mathbf{U} + \mathbf{U} \cdot \nabla_x \mathbf{U} + \frac{1}{r} \nabla_x p(r, \Theta) - \mathbf{g} \right) \cdot (\mathbf{u} - \mathbf{U}) \, dx \, dt \\ & = \int_0^\tau \int_{\Omega} \frac{\varrho}{r} \left(\operatorname{div}_x \mathbb{S}(\Theta, \nabla_x \mathbf{U}) + \operatorname{curl}_x \mathbf{H} \times \mathbf{H} \right) \cdot (\mathbf{U} - \mathbf{u}) \, dx \, dt \\ & = \int_0^\tau \int_{\Omega} \left(\frac{\varrho}{r} - 1 \right) \left(\operatorname{div}_x \mathbb{S}(\Theta, \nabla_x \mathbf{U}) + \operatorname{curl}_x \mathbf{H} \times \mathbf{H} \right) \cdot (\mathbf{U} - \mathbf{u}) \, dx \, dt \\ & \quad + \int_0^\tau \int_{\Omega} \left(\operatorname{div}_x \mathbb{S}(\Theta, \nabla_x \mathbf{U}) + \operatorname{curl}_x \mathbf{H} \times \mathbf{H} \right) \cdot (\mathbf{U} - \mathbf{u}) \, dx \, dt. \end{aligned} \quad (4.4)$$

Similarly, \mathbf{H} being the exact solution of the magnetic field equation,

$$\begin{aligned} & - \int_0^\tau \int_{\Omega} \left(\mathbf{B} \cdot \partial_t \mathbf{H} - (\mathbf{B} \times \mathbf{u}) \cdot \operatorname{curl}_x \mathbf{H} - \zeta(\vartheta) \operatorname{curl}_x \mathbf{B} \cdot \operatorname{curl}_x \mathbf{H} \right) \, dx \, dt + \int_0^\tau \int_{\Omega} \mathbf{H} \cdot \partial_t \mathbf{H} \, dx \, dt \\ & = \int_0^\tau \int_{\Omega} (\mathbf{B} - \mathbf{H}) \cdot \left(\operatorname{curl}_x (\mathbf{H} \times \mathbf{U}) + \operatorname{curl}_x (\zeta(\Theta) \operatorname{curl}_x \mathbf{H}) \right) \, dx \, dt \\ & \quad + \int_0^\tau \int_{\Omega} \left((\mathbf{B} \times \mathbf{u}) \cdot \operatorname{curl}_x \mathbf{H} + \zeta(\vartheta) \operatorname{curl}_x \mathbf{B} \cdot \operatorname{curl}_x \mathbf{H} \right) \, dx \, dt \end{aligned} \quad (4.5)$$

Regrouping the terms in (3.7), (4.4), (4.5) that do not contain magnetic dissipation, we get the expression

$$\begin{aligned} & - \int_{\Omega} (\mathbf{curl}_x \mathbf{B} \times \mathbf{B}) \cdot \mathbf{U} \, dx + \int_{\Omega} (\mathbf{curl}_x \mathbf{H} \times \mathbf{H}) \cdot (\mathbf{U} - \mathbf{u}) \, dx \\ & + \int_{\Omega} (\mathbf{B} - \mathbf{H}) \cdot \mathbf{curl}_x (\mathbf{H} \times \mathbf{U}) \, dx + \int_{\Omega} (\mathbf{B} \times \mathbf{u}) \cdot \mathbf{curl}_x \mathbf{H} \, dx. \end{aligned}$$

Now, we claim the identity

$$\begin{aligned} & - \int_{\Omega} (\mathbf{curl}_x \mathbf{B} \times \mathbf{B}) \cdot \mathbf{U} \, dx + \int_{\Omega} (\mathbf{curl}_x \mathbf{H} \times \mathbf{H}) \cdot (\mathbf{U} - \mathbf{u}) \, dx \\ & + \int_{\Omega} (\mathbf{B} - \mathbf{H}) \cdot \mathbf{curl}_x (\mathbf{H} \times \mathbf{U}) \, dx + \int_{\Omega} (\mathbf{B} \times \mathbf{u}) \cdot \mathbf{curl}_x \mathbf{H} \, dx \\ & = \int_{\Omega} (\mathbf{U} \times (\mathbf{B} - \mathbf{H})) \cdot \mathbf{curl}_x (\mathbf{B} - \mathbf{H}) \, dx + \int_{\Omega} ((\mathbf{U} - \mathbf{u}) \times (\mathbf{B} - \mathbf{H})) \cdot \mathbf{curl}_x \mathbf{H} \, dx \\ & + \int_{\Omega} \operatorname{div}_x ((\mathbf{B} - \mathbf{H}) \times (\mathbf{H} \times \mathbf{U})) \, dx. \end{aligned} \tag{4.6}$$

Indeed the integrals on the right-hand side can be treated as follows:

$$\begin{aligned} & \int_{\Omega} \left[(\mathbf{U} \times (\mathbf{B} - \mathbf{H})) \cdot \mathbf{curl}_x (\mathbf{B} - \mathbf{H}) + ((\mathbf{U} - \mathbf{u}) \times (\mathbf{B} - \mathbf{H})) \cdot \mathbf{curl}_x \mathbf{H} \right] \, dx \\ & = \int_{\Omega} (\mathbf{U} \times \mathbf{B}) \cdot \mathbf{curl}_x (\mathbf{B} - \mathbf{H}) \, dx - \int_{\Omega} (\mathbf{U} \times \mathbf{H}) \cdot \mathbf{curl}_x (\mathbf{B} - \mathbf{H}) \, dx \\ & \quad - \int_{\Omega} ((\mathbf{U} - \mathbf{u}) \times \mathbf{curl}_x \mathbf{H}) \cdot \mathbf{B} \, dx + \int_{\Omega} (\mathbf{curl}_x \mathbf{H} \times \mathbf{H}) \cdot (\mathbf{U} - \mathbf{u}) \, dx \\ & = - \int_{\Omega} (\mathbf{curl}_x \mathbf{B} \times \mathbf{B}) \mathbf{U} \, dx + \int_{\Omega} (\mathbf{curl}_x \mathbf{H} \times \mathbf{H}) \cdot (\mathbf{U} - \mathbf{u}) \, dx \\ & \quad - \int_{\Omega} (\mathbf{U} \times \mathbf{B}) \cdot \mathbf{curl}_x \mathbf{H} \, dx - \int_{\Omega} (\mathbf{U} \times \mathbf{H}) \cdot \mathbf{curl}_x (\mathbf{B} - \mathbf{H}) \, dx \\ & \quad - \int_{\Omega} (\mathbf{U} \times \mathbf{curl}_x \mathbf{H}) \cdot \mathbf{B} \, dx + \int_{\Omega} (\mathbf{u} \times \mathbf{curl}_x \mathbf{H}) \cdot \mathbf{B} \, dx \\ & = - \int_{\Omega} (\mathbf{curl}_x \mathbf{B} \times \mathbf{B}) \cdot \mathbf{U} \, dx + \int_{\Omega} (\mathbf{curl}_x \mathbf{H} \times \mathbf{H}) \cdot (\mathbf{U} - \mathbf{u}) \, dx \\ & \quad + \int_{\Omega} (\mathbf{B} \times \mathbf{u}) \cdot \mathbf{curl}_x \mathbf{H} \, dx + \int_{\Omega} (\mathbf{H} \times \mathbf{U}) \cdot \mathbf{curl}_x (\mathbf{B} - \mathbf{H}) \, dx \\ & = - \int_{\Omega} (\mathbf{curl}_x \mathbf{B} \times \mathbf{B}) \cdot \mathbf{U} \, dx + \int_{\Omega} (\mathbf{curl}_x \mathbf{H} \times \mathbf{H}) \cdot (\mathbf{U} - \mathbf{u}) \, dx \\ & \quad + \int_{\Omega} (\mathbf{B} \times \mathbf{u}) \cdot \mathbf{curl}_x \mathbf{H} \, dx + \int_{\Omega} (\mathbf{B} - \mathbf{H}) \cdot \mathbf{curl}_x (\mathbf{H} \times \mathbf{U}) \, dx \\ & \quad + \int_{\Omega} \operatorname{div}_x ((\mathbf{B} - \mathbf{H}) \times (\mathbf{H} \times \mathbf{U})) \, dx. \end{aligned}$$

Summing up the previous discussion, we may rewrite inequality (3.7) in the form

$$\begin{aligned} & \int_{\Omega} E(\varrho, \vartheta, \mathbf{u}, \mathbf{B} \mid r, \Theta, \mathbf{U}, \mathbf{H})(\tau, \cdot) \, dx \\ & + \int_0^\tau \int_{\Omega} \frac{\Theta}{\vartheta} \left(\mathbb{S} : \nabla_x \mathbf{u} - \frac{\mathbf{q} \cdot \nabla_x \vartheta}{\vartheta} + \zeta(\vartheta) |\mathbf{curl}_x \mathbf{B}|^2 \right) \, dx \, dt \\ & \leq - \int_0^\tau \int_{\Omega} \left(\varrho (\mathbf{u} - \mathbf{U}) \otimes (\mathbf{u} - \mathbf{U}) + p \mathbb{I} - \mathbb{S} \right) : \nabla_x \mathbf{U} \, dx \, dt \end{aligned}$$

$$\begin{aligned}
& + \int_0^\tau \int_\Omega \frac{\varrho}{r} (\mathbf{u} - \mathbf{U}) \cdot \nabla_x p(r, \Theta) \, dx \, dt \\
& - \int_0^\tau \int_\Omega \operatorname{div}_x \mathbb{S}(\Theta, \nabla_x \mathbf{U}) \cdot (\mathbf{u} - \mathbf{U}) \, dx \, dt \\
& - \int_0^\tau \int_\Omega \left(\varrho(s - s(r, \Theta)) \partial_t \Theta + \varrho(s - s(r, \Theta)) \mathbf{u} \cdot \nabla_x \Theta + \frac{\mathbf{q}}{\vartheta} \cdot \nabla_x \Theta \right) \, dx \\
& + \int_0^\tau \int_\Omega \left(\left(1 - \frac{\varrho}{r}\right) \partial_t p(r, \Theta) - \frac{\varrho}{r} \mathbf{u} \cdot \nabla_x p(r, \Theta) \right) \, dx \, dt \\
& + \int_0^\tau \int_\Omega (\mathbf{B} - \mathbf{H}) \cdot \operatorname{curl}_x (\zeta(\Theta) \operatorname{curl}_x \mathbf{H}) \, dx \, dt + \int_0^\tau \int_\Omega \zeta(\vartheta) \operatorname{curl}_x \mathbf{B} \cdot \operatorname{curl}_x \mathbf{H} \, dx \, dt \\
& + \int_0^\tau \int_\Omega \left(\frac{\varrho}{r} - 1 \right) \left(\operatorname{div}_x \mathbb{S}(\Theta, \nabla_x \mathbf{U}) + \operatorname{curl}_x \mathbf{H} \times \mathbf{H} \right) \cdot (\mathbf{U} - \mathbf{u}) \, dx \, dt \\
& + \int_0^\tau \int_\Omega \left(\mathbf{U} \times (\mathbf{B} - \mathbf{H}) \right) \cdot \operatorname{curl}_x (\mathbf{B} - \mathbf{H}) \, dx \, dt \\
& + \int_0^\tau \int_\Omega \left((\mathbf{U} - \mathbf{u}) \times (\mathbf{B} - \mathbf{H}) \right) \cdot \operatorname{curl}_x \mathbf{H} \, dx \, dt \\
& + \int_0^\tau \int_\Omega \operatorname{div}_x ((\mathbf{B} - \mathbf{H}) \times (\mathbf{H} \times \mathbf{U})) \, dx \, dt \tag{4.7}
\end{aligned}$$

In addition, we have

$$\begin{aligned}
\int_\Omega (\mathbf{B} - \mathbf{H}) \cdot \operatorname{curl}_x (\zeta(\Theta) \operatorname{curl}_x \mathbf{H}) \, dx &= \int_\Omega \zeta(\Theta) \operatorname{curl}_x \mathbf{H} \cdot \operatorname{curl}_x (\mathbf{B} - \mathbf{H}) \, dx \\
&+ \int_\Omega \operatorname{div}_x (\zeta(\Theta) (\mathbf{B} - \mathbf{H}) \cdot \operatorname{curl}_x \mathbf{H}) \, dx. \tag{4.8}
\end{aligned}$$

We claim

$$\begin{aligned}
& \int_0^\tau \int_\Omega \operatorname{div}_x ((\mathbf{B} - \mathbf{H}) \times (\mathbf{H} \times \mathbf{U})) \, dx \, dt + \int_\Omega \operatorname{div}_x (\zeta(\Theta) (\mathbf{B} - \mathbf{H}) \times \operatorname{curl}_x \mathbf{H}) \, dx \\
&= \int_0^\tau \int_{\partial\Omega} \left((\mathbf{B} - \mathbf{H}) \times (\mathbf{H} \times \mathbf{U}) \right) \cdot \mathbf{n} + \left(\zeta(\Theta) (\mathbf{B} - \mathbf{H}) \cdot \operatorname{curl}_x \mathbf{H} \right) \cdot \mathbf{n} \, d\sigma \, dt = 0. \tag{4.9}
\end{aligned}$$

Indeed either \mathbf{B} , \mathbf{H} satisfy the boundary condition (1.12) and then $(\mathbf{B} - \mathbf{H}) \times \mathbf{n} = 0$ or \mathbf{H} satisfies the flux condition (1.14). In both cases the surface integral in (4.9) vanishes.

Thus relation (4.7) takes the form

$$\begin{aligned}
& \int_\Omega E \left(\varrho, \vartheta, \mathbf{u}, \mathbf{B} \mid r, \Theta, \mathbf{U}, \mathbf{H} \right) (\tau, \cdot) \, dx \\
& + \int_0^\tau \int_\Omega \frac{\Theta}{\vartheta} \left(\mathbb{S}(\vartheta, \nabla_x \mathbf{u}) : \nabla_x \mathbf{u} - \frac{\mathbf{q}(\vartheta, \nabla_x \vartheta) \cdot \nabla_x \vartheta}{\vartheta} + \zeta(\vartheta) |\operatorname{curl}_x \mathbf{B}|^2 \right) \, dx \, dt \\
& \leq \int_0^\tau \int_\Omega \frac{\varrho}{r} (\mathbf{u} - \mathbf{U}) \cdot \nabla_x p(r, \Theta) \, dx \, dt - \int_0^\tau \int_\Omega p(\varrho, \vartheta) \operatorname{div}_x \mathbf{U} \, dx \, dt \\
& + \int_0^\tau \int_\Omega \mathbb{S}(\Theta, \nabla_x \mathbf{U}) : \nabla_x (\mathbf{u} - \mathbf{U}) \, dx \, dt + \int_0^\tau \int_\Omega \mathbb{S}(\vartheta, \nabla_x \mathbf{u}) : \nabla_x \mathbf{U} \, dx \, dt \\
& - \int_0^\tau \int_\Omega \left(\varrho(s(\varrho, \vartheta) - s(r, \Theta)) \partial_t \Theta + \varrho(s(\varrho, \vartheta) - s(r, \Theta)) \mathbf{u} \cdot \nabla_x \Theta + \frac{\mathbf{q}}{\vartheta} \cdot \nabla_x \Theta \right) \, dx \\
& + \int_0^\tau \int_\Omega \left(\left(1 - \frac{\varrho}{r}\right) \partial_t p(r, \Theta) - \frac{\varrho}{r} \mathbf{u} \cdot \nabla_x p(r, \Theta) \right) \, dx \, dt \\
& + \int_0^\tau \int_\Omega \zeta(\Theta) \operatorname{curl}_x (\mathbf{B} - \mathbf{H}) \cdot \operatorname{curl}_x \mathbf{H} \, dx \, dt + \int_0^\tau \int_\Omega \zeta(\vartheta) \operatorname{curl}_x \mathbf{B} \cdot \operatorname{curl}_x \mathbf{H} \, dx \, dt
\end{aligned}$$

$$\begin{aligned}
 & + \int_0^\tau \int_\Omega \left(\frac{\varrho}{r} - 1\right) \left(\operatorname{div}_x \mathbb{S}(\Theta, \nabla_x \mathbf{U}) + \operatorname{curl}_x \mathbf{H} \times \mathbf{H}\right) \cdot (\mathbf{U} - \mathbf{u}) \, dx \, dt \\
 & + \int_\Omega (\mathbf{U} \times (\mathbf{B} - \mathbf{H})) \cdot \operatorname{curl}_x (\mathbf{B} - \mathbf{H}) \, dx + \int_\Omega \left((\mathbf{U} - \mathbf{u}) \times (\mathbf{B} - \mathbf{H})\right) \cdot \operatorname{curl}_x \mathbf{H} \, dx \\
 & - \int_0^\tau \int_\Omega \left(\varrho(\mathbf{u} - \mathbf{U}) \otimes (\mathbf{u} - \mathbf{U})\right) : \nabla_x \mathbf{U} \, dx \, dt
 \end{aligned} \tag{4.10}$$

4.2. Pressure and Entropy

After a simple manipulation, we can rewrite (4.10) in the form

$$\begin{aligned}
 & \int_\Omega E(\varrho, \vartheta, \mathbf{u}, \mathbf{B} \mid r, \Theta, \mathbf{U}, \mathbf{H})(\tau, \cdot) \, dx \\
 & + \int_0^\tau \int_\Omega \frac{\Theta}{\vartheta} \left(\mathbb{S}(\vartheta, \nabla_x \mathbf{u}) : \nabla_x \mathbf{u} - \frac{\mathbf{q}(\vartheta, \nabla_x \vartheta) \cdot \nabla_x \vartheta}{\vartheta} + \zeta(\vartheta) |\operatorname{curl}_x \mathbf{B}|^2 \right) \, dx \, dt \\
 & \leq \int_0^\tau \int_\Omega \left(p(r, \Theta) - p(\varrho, \vartheta) \right) \operatorname{div}_x \mathbf{U} \, dx \, dt \\
 & - \int_0^\tau \int_\Omega \left(r(s(\varrho, \vartheta) - s(r, \Theta)) \partial_t \Theta + r(s(\varrho, \vartheta) - s(r, \Theta)) \mathbf{U} \cdot \nabla_x \Theta \right) \, dx \\
 & + \int_0^\tau \int_\Omega \left(1 - \frac{\varrho}{r} \right) \left(\partial_t p(r, \Theta) + \mathbf{U} \cdot \nabla_x p(r, \Theta) \right) \, dx \, dt \\
 & + \int_0^\tau \int_\Omega \mathbb{S}(\Theta, \nabla_x \mathbf{U}) : \nabla_x (\mathbf{u} - \mathbf{U}) \, dx \, dt + \int_0^\tau \int_\Omega \mathbb{S}(\vartheta, \nabla_x \mathbf{u}) : \nabla_x \mathbf{U} \, dx \, dt \\
 & + \int_0^\tau \int_\Omega \zeta(\Theta) \operatorname{curl}_x (\mathbf{B} - \mathbf{H}) \cdot \operatorname{curl}_x \mathbf{H} \, dx \, dt + \int_0^\tau \int_\Omega \zeta(\vartheta) \operatorname{curl}_x \mathbf{B} \cdot \operatorname{curl}_x \mathbf{H} \, dx \, dt \\
 & - \int_0^\tau \int_\Omega \frac{\mathbf{q}(\vartheta, \nabla_x \vartheta)}{\vartheta} \cdot \nabla_x \Theta \, dx \, dt + \int_0^\tau \mathcal{R}_1,
 \end{aligned} \tag{4.11}$$

with a quadratic remainder

$$\begin{aligned}
 \mathcal{R}_1 = & \int_0^\tau \int_\Omega (r - \varrho) \left(s(\varrho, \vartheta) - s(r, \Theta) \right) \partial_t \Theta \, dx \, dt \\
 & + \int_0^\tau \int_\Omega (r - \varrho) \left(s(\varrho, \vartheta) - s(r, \Theta) \right) \mathbf{U} \cdot \nabla_x \Theta \, dx \, dt \\
 & + \int_0^\tau \int_\Omega \varrho \left(s(\varrho, \vartheta) - s(r, \Theta) \right) (\mathbf{U} - \mathbf{u}) \cdot \nabla_x \Theta \, dx \, dt \\
 & + \int_0^\tau \int_\Omega \left(\frac{\varrho}{r} - 1\right) \left(\operatorname{div}_x \mathbb{S}(\Theta, \nabla_x \mathbf{U}) + \operatorname{curl}_x \mathbf{H} \times \mathbf{H} - \nabla_x p(r, \Theta)\right) \cdot (\mathbf{U} - \mathbf{u}) \, dx \, dt \\
 & + \int_\Omega (\mathbf{U} \times (\mathbf{B} - \mathbf{H})) \cdot \operatorname{curl}_x (\mathbf{B} - \mathbf{H}) \, dx + \int_\Omega \left((\mathbf{U} - \mathbf{u}) \times (\mathbf{B} - \mathbf{H})\right) \cdot \operatorname{curl}_x \mathbf{H} \, dx \\
 & - \int_0^\tau \int_\Omega \left(\varrho(\mathbf{u} - \mathbf{U}) \otimes (\mathbf{u} - \mathbf{U})\right) : \nabla_x \mathbf{U} \, dx \, dt \\
 & + \int_0^\tau \int_\Omega \left(\frac{\varrho}{r} - 1\right) (\mathbf{u} - \mathbf{U}) \cdot \nabla_x p(r, \Theta) \, dx \, dt.
 \end{aligned} \tag{4.12}$$

Using the identity

$$\frac{\partial s(r, \Theta)}{\partial \varrho} = -\frac{1}{r^2} \frac{\partial p(r, \Theta)}{\partial \vartheta}$$

we compute, exactly as in [5, Section 3.3],

$$\begin{aligned} & \left(1 - \frac{\varrho}{r}\right) (\partial_t p(r, \Theta) + \mathbf{U} \cdot \nabla_x p(r, \Theta)) + \operatorname{div}_x \mathbf{U} \left(p(r, \Theta) - p(\varrho, \vartheta)\right) \\ &= \operatorname{div}_x \mathbf{U} \left(p(r, \Theta) - \frac{\partial p(r, \Theta)}{\partial \varrho} (r - \varrho) - \frac{\partial p(r, \Theta)}{\partial \vartheta} (\Theta - \vartheta) - p(\varrho, \vartheta)\right) \\ & \quad - r(r - \varrho) \frac{\partial s(r, \Theta)}{\partial \varrho} (\partial_t \Theta + \mathbf{U} \cdot \nabla_x \Theta) - r(\Theta - \vartheta) \frac{\partial s(r, \Theta)}{\partial \vartheta} (\partial_t \Theta + \mathbf{U} \cdot \nabla_x \Theta) \\ & \quad - (\Theta - \vartheta) \operatorname{div}_x \left(\frac{\mathbf{q}(\Theta, \nabla_x \Theta)}{\Theta}\right) + \left(1 - \frac{\vartheta}{\Theta}\right) \left(\mathbb{S}(\Theta, \nabla_x \mathbf{U}) : \nabla_x \mathbf{U} - \frac{\mathbf{q}(\Theta, \nabla_x \Theta) \cdot \nabla_x \Theta}{\Theta}\right). \end{aligned}$$

Consequently, inequality (4.11) can be written as

$$\begin{aligned} & \int_{\Omega} E(\varrho, \vartheta, \mathbf{u}, \mathbf{B} \mid r, \Theta, \mathbf{U}, \mathbf{H})(\tau, \cdot) \, dx \\ & \quad + \int_0^\tau \int_{\Omega} \frac{\Theta}{\vartheta} \left(\mathbb{S}(\vartheta, \nabla_x \mathbf{u}) : \nabla_x \mathbf{u} - \frac{\mathbf{q}(\vartheta, \nabla_x \vartheta) \cdot \nabla_x \vartheta}{\vartheta} + \zeta(\vartheta) |\operatorname{curl}_x \mathbf{B}|^2\right) \, dx \, dt \\ & \leq - \int_0^\tau \int_{\Omega} (\Theta - \vartheta) \operatorname{div}_x \left(\frac{\mathbf{q}(\Theta, \nabla_x \Theta)}{\Theta}\right) \, dx \, dt \\ & \quad + \int_0^\tau \int_{\Omega} \left(1 - \frac{\vartheta}{\Theta}\right) \left(\mathbb{S}(\Theta, \nabla_x \mathbf{U}) : \nabla_x \mathbf{U} - \frac{\mathbf{q}(\Theta, \nabla_x \Theta) \cdot \nabla_x \Theta}{\Theta} + \zeta(\Theta) |\operatorname{curl}_x \mathbf{H}|^2\right) \, dx \, dt \\ & \quad + \int_0^\tau \int_{\Omega} \mathbb{S}(\Theta, \nabla_x \mathbf{U}) : \nabla_x (\mathbf{u} - \mathbf{U}) \, dx \, dt + \int_0^\tau \int_{\Omega} \mathbb{S}(\vartheta, \nabla_x \mathbf{u}) : \nabla_x \mathbf{U} \, dx \, dt \\ & \quad + \int_0^\tau \int_{\Omega} \zeta(\Theta) \operatorname{curl}_x (\mathbf{B} - \mathbf{H}) \cdot \operatorname{curl}_x \mathbf{H} \, dx \, dt + \int_0^\tau \int_{\Omega} \zeta(\vartheta) \operatorname{curl}_x \mathbf{B} \cdot \operatorname{curl}_x \mathbf{H} \, dx \, dt \\ & \quad - \int_0^\tau \int_{\Omega} \frac{\mathbf{q}(\vartheta, \nabla_x \vartheta)}{\vartheta} \cdot \nabla_x \Theta \, dx \, dt + \int_0^\tau \mathcal{R}_2, \end{aligned} \tag{4.13}$$

with a remainder

$$\begin{aligned} \mathcal{R}_2 &= \int_0^\tau \int_{\Omega} (r - \varrho) (s(\varrho, \vartheta) - s(r, \Theta)) \partial_t \Theta \, dx \, dt \\ & \quad + \int_0^\tau \int_{\Omega} (r - \varrho) (s(\varrho, \vartheta) - s(r, \Theta)) \mathbf{U} \cdot \nabla_x \Theta \, dx \, dt \\ & \quad + \int_0^\tau \int_{\Omega} \varrho (s(\varrho, \vartheta) - s(r, \Theta)) (\mathbf{U} - \mathbf{u}) \cdot \nabla_x \Theta \, dx \, dt \\ & \quad + \int_0^\tau \int_{\Omega} \left(\frac{\varrho}{r} - 1\right) \left(\operatorname{div}_x \mathbb{S}(\Theta, \nabla_x \mathbf{U}) + \operatorname{curl}_x \mathbf{H} \times \mathbf{H} - \nabla_x p(r, \Theta)\right) \cdot (\mathbf{U} - \mathbf{u}) \, dx \, dt \\ & \quad + \int_{\Omega} (\mathbf{U} \times (\mathbf{B} - \mathbf{H})) \cdot \operatorname{curl}_x (\mathbf{B} - \mathbf{H}) \, dx + \int_{\Omega} ((\mathbf{U} - \mathbf{u}) \times (\mathbf{B} - \mathbf{H})) \cdot \operatorname{curl}_x \mathbf{H} \, dx \\ & \quad - \int_0^\tau \int_{\Omega} (\varrho (\mathbf{u} - \mathbf{U}) \otimes (\mathbf{u} - \mathbf{U})) : \nabla_x \mathbf{U} \, dx \, dt \\ & \quad + \int_0^\tau \int_{\Omega} \left(\frac{\varrho}{r} - 1\right) (\mathbf{u} - \mathbf{U}) \cdot \nabla_x p(r, \Theta) \, dx \, dt \\ & \quad + \int_0^\tau \int_{\Omega} \operatorname{div}_x \mathbf{U} \left(p(r, \Theta) - \frac{\partial p(r, \Theta)}{\partial \varrho} (r - \varrho) - \frac{\partial p(r, \Theta)}{\partial \vartheta} (\Theta - \vartheta) - p(\varrho, \vartheta)\right) \, dx \, dt \\ & \quad + \int_0^\tau \int_{\Omega} r \left(s(r, \Theta) - \frac{\partial s(r, \Theta)}{\partial \varrho} (r - \varrho) - \frac{\partial s(r, \Theta)}{\partial \vartheta} (\Theta - \vartheta) - s(\varrho, \vartheta)\right) (\partial_t \Theta + \mathbf{U} \cdot \nabla_x \Theta) \, dx \, dt. \end{aligned} \tag{4.14}$$

4.3. Diffusive Terms

First, as ϑ, Θ coincide on $\partial\Omega$, we can integrate by parts:

$$-\int_0^\tau \int_\Omega (\Theta - \vartheta) \operatorname{div}_x \left(\frac{\mathbf{q}(\Theta, \nabla_x \Theta)}{\Theta} \right) dx dt = \int_0^\tau \int_\Omega \left(\frac{\mathbf{q}(\Theta, \nabla_x \Theta)}{\Theta} \right) \cdot (\nabla_x \Theta - \nabla_x \vartheta) dx dt$$

Putting all diffusive terms on the left-hand side of the relative energy inequality (4.13), we get

$$\begin{aligned} & \int_0^\tau \int_\Omega \left(\frac{\Theta}{\vartheta} \mathbb{S}(\vartheta, \nabla_x \mathbf{u}) : \nabla_x \mathbf{u} - \mathbb{S}(\vartheta, \nabla_x \mathbf{u}) : \nabla_x \mathbf{U} - \mathbb{S}(\Theta, \nabla_x \mathbf{U}) : \nabla_x (\mathbf{u} - \mathbf{U}) \right) dx dt \\ &= \int_0^\tau \int_\Omega \left(\left(\frac{\Theta}{\vartheta} - 1 \right) \mathbb{S}(\vartheta, \nabla_x \mathbf{u}) : \nabla_x \mathbf{u} + \left(\mathbb{S}(\vartheta, \nabla_x \mathbf{u}) - \mathbb{S}(\Theta, \nabla_x \mathbf{U}) \right) : \nabla_x (\mathbf{u} - \mathbf{U}) \right) dx dt. \end{aligned} \quad (4.15)$$

Similarly

$$\begin{aligned} & \int_0^\tau \int_\Omega \left(\frac{\Theta}{\vartheta} \zeta(\vartheta) |\operatorname{curl}_x \mathbf{B}|^2 - \zeta(\Theta) \operatorname{curl}_x (\mathbf{B} - \mathbf{H}) \cdot \operatorname{curl}_x \mathbf{H} - \zeta(\vartheta) \operatorname{curl}_x \mathbf{B} \cdot \operatorname{curl}_x \mathbf{H} \right) dx dt \\ &= \int_0^\tau \int_\Omega \left(\left(\frac{\Theta}{\vartheta} - 1 \right) \zeta(\vartheta) |\operatorname{curl}_x \mathbf{B}|^2 + \left(\zeta(\vartheta) \operatorname{curl}_x \mathbf{B} - \zeta(\Theta) \operatorname{curl}_x \mathbf{H} \right) \cdot \operatorname{curl}_x (\mathbf{B} - \mathbf{H}) \right) dx dt. \end{aligned} \quad (4.16)$$

Finally,

$$\begin{aligned} & -\int_0^\tau \int_\Omega \left(\frac{\Theta}{\vartheta} \frac{\mathbf{q}(\varrho, \nabla_x \vartheta) \cdot \nabla_x \vartheta}{\vartheta} \right) dx dt + \int_0^\tau \int_\Omega (\Theta - \vartheta) \operatorname{div}_x \left(\frac{\mathbf{q}(\Theta, \nabla_x \Theta)}{\Theta} \right) dx dt \\ & \quad + \int_0^\tau \int_\Omega \left(1 - \frac{\vartheta}{\Theta} \right) \left(\frac{\mathbf{q}(\Theta, \nabla_x \Theta) \cdot \nabla_x \Theta}{\Theta} \right) dx dt + \int_0^\tau \int_\Omega \frac{\mathbf{q}(\vartheta, \nabla_x \vartheta)}{\vartheta} \cdot \nabla_x \Theta dx dt \\ &= \int_0^\tau \int_\Omega \left(\left(1 - \frac{\Theta}{\vartheta} \right) \frac{\mathbf{q}(\vartheta, \nabla_x \vartheta) \cdot \nabla_x \vartheta}{\vartheta} + \left(1 - \frac{\vartheta}{\Theta} \right) \frac{\mathbf{q}(\Theta, \nabla_x \Theta) \cdot \nabla_x \Theta}{\Theta} \right) dx dt \\ & \quad + \int_0^\tau \int_\Omega \left(\frac{\mathbf{q}(\vartheta, \nabla_x \vartheta)}{\vartheta} - \frac{\mathbf{q}(\Theta, \nabla_x \Theta)}{\Theta} \right) \cdot (\nabla_x \Theta - \nabla_x \vartheta) dx dt \end{aligned} \quad (4.17)$$

Thus (4.13) can be rewritten in the final form

$$\begin{aligned} & \int_\Omega E(\varrho, \vartheta, \mathbf{u}, \mathbf{B} \mid r, \Theta, \mathbf{U}, \mathbf{H})(\tau, \cdot) dx \\ & \quad + \int_0^\tau \int_\Omega \left(\frac{\Theta}{\vartheta} - 1 \right) \mathbb{S}(\vartheta, \nabla_x \mathbf{u}) dx dt + \int_0^\tau \int_\Omega \left(\frac{\vartheta}{\Theta} - 1 \right) \mathbb{S}(\Theta, \nabla_x \mathbf{U}) dx dt \\ & \quad + \int_0^\tau \int_\Omega \left(\frac{\Theta}{\vartheta} - 1 \right) \zeta(\vartheta) |\operatorname{curl}_x \mathbf{B}|^2 dx dt + \int_0^\tau \int_\Omega \left(\frac{\vartheta}{\Theta} - 1 \right) \zeta(\Theta) |\operatorname{curl}_x \mathbf{H}|^2 dx dt \\ & \quad + \int_0^\tau \int_\Omega \left(\left(1 - \frac{\Theta}{\vartheta} \right) \frac{\mathbf{q}(\varrho, \nabla_x \vartheta) \cdot \nabla_x \vartheta}{\vartheta} + \left(1 - \frac{\vartheta}{\Theta} \right) \frac{\mathbf{q}(\Theta, \nabla_x \Theta) \cdot \nabla_x \Theta}{\Theta} \right) dx dt \\ & \quad + \int_0^\tau \int_\Omega \left(\mathbb{S}(\vartheta, \nabla_x \mathbf{u}) - \mathbb{S}(\Theta, \nabla_x \mathbf{U}) \right) : \nabla_x (\mathbf{u} - \mathbf{U}) dx dt \\ & \quad + \int_0^\tau \int_\Omega \left(\zeta(\vartheta) \operatorname{curl}_x \mathbf{B} - \zeta(\Theta) \operatorname{curl}_x \mathbf{H} \right) \cdot \operatorname{curl}_x (\mathbf{B} - \mathbf{H}) dx dt \\ & \quad + \int_0^\tau \int_\Omega \left(\frac{\mathbf{q}(\vartheta, \nabla_x \vartheta)}{\vartheta} - \frac{\mathbf{q}(\Theta, \nabla_x \Theta)}{\Theta} \right) \cdot (\nabla_x \Theta - \nabla_x \vartheta) dx dt \\ & \leq \int_0^\tau \mathcal{R}_2 dt, \end{aligned} \quad (4.18)$$

with \mathcal{R}_2 given in (4.14).

4.4. Refined Estimates

Similarly to [10, Chapter 4, Section 4.2.1], we introduce the “essential” and “residual” component of a measurable function. We set

$$\begin{aligned} \mathcal{O}_{\text{ess}} &= \left\{ (t, x) \in (0, T) \times \Omega \mid \frac{1}{2} \inf_{(0, T) \times \Omega} r \leq \varrho(t, x) \leq 2 \sup_{(0, T) \times \Omega} r \right. \\ &\quad \left. \frac{1}{2} \inf_{(0, T) \times \Omega} \Theta \leq \vartheta(t, x) \leq 2 \sup_{(0, T) \times \Omega} \Theta \right\} \\ \mathcal{O}_{\text{res}} &= \left((0, T) \times \Omega \right) \setminus \mathcal{O}_{\text{ess}}, \\ h_{\text{ess}} &= h \mathbb{1}_{\mathcal{O}_{\text{ess}}}, \quad h_{\text{res}} = h - h_{\text{ess}} = h \mathbb{1}_{\mathcal{O}_{\text{res}}}. \end{aligned}$$

The remaining part of the proof of weak–strong uniqueness principle follows essentially the arguments of [10, Chapter 4, Section 4], with the necessary modifications to accommodate the terms containing the magnetic field as well as the complete slip/Dirichlet boundary conditions. First, we recall the coercivity property of the relative energy:

$$\begin{aligned} E \left(\varrho, \vartheta, \mathbf{u}, \mathbf{B} \mid r, \Theta, \mathbf{U}, \mathbf{H} \right) &\geq c \left(\left| [\varrho - r]_{\text{ess}} \right|^2 + \left| [\vartheta - \Theta]_{\text{ess}} \right|^2 + \left| [\mathbf{u} - \mathbf{U}]_{\text{ess}} \right|^2 + \left| [\mathbf{B} - \mathbf{H}]_{\text{ess}} \right|^2 \right), \\ E \left(\varrho, \vartheta, \mathbf{u}, \mathbf{B} \mid r, \Theta, \mathbf{U}, \mathbf{H} \right) &\geq (1_{\text{res}} + [\varrho \mathbf{u}^2]_{\text{res}} + [\varrho e(\varrho, \vartheta)]_{\text{res}} + [\varrho |s(\varrho, \vartheta)]_{\text{res}} + \left| [\mathbf{B}^2]_{\text{res}} \right|), \end{aligned} \tag{4.19}$$

where the constant depend on $\inf r, \sup r, \inf \Theta, \sup \Theta$.

4.4.1. Temperature Gradient. The following crucial estimate was proved in [10, Chapter 4, Section 4.2.2] for $\beta = 3$. Here, we extend its validity to general $\beta \geq 3$.

Remark 4.4. Under the hypotheses of Theorem 4.1, there is a constant $c > 0$,

$$\begin{aligned} &\| \vartheta - \Theta \|_{W^{1,2}(\Omega; \mathbb{R}^3)}^2 \\ &\leq c \int_{\Omega} \left(\left(\frac{\Theta}{\vartheta} - 1 \right) \frac{\kappa(\vartheta) |\nabla_x \vartheta|^2}{\vartheta} + \left(\frac{\vartheta}{\Theta} - 1 \right) \frac{\kappa(\Theta) |\nabla_x \Theta|^2}{\Theta} \right) dx \\ &\quad + \int_{\Omega} \left(\frac{\kappa(\Theta) \nabla_x \Theta}{\Theta} - \frac{\kappa(\vartheta) \nabla_x \vartheta}{\vartheta} \right) \cdot (\nabla_x \Theta - \nabla_x \vartheta) dx + \xi \int_{\Omega} E \left(\varrho, \vartheta, \mathbf{u}, \mathbf{B} \mid r, \Theta, \mathbf{U}, \mathbf{H} \right) dx \end{aligned} \tag{4.20}$$

where $\xi > 0$ depends on $\Theta, \inf \Theta$, and $\| \vartheta \|_{L^4(\Omega)}$.

Remark 4.5. As we shall see below, the result holds whenever $\kappa(\vartheta) \approx 1 + \vartheta^\beta, \beta \geq 2$.

Proof. We consider first the essential part. Rewrite

$$\begin{aligned} &\left(\frac{\Theta}{\vartheta} - 1 \right) \frac{\kappa(\vartheta) |\nabla_x \vartheta|^2}{\vartheta} + \left(\frac{\vartheta}{\Theta} - 1 \right) \frac{\kappa(\Theta) |\nabla_x \Theta|^2}{\Theta} + \left(\frac{\kappa(\Theta) \nabla_x \Theta}{\Theta} - \frac{\kappa(\vartheta) \nabla_x \vartheta}{\vartheta} \right) \cdot (\nabla_x \Theta - \nabla_x \vartheta) \\ &\geq \Theta \frac{\kappa(\vartheta)}{\vartheta^2} |\nabla_x \vartheta - \nabla_x \Theta|^2 + 2 \frac{\kappa(\vartheta)}{\vartheta^2} (\Theta - \vartheta) (\nabla_x \vartheta - \nabla_x \Theta) \cdot \nabla_x \Theta \\ &\quad + \left(\frac{\kappa(\vartheta)}{\vartheta} - \frac{\kappa(\Theta)}{\Theta} \right) (\nabla_x \vartheta - \nabla_x \Theta) \cdot \nabla_x \Theta + \left(\frac{\kappa(\vartheta)}{\vartheta} - \frac{\kappa(\Theta)}{\Theta} \right) \left(\frac{\Theta - \vartheta}{\Theta} \right) |\nabla_x \Theta|^2. \end{aligned} \tag{4.21}$$

Applying Hölder’s inequality we deduce from (4.21)

$$\begin{aligned} &\int_{\Omega} \left| \nabla_x \vartheta - \nabla_x \Theta \right|_{\text{ess}}^2 dx \\ &\lesssim \left[\left(\frac{\Theta}{\vartheta} - 1 \right) \frac{\kappa(\vartheta) |\nabla_x \vartheta|^2}{\vartheta} + \left(\frac{\vartheta}{\Theta} - 1 \right) \frac{\kappa(\Theta) |\nabla_x \Theta|^2}{\Theta} + \left(\frac{\kappa(\Theta) \nabla_x \Theta}{\Theta} - \frac{\kappa(\vartheta) \nabla_x \vartheta}{\vartheta} \right) \cdot (\nabla_x \Theta - \nabla_x \vartheta) \right]_{\text{ess}} \end{aligned}$$

$$+ \xi \int_{\Omega} E \left(\varrho, \vartheta, \mathbf{u}, \mathbf{B} \mid r, \Theta, \mathbf{U}, \mathbf{H} \right)_{\text{ess}} dx, \tag{4.22}$$

where ξ depends on Θ . As a matter of fact, the only hypotheses to be imposed on κ is that $\kappa(\vartheta) > 0$ whenever $\vartheta > 0$.

As for the residual part, we first rewrite

$$\begin{aligned} & \left(\frac{\Theta}{\vartheta} - 1 \right) \frac{\kappa(\vartheta) |\nabla_x \vartheta|^2}{\vartheta} + \left(\frac{\vartheta}{\Theta} - 1 \right) \frac{\kappa(\Theta) |\nabla_x \Theta|^2}{\Theta} + \left(\frac{\kappa(\Theta) \nabla_x \Theta}{\Theta} - \frac{\kappa(\vartheta) \nabla_x \vartheta}{\vartheta} \right) \cdot (\nabla_x \Theta - \nabla_x \vartheta) \\ &= \frac{\Theta}{\vartheta} \frac{\kappa(\vartheta) |\nabla_x \vartheta|^2}{\vartheta} + \frac{\vartheta}{\Theta} \frac{\kappa(\Theta) |\nabla_x \Theta|^2}{\Theta} - \frac{\kappa(\vartheta) \nabla_x \vartheta}{\vartheta} \cdot \nabla_x \Theta - \frac{\kappa(\Theta) \nabla_x \Theta}{\Theta} \cdot \nabla_x \vartheta \end{aligned} \tag{4.23}$$

Obviously,

$$\left[\frac{\vartheta}{\Theta} \frac{\kappa(\Theta) |\nabla_x \Theta|^2}{\Theta} \right]_{\text{res}} \lesssim \int_{\Omega} E \left(\varrho, \vartheta, \mathbf{u}, \mathbf{B} \mid r, \Theta, \mathbf{U}, \mathbf{H} \right)_{\text{res}} dx.$$

Consequently, as $\inf \Theta > 0$, it remains to handle a quantity

$$\left[\frac{\kappa(\vartheta) |\nabla_x \vartheta|^2}{\vartheta^2} - C \kappa(\vartheta) |\nabla_x \Theta| \right]_{\text{res}}$$

where C is a positive constant depending on $\Theta, \inf \Theta$. Note that this step requires $\vartheta^2 \lesssim \kappa(\vartheta)$.

To proceed we consider $\bar{\vartheta} > 0$ to be fixed below. We write

$$\left[\frac{\kappa(\vartheta) |\nabla_x \vartheta|^2}{\vartheta^2} - C \kappa(\vartheta) |\nabla_x \Theta| \right]_{\text{res}} \mathbf{1}_{\vartheta < \bar{\vartheta}} + \left[\frac{\kappa(\vartheta) |\nabla_x \vartheta|^2}{\vartheta^2} - C \kappa(\vartheta) |\nabla_x \Theta| \right]_{\text{res}} \mathbf{1}_{\vartheta \geq \bar{\vartheta}}.$$

Obviously, there is $\xi = \xi(\bar{\vartheta}, \Theta)$ such that

$$\begin{aligned} & \int_{\Omega} |[\nabla_x \vartheta]_{\text{res}}|^2 \mathbf{1}_{\vartheta < \bar{\vartheta}} dx \\ & \lesssim \int_{\Omega} \left[\frac{\kappa(\vartheta) |\nabla_x \vartheta|^2}{\vartheta^2} - C \kappa(\vartheta) |\nabla_x \Theta| \right]_{\text{res}} \mathbf{1}_{\vartheta < \bar{\vartheta}} dx + \xi \int_{\Omega} E \left(\varrho, \vartheta, \mathbf{u}, \mathbf{B} \mid r, \Theta, \mathbf{U}, \mathbf{H} \right)_{\text{res}} dx. \end{aligned} \tag{4.24}$$

Thus it remains to handle the integral

$$\int_{\Omega} \left[\frac{\kappa(\vartheta) |\nabla_x \vartheta|^2}{\vartheta^2} - C \kappa(\vartheta) |\nabla_x \Theta| \right]_{\text{res}} \mathbf{1}_{\vartheta \geq \bar{\vartheta}} dx = \int_{\vartheta \geq \bar{\vartheta}} \left(\frac{\kappa(\vartheta) |\nabla_x \vartheta|^2}{\vartheta^2} - C \kappa(\vartheta) |\nabla_x \Theta| \right) dx.$$

As $\kappa(\vartheta) \approx (1 + \vartheta^\beta)$, our task reduces to

$$\int_{\vartheta \geq \bar{\vartheta}} \left(\frac{\kappa(\vartheta) |\nabla_x \vartheta|^2}{\vartheta^2} - C \kappa(\vartheta) |\nabla_x \Theta| \right) dx \approx \int_{\vartheta \geq \bar{\vartheta}} \left(|\nabla_x \vartheta^{\frac{\beta}{2}}|^2 - C \vartheta^\beta |\nabla_x \Theta| \right) dx.$$

First observe

$$\int_{\vartheta \geq \bar{\vartheta}} \vartheta^\beta |\nabla_x \Theta| dx \leq \int_{\vartheta \geq \bar{\vartheta}} \left(\vartheta^{\frac{\beta}{2}} - \bar{\vartheta}^{\frac{\beta}{2}} \right)^2 |\nabla_x \Theta| dx + \int_{\vartheta \geq \bar{\vartheta}} \bar{\vartheta}^\beta |\nabla_x \Theta| dx,$$

where

$$\int_{\vartheta \geq \bar{\vartheta}} \bar{\vartheta}^\beta |\nabla_x \Theta| dx \leq \zeta \int_{\Omega} E \left(\varrho, \vartheta, \mathbf{u}, \mathbf{B} \mid r, \Theta, \mathbf{U}, \mathbf{H} \right)_{\text{res}} dx.$$

Finally, it remains to handle the integral

$$\begin{aligned} & \int_{\vartheta \geq \bar{\vartheta}} \left(|\nabla_x \vartheta^{\frac{\beta}{2}}|^2 - C \left(\vartheta^{\frac{\beta}{2}} - \bar{\vartheta}^{\frac{\beta}{2}} \right)^2 |\nabla_x \Theta| \right) dx \\ &= \int_{\vartheta \geq \bar{\vartheta}} \left(\left| \nabla_x \left(\vartheta^{\frac{\beta}{2}} - \bar{\vartheta}^{\frac{\beta}{2}} \right) \right|^2 - C \left(\vartheta^{\frac{\beta}{2}} - \bar{\vartheta}^{\frac{\beta}{2}} \right)^2 |\nabla_x \Theta| \right) dx. \end{aligned}$$

On the one hand, using Sobolev–Poincaré inequality we get

$$\begin{aligned} \int_{\vartheta \geq \bar{\vartheta}} \left| \nabla_x \left(\vartheta^{\frac{\beta}{2}} - \bar{\vartheta}^{\frac{\beta}{2}} \right) \right|^2 dx &= \int_{\Omega} \left| \nabla_x \left[\vartheta^{\frac{\beta}{2}} - \bar{\vartheta}^{\frac{\beta}{2}} \right]^+ \right|^2 dx \\ &\gtrsim \left(\int_{\Omega} \left| \left[\vartheta^{\frac{\beta}{2}} - \bar{\vartheta}^{\frac{\beta}{2}} \right]^+ \right|^6 dx \right)^{\frac{1}{3}} = \left(\int_{\vartheta \geq \bar{\vartheta}} \left(\vartheta^{\frac{\beta}{2}} - \bar{\vartheta}^{\frac{\beta}{2}} \right)^6 dx \right)^{\frac{1}{3}}. \end{aligned} \tag{4.25}$$

On the other hand, by Hölder’s inequality,

$$\int_{\vartheta \geq \bar{\vartheta}} \left(\vartheta^{\frac{\beta}{2}} - \bar{\vartheta}^{\frac{\beta}{2}} \right)^2 |\nabla_x \Theta| dx \leq \left(\int_{\vartheta \geq \bar{\vartheta}} \left(\vartheta^{\frac{\beta}{2}} - \bar{\vartheta}^{\frac{\beta}{2}} \right)^4 dx \right)^{\frac{1}{2}} \left(\int_{\vartheta \geq \bar{\vartheta}} |\nabla_x \Theta|^2 dx \right)^{\frac{1}{2}}.$$

Next, by Jensen’s inequality,

$$\begin{aligned} \left(\int_{\vartheta \geq \bar{\vartheta}} \left(\vartheta^{\frac{\beta}{2}} - \bar{\vartheta}^{\frac{\beta}{2}} \right)^4 dx \right)^{\frac{3}{2}} &= \left(\int_{\Omega} \left(\left[\vartheta^{\frac{\beta}{2}} - \bar{\vartheta}^{\frac{\beta}{2}} \right]^+ \right)^4 dx \right)^{\frac{3}{2}} \\ &\lesssim \int_{\Omega} \left(\left[\vartheta^{\frac{\beta}{2}} - \bar{\vartheta}^{\frac{\beta}{2}} \right]^+ \right)^6 dx = \int_{\vartheta \geq \bar{\vartheta}} \left(\vartheta^{\frac{\beta}{2}} - \bar{\vartheta}^{\frac{\beta}{2}} \right)^6 dx; \end{aligned}$$

whence

$$\left(\int_{\vartheta \geq \bar{\vartheta}} \left(\vartheta^{\frac{\beta}{2}} - \bar{\vartheta}^{\frac{\beta}{2}} \right)^4 dx \right)^{\frac{1}{2}} \lesssim \left(\int_{\vartheta \geq \bar{\vartheta}} \left(\vartheta^{\frac{\beta}{2}} - \bar{\vartheta}^{\frac{\beta}{2}} \right)^6 dx \right)^{\frac{1}{3}}. \tag{4.26}$$

Our final observation is that as $\|\vartheta\|_{L^4(\Omega)}$ is bounded, for any $\delta > 0$, there is $\bar{\vartheta}(\delta)$ such that

$$\int_{\vartheta \geq \bar{\vartheta}} |\nabla_x \Theta|^2 dx \leq \delta.$$

Combining (4.25), (4.26) we may infer

$$\begin{aligned} &\int_{\Omega} |[\nabla_x \vartheta]_{\text{res}}|^2 \mathbf{1}_{\vartheta \geq \bar{\vartheta}} dx \\ &\lesssim \int_{\Omega} \left[\frac{\kappa(\vartheta) |\nabla_x \vartheta|^2}{\vartheta^2} - C\kappa(\vartheta) |\nabla_x \Theta| \right]_{\text{res}} dx + \xi \int_{\Omega} E(\varrho, \vartheta, \mathbf{u}, \mathbf{B} | r, \Theta, \mathbf{U}, \mathbf{H})_{\text{res}} dx \end{aligned} \tag{4.27}$$

Putting together (4.22), (4.24), and (4.27) we obtain (4.20). □

4.4.2. Viscous Stress and Magnetic Diffusion. Exactly as in [10, Chapter 4, Section 4.2.3], we estimate the viscous stress dissipation:

$$\begin{aligned} &\left\| \nabla_x(\mathbf{u} - \mathbf{U}) + \nabla_x^t(\mathbf{u} - \mathbf{U}) - \frac{2}{3} \text{div}_x(\mathbf{u} - \mathbf{U}) \mathbb{I} \right\|_{L^q(\Omega; \mathbb{R}^{3 \times 3})}^2 \\ &\leq c \left(\int_{\Omega} \left(\frac{\Theta}{\vartheta} - 1 \right) \mathbb{S}(\vartheta, \nabla_x \mathbf{u}) + \left(\frac{\vartheta}{\Theta} - 1 \right) \mathbb{S}(\Theta, \nabla_x \mathbf{U}) dx \right. \\ &\quad \left. + \int_{\Omega} \left(\mathbb{S}(\vartheta, \nabla_x \mathbf{u}) - \mathbb{S}(\Theta, \nabla_x \mathbf{U}) \right) : \nabla_x(\mathbf{u} - \mathbf{U}) dx + \int_{\Omega} E(\varrho, \vartheta, \mathbf{u}, \mathbf{B} | r, \Theta, \mathbf{U}, \mathbf{H}) dx \right), \end{aligned} \tag{4.28}$$

where

$$q = \frac{8}{5 - \alpha}. \tag{4.29}$$

Next, repeating the same arguments, we get

$$\|\text{curl}_x(\mathbf{B} - \mathbf{H})\|_{L^2(\Omega; \mathbb{R}^3)}^2 \leq c \left(\int_{\Omega} \left(\frac{\Theta}{\vartheta} - 1 \right) \zeta(\vartheta) |\text{curl}_x \mathbf{B}|^2 + \left(\frac{\vartheta}{\Theta} - 1 \right) \zeta(\Theta) |\text{curl}_x \mathbf{H}|^2 dx \right)$$

$$+ \int_{\Omega} \left(\zeta(\vartheta) \mathbf{curl}_x \mathbf{B} - \zeta(\Theta) \mathbf{curl}_x \mathbf{H} \right) \cdot \mathbf{curl}_x (\mathbf{B} - \mathbf{H}) \, dx + \int_{\Omega} E \left(\varrho, \vartheta, \mathbf{u}, \mathbf{B} \mid r, \Theta, \mathbf{U}, \mathbf{H} \right) \, dx \Big). \tag{4.30}$$

Finally, we recall the generalized Korn–Poincaré inequality [9, Chapter 11, Theorem 11.23]:

$$\|\mathbf{v}\|_{W^{1,q}(\Omega;R^3)} \leq c(\delta, q) \left(\left\| \nabla_x \mathbf{v} + \nabla_x^t \mathbf{v} - \frac{2}{3} \operatorname{div}_x \mathbf{v} \mathbb{I} \right\|_{L^q(\Omega;R^{3 \times 3})} + \int_{\Omega} \varrho |\mathbf{v}| \, dx \right) \tag{4.31}$$

whenever

$$1 < q < \infty, \varrho \geq 0, \int_{\Omega} \varrho \, dx \geq \delta, \int_{\Omega} \varrho^\gamma \, dx < \frac{1}{\delta}, \gamma > 1. \tag{4.32}$$

The previous estimates, together with (4.32), allow us to rewrite the inequality (4.18) in the form

$$\begin{aligned} & \int_{\Omega} E \left(\varrho, \vartheta, \mathbf{u}, \mathbf{B} \mid r, \Theta, \mathbf{U}, \mathbf{H} \right) (\tau, \cdot) \, dx \\ & + \int_0^\tau \left(\|\mathbf{u} - \mathbf{U}\|_{W^{1,q}(\Omega;R^3)}^2 + \|\vartheta - \Theta\|_{W^{1,2}(\Omega)}^2 + \|\mathbf{curl}_x (\mathbf{B} - \mathbf{H})\|_{L^2(\Omega;R^3)}^2 \right) dt \\ & \leq \int_0^\tau \mathcal{R}_2 \, dt + c \int_0^\tau \int_{\Omega} E \left(\varrho, \vartheta, \mathbf{u}, \mathbf{B} \mid r, \Theta, \mathbf{U}, \mathbf{H} \right) \, dx dt, \quad q = \frac{8}{5 - \alpha}, \end{aligned} \tag{4.33}$$

where the remainder \mathcal{R}_2 is specified in (4.14).

4.5. Completing the Proof of Weak–Strong Uniqueness

The rest of the proof of weak–strong uniqueness consists in absorbing all integrals appearing in the remainder \mathcal{R}_2 by the left-hand side of (4.33) and applying Gronwall’s argument. We omit the details as the whole procedure is described in [10, Chapter 4].

We have proved Theorem 4.1.

Remark 4.6. The Lipschitz regularity of the domain Ω required in Theorem 4.1 may not be sufficient for the *existence* of the strong solution to the MHD system.

5. Existence of Weak Solutions

The proof of existence of weak solutions can be carried out by means of the approximate scheme, *a priori* estimates, and compactness arguments specified in Chapter 5 and Chapter 12 of the monograph [10].

The magnetic field equations can be incorporated exactly as in [7] without any additional difficulties. Similarly to [10, Chapter 12], two additional assumptions must be imposed on the constitutive relations, specifically,

$$\lim_{Z \rightarrow \infty} \mathcal{S}(Z) = 0, \tag{5.1}$$

where \mathcal{S} is the function defining the entropy s_M . Moreover, we require

$$\beta > 6, \tag{5.2}$$

where β is the exponent in (2.8). While (5.1) corresponds to the Third Law of Thermodynamics, hypothesis (5.2) is purely technical to ensure the necessary *a priori* bounds.

5.1. Approximation Scheme

The existence of weak solutions can be shown by solving the following multi-level approximation scheme. The equation of continuity (1.1) is replaced by its *parabolic regularization*:

$$\partial_t \varrho + \operatorname{div}_x(\varrho \mathbf{u}) = \varepsilon \Delta_x \varrho \text{ in } (0, T) \times \Omega, \quad \varepsilon > 0, \tag{5.3}$$

supplemented by the Neumann boundary conditions

$$\nabla_x \varrho \cdot \mathbf{n}|_{\partial\Omega} = 0, \tag{5.4}$$

and the initial condition

$$\varrho(0, \cdot) = \varrho_{0,\delta}, \tag{5.5}$$

where $\varrho_{0,\delta} > 0$ is a smooth regularization of the initial density ϱ_0 . Here and hereafter, ε and δ are small parameters to be sent to zero successively in the limit passage.

The momentum equation (1.2) is replaced by a Galerkin approximation

$$\begin{aligned} & \int_0^\tau \int_\Omega \left[\varrho \mathbf{u} \cdot \partial_t \boldsymbol{\varphi} + \varrho \mathbf{u} \otimes \mathbf{u} : \nabla_x \boldsymbol{\varphi} + p_\delta(\varrho, \vartheta) \operatorname{div}_x \boldsymbol{\varphi} - \mathbb{S}_\delta(\vartheta, \nabla_x \mathbf{u}) : \nabla_x \boldsymbol{\varphi} \right] dx dt \\ & - \varepsilon \int_0^\tau \int_\Omega \nabla_x \varrho \cdot \nabla_x \mathbf{u} \cdot \nabla_x \boldsymbol{\varphi} dx dt + \int_0^\tau \int_\Omega \left[\operatorname{curl}_x \mathbf{B} \times \mathbf{B} \cdot \boldsymbol{\varphi} + \varrho \mathbf{g} \cdot \boldsymbol{\varphi} \right] dx dt \\ & = \int_\Omega \varrho \mathbf{u}(\tau, \cdot) \cdot \boldsymbol{\varphi}(\tau, \cdot) dx - \int_\Omega (\varrho \mathbf{u})_0 \cdot \boldsymbol{\varphi}(0, \cdot) dx \end{aligned} \tag{5.6}$$

for any test function $\boldsymbol{\varphi} \in C^1([0, T]; X_N)$, where X_N is a finite dimensional space spanned by the eigenvectors of the Lamé operator with the complete slip boundary conditions (1.10). Here,

$$p_\delta(\varrho, \vartheta) = p(\varrho, \vartheta) + \delta (\varrho^2 + \varrho^\Gamma), \quad \mathbb{S}_\delta = \mathbb{S}(\vartheta, \nabla_x \mathbf{u}) + \delta \vartheta \left(\nabla_x \mathbf{u} + \nabla_x^t \mathbf{u} - \frac{2}{3} \operatorname{div}_x \mathbf{u} \mathbb{I} \right), \quad \delta > 0. \tag{5.7}$$

Similarly, the induction equation is replaced by a Galerkin approximation. We look for solutions in the form

$$\mathbf{B} = \mathbf{B}_B + \mathbf{b},$$

where

$$\begin{aligned} & \int_0^\tau \int_\Omega \left[(\mathbf{B}_B + \mathbf{b}) \cdot \partial_t \boldsymbol{\varphi} - \left((\mathbf{B}_B + \mathbf{b}) \times \mathbf{u} \right) \cdot \operatorname{curl}_x \boldsymbol{\varphi} - \zeta(\vartheta) \operatorname{curl}_x (\mathbf{B}_B + \mathbf{b}) \cdot \operatorname{curl}_x \boldsymbol{\varphi} \right] dx dt \\ & = \int_\Omega (\mathbf{B}_B + \mathbf{b}) \cdot \boldsymbol{\varphi}(\tau, \cdot) dx - \int_\Omega \mathbf{B}_0 \cdot \boldsymbol{\varphi}(0, \cdot) dx \end{aligned} \tag{5.8}$$

for any $\boldsymbol{\varphi} \in C^1([0, T]; Y_N)$. The finite-dimensional space Y_N is chosen in accordance with the boundary conditions. The functions in Y_N are smooth, solenoidal,

$$\operatorname{div}_x \boldsymbol{\varphi} = 0 \text{ in } \Omega,$$

and $\boldsymbol{\varphi} \times \mathbf{n}|_{\partial\Omega} = 0$, $\cup_{N>0} Y_N$ is dense in $H_{0,\tau}$ in the case of boundary condition (1.12), and $\boldsymbol{\varphi} \cdot \mathbf{n}|_{\partial\Omega} = 0$, $\cup_{N>0} Y_N$ is dense in $H_{0,n}$ in the case of boundary conditions (1.13), (1.14).

Finally, the internal energy balance is regularized, adapted to the viscous approximation of the equation of continuity, and solved exactly,

$$\begin{aligned} & \partial_t (\varrho e_\delta(\varrho, \vartheta)) + \operatorname{div}_x (\varrho e_\delta(\varrho, \vartheta) \mathbf{u}) + \operatorname{div}_x \mathbf{q}_\delta(\vartheta, \nabla_x \vartheta) = \mathbb{S}_\delta(\vartheta, \nabla_x \mathbf{u}) + \zeta(\vartheta) |\operatorname{curl}_x \mathbf{B}|^2 \\ & - p(\varrho, \vartheta) \operatorname{div}_x \mathbf{u} + \varepsilon \delta (\varrho^{\Gamma-2} + 2) |\nabla_x \varrho|^2 + \delta \frac{1}{\varrho^2} - \varepsilon \vartheta^5, \end{aligned} \tag{5.9}$$

with the regularized initial/boundary conditions

$$\vartheta(0, \cdot) = \vartheta_{0,\delta}, \quad \vartheta|_{\partial\Omega} = \vartheta_B. \tag{5.10}$$

Here, we have denoted

$$\begin{aligned} \mathbf{q}_\delta &= \mathbf{q} - \delta \left(\vartheta^\Gamma + \frac{1}{\vartheta} \right) \nabla_x \vartheta, \\ e_\delta &= e + \delta \vartheta. \end{aligned} \tag{5.11}$$

5.2. Existence of Weak Solutions

The existence of weak solutions to the Navier–Stokes–Fourier system (without magnetic field) follows by solving the approximate scheme (5.3)–(5.7), (5.9)–(5.11). The proof in the absence of magnetic field and for the *Dirichlet boundary conditions* for the velocity field is given with full details in [10, Part II, Chapter 5]. The crucial observation made in [7] is that the basic energy estimate consists in considering the scalar product of (5.6) with \mathbf{u} and (5.8) with \mathbf{b} —a step compatible with the Galerkin approximation. The proof can be reproduced without any essential difficulties in the present setting as soon as the *complete slip boundary conditions* (1.10) are accommodated.

The main problem associated to the complete slip boundary conditions (1.10) is the absence of the Korn–Poincaré inequality necessary to control the term

$$\int_\Omega \varrho s \mathbf{u} \cdot \nabla_x \tilde{\vartheta} \, dx$$

or, more specifically, its radiation component

$$\int_\Omega \vartheta^3 \mathbf{u} \cdot \nabla_x \tilde{\vartheta} \, dx \tag{5.12}$$

in the ballistic energy inequality, where $\tilde{\vartheta}$ is the unique harmonic extension of the boundary temperature,

$$\Delta_x \tilde{\vartheta} = 0, \quad \tilde{\vartheta}|_{\partial\Omega} = \vartheta_B,$$

cf. [10, Chapter 12, Section 12.4.2].

To overcome this difficulty, an extra hypothesis is needed depending on the specific shape of the spatial domain Ω . Denote

$$\mathcal{N} = \left\{ \mathbf{w} \in W^{1,2}(\Omega; \mathbb{R}^3) \mid \underline{\mu} \left(\mathbb{D}_x \mathbf{w} - \frac{1}{3} \operatorname{div}_x \mathbf{w} \mathbb{I} \right) = 0, \quad \underline{\eta} \operatorname{div}_x \mathbf{w} = 0 \text{ in } \Omega, \quad \mathbf{w} \cdot \mathbf{n}|_{\partial\Omega} = 0 \right\},$$

where $\underline{\mu} > 0$, $\underline{\eta} \geq 0$ are the structural constants introduced in (2.7). If $\underline{\mu} > 0$, $\underline{\eta} = 0$, the set \mathcal{N} contains all conformal Killing vectors with vanishing normal trace, see Schirra [20, Proposition 2.5]. If, in addition, $\underline{\eta} > 0$, then the set \mathcal{N} consists of all *rigid motions* $\mathbf{w} = \boldsymbol{\omega} \times x + \mathbf{a}$ with zero normal trace. In the latter case, $\mathcal{N} \neq \{0\}$ if and only if Ω is rotationally symmetric. Consequently, a suitable hypothesis to handle the integral (5.12) reads:

$$\nabla_x \tilde{\vartheta}(t, \cdot) \cdot \mathbf{w} = 0 \text{ for any } \mathbf{w} \in \mathcal{N} \text{ and any } t \in [0, T]. \tag{5.13}$$

Note that (5.13) is satisfied in the case when Ω is an annulus bounded by two concentric spheres,

$$\Omega = \left\{ x \in \mathbb{R}^3 \mid 0 < r_1 < |x| < r_2 \right\} \text{ and } \vartheta_B = \Theta_1 \text{ if } |x| = r_1, \quad \vartheta_B = \Theta_2 \text{ if } |x| = r_2,$$

provided $\underline{\mu}, \underline{\eta} > 0$, and Θ_1, Θ_2 are possibly time dependent positive constants. Indeed we recall Korn–Poincaré inequality, see e.g. Lewintan, Müller, Neff [16]:

$$\|\mathbf{v} - \Pi_{\mathcal{N}} \mathbf{v}\|_{W^{1,q}(\Omega; \mathbb{R}^3)} \lesssim \left(\underline{\mu} \left\| \mathbb{D}_x \mathbf{v} - \frac{1}{3} \operatorname{div}_x \mathbf{v} \mathbb{I} \right\|_{L^q(\Omega; \mathbb{R}^{3 \times 3})} + \underline{\eta} \|\operatorname{div}_x \mathbf{v}\|_{L^q(\Omega)} \right), \quad 1 < q < \infty, \tag{5.14}$$

for any $\mathbf{v} \in W^{1,q}(\Omega; R^3)$, $\mathbf{v} \cdot \mathbf{n}|_{\partial\Omega} = 0$, where $\Pi_{\mathcal{N}}$ denotes a projection onto the space \mathcal{N} . Accordingly, the integral (5.12) can be written as

$$\int_{\Omega} \vartheta^3 \mathbf{u} \cdot \nabla_x \tilde{\vartheta} \, dx = \int_{\Omega} \vartheta^3 (\mathbf{u} - \Pi_{\mathcal{N}} \mathbf{u}) \cdot \nabla_x \tilde{\vartheta} \, dx,$$

and estimated via (5.14), exactly as in [10, Chapter 12, Section 12.4.2].

We may therefore state the following result concerning the existence of weak solutions to the compressible MHD system.

Theorem 5.1. (Existence of weak solution) *Let $\Omega \subset R^3$ be a bounded domain of class at least C^3 . Suppose the data $\vartheta_B, \mathbf{B}_B$ are twice continuously differentiable,*

$$\inf_{[0,T] \times \partial\Omega} \vartheta_B > 0, \quad \operatorname{div}_x \mathbf{B}_B = 0 \text{ in } (0, T) \times \Omega.$$

Let the thermodynamic functions p, e, s satisfy the hypotheses (2.1)–(2.6) and the transport coefficients $\mu, \eta, \kappa,$ and ζ comply with (2.7)–(2.9), where

$$\frac{1}{2} \leq \alpha \leq 1, \quad \beta > 6, \quad \lim_{Z \rightarrow \infty} \mathcal{S}(Z) = 0. \tag{5.15}$$

In addition, suppose the harmonic extension of the boundary temperature satisfies (5.13).

Then for any initial data

$$\begin{aligned} \varrho_0, \varrho_0 > 0, \quad \vartheta_0, \vartheta_0 > 0, \quad \mathbf{u}_0, \mathbf{B}_0, \quad \operatorname{div}_x \mathbf{B}_0 = 0, \\ \int_{\Omega} \left[\frac{1}{2} \varrho_0 |\mathbf{u}_0|^2 + \varrho_0 e(\varrho_0, \vartheta_0) + \frac{1}{2} |\mathbf{B}_0|^2 \right] dx < \infty, \end{aligned}$$

and any $\mathbf{g} \in L^\infty((0, T) \times \Omega; R^3)$, the compressible MHD system admits a weak solution in $(0, T) \times \Omega$ in the sense specified in Definition 2.1.

Remark 5.2. As a matter of fact, positivity of the initial density and temperature is not really necessary in the weak framework. Expressing the initial data in terms of the initial density ϱ_0 , the momentum \mathbf{m}_0 , the total entropy S_0 , and the magnetic field \mathbf{B}_0 , we can show global in time existence under the condition that the initial energy

$$\int_{\Omega} \left[\frac{1}{2} \frac{|\mathbf{m}_0|^2}{\varrho_0} + \varrho_0 e(\varrho_0, S_0) + \frac{1}{2} |\mathbf{B}_0|^2 \right] dx < \infty$$

is finite.

Remark 5.3. The weak solutions resulting from the approximate scheme specified above satisfy the differential version of the ballistic energy inequality (2.24).

Remark 5.4. The hypothesis $\lim_{Z \rightarrow \infty} \mathcal{S}(Z) = 0$ can be omitted and $\beta \geq 3$ allowed as soon as the boundary temperature ϑ_B is constant.

6. Concluding Remarks

The theory can be extended in a straightforward manner to other types of boundary conditions.

- The complete slip boundary conditions (1.10) can be replaced by the Dirichlet (no slip) boundary conditions

$$\mathbf{u}|_{\partial\Omega} = \mathbf{u}_B$$

as long as $\mathbf{u}_B \cdot \mathbf{n} = 0$. This enables to include problems of Taylor–Coutte type. The extension to general in/out flow boundary conditions for the velocity would require a more elaborated treatment in view of the presence of the magnetic field. For the in/out flow boundary conditions for the Navier–Stokes–Fourier system in the absence of magnetic field see [10].

- The Dirichlet boundary conditions for the temperature can be replaced by a flux type condition on a part of the boundary:

$$\partial\Omega = \Gamma_1 \cup \Gamma_2, \quad \vartheta|_{\Gamma_1} = \vartheta_B, \quad \mathbf{q} \cdot \mathbf{n}|_{\Gamma_2} = \mathbf{h}.$$

Note that this type of condition is particularly relevant in the modelling of stellar magnetoconvection, see Thompson and Christensen–Dalsgaard [23].

- The induction flux (electric field) can be prescribed in (1.14):

$$\left[\mathbf{B} \times \mathbf{u} + \zeta \operatorname{curl}_x \mathbf{B} \right] \times \mathbf{n}|_{\partial\Omega} = \mathbf{E}_B \times \mathbf{n}.$$

- Mixed type boundary conditions on *different* components of $\partial\Omega$ can be accommodated without essential difficulties.

Funding Funding was provided by Grantová Agentura České Republiky (Grant No. 21-02411S), Korea National Science Foundation (Grant No. NRF2022R1F1A1073801). Narodowe Centrum Badań i Rozwoju (Grant No. 2021/43/B/ST1/02851).

Data Availability Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

Conflict of interest The authors declare that they have no conflict of interest.

Open Access. This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

References

- [1] Alekseev, G.V.: Solvability of control problems for stationary equations of the magnetohydrodynamics of a viscous fluid. *Sibirsk. Mat. Zh.* **45**(2), 243–263 (2004)
- [2] Alexander, J.C., Auchmuty, G.: l^2 -well-posedness of 3d div-curl boundary value problems. *Quart. Appl. Math.* **63**(3), 479–508 (2005)
- [3] Battaner, A.: *Astrophysical Fluid Dynamics*. Cambridge University Press, Cambridge (1996)
- [4] Buckmaster, T., Cao-Labora, G., Gómez-Serrano, J.: Smooth self-similar imploding profiles to 3d compressible Euler. arxiv preprint No. [arXiv:2301.10101](https://arxiv.org/abs/2301.10101) (2023)
- [5] Chaudhuri, N., Feireisl, E.: Navier–Stokes–Fourier system with Dirichlet boundary conditions. *Appl. Anal.* **101**(12), 4076–4094 (2022)
- [6] Davidson, P.A.: *Turbulence: An Introduction For Scientists and Engineers*. Oxford University Press, Oxford (2004)
- [7] Ducomet, B., Feireisl, E.: The equations of magnetohydrodynamics: on the interaction between matter and radiation in the evolution of gaseous stars. *Commun. Math. Phys.* **266**, 595–629 (2006)
- [8] Feireisl, E., Novotný, A.: Weak-strong uniqueness property for the full Navier–Stokes–Fourier system. *Arch. Ration. Mech. Anal.* **204**, 683–706 (2012)
- [9] Feireisl, E., Novotný, A.: *Singular Limits in Thermodynamics of Viscous Fluids*. *Advances in Mathematical Fluid Mechanics*, 2nd edn. Birkhäuser, Cham (2017)
- [10] Feireisl, E., Novotný, A.: *Mathematics of Open Fluid Systems*. Birkhäuser, Basel (2022)
- [11] Feireisl, E., Petzeltová, H.: On the long-time behaviour of solutions to the Navier–Stokes–Fourier system with a time-dependent driving force. *J. Dyn. Differ. Equ.* **19**(3), 685–707 (2007)
- [12] Gough, D.: Towards understanding solar convection and activity. *Solar Phys.* **192**, 3–26 (2000)
- [13] Kotschote, M.: Strong solutions to the compressible non-isothermal Navier–Stokes equations. *Adv. Math. Sci. Appl.* **22**(2), 319–347 (2012)

- [14] Ladyženskaja, O.A., Solonnikov, V.A.: Solution of some non-stationary problems of magnetohydrodynamics for a viscous incompressible fluid. *Trudy Mat. Inst. Steklov* **59**, 115–173 (1960)
- [15] Lassner, G.: Über ein Rand-Anfangswertproblem der Magnetohydrodynamik. *Arch. Ration. Mech. Anal.* **25**, 388–405 (1967)
- [16] Lewintan, P., Müller, S., Neff, P.: Korn inequalities for incompatible tensor fields in three space dimensions with conformally invariant dislocation energy. *Calc. Var. Partial Differ. Equ.* **60**(4), 150, 46 (2021)
- [17] Matsumura, A., Nishida, T.: The initial value problem for the equations of motion of compressible and heat conductive fluids. *Commun. Math. Phys.* **89**, 445–464 (1983)
- [18] Merle, F., Raphaël, P., Rodnianski, I., Szeftel, J.: On the implosion of a compressible fluid I: smooth self-similar inviscid profiles. *Ann. Math. (2)* **196**(2), 567–778 (2022)
- [19] Merle, F., Raphaël, P., Rodnianski, I., Szeftel, J.: On the implosion of a compressible fluid II: singularity formation. *Ann. Math. (2)* **196**(2), 779–889 (2022)
- [20] Schirra, O.D.: New Korn-type inequalities and regularity of solutions to linear elliptic systems and anisotropic variational problems involving the trace-free part of the symmetric gradient. *Calc. Var. Partial Differ. Equ.* **43**(1–2), 147–172 (2012)
- [21] Ströhmer, G.: About an initial-boundary value problem from magnetohydrodynamics. *Math. Z.* **209**(3), 345–362 (1992)
- [22] Tao, L., Weiss, N.O., Brownjohn, D.P., Proctor, M.R.E.: Flux separation in stellar magnetoconvection. *Astronom. J.* **496**, L39–L42 (1998)
- [23] Thompson, M.J., Christensen-Dalsgaard, J.: *Stellar Astrophysical Fluid Dynamics*. Cambridge University Press, Cambridge (2003)
- [24] Valli, A.: Periodic and stationary solutions for compressible Navier–Stokes equations via a stability method. *Ann. Sc. Norm. Sup. Pisa* **10**(1), 607–646 (1983)
- [25] Valli, A., Zajaczkowski, M.: Navier–Stokes equations for compressible fluids: Global existence and qualitative properties of the solutions in the general case. *Commun. Math. Phys.* **103**, 259–296 (1986)
- [26] Weiss, N.O., Proctor, M.R.E.: *Magnetoconvection*. Cambridge Monographs on Mechanics. Cambridge University Press, Cambridge (2014)
- [27] Yang, Y., Wan, M., Matthaeus, W.M., Chen, S.: Energy budget in decaying compressible MHD turbulence. *J. Fluid Mech.* **916**, A4-1-A4-29 (2021)

Eduard Feireisl
Institute of Mathematics of the Academy of Sciences of the
Czech Republic
Žitná 25
115 67 Praha 1
Czech Republic

Piotr Gwiazda
Institute of Mathematics of Polish Academy of Sciences
Śniadeckich 8
00-956 Warszawa
Poland

Young-Sam Kwon
Department of Mathematics
Dong-A University
Busan 49315
Republic of Korea

Agnieszka Świerczewska-Gwiazda
Institute of Applied Mathematics and Mechanics
University of Warsaw Banacha 2
02-097 Warsaw
Poland
e-mail: aswiercz@mimuw.edu.pl

(accepted: August 28, 2023; published online: October 7, 2023)