



On Perturbations of Extreme Kerr–Newman Black Holes and their Evolution

Martin Reiris

Abstract. Using black hole inequalities and the increase of the horizon's areas, we show that there are arbitrarily small electro-vacuum perturbations of the standard initial data of the extreme Reissner–Nordström black hole that (by contradiction) cannot decay in time into any extreme Kerr–Newman black hole. This proves that, in a formal sense, the reduced family of the extreme Kerr–Newman black holes is unstable. It remains of course to be seen whether the whole family of charged black holes, including those extremes, is stable or not.

1. Introduction

In this article it is proved that there are arbitrarily small electrovacuum perturbations of the standard initial data of the extreme Reissner–Nordström black hole that cannot decay in time into any extreme Kerr–Newman (EKN) black hole. Strictly speaking, this says that the reduced family of EKN black holes is unstable. One must make it clear however that this does not prove the *instability*, nor the *stability*, in the way that they are currently formulated in the literature and which are, for their physical significance, the relevant notions than one should consider. Roughly speaking (see [15]), the eKN black holes are called stable if small generic perturbations of them settle eventually into extreme or near-extreme Kerr–Newman black holes. Otherwise, they are called *unstable*. As discussed later, the results presented in this article will favor the stability (in this last sense), rather than the instability.

To bring more accuracy to this introduction, let us start reviewing the mathematics and the qualitative properties of the extreme black holes. The Lorentzian metric of the EKN space-time of electric charge Q_E , magnetic charge Q_M , angular momentum J and mass $m^2 = (Q^2 + \sqrt{4J^2 + Q^4})/2 \neq 0$, ($Q^2 = Q_E^2 + Q_M^2$), is given by

$$\mathbf{g} = -\frac{\Delta - a^2 \sin^2 \theta}{\Sigma} dt^2 - \frac{2a \sin^2 \theta}{\Sigma} (r^2 + a^2 - \Delta) dt d\phi + \frac{(r^2 + a^2)^2 - \Delta a^2 \sin^2 \theta}{\Sigma} \sin^2 \theta d\phi^2 + \frac{\Sigma}{\Delta} dr^2 + \Sigma d\theta^2, \tag{1}$$

where $a = J/m$, $\Sigma = r^2 + a^2 \cos^2 \theta$, and $\Delta = r^2 + a^2 + Q^2 - 2mr$ (see for instance [5]). The coordinate t ranges in $(-\infty, \infty)$, r in (m, ∞) and (θ, φ) are the standard coordinates of the unit sphere \mathbb{S}^2 . The space-time \mathbf{M} is therefore diffeomorphic to $\mathbb{R} \times \mathbb{R} \times \mathbb{S}^2$.

The electromagnetic potential \mathbf{A} is given explicitly by

$$\mathbf{A} = -\frac{Q_E r}{\Sigma} (dt - a \sin^2 \theta d\phi) + \frac{Q_M \cos \theta}{\Sigma} (a dt - (r^2 + a^2) d\phi)$$

and recall that the electromagnetic tensor is $\mathbf{F}_{ab} = \nabla_a \mathbf{A}_b - \nabla_b \mathbf{A}_a$.¹ The solution is rotational symmetric and stationary. Of particular interest for this article are the EKN solutions with $J = 0$, $Q_M = 0$ but $Q_E \neq 0$, which are called extreme Reissner–Nordström (ERN). When $Q_E = 1$ the ERN metric (from now on ERN₁) takes the synthetic form

$$\mathbf{g} = -(1 - 1/r)^2 dt^2 + \frac{1}{(1 - 1/r)^2} dr^2 + r^2 d\Omega^2, \tag{2}$$

($d\Omega^2$ is the line element of the round sphere), and the electromagnetic potential simplifies to $\mathbf{A} = -dt/r$. Over the Cauchy hypersurface $\{t = 0\}$ the electric field is $E_a = \mathbf{F}_{ab} \mathbf{n}^b = \partial_r / (r^2 |\partial_r|)$ and the magnetic field is zero, i.e., $B_a = \star \mathbf{F}_{ab} \mathbf{n}^b = 0$. Here \mathbf{n} is the time-like unit normal to $\{t = 0\}$. The solution is time symmetric and therefore the second fundamental form K of the slice $\{t = 0\}$ is zero. Finally, the solution is spherically symmetric and static. For future reference the data set over $\Sigma_0 := \{t = 0\}$ will be called the *standard initial data of the ERN₁ solution* and denoted by $(\Sigma_0; g_0, K_0; E_0, B_0)$.

The EKN solutions form part of the larger family of Kerr–Newman (KN) space-times and lie exactly between those KN space-times representing black holes and those exhibiting naked singularities. Due to their special properties, the EKN solutions have played a peculiar role in the mathematical and physical analysis of black holes. Some of their most noticeable features are the following. The past and the future null infinity of the ERN space-time can be reached from any of its space-time points. Yet the ERN space-time is geodesically incomplete and exhibits future and past Cauchy horizons. Each Cauchy horizon is diffeomorphic to $\mathbb{R} \times \mathbb{S}^2$, has complete null generators and the area of any spherical section is

$$A = 4\pi \sqrt{4|J| + Q^2}$$

In particular, if an extreme solution has $Q_E = 1$ then to be the one with $Q_E = 1$, $Q_M = 0$ and $J = 0$ it is necessary and sufficient that $A = 4\pi$. Moreover, the “initial” Cauchy hypersurface $\{t = 0\}$ is maximal and complete (as a Riemannian manifold), and possess no trapped region. This hypersurface is diffeomorphic to $\mathbb{R} \times \mathbb{S}^2$ and has one cylindrical end and one asymptotically

¹ Note that \mathbf{A} is not smooth at $\{\theta = 0\} \cup \{\theta = \pi\}$. In this article smooth means C^∞ .




FIGURE 1. Picture of the (half) Penrose diagram of the EKN black-holes. The picture shows also a visualization of the geometry of the standard initial data and the future Cauchy horizon

flat (AF) end (see Fig. 1). Of special interest to us is the cylindrical space-time of the ERN_1 solution (Bertotti’s space-time). It is found by taking a sequence $r_i \rightarrow 1$, making then the change of variables $\bar{x} = \ln((r-1)/(r_i-1))$, $\bar{t} = (r_i-1)t$ in (2), and finally taking the limit as $r_i \rightarrow 1$. This gives the result

$$\check{g} = -e^{2\bar{x}} d\bar{t}^2 + d\bar{x}^2 + d\Omega^2 \tag{3}$$

The three-metric over $\{\bar{t} = 0\}$ is then $\check{g}_0 = d\bar{x}^2 + d\Omega^2$, that is, that of the metric product $\mathbb{R} \times \mathbb{S}^2$, hence cylindrical. For future reference, over this slice the electric field is $\check{E}_0 = \partial_{\bar{x}}$ and the magnetic field \check{B}_0 and the second fundamental form \check{K}_0 are zero. The data set $(\mathbb{R} \times \mathbb{S}^2; \check{g}_0, \check{K}_0; \check{E}_0, \check{B}_0)$ will be called the *standard initial data of the extreme RN_1 throat* ($ERNT_1$).

It is fundamentally the presence of these peculiar Cauchy horizons what makes extreme solutions so special. Are extreme black holes physically realistic solutions? Are they stable under small perturbations of the initial data? What occurs to their horizons under such perturbations?

A revitalised interest in these old questions reappeared in the last years as a part of new and larger mathematical investigations on the stability of black hole space-times, [1, 3, 11, 14, 17] (to mention some). Most of these theoretical developments are characterized by the use of linear techniques over the otherwise unperturbed ERN background. As a contribution to the ongoing discussion we prove here that there are arbitrarily small perturbations of the standard ERN_1 initial data whose evolution cannot decay in any way into any EKN solution. The proof is satisfactory to us in that it is the result of combining black hole inequalities [10, 12], and the ubiquitous law of area increase of event horizons [7], and does not rely in any linear or linearization technique. In a sense, our argument belongs to a class of natural procedures to prove

instabilities that was used in the literature during the last years² and which consists in finding certain inequalities at the level of the perturbed initial data that are shown to be propagated along the evolution and that are incompatible with the stationary states that one wants to rule out as the long time limit of the evolution (see for instance [13] and references therein).

Before we pass to explain the generalities behind the proof, let us explain in precise terms the main statement to be proved. We first introduce the notion of “perturbation” of the standard initial data $(\Sigma_0; g_0, K_0; E_0, B_0)$ of the ERN₁ space-time.

Definition 1. Let $(\Sigma; g, K; E, B)$ be a smooth an maximal electro-vacuum data set and let k be an integer greater or equal than 1. We say that the data set is ε -close in C^k to the ERN₁ standard initial data iff there is a diffeomorphism $\varphi : \Sigma_0 \rightarrow \Sigma$ such that for any (U, U_0) equal to either (g, g_0) , (K, K_0) , (E, E_0) or (B, B_0) we have

$$\| \varphi^*U - U_0 \|_{C^k_{g_0}(\Sigma_0)} \leq \varepsilon.$$

The $C^k_{g_0}$ norm of a tensor W (no matter its valence) is defined as usual by

$$\| W \|_{C^k_{g_0}(\Sigma_0)}^2 = \sup_{p \in \Sigma_0} \left[\sum_{j=0}^{j=k} |(\nabla^{(j)} W)(p)|_{g_0}^2 \right]$$

The Definition 1 is satisfactory, but we need to make sure that the perturbation “falls off” along the asymptotically cylindrical end and that the “cylindrical asymptotic” is preserved. To be concrete, we will work with perturbations that “fall off exponentially along the cylindrical end into the ERN₁ standard initial data”. Precisely, we say that a data set $(\Sigma; g, K; E, B)$, ε -close in C^k to $(\Sigma_0; g_0, K_0; E_0, B_0)$, falls off exponentially into $(\Sigma_0; g_0, K_0; E_0, B_0)$ along the cylindrical end iff there is $\Lambda > 0$ such that for any (U, U_0) equal to either (g, g_0) , (K, K_0) , (E, E_0) or (B, B_0) we have

$$\lim_{r(p) \rightarrow 1} e^{\Lambda \ln(r-1)} \left[\sum_{j=0}^{j=k} |(\nabla^{(j)} (\varphi^*U - U_0))(p)|_{g_0}^2 \right] = 0,$$

where φ^* is the pull-back by the diffeomorphism $\varphi : \Sigma_0 \rightarrow \Sigma$ (note that $r(p) \rightarrow 1$ means that “ p ” diverges along the cylindrical end).

With all these definitions at hand we can state our main result as follows.

Theorem 1. For any $\bar{\varepsilon} > 0$ and integer $k \geq 1$ there is a smooth and maximal electro-vacuum data set $(\bar{\Sigma}; \bar{g}, \bar{K}; \bar{E}, \bar{B})$, $\bar{\varepsilon}$ -close in C^k to the standard ERN₁ initial data and falling into it exponential along the cylindrical end, which cannot decay, towards the future or the past, into any EKN solution.

Let us overview now the basics behind the proof. A technical proof has to be found inside the text. The argument that follows can be done in any time direction. The idea is to construct (arbitrarily small) axisymmetric perturbations of the standard ERN₁ initial data and do so with sufficiently control to

² I would like to thank Piotr Chruściel for making this remark to me.




FIGURE 2. Diagram of the initial data used in this article

be able to prove that a Marginally Outer Trapped Surface (MOTS) forms separating the two ends (see Fig. 2). In addition, the perturbation is done keeping $Q_E = 1$, $Q_M = 0$ and $J = 0$. In particular, and because the electromagnetic charges and the angular momentum are conserved, if the perturbation evolves into an EKN space-time in the long-time, then it must be one with $Q_E = 1$, $Q_M = 0$ and $J = 0$, that is, it has to be the ERN that is being perturbed.³ Moreover, due to presence of a MOTS which acts as a barrier, the event horizon must intersect the initial Cauchy hypersurface somewhere between the MOTS and the asymptotically flat end. In parallel to all this it is shown that every surface S embedded in the initial hypersurface and separating the two ends has area strictly greater than 4π . In particular, the intersection of the event horizon and the initial hypersurface must have area strictly greater than 4π . As the areas of sections of the event horizon are non-decreasing in time, we conclude that the initial data cannot evolve into the ERN_1 solution because its horizon has area exactly 4π . The perturbed data set is depicted in Fig. 2 and the (presumed) evolution in Fig. 3.

This argumentation essentially contains all the ingredients of the proof, but as the reader may have wondered, it assumes implicitly a particular “convergence” of the dynamical horizon to the smooth one of ERN_1 . This is not entirely satisfying because horizons are found out of global information and their local regularity is not tied up necessarily to the local regularity of the space-time. For this reason one could imagine a space-time decaying into a EKN but with a highly irregular horizon whose areas (due to this irregularity) do not approach the area of the horizon of the EKN but rather some bigger value. To avoid this objection the definition that we adopt for “decaying into a EKN solution” does not make any hypothesis about the horizon but instead assumes only convergence to a EKN strictly outside of it. The statement is given in Definition 4 and is postponed until Sect. 7 (when we use it for the first time) as introducing it here would cause much disruption. The actual

³ To be certain here, the charges and the angular momentum are not only conserved at null infinity, they take also the same values over any embedded sphere isotopic to a “sphere” at “spatial infinity”. This is explained in Sect. 2.




FIGURE 3. Picture of the geometric construction in the argument by contradiction of the proof of Theorem 1

proof of the Theorem 1 modifies slightly the argument given above to stick to the actual definition.

Like any argument by contradiction, our proof of Theorem 1 does not say what indeed occurs during the time evolution. It just says something of what cannot happen. Nevertheless, the presence of the mentioned MOTS in the perturbed initial data suggests that it must decay in the long-time into a non-extremal RN black hole. For this reason, it is expected also that whatever occurs to the “old” horizon of the ERN_1 , that part of the space-time stays hidden inside the new black hole region. Regardless of that, this work does not yield any light about the fate of the ERN horizon under perturbations. In this sense it does not make previous investigations about the ERN horizon less interesting.

In principle, with further work but following a similar argument, one should be able to prove that there are arbitrarily small perturbations of any EKN that cannot decay in any way into an EKN black hole. What makes the use of the ERN and not of any other EKN solution more useful is that the perturbations can be made time-symmetric and for this reason proving the existence of a MOTS reduces to proving the existence of a minimal surface which is technically more accessible.⁴

The organization of this article is the following. In Sect. 2, we recall the basic material to be used about electro-vacuum space-times. In Sect. 3, we discuss black hole inequalities on data sets that we call of the ERN_1 “type” and that are introduced in Definition 3. Roughly speaking, such data sets are defined to share the topology and the asymptotic geometry of the standard initial data of the ERN_1 solution. Not surprisingly, the perturbations of the standard initial data of the ERN_1 solution that we are going to use are of the ERN_1 type. The main result of this section is to prove that the area of any (compact, boundaryless and embedded) surface separating the two ends

⁴ I would like to thank Sergio Dain for pointing this out.

of any data set of the ERN_1 type is strictly greater than 4π . The analysis in this section shares many elements with [19]. In Sect. 4, we construct the mentioned initial perturbations using the conformal method. The existence of solutions of the conformal equations is proved following standard barrier methods [8] which give good control on the solutions. In Sect. 5, we show the rigidity of the $ERNT_1$ initial data which will be necessary in Sect. 6 to show that one can make arbitrarily small perturbations containing MOTS. It is worth mentioning that the rigidity of the $ERNT_1$ initial data is of interest in itself. In particular, the formation of extreme RN throats along sequence of data sets can be studied in the same way as was done in [19] with the formation of extreme Kerr-throats. The proof of the main result following the lines explained above is made formally and finally in Sect. 7.

2. Background Material

In this section, we recall succinctly and with certain formality those notions, like that of electric and magnetic charges, that will be necessary throughout the article. The formal treatment is justified by the mathematical nature of the paper.

We will be working with smooth (C^∞) electro-vacuum space-times $(\mathbf{M}; \mathbf{g}; \mathbf{F})$, where $(\mathbf{M}; \mathbf{g})$ is an orientable and time orientable Lorentzian manifold. We will assume that an orientation on \mathbf{M} was chosen and that a future direction was assigned. Let Σ be a space-like hyper-surface and \mathbf{n} a future unit normal to Σ . As usual, the orientation on \mathbf{M} and the field \mathbf{n} provide an orientation on Σ , more precisely: $\{e_1(p), e_2(p), e_3(p)\}$ is a positive basis of $T_p\Sigma$ iff $\{\mathbf{n}(p), e_1(p), e_2(p), e_3(p)\}$ is a positive basis of $T_p\mathbf{M}$. Space-times tensors, like the Ricci curvature \mathbf{Ric} of \mathbf{g} , will be boldfaced.

(i) The Einstein–Maxwell system.

In coordinate-independent form the Einstein–Maxwell equations are

$$\mathbf{Ric} - \frac{1}{2} \mathbf{R} \mathbf{g} = 8\pi \mathbf{T}, \quad d\mathbf{F} = 0, \quad \text{and} \quad d\star\mathbf{F} = 0 \tag{4}$$

where d is the exterior derivative and \star is the \mathbf{g} -Hodge star, namely $\star\mathbf{F}_{ab} = \epsilon_{abcd}\mathbf{F}^{cd}/2$. The electromagnetic energy-momentum tensor \mathbf{T} appearing in (4) is

$$\mathbf{T}_{ab} = \frac{1}{4\pi} \left(\mathbf{F}_{ac}\mathbf{F}_b{}^c - \frac{1}{4}\mathbf{F}_{cd}\mathbf{F}^{cd}\mathbf{g}_{ab} \right).$$

The 3+1 picture of (4) will be also used during the article. We recall it in what follows [6]. Let Σ_0 be a space-like hyper-surface (possibly with boundary) and \mathbf{V} a nowhere zero time-like vector field defined on an open neighborhood of Σ_0 . By moving Σ_0 along \mathbf{V} one obtains a flow of space-like hypersurfaces Σ_t (at least for a short time). Coordinates charts (x^1, x^2, x^3) are propagated by \mathbf{V} to every Σ_t and any two Σ_t and $\Sigma_{t'}$ are naturally diffeomorphic. In this way one obtains a flow $(g_{ij}(t), K_{ij}(t))$ of induced three-metrics and second fundamental forms on the fixed manifold Σ_0 . Writing $\mathbf{V}|_{\Sigma_t} = N(t)\mathbf{n} + X^i(t)\partial_i$, where \mathbf{n} is

a future unit normal to Σ_t , one obtains also a flow of lapse functions $N(t)$ and shift vectors $X(t) = X^i \partial_i$. In this 3 + 1 setup the Einstein equation (first Eq. in (4)) is

$$\begin{cases} \dot{g}_{ij} = -2NK_{ij} + \mathcal{L}_X g_{ij}, \\ \dot{K}_{ij} = -\nabla_i \nabla_j N + N(Ric_{ij} + kK_{ij} - 2K_{il}K^l_j) \\ \quad + \mathcal{L}_X K_{ij} - 8\pi N(\mathbf{T}_{ij} + \frac{1}{2}(\mathbf{T}_{ab} \mathbf{g}^{ab})g_{ij}), \\ R = |K|^2 - k^2 + 16\pi \mathbf{T}_{00}, \\ \nabla^i K_{ij} - \nabla_j k = 8\pi \mathbf{T}_{0i}, \end{cases}$$

where $\mathbf{T}_{00} = \mathbf{T}(\mathbf{n}, \mathbf{n})$ and $\mathbf{T}_{0i} = \mathbf{T}(\mathbf{n}, \partial_i)$, ∇ is the g -covariant derivative, $k = \text{tr}_g K$ is the mean curvature and \mathcal{L} is the Lie-derivative. The space-time metric is written in the form

$$\mathbf{g} = -(N^2 - X_i X^i) dt^2 + X_i (dt \otimes dx^i + dx^i \otimes dt) + g_{ij} dx^i dx^j$$

At every slice Σ_t , the electric and magnetic fields E and B , are defined by $E^i = \mathbf{F}^i_a \mathbf{n}^a$ and $B^i = \star \mathbf{F}^i_a \mathbf{n}^a$. In terms of them the electro-vacuum constraint equations are

$$\begin{cases} R = |K|^2 - k^2 + 2(|E|^2 + |B|^2), \\ \nabla^i K_{ij} - \nabla_j k = 2(E \times B)_j, \\ \nabla^i E_i = 0, \\ \nabla^i B_i = 0 \end{cases} \tag{5}$$

where $(E \times B)_j = \epsilon_{ijk} E^j B^k$. A set $(g, K; E, B)$ satisfying the constraint equations (5) on a manifold Σ is called an *electro-vacuum data set*. The data is maximal if $k = \text{tr}_g K = 0$.

(ii) The electric and magnetic charges.

Let $[S]$ be an oriented, compact and boundaryless surface S embedded in \mathbf{M} . The bracket $[\]$ signifies that an orientation on S has been assigned. Then $Q_E([S])$ and $Q_M([S])$ are defined by

$$Q_E([S]) = -\frac{1}{4\pi} \int_{[S]} \star \mathbf{F} \quad \text{and} \quad Q_M([S]) := -\frac{1}{4\pi} \int_{[S]} \mathbf{F}$$

As $d\mathbf{F} = 0$ and $d\star \mathbf{F} = 0$ then $Q_E([S])$ and $Q_M([S])$ depend only on the homology class of $[S]$. We will be referring this fact as the *conservation of charge*. If S is embedded in a space-like hypersurface Σ then $Q_E([S])$ and $Q_M([S])$ take the more familiar expressions

$$Q_E([S]) := \frac{1}{4\pi} \int_S \langle E, \zeta \rangle dA \quad \text{and} \quad Q_M([S]) := \frac{1}{4\pi} \int_S \langle B, \zeta \rangle dA \tag{6}$$

where $\langle E, \zeta \rangle = E^i \zeta^j g_{ij}$ and where ζ the unit normal field to S in Σ such that if $\{e_2(p), e_3(p)\}$ is a positive basis for $T_p S$ then $\{n(p), \zeta(p), e_2(p), e_3(p)\}$ is a positive basis for \mathbf{M} . Observe that if $[S]$ and $[S']$ are homologous in Σ (and therefore in \mathbf{M}) then the conservations $Q_E([S]) = Q_E([S'])$ and $Q_M([S]) =$

$Q_M([S'])$ can be seen also as a consequence of the laws $\operatorname{div} E = 0$ and $\operatorname{div} B = 0$ ($\operatorname{div} U = \nabla^i U_i$).

In this context, the total charges Q_E and Q_M that show up in the metric expression (1) of the EKN solutions are of course the electric and magnetic charges of any sphere with t and r constant and oriented using the outgoing normal $\zeta = \partial_r/|\partial_r|$.⁵

It is the case that the normal ζ will be given from the context (or simply will not matter). For this reason we will often write $Q_E(S)$ and $Q_M(S)$.

(iii) Angular momentum in electro-vacuum space-times.

Suppose now that the electro-vacuum space-time $(M; g; F)$ is axisymmetric and that $F = dA$ with the potential A axisymmetric.⁶ Denote by ξ the axisymmetric Killing field. Then the angular momentum of an oriented and axisymmetric (compact and boundaryless) surface $[S]$ is [5]

$$J([S]) := \frac{1}{8\pi} \int_{[S]} \star(\nabla_a \xi_b) + \frac{1}{4\pi} \int_{[S]} (A^a \xi_a) \star F \tag{7}$$

The angular momentum is conserved too [5]. Namely if $[\Sigma]$ is an oriented compact and axisymmetric hypersurface of M and $\partial[\Sigma] = [S] - [S']$, then $J([S]) = J([S'])$.

If S is embedded in an axisymmetric Cauchy hypersurface Σ , then the first term in (7) (which is the Komar angular momentum) reduces to the standard form $(\int_S K(\xi, \zeta) dA)/8\pi$ and is therefore zero when $K = 0$. If in addition $B = 0$ over Σ then the second term in (7) is also zero. To see this use the axisymmetry of A to get $\xi^a F_{ai} = \nabla_i A(\xi)$ and to conclude that $A(\xi)$ must be a constant over Σ . When S is in addition a sphere then the constant must be zero because $A(\xi)$ must vanish at the axes. This information shows that the perturbations constructed in Sect. 4, which have $K = 0$ and $B = 0$, also have total angular momentum J equal to zero.

(iv) The stability inequality of minimal surfaces embedded in maximal data sets.

Let $(\Sigma; g, K; E, B)$ be an electro-vacuum data set and suppose that S is a (compact, boundary-less and orientable) minimal surface embedded in Σ . Recall that a surface S is said minimal inside $(\Sigma; g)$ if its mean curvature is identically zero. Let ζ be a unit normal vector field to S in Σ and let $\alpha : S \rightarrow \mathbb{R}$ be a smooth function. The first variation of area when S is deformed along $\alpha\zeta$ is zero by minimality. Instead, the second variation is [9]

$$A''_\alpha(S) := \int_S [|\nabla\alpha|^2 - (|\Theta|^2 + \operatorname{Ric}(\zeta, \zeta)) \alpha^2] dA, \tag{8}$$

where here Θ is the second fundamental form of S . The surface S is said to be stable if $A''_\alpha(S) \geq 0$ for all α . In dimension three the r.h.s of (8) is simplified

⁵ Assume $\{\partial_t, \partial_r, \partial_\theta, \partial_\varphi\}$ is positive for M .

⁶ If F is exact then an axisymmetric potential A can always be found by averaging any potential by the rotational group $U(1)$. Observe too that F is exact iff all the magnetic charges (i.e., $Q_M([S]) = 0$ for all S) are zero.

due to the identity $2\kappa = (\text{tr}_h \Theta)^2 - |\Theta|^2 + R - 2\text{Ric}(\zeta, \zeta)$, where κ is the Gaussian curvature of S (with its induced metric). Using this expression, the minimality of S (i.e., $\text{tr}_h \Theta = 0$) and the energy constraint, we deduce that if S is stable then for any α we have

$$\int_S (|\nabla \alpha|^2 + \kappa \alpha^2) \, dA \geq \frac{1}{2} \int_S (2|E|^2 + 2|B|^2 + |K|^2 + |\Theta|^2 - k^2) \alpha^2 \, dA. \tag{9}$$

3. Black Holes Inequalities in Maximal Data Sets

Definition 2. We say that a sphere S embedded in a maximal electro-vacuum data set $(\Sigma; g, K; E, B)$ is a (normalized) extreme RN sphere if over S we have

$$\kappa = 1, \quad E = \zeta, \quad B = 0, \quad \Theta = 0, \quad \text{and} \quad K = 0, \tag{10}$$

where κ is the Gaussian curvature, ζ is a unit normal to S in Σ and Θ is the second fundamental form of S in $(\Sigma; g)$.

Normalized extreme RN spheres S are totally geodesic and have $|Q_E(S)| = 1$, $Q_M(S) = 0$ and $A(S) = 4\pi$.

The following lemma discusses the equality case in the general inequality $A \geq 4\pi Q_E^2$ and that was not treated in [12].

Lemma 1. *Let S be a stable (compact, boundaryless and orientable) minimal surface embedded in a maximal electro-vacuum data set and having $A(S) = 4\pi$ and $|Q_E(S)| = 1$. Then, S is a (normalized) extreme RN sphere.*

Proof. Recall from (9) that the stability inequality of the area implies

$$\int_S (|\nabla \alpha|^2 + \kappa \alpha^2) \, dA \geq \int_S \left(|E|^2 + |B|^2 + \frac{|K|^2}{2} + \frac{|\Theta|^2}{2} \right) \alpha^2 \, dA \tag{11}$$

for all $\alpha : S \rightarrow \mathbb{R}$. As $|Q_E(S)| = 1$ we can select the unit normal field ζ to S such that $\frac{1}{4\pi} \int_S \langle E, \zeta \rangle \, dA = |Q_E(S)| = 1$. Choosing $\alpha = 1$ in (11) and using then Gauss–Bonnet and that

$$1 = |Q_E(S)| = \frac{1}{4\pi} \left| \int_S \langle E, \zeta \rangle \, dA \right| \leq \frac{1}{(4\pi)^{1/2}} \left(\int_S |E|^2 \, dA \right)^{1/2} \tag{12}$$

we obtain

$$4\pi \geq 4\pi + \int_S \left(|B|^2 + \frac{|K|^2}{2} + \frac{|\Theta|^2}{2} \right) \, dA$$

This shows that $B = 0$, $K = 0$ and $\Theta = 0$ and that equality must hold. Therefore equality must hold also in (12) which implies (by Cauchy–Schwarz) that $E = \zeta$. It remains to see that $\kappa = 1$, i.e., that S has a round metric. Let us show this below.

Using $B = 0$, $K = 0$, $\Theta = 0$ and $E = \zeta$ in (11) we obtain

$$\int_S (|\nabla \alpha|^2 + (\kappa - 1)\alpha^2) \, dA \geq 0$$

for all functions α . This implies that the first eigenvalue λ of the operator $\alpha \rightarrow -\Delta \alpha + (\kappa - 1)\alpha$ must be non-negative. Denote by α_λ its eigenfunction

(which is unique up to a constant and that is well known to be nowhere zero). Then we have

$$-\Delta\alpha_\lambda + (\kappa - 1)\alpha_\lambda = \lambda\alpha_\lambda \tag{13}$$

Multiplying by $1/\alpha_\lambda$ and integrating over S we obtain

$$-\int_S |\nabla \ln \alpha_\lambda|^2 dA = 4\pi\lambda \geq 0$$

This implies that $\lambda = 0$ and that α_λ is a constant. Using this information in (13) we obtain $\kappa = 1$ as wished. \square

Definition 3. A maximal electro-vacuum data set $(\Sigma; g, K; E, B)$ is said to be of the ERN_1 type if there is a (smooth) diffeomorphism $\varphi : \Sigma_0 \rightarrow \Sigma$ such that

$$\lim_{p \rightarrow \mathcal{E}nd} |(\varphi^*U)(p) - U_0(p)|_{g_0} = 0$$

where (U, U_0) is any of the pairs $(g, g_0), (K, K_0), (E, E_0), (B, B_0)$ and $p \rightarrow \mathcal{E}nd$ means “as p diverges along the cylindrical end or the asymptotically flat end”.

Observe that we require that $(\varphi^*g, \varphi^*K; \varphi^*E, \varphi^*B)$ converges to $(g_0, K_0; E_0, B_0)$ along the ends only in C^0 . For this reason the ADM masses of both data sets are not necessarily equal. However, the total electric and magnetic charges must stay the same as they can be calculated from the formulas (6) along the divergent sequence of spheres $S_{r_i} = \{r = r_i\}$ on the cylindrical end. That is, any data set of the ERN_1 type has total charges $|Q_E| = 1$ and $Q_M = 0$.

The next proposition is essentially a particular case of the results in [12]. We include a proof for a more convenient exposition.

Proposition 1. *Let $(\Sigma; g, K; E, B)$ be a maximal electro-vacuum data set of ERN_1 type. Then every (compact, boundaryless and orientable) embedded surface S which is non-contractible inside Σ has*

$$|Q_E(S)| = 1 \quad \text{and} \quad A(S) \geq 4\pi. \tag{14}$$

Proof. We prove first that $|Q_E(S)| = 1$. Think Σ as $\mathbb{R}^3 \setminus \{o\}$ and S as a surface embedded in it. Then recall that any compact, boundary-less and orientable surface embedded in \mathbb{R}^3 divides \mathbb{R}^3 into two connected components one of which is necessarily unbounded. As S is non-contractible inside $\mathbb{R}^3 \setminus \{o\}$ then the bounded component of $\mathbb{R}^3 \setminus S$ must contain the origin o . That is, S separates the two ends of Σ and the electric charge of S (with an appropriate normal) must be that of the asymptotically flat end, i.e., $|Q_E(S)| = 1$.

We prove now that $A(S) \geq 4\pi$. Assume by contradiction the existence of an S with $A(S) < 4\pi$. Let $\underline{A}(S) = \inf\{A(S'), S' \text{ isotopic to } S\}$. Then obviously we have $4\pi > \underline{A}$. We claim that we also have $\underline{A}(S) > 0$. In fact, if there is a sequence S'_j of surfaces isotopic to S such that $A(S'_j) \rightarrow 0$ then

$$1 = |Q_E(S'_j)| = \frac{1}{4\pi} \left| \int_{S'_j} \langle E, \zeta \rangle dA \right| \leq \frac{1}{4\pi} \|E\|_{L^\infty_g} A(S'_j) \rightarrow 0$$

which would show a contradiction.

Now, following [16] (THEOREM 1')⁷ there is a (non-empty) set of compact boundary-less and non-contractible (inside Σ) minimal surfaces $\{S_1, \dots, S_l\}$ embedded in Σ and a set of positive integers $\{n_1, \dots, n_l\}$ such that

$$\underline{A}(S) = \sum_{i=1}^{i=l} n_i A(S_i)$$

As Σ is diffeomorphic to $\mathbb{R}^3 \setminus \{o\}$ then all the S_i 's must be orientable and therefore stable minimal surfaces [16]. Consider now S_1 and note that $A(S_1) \leq \underline{A}(S) < 4\pi$. We show now that in addition to this it must also be $A(S_1) \geq 4\pi$, which is a contradiction. To show $A(S_1) \geq 4\pi$ we recall (as was shown before) that $|Q_E(S_1)| = 1$. Therefore plugging $\alpha = 1$ in (11) we have

$$4\pi \geq \int_{S_1} |E|^2 dA \geq \frac{1}{A(S_1)} \left(\int_{S_1} |\langle E, \zeta \rangle| dA \right)^2 \geq \frac{(4\pi |Q_E(S_1)|)^2}{A(S_1)} = \frac{(4\pi)^2}{A(S_1)} \tag{15}$$

as wished. □

The following crucial refinement of Proposition 1 shows that equality in the second equation of (14) cannot be achieved. The proof is based in similar arguments to those in [19].

Proposition 2. *Let $(\Sigma; g, K; E, B)$ be a maximal electro-vacuum data set of ERN_1 type. Then every (compact, boundary-less and orientable) embedded surface S which is non-contractible inside Σ has*

$$A(S) > 4\pi.$$

Proof. By Proposition 1 it is enough to show that equality in (14) cannot be achieved. Proceeding by contradiction assume then that there is S_0 with $A(S_0) = 4\pi$. Then observe that if S is isotopic to S_0 then S is also non-contractible inside Σ . Therefore, again by Proposition 1, we have $A(S) \geq 4\pi$ for any surface S isotopic to S_0 . This implies that S_0 is minimal and stable.⁸ By Lemma 1 S_0 is an extreme RN sphere.

Let \mathcal{Y}_0 be a large and strictly convex sphere (w.r.t the outer normal) over the asymptotically flat end. Denote by Ω_0 the region enclosed by it and the

⁷ There is a caveat here. Strictly speaking THEOREM 1' applies to manifolds with convex boundary which is not the case here (instead we have an AF end and a Cylindrical end $\sim \mathbb{R} \times \mathbb{S}^2$). To apply THEOREM 1' one can work between two spheres, one convex and far away in the AF end and another far away on the cylindrical end where in a neighborhood of it one modifies slightly the metric to have also a convex boundary. Apply THEOREM 1' and then show that the minimizer does not intersect the deformed region. The reader can see how this type of argument works when we use as similar one in the proof of Aux-Proposition 3.

⁸ More explicitly, for any smooth $F : [-\varepsilon, \varepsilon] \times S_0 \rightarrow \Sigma$ with $F(0, -) = \text{Id}(-)$ and ε small to have $F(x, -) : S_0 \rightarrow \Sigma$ a smooth embedding, the real function $\lambda \rightarrow A(F(\lambda, S_0))$, (which is greater or equal than 4π for all λ), must have an absolute minimum at $\lambda = 0$. It follows that the first λ -derivative is zero and the second is non-negative. As this is valid for all F then the surface is minimal and stable.

cylindrical end and assume that $S_0 \subset \text{Int}(\Omega_0)$. In what follows we are going to use this region Ω_0 together with a positive solution $N = N_0$ of

$$\Delta N - |E|^2 N = 0 \tag{16}$$

over Ω_0 , asymptotically vanishing over the cylindrical end and not-identical to a constant over S_0 . The existence of such N_0 is proved as follows. Take any two linearly independent smooth positive functions f_1 and f_2 over \mathcal{Y}_0 . For $i = 1, 2$, let \tilde{N}_i be the solution to (16) on Ω_0 with the boundary condition $\tilde{N}_i|_{\mathcal{Y}_0} = f_i$ and asymptotically vanishing over the cylindrical end of Ω_0 . By the maximum principle we have $\tilde{N}_i > 0$ for $i = 1, 2$. If both solutions are constant over S_0 then one can take a linear combination $\tilde{N} := \alpha_1 \tilde{N}_1 + \alpha_2 \tilde{N}_2$ vanishing exactly over S_0 but with $\alpha_1 \neq 0$ and $\alpha_2 \neq 0$. As \tilde{N} asymptotically vanishes over the cylindrical end of Ω_0 and is zero over S_0 then, by the uniqueness of solutions to (16), the combination has to be zero all over the set enclosed by S_0 and the cylindrical end. Then, the unique continuation principle [2] tells that \tilde{N} has to be zero all over Ω_0 which is not possible because f_1 and f_2 were chosen to be linearly independent.

The reason why we take such N_0 is twofold and will be explained adequately during the argumentation below.

In the space-time generated by the initial data consider the future-pointing congruence $\{\gamma(p, \tau)\}$ of time-like geodesics $\gamma(p, \tau)$ starting perpendicularly to Ω_0 at $p \in \Omega_0$ and parametrized by proper time τ . We are going to move Ω_0 with the help of this congruence and obtain a foliation $\{\Omega_t\}$.⁹ The leaves Ω_t of the foliation are defined, for every given t , as the image of the map

$$F_t : p \in \Omega_0 \rightarrow \gamma(p, N_0(p)t) \in \Omega_t$$

This map in turn induces Lapse and Shifts, N_t, X_t over each Ω_t with the property that $N_{t=0} = N_0$ and $X_0 = 0$. Of course the result of moving a point $p \in \Omega_0$ through the space-time vector field $N_\tau \mathbf{n}_\tau + X_\tau$ and for a lapse of time t is the same as $F_t(p)$. The leaves Ω_t are naturally identified to Ω_0 and thus the space-time metric together with the electromagnetic tensor are described by a flow $(g_t, K_t; N_t, X_t; E_t, B_t)$ over Ω_0 (c.f. Sect. 2 item (i); note also that we are changing notation from $(g(t), K(t); N(t), X(t); E(t), B(t))$ to $(g_t, K_t; N_t, X_t; E_t, B_t)$ which makes the writing clearer in this part).

To simplify notation below, when we omit the subindex t we mean $t = 0$.

We can comment now on one of the reasons why we chose N_0 satisfying (16). In general, the time derivative of the mean curvature k_t of the leaves of a space-like foliation $\{\Omega_t\}$ with Lapse N_t and Shift X_t is given by

$$\partial_t k_t - \mathcal{L}_X k_t = -\Delta_{g_t} N_t + (4\pi(\mathbf{T}_{00} + \mathbf{T}_{ij}g_t^{ij}) + |K_t|^2)N_t$$

In our case we have $k_t|_{t=0} = 0$ and thus $\mathcal{L}_X k_{t=0} = 0$. Also at time t equal zero we have $(4\pi(\mathbf{T}_{00} + \mathbf{T}_{ij}g^{ij}) + |K|^2) = |E|^2$ (use $\mathbf{T}_{00} = \mathbf{T}_{ij}g^{ij}$ and $8\pi\mathbf{T}_{00} = |E|^2 + |B|^2$) and hence $\partial_t k_t|_{t=0} = 0$. Thus we obtain $k_t = (\partial_t^2 k_t|_{t=0})t^2/2 + O(t^3)$ in short times. Having this quadratic behavior of k_t in short times was one of the reasons behind the choice of N_0 and will be crucial later.

⁹ Of course is a foliation of a piece of the space-time.

Define $S_t = F_t(S_0) \subset \Omega_t$, the translation of S_0 by F_t . Recall that we are identifying Ω_t to Ω_0 through F_t . In this identification the surface S_t is identified to S_0 . In this sense the area of S_t is the same as $A_{g_t}(S_0)$, a notation that we keep using below.

We claim that

$$\ddot{A}_{g_t}(S_0) \Big|_{t=0} = -A''_{N_0}(S_0) \tag{17}$$

where the double dot means twice the t -derivative of $A_{g_t}(S_0)$ and $A''_{N_0}(S_0)$ is, following the notation introduced before, the second variation of area of S_0 along $N_0\zeta$. We prove this claim in what follows. As was calculated in Proposition 3 in [19] we have

$$\begin{aligned} \ddot{A}_{g_t}(S_0) \Big|_{t=0} &= \int_{S_0} [N_0 \nabla_A \nabla_B N_0 - N_0^2 (Ric_{AB} - 2K_{Ai} K^i_B)] h^{AB} dA \\ &\quad + \int_{S_0} 8\pi N_0^2 [\mathbf{T}_{AB} - \frac{1}{2}(\mathbf{T}_{ij} g^{ij} - \mathbf{T}_{00}) g_{AB}] h^{AB} dA \end{aligned} \tag{18}$$

where we included here the term involving \mathbf{T} that was omitted in [19] as in there only vacuum solutions were considered.¹⁰ In the previous formula Ric is the Ricci curvature of $g = g_0$ and ∇ its covariant derivative. We note then that:

1. The electromagnetic stress-energy is traceless and therefore $\mathbf{T}_{ij} g^{ij} - \mathbf{T}_{00} = 0$,
2. $Ric_{AB} h^{AB} = R - Ric(\zeta, \zeta) = 2|E|^2 - Ric(\zeta, \zeta)$,
3. And finally, because S_0 has the geometry of an extreme RN-horizon the conditions (10) hold and we have

$$\begin{aligned} 8\pi \mathbf{T}_{AB} h^{AB} &= 2|E|^2, \quad K_{Ai} K^i_B h^{AB} = 0, \quad \text{and,} \\ \int_{S_0} N_0 (\nabla_A \nabla_B N_0) h^{AB} dA &= - \int_{S_0} |\nabla N_0|^2 dA \end{aligned}$$

where in the last formula the gradient of N_0 is taken over S_0 .

Combining this information in (18) and after a crucial cancelation of the terms involving $|E|^2$ we obtain

$$\ddot{A}_{g_t}(S_0) \Big|_{t=0} = - \int_{S_0} (|\nabla N_0|^2 - Ric(\zeta, \zeta) N_0^2) dA = -A''_{N_0}(S_0)$$

where to deduce the second equality we have used (8) and that $\Theta = 0$ over S_0 . We can comment now on the second reason for our particular selection of N_0 . If N_0 is not exactly the constant function one over S_0 , as we are assuming, then $A''_{N_0}(S_0) > 0$ and therefore $\ddot{A}_{g_t}(S_0)|_{t=0} < 0$. This is our second reason and will be also crucial below.

¹⁰ More precisely, in the second formula of Proposition 3 use $\dot{K}_{ij} = -\nabla_i \nabla_j N + N(Ric_{ij} - 2K_{il} K^l_j) - 8\pi N(\mathbf{T}_{ij} + \frac{1}{2}(\mathbf{T}_{lm} g^{lm} - \mathbf{T}_{00}) g_{ij})$ instead of just $\dot{K}_{ij} = -\nabla_i \nabla_j N + N(Ric_{ij} - 2K_{il} K^l_j)$ (recall that the data at the initial time is maximal, that is $k = 0$).

The space-time vector field \mathbf{V} which moves Ω_0 to Ω_t and which generates the flow g_t , is, at a space-time point $q = \gamma(p, N_0(p)t)$, given by

$$\mathbf{V}(q) = \frac{dF(p, N_0(p)t)}{dt} = N_0(p) \frac{d\gamma(p, \tau)}{d\tau} \Big|_{\tau=N_0(p)t} = N_0(p)\gamma'(q)$$

Recalling that N_0 tends to zero (indeed exponentially) over the asymptotically cylindrical end of (Ω_0, g_0) we conclude that \mathbf{V} tends to zero over the asymptotically cylindrical end and for this reason the evolution of g_t over the end freezes up. Thus the metrics g_t inherit exactly the same cylindrical asymptotic for every t , that is, that of the metric product of the unit two-sphere and the half-real line.

Take (by continuity) $t^* > 0$ small enough such that for all $t \in [0, t^*]$, the boundary of (Ω_0, g_t) is still strictly convex. Assume that t^* was chosen small enough that $A_{g_t}(S_0) < 4\pi$ for every $t \in [0, t^*]$. Then, again based on general results on minimal surfaces [16] we can guarantee, for every $t \in [0, t^*]$, the existence of a stable minimal sphere¹¹ \hat{S}_t in Ω_0 of area less or equal than $A_{g_t}(S_0)$, non-contractible inside Ω_0 and thus of electric charge one.

We proceed now to gather conveniently all the information obtained so far and use it thereafter to reach a contradiction.

1. From $k_t = (\partial_t^2 k_t|_{t=0})t^2/2 + O(t^3)$ we have, for all $t \in [0, t^*]$ (chose t^* smaller if necessary),

$$k_t^2 \leq 2c_1^2 t^4 \quad \text{where} \quad c_1 = \sup \left\{ \frac{|\partial_t^2 k_t(p)|_{t=0}}{2}, p \in \Omega_0 \right\}, \tag{19}$$

2. From $A_{g_t}(S_0) = 4\pi - A''_{N_0}(S_0)t^2/2 + O(t^3)$ we have, for all $t \in [0, t^*]$ (chose t^* smaller if necessary),

$$A_{g_t}(S_0) \leq 4\pi - \frac{c_2}{2}t^2 \leq 4\pi \quad \text{where} \quad c_2 = \frac{A''_{N_0}(S_0)}{2} > 0 \tag{20}$$

3. For every $t \in [0, t^*]$ there is a stable minimal sphere \hat{S}_t with $Q_E(\hat{S}_t) = 1$ and $A_{g_t}(\hat{S}_t) \leq A_{g_t}(S_0)$.

Now, the stability inequality at \hat{S}_t with trial function $\alpha = 1$ gives

$$4\pi \geq \int_{\hat{S}_t} |E_t|^2 dA_t - \int_{\hat{S}_t} \frac{k_t^2}{2} dA_t$$

Use then (19) and that $\int_{\hat{S}_t} |E_t|^2 dA_t \geq (4\pi)^2/A(\hat{S}_t)$ (because $Q_E(\hat{S}_t) = 1$) to transform this equation into

$$4\pi \geq \frac{(4\pi)^2}{A(\hat{S}_t)} - c_1^2 t^4 A(\hat{S}_t)$$

Multiply this equation by $A(\hat{S}_t)/4\pi$ and then use that $A(\hat{S}_t) \leq A(S_0)$ and (20) to deduce $4\pi - c_2 t^2/2 \geq 4\pi - 4\pi c_1^2 t^4$ or, the same, $8\pi c_2^2 t^4 \geq c_1 t^2$, which is impossible for small t . □

¹¹ That the limit is connected and is a sphere follows from the genus bounds (1.4) of THEOREM 1 in [16].

4. A Family of Perturbations of the ERN₁ Initial Data

Recall that the metric of the ERN₁ space-time is

$$g = -(1 - 1/r)^2 dt^2 + \frac{1}{(1 - 1/r)^2} dr^2 + r^2 d\Omega^2 \tag{21}$$

and that on the hypersurface $\Sigma_0 = \{t = 0\}$ we have $K_0 = 0$, $B_0 = 0$ and that the electric field is radial and takes the form $E_0 = \zeta/r^2$ where $\zeta = \partial_r/|\partial_r|$ is the unit normal to the radial spheres $S_{\bar{r}} = \{r = \bar{r}\}$. Now, the constraint equations (5) of an electro-vacuum data set $(g, K; E, B)$ with $K = 0$ and $B = 0$ reduce to

$$\begin{cases} R = 2|E|^2, \\ \operatorname{div} E = 0 \end{cases} \tag{22}$$

Because of this the scalar curvature R_0 of the metric g_0 of the ERN₁ standard initial data is $R_0 = 2/r^4$.

In the argumentation given below we will make use of an expression for the three-Laplacian Δ_{g_0} acting on radial functions $\phi = \phi(r)$ of Σ_0 . A direct calculation using the general formula $\Delta\phi = \partial_r(\sqrt{g}g^{rr}\partial_r\phi)/\sqrt{g}$ gives, when $\phi = \phi(r)$, the expression

$$\Delta_{g_0} \phi = \frac{r(r-1)}{r^4} \frac{d}{dr} \left[r(r-1) \frac{d}{dr} \phi \right]$$

This formula is simplified if we use the harmonic radial coordinate $x = \ln(1 - 1/r)$ instead of r (harmonic means $\Delta_{g_0} x = 0$). With this definition the range of x is $(-\infty, 0)$. In this new coordinate the Laplacian acting on radial functions reads

$$\Delta_{g_0} \phi = \frac{\phi''}{r^4}, \tag{23}$$

where here $\phi'' = d^2\phi/dx^2$. Note then that $\Delta_{g_0} \phi = |E_0|^2 \phi''$.

We proceed now to construct the bi-parametric family of axisymmetric “perturbations” of the initial data on Σ_0 . The axisymmetric Killing field will be ∂_φ , which, note, is also axisymmetric Killing for the background data set. The two parameters of the family will be $\hat{\epsilon}$ and \hat{x} . Roughly speaking the variable $\hat{\epsilon}$ represents the “strength” of the perturbation while \hat{x} marks the sphere around which the perturbation “concentrates”. This interpretation will be clear as the construction progresses. To explain the construction let us recall in what follows the *conformal method* to solve the constraint equations but for the situation that is of interest here, namely when the data set to be found is time symmetric and has no magnetic field. Let (Σ, g) be a Riemannian manifold of scalar curvature R . On it let \hat{E} be a g -divergence-less vector field. If for $\phi > 0$ we have

$$\Delta \phi = R\phi - 2|\hat{E}|^2 \phi^{-3}, \tag{24}$$

then $\bar{g} = \phi^4 g$ and $\bar{E} = \phi^{-6} \hat{E}$ satisfy the constrain equations (22). We will use this method below with $(\Sigma, g) = (\Sigma_0, g_0)$ and $\hat{E} = E_{\hat{\epsilon}, \hat{x}}$ suitably chosen.

In what follows we will identify Σ_0 to $(-\infty, 0] \times \mathbb{S}^2$ where the factor $(-\infty, 0]$ is the range of the coordinated x introduced before. From now on the parameter \hat{x} is set to vary in $(-\infty, -2]$ and $\hat{\epsilon}$ in $(0, 1/16)$. Fix a smooth and non-zero axisymmetric two-form ω supported on $(-3, -1) \times \mathbb{S}^2 \subset (-\infty, -1) \times \mathbb{S}^2$. This form is set to be fixed from now on and will not be adjusted anymore. For every \hat{x} let $\chi_{\hat{x}}^* \omega$ be the pull-back of ω to $[\hat{x} - 1, \hat{x} + 1] \times \mathbb{S}^2$ under the transformation $\chi_{\hat{x}} : [\hat{x} - 1, \hat{x} + 1] \times S^2 \rightarrow [-3, -1] \times S^2$ given by $(x, \theta, \varphi) \rightarrow (x - \hat{x} - 2, \theta, \varphi)$. Then, for every \hat{x} and $\hat{\epsilon}$ define

$$\hat{E}_{\hat{x}, \hat{\epsilon}} = E_0 + \hat{\lambda} (\star d \star (\chi_{\hat{x}}^* \omega))^\# \tag{25}$$

where $\hat{\lambda} = \hat{\lambda}_{\hat{x}, \hat{\epsilon}}$ is a factor chosen to have $\hat{\epsilon} = \sup |1 - |\hat{E}_{\hat{x}, \hat{\epsilon}}|^2 / |E_0|^2|$ (here $|\dots| = |\dots|_{g_0}$), the star \star in $\star d \star$ is the g_0 -Hodge star and $(\star d \star (\chi_{\hat{x}}^* \omega))^\#$ is the g_0 -dual vector field of the form $\star d \star (\chi_{\hat{x}}^* \omega)$. In this way $\hat{E}_{\hat{x}, \hat{\epsilon}}$ comprises a bi parametric family of divergence-less axisymmetric vector fields which are equal to the background field E_0 outside $[\hat{x} - 1, \hat{x} + 1] \times \mathbb{S}^2$ but otherwise not very different from it.

In what follows and to simplify notation we keep using $|\dots| = |\dots|_{g_0}$ and make also $\hat{E} = \hat{E}_{\hat{x}, \hat{\epsilon}}$.

We pass now to show that for every \hat{E} we can find an axisymmetric solution to the Lichnerowitz equation (24) (L-equation from now on) with good geometric properties. To this extent we use the method of sub and super-solutions. Namely, if for axisymmetric functions (barriers) $\phi_+ > 0$ and $\phi_- > 0$ with $\phi_+ > \phi_-$ we have

$$\begin{cases} \Delta_{g_0} \phi_+ \leq 2|E_0|^2 \phi_+ - 2|\hat{E}|^2 \phi_+^{-3}, \\ \Delta_{g_0} \phi_- \geq 2|E_0|^2 \phi_- - 2|\hat{E}|^2 \phi_-^{-3} \end{cases} \tag{26}$$

(recall $R_0 = 2|E_0|^2$) then there is an axisymmetric solution $\phi > 0$ to (24) with $\phi_- \leq \phi \leq \phi_+$, (for a proof of this fact in this context see [8]¹²). We explain now how to find ϕ_- which will be a radial function, i.e., $\phi_- = \phi_-(x)$. In (I) below we define $\phi_-(x)$ over $(-\infty, -1]$ and in (II) over $[-1, 0)$. The global function defined by (I) and (II) will be smooth over the separate domains $(-\infty, -1)$ and $(-1, 0)$ but will be just C^0 at $x = -1$. For this reason, to check that such global function is a barrier in the distributional sense [8] it will be necessary to check that its left derivative at $x = -1$ is less than its right derivative.¹³ This will be done after (I) and (II) below.

(I) *Defining $\phi_-(x)$ on $(-\infty, -1]$.* Make $\psi_- = \phi_- - 1$ and recall that $\Delta \psi_- = |E_0|^2 \psi_-''$. With this information and after a simple manipulation the second equation in (26) can be displayed in the form

$$\psi_-'' \geq 2 [1 + \phi_-^{-1} + \phi_-^{-2} + \phi_-^{-3}] \psi_- + 2 \left[1 - \frac{|\hat{E}|^2}{|E_0|^2} \right] \phi_-^{-3} \tag{27}$$

¹² To get an axisymmetric solution out of the method of barriers just work inside the family of axisymmetric functions all the time in [8].

¹³ Alternatively, a smooth barrier can be easily found by rounding off the global function constructed by (I) and (II).

Now, it can be easily checked that for any real number γ such that $|\gamma - 1| \leq 1/8$ we have

$$3 \leq 1 + \gamma^{-1} + \gamma^{-2} + \gamma^{-3} \leq 5, \quad \text{and} \quad \frac{1}{2} \leq \gamma^{-3} \leq 2 \tag{28}$$

Thus, if we can find $\psi_-(x)$ with $-1/8 < \psi_- < 0$ and satisfying

$$\psi_-'' \geq 6\psi_- + 4\hat{E}I \tag{29}$$

where $\hat{I} = \hat{I}(x)$ is the *indicator function* on $[\hat{x} - 1, \hat{x} + 1]$, (i.e., equal to one on $[\hat{x} - 1, \hat{x} + 1]$ and zero otherwise), then $\phi_- = 1 + \psi_-$ will verify (27) because, in this case, we would have

$$6\psi_- + 4\hat{E}I \geq 2 [1 + \phi_-^{-1} + \phi_-^{-2} + \phi_-^{-3}] \psi_- + 2 \left[1 - \frac{|\hat{E}|^2}{|E_0|^2} \right] \phi_-^{-3}$$

due to (28) (with $\gamma = \phi_-$) and because, by construction, we have $1 - |\hat{E}|^2/|E_0|^2 \leq \hat{\epsilon}$ point-wise. The function

$$\psi_-(x) = \frac{-4\hat{\epsilon}}{\cosh(x - \hat{x})}$$

verifies $-1/8 < \psi_-(x) < 0$ because $\hat{\epsilon} < 1/16$. To see that it also satisfies (29) on $(-\infty, -1]$ we argue as follows. First we compute $\psi_-'' = 4\hat{\epsilon}(1 - 2 \sinh^2(x - \hat{x})/\cosh^2(x - \hat{x}))/\cosh(x - \hat{x})$ and, after plugging this inside (29) and after a simple manipulation we conclude that to verify (29) it is enough to verify the inequality $7 - 2 \sinh^2(x - \hat{x})/\cosh^2(x - \hat{x}) \geq (\cosh(x - \hat{x})) \hat{I}(x)$ for all $x \in (-\infty, -1]$. This is easily seen because the l.h.s of this expression is greater than five and the r.h.s is less or equal than $\cosh 1$ which is less than e . Summarizing, $\phi_- = \psi_- + 1$ is a sub-solution on this range of x . Note that as $\psi_- < 0$ then it is $\phi_- < 1$.

(II) *Defining $\phi_-(x)$ on $[-1, 0)$.* On $[-1, 0)$ define $\phi_-(x)$ by $\phi_-(x) = 1 + \psi_-(x)$ where

$$\psi_-(x) = \frac{4\hat{\epsilon}x}{\cosh(-1 - \hat{x})}$$

To see that ϕ_- is a sub-solution it is necessary to check (27). Firstly, as $\phi_-(x)$ is linear in x the l.h.s of (27) is zero. Secondly, the second term on the r.h.s of (27) is zero because when $x \in [-1, 0)$ it is $|E_0|^2 = |\hat{E}|^2$. The inequality (27) then follows because $\psi_- < 0$ and so is the first term on the r.h.s of (27).

So far we have defined ϕ_- and proved that it is a sub-solution when restricted to the intervals $(-\infty, -1)$ and $(-1, 0)$. It remains to prove that it is also a sub-solution in the neighborhood of $x = -1$. As said, to see this it is enough to check that the left-sided derivative of ϕ_- at $x = -1$ is less than its right-sided derivative. The left-sided derivative at $x = -1$ is $4\hat{\epsilon} \sinh(-1 - \hat{x})/\cosh^2(-1 - \hat{x})$ while the right-sided is $4\hat{\epsilon}/\cosh(-1 - \hat{x})$ and the desired inequality follows.




FIGURE 4. Picture of the barriers ϕ_- and ϕ_+

Summarizing, the sub-solution is

$$\phi_-(x) = \begin{cases} 1 - \frac{4\hat{\epsilon}}{\cosh(x - \hat{x})} & \text{if } x \in (-\infty, -1], \\ 1 + \frac{4\hat{\epsilon}x}{\cosh(-1 - \hat{x})} & \text{if } x \in [-1, 0) \end{cases} \tag{30}$$

A graph of ϕ_- is presented in Fig. 4. Reproducing the argument that lead to ϕ_- , it is found that $\phi_+(x)$, defined by

$$\phi_+(x) = \begin{cases} 1 + \frac{4\hat{\epsilon}}{\cosh(x - \hat{x})} & \text{if } x \in (-\infty, -1], \\ 1 - \frac{4\hat{\epsilon}x}{\cosh(-1 - \hat{x})} & \text{if } x \in [-1, 0) \end{cases} \tag{31}$$

is a super-solution. We conclude that there is $\phi > 0$, solution of (24), and satisfying $\phi_- \leq \phi \leq \phi_+$. The metric $\bar{g} = \phi^2 g_0$ and the electric field $\bar{E} = \hat{E}\phi^{-6}$ satisfy the constraint equations (22).

Summarizing, from the explicit form of the sub and super-solutions we observe that $\phi - 1$ “concentrates“ around \hat{x} and decays exponentially to zero in both directions of x starting from \hat{x} . In the direction of increasing x the exponential decay however stops at $x = -1$ and after that it is linear in x , namely of the order $1/r$ in the r -coordinate. Observe, to be recalled later, that the exponential decay of ϕ in the asymptotically cylindrical end implies by standard elliptic estimates that the perturbed data sets $(\bar{g}, \bar{K}; \bar{E}, \bar{B})$ decay exponentially as defined in the introduction.

5. Rigidity of the ERNT₁ Initial Data

The next lemma shows the rigidity of the ERNT space-time and has interest in itself. It will be used in the proof of Proposition 4.

Lemma 2. *Let $(\Sigma; g, K; E, B)$ be a smooth complete and maximal electrovacuum data set where Σ is diffeomorphic to $\mathbb{R} \times \mathbb{S}^2$. Let $S_0 := 0 \times \mathbb{S}^2$ and suppose that $|Q_E(S_0)| = 1$. Suppose too that for any (compact, boundaryless and embedded) surface S non-contractible inside Σ we have $A(S) \geq 4\pi$, and that there is at least one such S with $A(S) = 4\pi$. Then the data set is the standard ERNT₁ initial data.*




FIGURE 5. Representation of the manifolds $\mathcal{V}_1, \mathcal{V}_2, \mathcal{W}_1$ and \mathcal{W}_2

For expository reasons it is better to divide the proof into three Auxiliary Propositions. In every one of them we let \mathcal{F} be the set of (compact, boundaryless and embedded) surfaces of area 4π and which are non-contractible inside Σ .

Aux-Proposition 1. *Assume the hypothesis of Lemma 2. Then, each $S \in \mathcal{F}$ is a (normalized) ERN sphere and every two different spheres in \mathcal{F} are disjoint. Moreover, the set $\bigcup_{S \in \mathcal{F}} \{S\}$ is closed as a set in Σ .*

Aux-Proposition 2. *Assume the hypothesis of Lemma 2. If $\bigcup_{S \in \mathcal{F}} \{S\} = \Sigma$ then the data set is the standard ERNT₁ initial data $(\tilde{\Sigma}; \check{g}_0, \check{K}_0; \check{E}_0, \check{B}_0)$.*

Aux-Proposition 3. *Assume the hypothesis of Lemma 2. Then, $\bigcup_{S \in \mathcal{F}} \{S\} = \Sigma$.*

The proofs of the three propositions are presented consecutively.

Proof of Aux-Proposition 1. By the hypothesis of Lemma 2 every non-contractible surface has area greater or equal than 4π . Therefore the surfaces in \mathcal{F} , which have area equal to 4π , must be minimal and stable (see footnote 8). By Lemma 1 they are (normalized) ERN spheres. We show next that two different spheres S_1 and S_2 in \mathcal{F} (in case \mathcal{F} has more than one element) must be disjoint. If $S_1 \cap S_2 \neq \emptyset$ then, being minimal surfaces, they must intersect transversely. We will think the surfaces $S_i, i = 1, 2$ as embedded in $(\mathbb{R}^3 \setminus \{o\}) \sim \Sigma$. As the $S_i, i = 1, 2$ are non-contractible inside $\mathbb{R}^3 \setminus \{o\}$ then there are open balls B_1 and B_2 in \mathbb{R}^3 containing the origin o and such that $\partial \bar{B}_i = S_i$ for $i = 1, 2$. Define the manifolds

$$\mathcal{V}_1 := S_1 \cap \text{Int}(B_2^c), \quad \mathcal{V}_2 := S_2 \cap \text{Int}(B_1^c) \quad \text{and} \quad \mathcal{W}_1 := S_1 \cap B_2, \quad \mathcal{W}_2 := S_2 \cap B_1$$

where $\text{Int}(B_i^c)$ is the interior of the complement of B_i (see Fig. 5). The manifolds $\mathcal{V}_1, \mathcal{V}_2, \mathcal{W}_1$ and \mathcal{W}_2 are pairwise disjoint and their closures have the same boundary. We will denote such boundary (a union of embedded circles indeed) by \mathcal{B} . We have $S_1 = \bar{\mathcal{V}}_1 \cup \bar{\mathcal{W}}_1$ and $S_2 = \bar{\mathcal{V}}_2 \cup \bar{\mathcal{W}}_2$ and for this reason it is

$$4\pi = A(\bar{\mathcal{V}}_1) + A(\bar{\mathcal{W}}_1) \quad \text{and} \quad 4\pi = A(\bar{\mathcal{V}}_2) + A(\bar{\mathcal{W}}_2). \tag{32}$$

The manifolds

$$\mathcal{V} := \overline{\mathcal{V}}_1 \cup \overline{\mathcal{V}}_2 \quad \text{and} \quad \mathcal{W} := \overline{\mathcal{W}}_1 \cup \overline{\mathcal{W}}_2,$$

are embedded and smooth except at \mathcal{B} , where they have necessarily corners. Note that \mathcal{V} and \mathcal{W} are not necessarily connected (see Fig. 5). Moreover, we have $\mathcal{V} = \partial(\overline{B_1 \cup B_2})$ and $\mathcal{W} = \partial(\overline{B_1 \cap B_2})$. Therefore, as $o \in B_1 \cup B_2$ and $o \in B_1 \cap B_2$, then at least one of the connected component of \mathcal{V} and at least one of \mathcal{W} divide $\mathbb{R}^3 \setminus \{o\}$ into two connected components and are consequently non-contractible inside Σ . By (32) if $A(\mathcal{V}_2) \leq A(\mathcal{W}_1)$ then $A(\mathcal{V}) \leq 4\pi$, while if $A(\mathcal{V}_2) \geq A(\mathcal{W}_1)$ then $A(\mathcal{W}) \leq 4\pi$. In any case, we can round off the corners at \mathcal{B} of either the manifold \mathcal{V} or the manifold \mathcal{W} to obtain one of area less than 4π and having at least one connected component non-contractible inside Σ . This is against hypothesis and therefore the surfaces S_1 and S_2 have to be disjoint.

It remains to be proved that the set $\bigcup_{S \in \mathcal{F}} \{S\}$ is closed in Σ . But if $p_i (\in S_i \in \mathcal{F})$ is a sequence of points in $\bigcup_{S \in \mathcal{F}} \{S\}$ with limit point p_∞ , then the sequence S_i of (normalized) ERN spheres, and therefore of stable and area minimizing minimal surfaces, has a subsequence converging (in C^k for every $k \geq 1$) to a limit stable minimal sphere $S_\infty \ni p_\infty$, [18].¹⁴ The sphere S_∞ cannot be contractible inside Σ , otherwise the S_i s would be contractible for sufficiently big i . We have $4\pi = \lim A(S_i) = A(S_\infty)$, thus $S_\infty \in \mathcal{F}$ and therefore $p_\infty \in \bigcup_{S \in \mathcal{F}} \{S\}$. \square

Proof of Aux-Proposition 2. Assume at the moment that the foliation \mathcal{F} is smooth (see the definition of smooth foliation in [4]). We will be proving this later. Fix a sphere S^* in \mathcal{F} and denote by Σ_L^* and Σ_R^* the connected components of $\Sigma \setminus S^*$. For any $p \in \Sigma$ let $S(p)$ be the sphere in \mathcal{F} containing p and denote by $\Omega(p)$ the region enclosed by S^* and $S(p)$. Then define the (smooth) function $\tilde{x} : \Sigma \rightarrow \mathbb{R}$ as

$$\tilde{x}(p) = \begin{cases} \text{Vol}(\Omega(p)) & \text{if } p \in \Sigma_R^*, \\ -\text{Vol}(\Omega(p)) & \text{if } p \in \Sigma_L^* \end{cases}$$

This function is constant over every leaf and has nowhere zero gradient.¹⁵ Let $Y = \nabla \tilde{x} / |\nabla \tilde{x}|^2$ and note that as $Y(\tilde{x}) = 1$ the flow induced by Y carries leaves (of \mathcal{F}) into leaves (of \mathcal{F}). Fix an isometry $\psi : \mathbb{S}^2 \rightarrow S^*$ and define the diffeomorphism $\Phi : \mathbb{R} \times \mathbb{S}^2 \rightarrow \Sigma$ by sending a pair (t, s) into the translation of $\psi(s)$ through the flow induced by Y and by a parametric time t . Of course we have $\Phi_* \partial t = Y$. On the other hand, if we denote by $h_{\tilde{x}}$ the induced metric on the leaves, then we have $\mathcal{L}_Y h_{\tilde{x}} = 0$ because each leaf is totally geodesic (here \mathcal{L} is the Lie-derivative). Therefore we can write

$$\Phi^* g = |\nabla \tilde{x}|^2 d\tilde{x}^2 + d\Omega^2$$

¹⁴ Precisely there are embeddings $f_i : \mathbb{S}^2 \rightarrow S_i$ converging in C^k to a covering immersion $f_\infty : \mathbb{S}^2 \rightarrow S_\infty$. But in our case $\Sigma \sim \mathbb{R}^3 \setminus \{o\}$ and therefore S_∞ must be orientable, hence a sphere and f_∞ an embedding.

¹⁵ This can be easily seen from the fact that \mathcal{F} is assumed smooth.

We show now that $|\nabla\bar{x}|$ is constant over every leaf. Indeed, as the areas of the spheres of \mathcal{F} is 4π then the second variation of area of any sphere in \mathcal{F} along Y is zero, i.e., $A''_Y(S) = 0$. This implies that $|Y| = 1/|\nabla\bar{x}|$ is constant over every sphere (see the proof of Lemma 1). The metric (3) is recovered by making a simple change of variables $\bar{x} = \bar{x}(\hat{x})$, with $|\nabla\bar{x}| = d\bar{x}/d\hat{x}$. Finally by Lemma 1 we have $B = 0, K = 0$ and $E = \zeta$ with ζ a normal field to the leaves of \mathcal{F} (i.e., either $\partial_{\bar{x}}$ or $-\partial_{\bar{x}}$). Hence we have $(g, K; E, B) = (\hat{g}, \hat{K}; \hat{E}, \hat{B})$ as claimed.

It remains to prove that the foliation \mathcal{F} is smooth. We will show that the 1-distribution of lines perpendicular to the leaves of \mathcal{F} is smooth. This implies that the distribution of the tangent planes to the leaves of \mathcal{F} is smooth and the smoothness of \mathcal{F} is then direct from Frobenius’s theorem [4]. Let S be a sphere of \mathcal{F} , let ζ be a normal field to it and let h be the induced two-metric. We will show that the Ricci curvature Ric of g over S has the following form: $Ric(\zeta, \zeta) = 0$ and for any $v, w \in TS$ we have $Ric(\zeta, v) = 0$ and $Ric(v, w) = h(v, w)$. The 1-distribution of normal directions to \mathcal{F} is then uniquely characterized by the null space of Ric (i.e., $\{v \in TS, Ric(v, v) = 0\}$), and is easily seen to be smooth because Ric is smooth.

Again let S be a surface in \mathcal{F} and ζ a unit normal field to it. Let $\{\gamma_q(\tau), q \in S, 0 \leq \tau \leq \tau_0\}$ be the congruence of geodesics in Σ starting at $\tau = 0$ perpendicularly to S in the direction of ζ and parametrized by the arc-length τ . We will move S by the vector field $V = \partial_\tau \gamma_q(\tau)$ and obtain a smooth one-parametric family of surfaces $S(\tau)$. We assume that τ_0 is small enough that the surfaces $S(\tau)$ are embedded (and smooth).

In the forthcoming equations, but inside this proof, we will denote the mean curvature $\text{tr}_h \Theta$ by μ . Recall from Lemma 1 that over S we have $\kappa = 1, R = 2$ and $\Theta = 0$. Therefore from the general identity

$$2\kappa - |\Theta|^2 + \mu^2 = R - 2Ric(\zeta, \zeta) \tag{33}$$

we obtain $Ric(\zeta, \zeta) = 0$. Also from $\text{div } \Theta - d\mu = Ric(\zeta, -)$ we obtain $Ric(\zeta, v) = 0$ for any $v \in TS$. To show that for any $v, w \in TS$ we have $Ric(v, w) = h(v, w)$ it is enough to prove that $\mathcal{L}_V \Theta = \dot{\Theta} = 0$ because of the general identity (on TS)

$$\dot{\Theta} = -\mu\Theta + 2\Theta \circ \Theta + \kappa h - Ric$$

which gives $\dot{\Theta}(0) = h - Ric$ at $\tau = 0$. Now, at any time $\tau \in (0, \tau_0)$ we have

$$\begin{aligned} \ddot{A}(S(\tau)) &= \int_{S(\tau)} (\dot{\mu} + \mu^2) dA \\ &= \int_{S(\tau)} \left(-\frac{|\Theta|^2}{2} + \frac{\mu^2}{2} + \kappa - |E|^2 \right) dA \\ &= \int_{S(\tau)} \left(-\frac{|\Theta|^2}{2} + \frac{\mu^2}{2} \right) dA + \left[4\pi - \int_{S(\tau)} |E|^2 dA \right] \\ &\leq \underbrace{\int_{S(\tau)} \left(-\frac{|\Theta|^2}{2} + \frac{\mu^2}{2} \right) dA}_{U(\tau)} \end{aligned} \tag{34}$$

where: (i) to obtain the first inequality we use $d\dot{A} = \mu dA$, (ii) to pass from the second to the third line we use the focussing (Riccati) equation $\dot{\mu} = -|\Theta|^2 - Ric(\zeta, \zeta)$ in conjunction with (33) and $R \geq 2|E|^2$, (iii) to pass from the second to the third line we use Gauss–Bonnet and (iv) from the third to the fourth we use (12). On the other hand, we can express $A(S(\tau))$ as

$$A(S(\tau)) = 4\pi + \int_0^\tau R\tilde{\tau} \int_0^{\tilde{\tau}} \ddot{A}(S(\tilde{\tau})) d\tilde{\tau}$$

and we have $\ddot{A}(S(\tau)) \leq U(\tau) = U(0) + U'(0)\tau + U''(0)\tau^2/2 + O(\tau^3)$ with $U(0) = 0, U'(0) = 0$ and

$$U''(0) = -\frac{1}{2} \int_{S(0)} |\dot{\Theta}(0)|^2 dA$$

as can be easily seen using $\mu(0) = 0, \dot{\mu}(0) = 0, \Theta(0) = 0$ and $\dot{\Theta}(0) = h - Ric$. Therefore, if $\dot{\Theta}(0) \neq 0$ then we would have $A(S(\tau)) < 4\pi$ for small τ which is against the hypothesis. This finishes the proof. \square

Proof of Aux-Proposition 3. We will proceed by contradiction and assume that $\bigcup_{S \in \mathcal{F}} \{S\} \neq \Sigma$. As by Aux-Proposition 1 the set $\bigcup_{S \in \mathcal{F}} \{S\}$ is closed, then every connected component of $\Sigma \setminus \bigcup_{S \in \mathcal{F}} \{S\}$ is either an open region enclosed by two spheres in \mathcal{F} or an open region enclosed by a sphere in \mathcal{F} and one of the two ends of Σ . Thus, if there is only one connected component of $\Sigma \setminus \bigcup_{S \in \mathcal{F}} \{S\}$ then $\bigcup_{S \in \mathcal{F}} \{S\}$ must at least contain a closed region enclosed by a sphere in \mathcal{F} and one end of Σ . As in Aux-Proposition 2 the data set over such region must be $ERNT_1$. Because of this one can cut out such region and “double” the remaining one to construct a new data set $(\Sigma'; g', K'; E', B')$ in the hypothesis of Lemma 2 but with two connected components of $\Sigma' \setminus \bigcup_{S \in \mathcal{F}} \{S\}$.

Assume then without loss of generality that there are at least two connected components of $\Sigma \setminus \bigcup_{S \in \mathcal{F}} \{S\}$. We want to prove that such data set cannot exist. This will be done exactly as in Proposition 2. For this reason the paragraphs below are first dedicated to construct a setup similar to the one in the proof of Proposition 2.

For the discussion that follows the Fig. 6 could be of great help. Denote two of the connected components of $\bigcup_{S \in \mathcal{F}} \{S\}$ by Ω_L and Ω_R (L for “Left” and R for “Right”). Let S_L and S_R be any two spheres embedded in Ω_L and Ω_R , respectively, and non-contractible inside Σ . Denote by Ω_{LR} the region enclosed by them and including them, and by Σ_L^- (resp. Σ_R^+) the connected component of $\Sigma \setminus S_L$ (resp. $\Sigma \setminus S_R$) not containing S_R (resp. S_L). Also let $D^* > 0$ be small enough such that

1. if $p \in \Sigma_L^-$ (resp. $p \in \Sigma_R^+$) and $\text{dist}(p, S_L) \leq D^*$ (resp. $\text{dist}(p, S_R) \leq D^*$) then $p \in \Omega_L$ (resp. $p \in \Omega_R$), and
2. for any $0 < D \leq D^*$ the set $\{p \in \Sigma_L^-, \text{dist}(p, S_L) = D\}$ (resp. $\{p \in \Sigma_R^+, \text{dist}(p, S_R) = D\}$) is a smooth and embedded sphere.

In this context define the sphere S_L^* (resp. S_R^*) as $S_L^* := \{p \in \Sigma_L^-, \text{dist}(p, S_L) = D^*\}$ (resp. $S_R^* := \{p \in \Sigma_R^+, \text{dist}(p, S_R) = D^*\}$) and let Ω_{LR}^* be the set enclosed by S_L^* and S_R^* including them. As the components Ω_L and Ω_R are different




FIGURE 6. Representation of the construction in the proof of Aux-Proposition 3

there is at least one sphere $S_0 \in \mathcal{F}$ embedded in Ω_{LR} and therefore in Ω_{LR}^* . Now, on Ω_{LR}^* consider a positive solution $N = N_0$ of the maximal lapse equation

$$\Delta N - (4\pi(\mathbf{T}_{00} + \mathbf{T}_{ij}g^{ij}) + |K|^2)N = \Delta N - (|E|^2 + |B|^2 + |K|^2)N = 0$$

and that is not identically to a constant over S_0 . The existence of such N_0 is shown in the same way as was done in Proposition 2 and is left to the reader. Also in the same way as in Proposition 2 construct from N_0 a time-like vector field \mathbf{V} and from it a flow $(g_t, K_t; E_t, B_t)$ over Ω_{LR}^* , with $0 \leq t \leq t^*$ and for some t^* small. As in Proposition 2 now we have $\dot{A}_{g_t}(S_0) = 0$ and $\ddot{A}_{g_t}(S_0) < 0$. Therefore $A_{g_t}(S_0) < 4\pi$ in short times t .

Instead of g_t we are going to consider a modified flow of metrics \tilde{g}_t conformally related to g_t . This will help to guarantee the existence of certain stable minimal spheres. To the purpose of defining \tilde{g}_t consider the following function of $z \in [0, D^*]$,

$$\Psi_\delta(z) = 1 + e^{-1/z} + 1/(D^* + \delta - z)$$

where δ is a constant to be fixed soon below. Observe that $\Psi_\delta(0) = 1$ and that all the right-sided derivatives of Ψ_δ are zero at $z = 0$. Observe too that $\Psi_\delta \geq 1$. We then define \tilde{g}_t by

$$\tilde{g}_t(p) = \begin{cases} g_t(p) & \text{if } p \in \Omega_{LR}, \\ \Psi_\delta(d(p, S_L)) g_t(p) & \text{if } p \in \Omega_{LR}^* \cap \Sigma_L^-, \\ \Psi_\delta(d(p, S_R)) g_t(p) & \text{if } p \in \Omega_{LR}^* \cap \Sigma_R^+ \end{cases}$$

Now chose t^* and $\delta > 0$ small enough that the boundaries of $(\Omega_{LR}^*, \tilde{g}_t)$ are strictly mean convex (in the outgoing directions) for any $0 \leq t \leq t^*$. Once this is granted we can consider for every $t \leq t^*$ a sphere \tilde{S}_t minimizing the \tilde{g}_t -area among all the spheres embedded in Ω_{LR}^* and isotopic to S_0 [16].

Until now we have not done any particular progress in the proof. The key point of the proof lies in showing that one chose t^* smaller if necessary in such a way that the area minimizing spheres \tilde{S}_t are embedded in $\text{Int}(\Omega_{LR})$ and therefore do not intersect the regions where the metric g_t was conformally

modified. Once this is shown a contradiction is proved following exactly the same argument as in Proposition 2 and will not be repeated here.

Suppose then that there is a sequence of times $t_i \downarrow 0$ such that for each t_i the minimal and stables sphere \tilde{S}_{t_i} is not strictly embedded in $\text{Int}(\Omega_{LR})$. Take then a subsequence (indexed again by “i”) such that \tilde{S}_{t_i} converges to a stable minimal sphere \tilde{S}_0 intersecting $\Omega_{LR}^* \setminus \text{Int}(\Omega_{LR})$. As the \tilde{S}_{t_i} are non-contractible inside Σ then neither is \tilde{S}_0 . Moreover, as $A_{\tilde{g}_{t_i}}(\tilde{S}_{t_i}) \leq A_{\tilde{g}_{t_i}}(S_0) < 4\pi$ then $A_{\tilde{g}_{t_i}}(\tilde{S}_0) \leq 4\pi$. But $A_{\tilde{g}_0}(\tilde{S}_0) \geq A_g(\tilde{S}_0)$ because the conformal factor is greater or equal than one. Then, the sphere $\tilde{S}_0 \subset \Sigma$ has $A(\tilde{S}_0) \leq 4\pi$. So it must be $A(\tilde{S}_0) = 4\pi$ by the hypothesis of Lemma 2 and by Lemma 1 it must be a (normalized) ERN sphere. Thus $\tilde{S}_0 \in \mathcal{F}$. But this is a contradiction as the set $\Omega_{LR}^* \setminus \text{Int}(\Omega_{LR})$ does not contain any point of $\bigcup_{S \in \mathcal{F}} \{S\}$. \square

6. Perturbations Containing MOTS

The following proposition is direct from standard elliptic estimates and is left to the reader (use $\varphi = \text{id}$ and recall that $\tilde{g} = \phi^2 g_0$, $\bar{K} = 0$, $\bar{E} = \hat{E}_0 \phi^{-6}$ and $\bar{B} = 0$). It says that the data sets constructed in Sect. 4 are small in the sense of Definition 1.

Proposition 3. *Given $0 < \hat{\epsilon} < 1/16$ and integer $k \geq 1$ there is $\varepsilon = \varepsilon(\hat{\epsilon}, k) > 0$ such that for any $\hat{x} \in (-\infty, -1]$ the data set $(\tilde{g}, \bar{K}; \bar{E}, \bar{B})$ constructed in Sect. 4 out of $\hat{\epsilon}$ and \hat{x} , is ε -close in C^k to the standard ERN_1 initial data. Moreover, $\varepsilon \rightarrow 0$ if we fix k and let $\hat{\epsilon} \rightarrow 0$.*

In what follows we explain a *pointed convergence* that will be useful inside the proof of Proposition 4. We keep identifying Σ_0 to $(-\infty, 0) \times \mathbb{S}^2$ as we did before, in particular the factor $(-\infty, 0)$ is the range of x . Let \hat{x}_i be a sequence diverging to minus infinity, i.e., $\lim \hat{x}_i = -\infty$ and let s_0 be a fixed point in \mathbb{S}^2 . If we “follow” the ERN_1 metric g_0 around the sequence of points $\hat{x}_i \times s_0$ then, as we know, it converges to the metric \check{g}_0 of the standard ERN_1 initial data. The standard mathematical way of saying this is that the pointed sequence $(\Sigma_0; g_0; \hat{x}_i \times s_0)$ converges smoothly to $(\mathbb{R} \times \mathbb{S}^2; \check{g}_0, 0 \times s_0)$. We write this convergence by saying that for any integers $n \geq 1$ and $k \geq 1$ we have

$$\lim_{i \uparrow \infty} \left\| \varphi_{n,i}^* g_0 - \check{g}_0 \right\|_{C_{\check{g}_0}^k([-n,n] \times \mathbb{S}^2)} = 0, \tag{35}$$

where $\varphi_{n,i} : [-n, n] \times \mathbb{S}^2 \rightarrow [-n + \hat{x}_i, n + \hat{x}_i] \times \mathbb{S}^2 (\subset \Sigma_0)$ is the map $\varphi_{n,i}(x, s) = (x + \hat{x}_i, s)$ (note that $\varphi_{n,i}(0 \times s_0) = \hat{x}_i \times s_0$ for all i). More generally, the pointed sequence of initial data $(\Sigma_0; g_0; E_0; \hat{x}_i \times s_0)$ converges smoothly to $(\mathbb{R} \times \mathbb{S}^2; \check{g}_0; \check{E}_0; 0 \times s_0)$ because in addition to (35) we have

$$\lim_{i \uparrow \infty} \left\| \varphi_{n,i}^* E_0 - \check{E}_0 \right\|_{C_{\check{g}_0}^k([-n,n] \times \mathbb{S}^2)} = 0, \tag{36}$$

for any $n \geq 1$ and $k \geq 1$. Fix now $0 < \hat{\epsilon} < 1/16$ and consider the sequence of vector fields $\hat{E}_i := \hat{E}_{\hat{x}_i, \hat{\epsilon}}$ given in (25) out of $\hat{x} = \hat{x}_i$, $\hat{\epsilon}$ and ω . In the same way as before, this sequence converges smoothly to $\hat{E}_\infty := \check{E}_0 + \hat{\lambda}_\infty (\star d \star \omega_\infty)^\#$

where ω_∞ is the pull-back of ω by the map $(x, s) \rightarrow (x - 2, s)$ from $[-1, 1] \times \mathbb{S}^2$ into $[-3, -1] \times \mathbb{S}^2$ and $\hat{\lambda}_\infty$ is a constant such that

$$\sup \left| 1 - \frac{|\hat{E}_\infty|_{\hat{g}_0}^2}{|\check{E}_0|_{\check{g}_0}^2} \right| = \hat{\epsilon} \tag{37}$$

As before this convergence is expressed by the limit

$$\lim_{i \uparrow \infty} \left\| \varphi_{n,i}^* \hat{E}_i - \hat{E}_\infty \right\|_{C_{\hat{g}_0}^k([-n,n] \times \mathbb{S}^2)} = 0, \tag{38}$$

for any $n \geq 1$ and $k \geq 1$. Note that as ω_∞ has support in $[-1, 1] \times \mathbb{S}^2$ then $\hat{E}_\infty = \check{E}_0$ outside $[-1, 1] \times \mathbb{S}^2$. In particular, $|\hat{E}_\infty|_{\hat{g}_0} = 1$ outside $[-1, 1] \times \mathbb{S}^2$ because $|\check{E}_0|_{\check{g}_0} = 1$.

Now, let ϕ_i be the sequence of conformal factors constructed in Sect. 4 out of \hat{x}_i and the fixed $\hat{\epsilon}$. Using standard elliptic estimates and the barrier bounds (30)–(31) one easily shows that the sequence ϕ_i has a subsequence (indexed again by “ i ”) converging smoothly to a limit smooth conformal factor $\phi_\infty > 0$. Namely,

$$\lim_{i \uparrow \infty} \left\| \varphi_{n,i}^* \phi_i - \phi_\infty \right\|_{C_{\check{g}_0}^k([-n,n] \times \mathbb{S}^2)} = 0, \tag{39}$$

for any $n \geq 1$ and $k \geq 1$. Moreover, because of (35), (36) and (38) the limit conformal factor ϕ_∞ satisfies the limit L-equation

$$\Delta_{\check{g}_0} \phi_\infty = 2|\check{E}_0|_{\check{g}_0}^2 \phi_\infty - 2|\hat{E}_\infty|_{\hat{g}_0}^2 \phi_\infty^{-3}. \tag{40}$$

The convergences (35), (36), (38) and (39) also show that the pointed subsequence $(\bar{g}_i = \phi_i^4 g_0; \bar{E}_i = \hat{E}_i = \phi^{-6} \hat{E}_i; \hat{x}_i \times s_0)$ converges smoothly to $(\bar{g}_\infty := \phi_\infty^4 \check{g}_0; \bar{E}_\infty := \phi_\infty^{-6} \hat{E}_\infty, 0 \times s_0)$.

It is an important fact that the limit data set $(\mathbb{R} \times \mathbb{S}^2; \bar{g}_\infty; \bar{E}_\infty)$ is never the ERNT₁ initial data. If this were the case then we would have $|\bar{E}_\infty|_{\bar{g}_\infty}^2 = 1$ and therefore $|\hat{E}_\infty|_{\hat{g}_0}^2 \phi_\infty^{-8} = 1$. Plugging this in (40) and recalling that $|\check{E}_0|_{\check{g}_0} = 1$ we would obtain

$$\Delta_{\check{g}_0} \phi_\infty = 2\phi_\infty - 2\phi_\infty^5.$$

Then observe that as $|\hat{E}_\infty|_{\hat{g}_0} = 1$ outside $[-1, 1] \times \mathbb{S}^2$ we would have $\phi_\infty = 1$ also outside $[-1, 1] \times \mathbb{S}^2$. Then, as the constant function one is a solution of (40) we must have $\phi_\infty = 1$ everywhere by the unique continuation principle. Thus, it would be $|\hat{E}_\infty|_{\hat{g}_0} = 1$ everywhere, contradicting (37).

Observe that any non-contractible surface S embedded in $(\mathbb{R} \times \mathbb{S}^2; \bar{g}_\infty)$ must have \bar{g}_∞ -area greater or equal than 4π . To see this use Proposition 1 to have $A_{\bar{g}_\infty}(S) = \lim A_{\bar{g}_i}(\varphi_{n,i}(S)) \geq 4\pi$. Similarly, we have $|Q_E(S)| = \lim |Q_E(\varphi_{n,i}(S))| = 1$. We can now use this information together with Lemma 2 and the fact that the limits $(\bar{g}_\infty; \bar{E}_\infty)$ are not the ERNT₁ initial data, to conclude that for any non-contractible embedded S we have $A_{\bar{g}_\infty}(S) > 4\pi$. This will be crucially used in the following proposition.

Proposition 4. *Let $0 < \hat{\epsilon} < 1/16$. Then there is $\hat{x}_0 = \hat{x}_0(\hat{\epsilon})$ such that for any $\hat{x} \leq \hat{x}_0$ the data set $(\bar{g}, \bar{K}; \bar{E}, \bar{B})$ constructed in Sect. 4 out of $\hat{\epsilon}$ and \hat{x} possess*

an embedded minimal and stable sphere M separating the two ends. Because $\bar{K} = 0$ such sphere is also a MOTS (to the past and to the future).

Proof. We will proceed by contradiction. Assume therefore that there is $0 < \hat{\epsilon} < 1/16$ and a sequence $\hat{x}_i \rightarrow -\infty$ such that, if we denote by $((-\infty, 0) \times \mathbb{S}^2; \bar{g}_i; \bar{E}_i)$ the data sets constructed out of $\hat{\epsilon}$ and \hat{x}_i , then none of the manifolds $((-\infty, 0) \times \mathbb{S}^2; \bar{g}_i)$ possess a stable minimal sphere M separating the two ends. We will see that this leads to a contradiction.

Firstly, as commented before, one can take a subsequence of the pointed sequence $((-\infty, 0) \times \mathbb{S}^2; \bar{g}_i; \bar{E}_i; \hat{x}_i \times s_0)$ converging (in the pointed sense) to a smooth data set $((-\infty, \infty) \times \mathbb{S}^2; \bar{g}_\infty, \bar{E}_\infty)$. Moreover, as was commented right before the statement of the proposition, we know that for any embedded sphere S isotopic to $S_0 := 0 \times \mathbb{S}^2$ we have $A_{\bar{g}_\infty}(S) > 4\pi$.

Secondly, let $\psi_\delta(z)$ be the smooth real function of the one variable $z \in [-1, \infty]$ defined as

$$\psi_\delta(z) = \begin{cases} 1 + e^{1/z + 1/(z + 1 + \delta)} & \text{if } z \in [-1, 0], \\ 1 & \text{if } z \in [0, \infty) \end{cases}$$

With this function define the metric $\tilde{g}_i = [\psi_\delta(x - \hat{x}_i)]\bar{g}_i$ on the manifold $[-1 + \hat{x}_i, 0) \times \mathbb{S}^2$ and set $\delta > 0$ small enough that the boundary $(-1 + \hat{x}_i) \times \mathbb{S}^2$ of $[-1 + \hat{x}_i, 0) \times \mathbb{S}^2$ is strictly mean convex (in the direction of decreasing x) for all i . Of course the pointed sequence $([-1 + \hat{x}_i, 0) \times \mathbb{S}^2; \tilde{g}_i; \hat{x}_i \times s_0)$ converges to $([-1, \infty) \times \mathbb{S}^2; \tilde{g}_\infty)$ where $\tilde{g}_\infty = \psi_\delta \bar{g}_\infty$ and because $\psi_\delta \geq 1$ we have $A_{\tilde{g}_\infty}(S) \geq A_{\bar{g}_\infty}(S) > 4\pi$ for any embedded sphere isotopic to S_0 .

Thirdly, recall that $\bar{g}_i = \phi_i^2 g_0$ where ϕ_i is a solution to the L-equation enjoying the upper and lower bounds $\phi_{i,-} \leq \phi_i \leq \phi_{i,+}$ where $\phi_{i,\pm}$ are given by (30)–(31) with $\hat{x} = \hat{x}_i$. In particular, the conformal factor ϕ_i restricted to the spheres $S_{\hat{x}_i/2} := \{x = \hat{x}_i/2\}$ is bounded below by $1 - 4\hat{\epsilon}/\cosh(\hat{x}_i/2)$ and above by $1 + 4\hat{\epsilon}/\cosh(\hat{x}_i/2)$. This implies that $A_{\bar{g}_i}(S_{\hat{x}_i/2}) \rightarrow 4\pi$ and therefore that $A_{\tilde{g}_i}(S_{\hat{x}_i/2}) \rightarrow 4\pi$. Let $\tilde{S}_i \subset [-1 + \hat{x}_i, 0) \times \mathbb{S}^2$ be the embedded sphere minimizing the \tilde{g}_i -area among all spheres embedded in $[-1 + \hat{x}_i, 0) \times \mathbb{S}^2$ and isotopic to S_0 . Such sphere always exists because $([-1 + \hat{x}_i, 0) \times \mathbb{S}^2, \tilde{g}_i)$ has strictly mean convex boundary and is asymptotically flat [16]. Moreover, as $A_{\tilde{g}_i}(S_{\hat{x}_i/2}) \rightarrow 4\pi$ and as $A_{\tilde{g}_i}(\tilde{S}_i) \geq 4\pi$ for all i then we must have $A_{\tilde{g}_i}(\tilde{S}_i) \rightarrow 4\pi$.

On the other hand, every surface \tilde{S}_i must intersect $[-1 + \hat{x}_i, \hat{x}_i] \times \mathbb{S}^2$, which is the domain where \tilde{g}_i differs from \bar{g}_i , otherwise \tilde{S}_i would be \bar{g}_i -minimal and stable which is against the assumption. Now, take another subsequence if necessary in such a way that \tilde{S}_i converges to a \tilde{g}_∞ -minimal and stable sphere intersecting $[-1, 0] \times \mathbb{S}^2$ (inside the limit space) and isotopic to $S_0 := 0 \times \mathbb{S}^2$. As discussed before we must have $A_{\tilde{g}_\infty}(\tilde{S}_\infty) > 4\pi$ and at the same time $A_{\tilde{g}_\infty}(\tilde{S}_\infty) = \lim A_{\tilde{g}_i}(\tilde{S}_i) = 4\pi$ which is a contradiction. \square

7. Proof of the Main Result

We are ready to prove the main result of this article. For the convenience of the reader we restate it below.

Theorem 1. *For any $\bar{\varepsilon} > 0$ and integer $k \geq 1$ there is a smooth and maximal electro-vacuum data set $(\bar{\Sigma}; \bar{g}, \bar{K}; E, B)$, $\bar{\varepsilon}$ -close in C^k to the standard ERN initial data and falling into it exponential along the cylindrical end, which cannot decay, towards the future or the past, into any EKN solution.*

The precise definition of decaying into a EKN that we adopt, and that we provide below, is the laxest one can imagine. In very rough terms it says that, as one moves conveniently to the future but remaining outside the future black hole (if any), the space-time becomes “more and more similar” to a EKN. This is quantified by demanding that the data sets induced on certain domains of a sequence of Cauchy hypersurfaces, (the domains outside the black hole if any), converges to the data set induced on a future Cauchy hypersurface of the EKN in an appropriate norm.

To alleviate the statement of the definition, we introduce before some preliminary terminology. Let $(\mathbf{M}; \mathbf{g})$ be a globally hyperbolic space-time. Then,

1. A sequence of Cauchy hypersurfaces Σ_i exhaust \mathbf{M} to the future if (i) $D^+(\Sigma_{i'}) \subset D^+(\Sigma_i)^\circ$ when $i' > i$, and (ii) $\bigcap D^+(\Sigma_i) = \emptyset$.
2. A space-like hypersurface \mathcal{C} is a future Cauchy hypersurface if $\mathbf{M} \setminus \mathcal{C}$ has two connected components one of which is $D^+(\mathcal{C}) \setminus \mathcal{C}$.
3. A sequence of open sets \mathcal{V}_i of a future Cauchy surface \mathcal{C} is an exhaustion if (i) $\mathcal{V}_i \subset \mathcal{V}_{i'}$ when $i' \geq i$, and (ii) $\bigcup \mathcal{V}_i = \mathcal{C}$.

With this terminology at hand, we give now the definition we need. We comment on it thereafter.

Definition 4. A globally hyperbolic electro-vacuum space-time $(\mathbf{M}, \mathbf{g}; \mathbf{F})$ decays (to the future) into an eKN space-time $(\mathbf{M}_e; \mathbf{g}_e; \mathbf{F}_e)$ iff

1. $(\mathbf{M}; \mathbf{g})$ is asymptotically flat at spatial and null infinity ([21], p. 276), and,
2. There are diffeomorphisms into the images

$$\varphi_i : \mathcal{V}_i \longrightarrow I^-(\mathcal{S}^+) \cap \Sigma_i,$$

from the opens \mathcal{V}_i of an exhaustion of a future Cauchy hypersurface \mathcal{C}_e of $(\mathbf{M}_e; \mathbf{g}_e)$ into $I^-(\mathcal{S}^+) \cap \Sigma_i$, where the sequence Σ_i exhausts $(\mathbf{M}; \mathbf{g})$ to the future, and such that for (U_i, U_e) equal either $(\varphi_i^* g_i, g_e)$, $(\varphi_i^* K_i, K_e)$, $(\varphi_i^* E_i, E_e)$ or $(\varphi_i^* B_i, B_e)$ we have

$$\lim_{i \rightarrow \infty} \|U_i - U_e\|_{C^1_{g_e}(\mathcal{V}_i)} = 0. \tag{41}$$

The hypersurface \mathcal{C}_e is assumed spherically axisymmetric when $(\mathbf{M}_e; \mathbf{g}_e; \mathbf{F}_e)$ is ERN, and axisymmetric when $(\mathbf{M}_e; \mathbf{g}_e; \mathbf{F}_e)$ is any other EKN. Finally if $(\mathbf{M}, \mathbf{g}; \mathbf{F})$ is axisymmetric then the Σ_i are also assumed axisymmetric.

A few comments are in order. Firstly, *item 1* in Definition 4 imposes a basic global condition which allows us to speak of future null-infinity \mathcal{S}^+ and therefore of black holes. Secondly, the use of the maps $\varphi_i : \mathcal{V}_i \longrightarrow I^-(\mathcal{S}^+) \cap \Sigma_i$ in *item 2* guarantees that the notion of “decaying into a EKN” deals only with the strict exterior of the black hole and does not hypothesize about the nature

of the event horizon as was discussed at length in the introduction. Thirdly, the convergence in (41) is only in C^1 which is very weak.

On these grounds it is hard to imagine a more adequate definition of “decaying into a EKN” than this. In any case, this is the one we use in Theorem 1 that we now prove.

Proof of Theorem 1. Set $\hat{\epsilon}$ be small enough in such a way that the $\varepsilon(\hat{\epsilon}, k)$ provided by Proposition 3 is less or equal than $\bar{\varepsilon}$. Let then \hat{x} be any number less or equal than the $\hat{x}(\hat{\epsilon})$ provided by Proposition 4 and let $(\bar{g}, \bar{K} = 0; \bar{E}, \bar{B} = 0)$ be the axisymmetric and time symmetric data set constructed in Sect. 4 out of $\hat{\epsilon}$ and \hat{x} . By Proposition 3 such data set is $\bar{\varepsilon}$ -close in C^k to the standard ERN initial data. Its total electromagnetic charges are $Q_E = 1, Q_M = 0$ and the total angular momentum is $J = 0$ (see Sect. 2 (iii)). Moreover, the data set falls off exponentially towards the background data set $(g_0, K_0; E_0, B_0)$ along the cylindrical end as explained at the end of Sect. 4. Also, by Proposition 4, such data set possess a stable minimal surface M separating the two ends, which is therefore a future and past MOTS. For this reason the following argument applies equally to the future and to the past. Here we will argue only to the future. The future globally hyperbolic development of the initial data will be denoted by $(M^+; g)$.

Suppose now that the future evolution of the initial data decays into an extreme EKN solution according to Definition 4. In this situation the EKN limit must be necessarily ERN_1 because the electromagnetic charges and the angular momentum are everywhere conserved. Also, because of the presence of the MOTS M acting as a barrier, the set $I^-(S^+) \cap \bar{\Sigma}$ lies in the region inside $\bar{\Sigma}$ enclosed by M and the asymptotically flat end. Note that $I^-(S^+) \cap \bar{\Sigma}$ is the exterior black hole region at the “instant” $\bar{\Sigma}$.

Let \mathcal{C}_e be the spherically symmetric future Cauchy surface in ERN_1 and let $\{\mathcal{V}_j\}$ be the exhaustion, as granted by Definition 4. Then observe that no matter which is the \mathcal{C}_e at hand, one can find a sequence of spheres $S_j \subset \mathcal{C}_e$ with strictly positive outer null expansions $\theta^+|_{S_j}$ and whose areas tend to 4π as $j \rightarrow \infty$. One can take for instance any sequence of spheres of constant coordinates r and t , “going towards” one of the ends of \mathcal{C}_e (where the future Cauchy horizon of ERN_1 “is”). Hence, appealing now to (41) in Definition 4, one can consider a subsequence $\{\mathcal{V}_{i_j}\}$ of $\{\mathcal{V}_j\}$ such that $S_j \subset \mathcal{V}_{i_j}$ for all j and such that the spheres $\varphi_{i_j}(S_j)$ have strictly positive outer null expansion $\theta^+|_{\varphi_{i_j}(S_j)}$ in $(M^+; g)$.

Let $\Sigma_{i_j}^*$ be the closure of the connected component of $\Sigma_{i_j} \setminus \varphi_{i_j}(S_j)$ containing the asymptotically flat end. Consider now the (non-necessarily smooth) null boundary

$$N := \left[\partial(J^-(\Sigma_{i_j}^*) \cap M^+) \right] \setminus \left(\Sigma_{i_j} \cup \bar{\Sigma} \right).$$

Then observe that every null generator of N ends at a point in $\varphi_{i_j}(S_j)$ and as $\varphi_{i_j}(S_j) \subset I^-(S^+)$ observe that $N \subset I^-(S^+)$. Also note that at any end point of a null generator of N the expansion of the null congruence is positive because

it is (proportional to) $\theta^+|_{\varphi_{i_j}(S_j)}$ which is positive. As by the Rauchadhury equation the expansion of N is non-increasing along its null generators, we deduce that it is also positive at every smooth point in N . In particular the area of $\varphi_{i_j}(S_j)$ is bigger than the area of the two-rectifiable set $H_j := \partial(J^-(\Sigma_{i_j}^*) \cap \bar{\Sigma})$ in the initial slice $\bar{\Sigma}$, [7]. Hence $\limsup A_{\bar{g}}(H_j) \leq 4\pi$.

By the *item 1* in Definition 4 the set $J^-(\Sigma_{i_j}^*) \cap \bar{\Sigma}$ contains (obviously) a large sphere over the asymptotically flat end of $\bar{\Sigma}$ and its exterior region. For this reason H_j is homologous to that sphere and therefore to M , [20]. Take now another sphere \tilde{M} far away along the cylindrical end of $\bar{\Sigma}$ in such a way that the region $\tilde{\Sigma}$ enclosed by it and the asymptotically flat end contains the MOTS M . Then, as we did for instance in the proof of Proposition 4, deform the metric \bar{g} around $\tilde{M} = \partial\tilde{\Sigma}$ in $\tilde{\Sigma}$ to a metric \tilde{g} by means of a conformal factor greater or equal than one, in such a way that the outward mean curvature of $\tilde{M} = \partial\tilde{\Sigma}$ in $(\tilde{\Sigma}, \tilde{g})$ is strictly positive. The conformal factor is assumed to be one on the region enclosed by M and the asymptotically flat end, therefore $\bar{g} = \tilde{g}$ in there. Then, based on general grounds of geometric measure theory [20], one can find a smooth \tilde{g} -area-minimizer among all the two-rectifiable sets in $\tilde{\Sigma}$ homologous to M . As $\limsup A_{\tilde{g}}(H_j) = \limsup A_{\bar{g}}(H_j) \leq 4\pi$, we conclude that the \tilde{g} -area of such minimizer (which is homologous to M and non-empty) is less or equal than 4π . As $\tilde{g} \geq \bar{g}$ we conclude that the \bar{g} -area of the minimizer is less or equal than 4π contradicting Proposition 2. \square

Acknowledgements

We would like to thank Sergio Dain for useful suggestions in the early stages of this article.

References

- [1] Aretakis, S.: Horizon instability of extremal black holes (2012) [arXiv:1206.6598](https://arxiv.org/abs/1206.6598)
- [2] Aronszajn, N.: A unique continuation theorem for solutions of elliptic partial differential equations or inequalities of second order. *J. Math. Pures Appl.* (9) **36**, 235–249 (1957)
- [3] Bizon, P., Friedrich, H.: A remark about wave equations on the extreme Reissner–Nordström black hole exterior. *Class. Quant. Grav.* **30**, 065001 (2013)
- [4] Camacho, C., Lins Neto, A.: *Geometric Theory of Foliations*. Birkhäuser Boston Inc., Boston (1985)
- [5] Carter, B.: Black hole equilibrium states. In: *Black holes/Les astres occlus (École d’Été Phys. Théor. Les Houches, 1972)*, pp. 57–214. Gordon and Breach, New York (1973)
- [6] Choquet-Bruhat, Y., York, J.W. Jr.: The Cauchy problem. In: *General Relativity and Gravitation*, vol. 1, pp. 99–172. Plenum, New York (1980)
- [7] Chrusciel, P.T., Delay, E., Galloway, G.J., Howard, R.: The Area theorem. *Annales Henri Poincaré* **2**, 109–178 (2001)

- [8] Chrusciel, P.T., Mazzeo, R.: Initial data sets with ends of cylindrical type: I. The Lichnerowicz Equation. [arXiv:1201.4937](https://arxiv.org/abs/1201.4937)
- [9] Colding, T.H., Minicozzi, W.P. II.: A course in minimal surfaces. In: Graduate Studies in Mathematics, vol. 121. American Mathematical Society, Providence, RI (2011)
- [10] Dain, S.: Geometric inequalities for axially symmetric black holes. *Class. Quant. Grav.* **29**, 073001 (2012)
- [11] Dain, S., Dotti, G.: The wave equation on the extreme Reissner–Nordström black hole. *Class. Quant. Grav.* **30**, 055011 (2013)
- [12] Dain, S., Jaramillo, J.L., Reiris, M.: Area-charge inequality for black holes. *Class. Quant. Grav.* **29**, 035013 (2012)
- [13] Figueras, P., Murata, K., Reall, H.S.: Black hole instabilities and local Penrose inequalities. *Class. Quant. Grav.* **28**, 225030 (2011)
- [14] Lucietti, J., Murata, K., Reall, H.S., Tanahashi, N.: On the horizon instability of an extreme Reissner–Nordström black hole. *JHEP* **1303**, 035 (2013)
- [15] Lucietti, J., Reall, H.S.: Gravitational instability of an extreme Kerr black hole. *Phys. Rev. D* **86**, 104030 (2012)
- [16] Meeks, W. III., Simon, L., Yau, S.T.: Embedded minimal surfaces, exotic spheres, and manifolds with positive Ricci curvature. *Ann. Math. (2)* **116**(3), 621–659 (1982)
- [17] Murata, K., Reall, H.S., Tanahashi, N.: What happens at the horizon(s) of an extreme black hole? *Class. Quant. Grav.* **30**, 235007 (2013)
- [18] Nakauchi, N.: Compactness of the space of incompressible stable minimal surfaces without boundary. *J. Math. Kyoto Univ.* **30**(2), 343–346 (1990)
- [19] Reiris, M.: On extreme Kerr-throats and zero temperature black-holes. *Class. Quant. Grav.* **31**, 025001 (2014)
- [20] Simon, L.: Lectures on geometric measure theory. In: Proceedings of the Centre for Mathematical Analysis, Australian National University, vol. 3. Australian National University Centre for Mathematical Analysis, Canberra (1983)
- [21] Wald, R.M.: General Relativity. University of Chicago Press, Chicago (1984)

Martin Reiris
Max Planck Institute für Gravitationsphysik
Golm, Germany
e-mail: martin@aei.mpg.de

Communicated by James A. Isenberg.

Received: January 15, 2014.

Accepted: July 2, 2014.