Born–Oppenheimer Approximation for an Atom in Constant Magnetic Fields

Sohei Ashida

Abstract. We obtain a reduction scheme for the study of the quantum evolution of an atom in constant magnetic fields using the method developed by Martinez, Nenciu and Sordoni based on the construction of almost invariant subspace. In Martinez and Sordoni (Mem AMS 936, 2009) such a case is also studied but their reduced Hamiltonian includes the vector potential terms. In this paper, using the center of mass coordinates and constructing the almost invariant subspace different from theirs, we obtain the reduced Hamiltonian which does not include the vector potential terms of the nucleus. Using the reduced evolution, we also obtain the asymptotic expansion of the evolution for a specific localized initial data, which verifies the straight motion of an atom in constant magnetic fields.

1. Introduction

In this paper, we study the evolution of the system of one nucleus and some electrons which is neutral, i.e., the total electric charge is zero, in constant magnetic fields and external electric potentials. We construct the reduction scheme of the evolution of the system to the evolution of center of mass of the particles with small error terms which is regarded as Born–Oppenheimer approximation. It has been shown that the reduced evolution of the center of mass is not influenced by the magnetic field directly, which implies the straight motion of the center of mass when the external potential does not exists.

The Hamiltonian of an atom with a nucleus and N electrons moving in constant magnetic fields has the form

$$\begin{aligned} \hat{P} = & \frac{1}{2m}(D_{x_1} - e_1 A(x_1))^2 + \sum_{i=2}^{N+1} \frac{1}{2m_e}(D_{x_i} - eA(x_i))^2 \\ & + \sum_{i < j} V_{ij}(x_i - x_j) + \sum_{i=1}^{N+1} V_i(x_i). \end{aligned} \quad (1.1)$$

Here $x_1 \in \mathbb{R}^3$ (resp., m) denotes the position (resp., the mass) of the nucleus, $x_j, j \geq 2$ (resp., m_e) denote the position (resp., the mass) of electrons and V_{ij} (resp., V_i) are interaction (resp., external) potentials. e_1 (resp., e) denotes the charge of the nucleus (resp., electrons) and A denotes the vector potential.

There is a lot of literature on the quantum many-body problem. Especially the large time asymptotics of the solution of the Schrödinger equation has been studied intensively and the complete classification of the asymptotic behavior called asymptotic completeness was proved for a large variety of potentials (see Dereziński and Gérard [4]). We do not deal with such a problem here, but we study the phenomenon which happens when there are differences of the masses of the particles instead. Roughly speaking the Born–Oppenheimer approximation says the following. Since the electrons are lighter than the nuclei, they move rapidly and adjust their state adiabatically as the nuclei move more slowly. To develop the mathematical theory of the Born–Oppenheimer approximation, the ratio of electronic and nuclear mass is denoted by h^2 and regarded as a small parameter. Before we consider the case with magnetic fields, it is instructive to recall the results for the case without magnetic fields. In this case, the Hamiltonian of some nuclei and electrons is written as

$$P(h) = -h^2 \Delta_x - \Delta_y + V(x, y),$$

where we denote the coordinates of the electrons by y and that of the nuclei by x . Our purpose is to study the asymptotics of the solution to the Schrödinger equation $ih\partial_t\varphi = P(h)\varphi$ as $h \rightarrow 0$. From the intuitive description of the Born–Oppenheimer approximation above, if the electrons are in bound states for the fixed nuclei, i.e., the bound states for $P_e(x) := -\Delta_y + V(x, y)$ at the initial time, we expect the electrons remain in the bound states even after time passes. This suggests there is an almost invariant subspace close to the electronic bound states under the evolution $e^{-itP(h)}$.

The almost invariant subspaces are described by the projections (see Nenciu [19, 20]). If an orthogonal projection Π satisfies $[P(h), \Pi] = \mathcal{O}(h^\infty)$, then $e^{-itP}\Pi = \Pi e^{-itP} + \mathcal{O}(h^\infty|t|)$ holds which means $\text{Ran}\Pi$ is the almost invariant subspace. We expect there exists such a projection Π such that $\Pi - \Pi_0 = \mathcal{O}(h)$ where $\Pi_0 = \int^\oplus \Pi_0(x)dx$ and $\Pi_0(x)$ is the spectral projection onto an arbitrarily chosen part of the discrete spectrum of $P_e(x)$ separated from the other part of the spectrum. In Nenciu [19, 20] the projections onto almost invariant subspaces were constructed by a recurrence formula and another construction by pseudodifferential calculus was introduced by Helffer and Sjöstrand [13] and Sjöstrand [23]. In the case of Born–Oppenheimer approximation, the almost invariant subspace does not seem to exist according to the physical intuition saying that the adiabatic decoupling becomes weaker and weaker when the energy increases. However for any cutoff function χ a projection Π which satisfies $[P(h), \Pi]\chi(P) = \mathcal{O}(h^\infty)$ is constructed by Sordani [24]. Using the projection Π the quantum evolution $e^{-itP/h}$ of the molecule is reduced to the evolution of the nuclei $e^{-itG/h}$ where G is a $k \times k$ matrix of semiclassical pseudodifferential operators $H^2(\mathbb{R}_x^n) \rightarrow L^2(\mathbb{R}_x^n)$, of the nuclear variables, k being the

rank of Π_0 (see Martinez and Sordoni [17, 18]). The symbol of G is written as $g = \xi^2 I_k + \mu(x) + \sum_{j=1}^{\infty} g_j h^j$, where I_k is the k -dimensional identity matrix and $\mu(x)$ is a matrix of $\Pi_0 P_e(x)$ in a basis of $\text{Ran} \Pi_0$. The first results of the reduction to the Hamiltonian of nuclear variables were obtained by Spohn and Teufel [25] (see also Teufel [26]). In [25] they proved that the difference between the evolution of the reduced Hamiltonian and the original one is of order $\mathcal{O}(h)$ and semiclassics for observables are also obtained.

Let f be a generalized coherent state of nuclei and ψ be a bound state of electrons. The initial state of the form $f(x)\psi(x, y) + \mathcal{O}(h)$ is used by Hagedorn (see Hagedorn [7–9] and Hagedorn and Joye [12]). In semiclassical limit the quantum evolution of generalized coherent states admits an asymptotic expansion each term of which is a generalized coherent state centered at the point reached by the classical flow (see Combes and Robert [3] and Combes [2]). In [18] Martinez and Sordoni applied this expansion to G to obtain the evolution like that of Hagedorn.

We now return to the case with constant magnetic fields. Its coordinates perpendicular to the magnetic field stay in a bounded region as in the classical mechanics. When there are N particles, the center of mass of the particles does not move freely, so that we cannot divide the motion of the particles into the internal and external motion. However, if the total charge of the particles is zero, there is a subspace $\mathcal{H}_{\text{bound}}$ of $L^2(\mathbb{R}^{3N})$ such that internal coordinates of the particles in the state in $\mathcal{H}_{\text{bound}}$ stay in a bounded region and the particles travel to infinity across the magnetic field (see Gérard and Laba [5]). This would occur from the viewpoint of the classical mechanics because the Lorentz force on the particles with the opposite charge has the opposite direction and the particles interact with the interaction potential. Thus we expect when there is an atom, the electrons around the nucleus offset the influence of the magnetic fields. Hence it seems to be natural that the vector potential terms do not appear in the reduced Hamiltonian G . The Born–Oppenheimer approximation with magnetic fields is also dealt with by Martinez and Sordoni [18] but in their construction the term $\hbar^2(D_x - eA(x))^2$ remains in G . This is because in the coordinates where independent variables are positions of the particles, the effects of magnetic fields on the nuclei and the electrons are treated separately and therefore even if we restrict the Hamiltonian P on $\text{Ran} \Pi_0$, the effect of magnetic fields does not disappear. Our purpose is to construct a reduction scheme from $\hat{P}$ in (1.1) to G without vector potential terms.

To obtain such G we change the coordinates $(x_1, \dots, x_{N+1})$ to the new coordinates where the independent variables are the center of mass x and the relative position $y = (y_2, \dots, y_{N+1})$ of the electrons from the nucleus (see (2.1)). Then the Hamiltonian $\hat{P}$ is transformed into a certain operator $\tilde{P}$. Next we transform $\tilde{P}$ by a unitary transformation $\mathcal{V} = \exp(-ieA(x) \sum_{i=2}^{N+1} y_i)$ as $P := \mathcal{V} \tilde{P} \mathcal{V}^*$. The transformed Hamiltonian is written in the following form:

$$P = \hbar^2 D_x^2 + P_e(x) - 4\hbar^2 e \sum_{i=2}^{N+1} A(y_i) D_x + \hbar^2 Q, \quad P_e(x) = \sum_{i=2}^{N+1} L_i^2 + V_0(x, y).$$

Here $L_i = D_{y_i} - eA(y_i)$, V_0 is the zeroth order term with respect to h , and Q is a certain operator on $L^2(\mathbb{R}_y^{3N})$. We regard $P_e(x)$ as the electric Hamiltonian and denote the spectral projection onto the chosen part of the discrete spectrum of $P_e(x)$ by $\Pi_0(x)$.

Our main results, Theorems 2.5–2.9 in Sect. 2 can be illustrated as follows. For any cutoff function $\chi \in C_0^\infty(\mathbb{R})$ there exists an orthogonal projection $\Pi(h)$ onto the almost invariant subspace close to Π_0 , that is $[P, \Pi]\chi(P) = \mathcal{O}(h^\infty)$ and $\Pi - \Pi_0 = \mathcal{O}(h)$. There also exists a reducing transformation

$$W: L^2(\mathbb{R}^{3(N+1)}) \rightarrow (L^2(\mathbb{R}_x^3))^{\oplus k},$$

where $k = \text{Rank} \Pi_0$, and reduced Hamiltonian G which is $k \times k$ self-adjoint matrix of operators on $L^2(\mathbb{R}_x^3)$ without vector potential term such that the restriction U of W to $\text{Ran} \Pi$ is a unitary operator satisfying

$$e^{-itP/h} \Pi \varphi_0 = U^* e^{-itG/h} U \Pi \varphi_0 + \mathcal{O}(|t|h^\infty \|\varphi_0\|),$$

for $\varphi_0 \in \text{Ran} \chi(P)$. This means that the motions of all the particles are reduced to the motion of the center of mass. Moreover, if $k = 1$, we study the case of an initial data which is a wave packet belonging to the range of $\mathcal{V}^* \Pi$. We give in Theorem 2.9 below a description of its evolution through $e^{-it\tilde{P}/h}$ and show that it conserves its wave packet structure. We refer to the precise statements of Sect. 2.

The result of Gérard and Laba [5] mentioned above also indicates that if the total charge is zero and the initial data is in a certain subspace of $L^2(\mathbb{R}^N)$, the particles can travel across the magnetic field. This result holds without any approximation but in our Born–Oppenheimer framework the localization is clearer, and although Gérard and Laba [5] considered the case without external potentials in which the pseudomomentum of the center of mass commutes with the Hamiltonian, our result covers the case with smooth bounded external potentials.

The content of this paper is as follows. In Sect. 2, we transform the Hamiltonian into the form without vector potential term of nuclear variable, introduce our assumptions and state our main results. In Sect. 3 we prove the main results. In the “Appendix” we collect the results we need on the pseudodifferential operators with operator valued symbols.

2. Some Preliminaries and Main Results

We suppose the magnetic field is parallel to the third axis, so that the vector potential is written as

$$A(x) = \begin{pmatrix} 0 & -b & 0 \\ b & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} x$$

where $b > 0$ is a constant. We also suppose that the the total charge of the particles is zero, i.e., $e_1 + Ne = 0$. Setting the mass of electrons to 1 we introduce new coordinates $(x, y_2, \dots, y_{N+1}) = (x, y) \in \mathbb{R}^3 \times \mathbb{R}^{3N}$ by setting

$$\begin{aligned}
 x &= \frac{1}{M} \left(mx_1 + \sum_{i=2}^{N+1} x_i \right), \quad M = m + N \\
 y_i &= x_i - x_1, \quad 2 \leq i \leq N+1.
 \end{aligned} \tag{2.1}$$

Here M is the total mass, x is the position of center of mass and y_i is the position of the electrons relative to the nucleus. With this choice of coordinates, we have the Hamiltonian

$$\tilde{P} = h^2 D_x^2 - 2h^2 e \sum_{i=2}^{N+1} A(y_i) D_x + \sum_{i=2}^{N+1} \tilde{L}_i(x)^2 + h^2 \tilde{Q} + V(x, y)$$

where $h^2 = \frac{1}{M}$, $\tilde{L}_i(x) = D_{y_i} - eA(y_i + x)$, and writing $f = \sum_{i=2}^{N+1} y_i$, $V(x, y)$ and $\tilde{Q} = \tilde{Q}_1 + \tilde{Q}_2$ are as follows:

$$\begin{aligned}
 V(x, y; h) &= \sum_{i=2}^{N+1} V_{1i}(y_i) + \sum_{2 \leq i < j} V_{ij}(y_i - y_j) \\
 &\quad + V_1(x - h^2 f) + \sum_{i=2}^{N+1} V_i(x + y_i - h^2 f), \\
 \tilde{Q}_1 &= \frac{1}{1 - Nh^2} \left(\sum_{i=2}^{N+1} \tilde{L}_i(x) \right)^2, \\
 \tilde{Q}_2 &= \frac{1}{1 - Nh^2} \left[2 \sum_{i=2}^{N+1} \tilde{L}_i(x) \left(\sum_{i=2}^{N+1} eA(y_i + h^2 f) \right) \right. \\
 &\quad \left. + \left(\sum_{i=2}^{N+1} eA(y_i + h^2 f) \right)^2 \right] + 2 \sum_{i=2}^{N+1} e\tilde{L}_i(x)A(f) + h^2 e^2 A(f)^2.
 \end{aligned}$$

Here the terms which consist of $A(x)D_x$ canceled by the assumption $e_1 + Ne = 0$.

We apply the unitary transformation $\mathcal{V} := \exp(-ieA(x) \sum_{i=2}^{N+1} y_i)$ and its inverse to $\tilde{P}$ and obtain

$$\begin{aligned}
 P &= \mathcal{V} \tilde{P} \mathcal{V}^* \\
 &= h^2 D_x^2 + P_e(x) - 4h^2 e \sum_{i=2}^{N+1} A(y_i) D_x + h^2 Q.
 \end{aligned}$$

Here,

$$P_e(x) = \sum_{i=2}^{N+1} L_i^2 + V_0(x, y), \tag{2.2}$$

and $Q = Q_1 + Q_2 + h^{-2}(V(x, y; h) - V_0(x, y))$, where $L_i = D_{y_i} - eA(y_i)$ and

$$\begin{aligned} Q_1 &= \frac{1}{1 - Nh^2} \left(\sum_{i=2}^{N+1} L_i \right)^2, \\ Q_2 &= \frac{1}{1 - Nh^2} \left[2 \sum_{i=2}^{N+1} L_i \left(\sum_{i=2}^{N+1} eA(y_i + h^2 f) \right) + \left(\sum_{i=2}^{N+1} eA(y_i + h^2 f) \right)^2 \right] \\ &\quad + 2 \sum_{i=2}^{N+1} eL_i A(f) + 2h^2 e^2 A(f)^2, \\ V_0(x, y) &= \sum_{i=2}^{N+1} V_{1i}(y_i) + \sum_{2 \leq i < j} V_{ij}(y_i - y_j) + V_1(x) + \sum_{i=2}^{N+1} V_i(x + y_i). \end{aligned}$$

Note that $V_0(x, y)$ is the zeroth order terms with respect to h of formal Taylor expansion of $V(x, y)$. We regard $P_e(x)$ as the electric Hamiltonian.

Remark 2.1. If φ is a solution of $ih\partial_t\varphi = P\varphi$, then $\mathcal{V}^*\varphi$ is a solution of $ih\partial_t\varphi = \tilde{P}\varphi$. Since $\mathcal{V}^*$ is multiplication of the complex number with modulus 1, $\mathcal{V}^*\varphi$ has the same modulus as φ .

We suppose interaction potentials V_{ij} and the external potentials V_i satisfy the following assumptions.

- (H1) (i) V_{ij} are real valued function Δ -bounded with relative bound smaller than 1.
(ii) $V_i \in C^\infty$ are real valued function and for any $\alpha \in \mathbb{N}^3$ there exists a constant C_α such that

$$|\partial^\alpha V_i(r)| \leq C_\alpha.$$

Remark 2.2. The assumption (H1)(i) covers the interaction potentials V_{ij} with Coulomb singularity. This is because after the transformation of the Hamiltonian, V_{ij} is a function of only internal coordinates y . However, since we use the global pseudodifferential calculus, the external potentials V_i must be smooth and are not allowed having such singularity.

By (H1) $\hat{P}$ is well defined as a self-adjoint operator with the domain $\{u \in L^2(\mathbb{R}^{3(N+1)}): ((D_{x_1} - e_1 A(x_1))^2 + \sum_{i=2}^{N+1} (D_{x_i} - eA(x_i))^2)u \in L^2(\mathbb{R}^{3(N+1)})\}$, so that P is also self-adjoint with some domain. $P_e(x)$ can also be regarded as a self-adjoint operator with the domain $\{u \in L^2(\mathbb{R}^{3N}): \sum_{i=2}^{N+1} L_i^2 u \in L^2(\mathbb{R}^{3N})\}$. We also suppose that

- (H2) The spectrum $\sigma(P_e(x))$ is the union of two disjoint components $\sigma_j(x)$, $j = 0, 1$, such that $\sigma_0(x)$ is a part of discrete spectrum of $P_e(x)$ with the corresponding subspace of $L^2(\mathbb{R}^{3N})$ being finite dimensional and there exists a number $d > 0$ such that

$$\inf_{x \in \mathbb{R}^3} \text{dist}(\sigma_0(x), \sigma_1(x)) \geq d.$$

Remark 2.3. By (H2) it is easy to see that there exist a continuous family of loops $\gamma(x)$ which encloses $\sigma_0(x)$ for each x and $\inf_{x \in \mathbb{R}^3} \text{dist}(\gamma(x), \sigma(P_e(x))) > d'$ for some $d' > 0$.

We denote by $\Pi_0(x)$ the spectral projection of $P_e(x)$ corresponding to σ_0 . We also suppose the following assumption.

(H3) $\text{Ran} \Pi_0(x)$ is spanned by a orthonormal basis $(u_1(x, y), \dots, u_k(x, y)) \in C^\infty(\mathbb{R}^3; L_y^2)$ such that

$$\begin{aligned} \int_{\mathbb{R}^{3N}} |u_i(x, y)|^2 e^{2\alpha|y|} dy &< C, \\ \int_{\mathbb{R}^{3N}} |\partial_x^\beta u_i(x, y)|^2 dy &< C_\beta, \quad \forall \beta \in \mathbb{N}^3 \end{aligned} \quad (2.3)$$

where constants $C > 0$, $\alpha > 0$ and C_β do not depend on x .

Remark 2.4. (1) The assumption (2.3) is needed for the symbol g in Theorem 2.7 to be in the symbol class $S^2(\mathbb{R}^3 \times \mathbb{R}^3)$.

(2) If $V_i(r) \equiv 0$, $1 \leq i \leq N+1$, then $\sigma_0(x)$ and $\Pi_0(x)$ do not depend on x , so that the assumptions (H2) and (H3) are obviously satisfied. If the eigenvalues in σ_0 are simple and the fluctuation of V_i is small enough, by the upper semicontinuity of the spectrum the eigenvalues in σ_0 do not cross each other, so that the corresponding eigenfunctions are smooth orthonormal basis. Moreover if V_i are periodic, the eigenfunctions decay uniformly with respect to x since Ω_ϵ in the proof of Agmon [1, Theorem 4.1] can be chosen uniformly in x .

Our main results are concerned with the almost invariant subspace which is close to electronic eigenspace. In the following theorem, $a = \mathcal{O}(h^\infty)$ means $a = \mathcal{O}(h^K)$ for any $K \in \mathbb{N}$.

Theorem 2.5. *Assume (H1)–(H3) hold true. Then for any $\Phi \in C_0^\infty(\mathbb{R})$ there exists a orthogonal projection $\Pi(h)$ on $L^2(\mathbb{R}^{3(N+1)})$ such that*

$$\|\Pi - \Pi_0\|_{L^2(\mathbb{R}^{3(N+1)})} = \mathcal{O}(h),$$

and, for any $\chi \in C_0^\infty(\mathbb{R})$ such that $\chi\Phi = \chi$, we have

$$\|\chi(P)[\Pi, P]\|_{L^2(\mathbb{R}^{3(N+1)})} + \|[\Pi, P]\chi(P)\|_{L^2(\mathbb{R}^{3(N+1)})} = \mathcal{O}(h^\infty).$$

The difficulty in our construction of Π is that when we expand the symbol of the resolvent $(P - z)^{-1}$ in formal power series in h as $\sum_{j=0}^\infty q_j(x, \xi; z)h^j$ where q_j are operators on $L^2(\mathbb{R}_y^{3N})$, the power of y contained in $q_j(z)$ becomes of higher order as j increases (cf. Remark 3.3). This difficulty is overcome by the exponential decay of the eigenfunctions of P_e and the fact that the line integral $\hat{\pi}_j := \oint_\Gamma q_j(z)dz$ where Γ is a certain curve can be written as the sum of the terms containing Π_0 . To prove the boundedness of $\chi(P)\hat{\Pi}_j$ where $\hat{\Pi}_j := \mathcal{O}p_h^w(\hat{\pi}_j)$, we also use the similar but slightly different expansion of the symbol of $(P - i)^{-1}$ (cf. Remark 3.6). Since in Martinez and Sordani [18] they consider only the localized initial data and use the position of the particles as the coordinates these difficulties do not appear.

Using Π we have the following theorem.

Theorem 2.6. *If φ is the solution of $ih\partial_t\varphi = P\varphi$ with initial data φ_0 satisfying $\chi(P)\varphi_0 = \varphi_0$ for some $\chi \in C_0^\infty(\mathbb{R})$ such that $\chi\Phi = \chi$, then*

$$\varphi = e^{-itP_1/h}\Pi\varphi_0 + e^{-itP_2/h}(1 - \Pi)\varphi_0 + \mathcal{O}(|t|h^\infty\|\varphi_0\|), \quad (2.4)$$

where $P_1 := \Pi P \Pi$ and $P_2 := (1 - \Pi)P(1 - \Pi)$ are self-adjoint on a domain containing $D(P)$.

We can reduce the evolution $e^{-itP_1/h}$ to an evolution on the L^2 space of only nuclear variables. To state the next result, we use h -admissible operators. For the definitions of pseudodifferential and h -admissible operators see the the “Appendix”.

Theorem 2.7. *There exists a h -admissible operator*

$$W: L^2(\mathbb{R}^{3(N+1)}) \rightarrow (L^2(\mathbb{R}_x^3))^{\oplus k}$$

with operator valued symbol and $k \times k$ self-adjoint matrix G of h -admissible operators on $L^2(\mathbb{R}_x^3)$ such that the restriction U of W to $\text{Ran}\Pi$:

$$U: \text{Ran}\Pi \rightarrow (L^2(\mathbb{R}_x^3))^{\oplus k}$$

is a unitary operator which satisfies

$$UP_1\Pi = GU\Pi,$$

so that $e^{-itP_1/h}\Pi = U^*e^{-itG/h}U\Pi$. The symbol $g(x, \xi)$ of G has the following form:

$$g(x, \xi) = \xi^2 I_k + \mu(x) + \sum_{j \geq 1} h^j g_j(x, \xi), \quad (2.5)$$

where I_k is the k -dimensional identity matrix and $\mu(x)$ is the matrix of $\Pi_0(x)P_e(x)$ in $(u_1(x), \dots, u_k(x))$. The component in the j th row and m th column of g_1 is

$$\begin{aligned} (g_1)_{jm} = & -2i \sum_{n=1}^3 (u_j(x), \partial_{x_n} u_m(x))_{L_y^2} \xi_n \\ & - 4e \sum_{n=1}^3 (u_j(x), (\sum_{\ell=2}^{N+1} A(y_\ell))_n u_m(x))_{L_y^2} \xi_n, \end{aligned} \quad (2.6)$$

where $(\sum_{\ell=2}^{N+1} A(y_\ell))_n$ is the n th component of $\sum_{\ell=2}^{N+1} A(y_\ell)$, and $g_j \in S^0(\mathbb{R}^3 \times \mathbb{R}^3)$, $j \geq 2$, so that $Op_h^w(g_j) \in \mathcal{L}(L^2(\mathbb{R}^{3(N+1)}))$, $j \geq 2$.

Remark 2.8. (1) The factor $(u_j(x), \partial_{x_n} u_m(x))$ in the first term of (2.6) corresponds to the additional vector potential A_{geo} in [25] which appears even if there is no magnetic field.

(2) Terms including the vector potential of the nucleus such as $-2heA(x)\xi$ do not appear in g . In particular if the external potentials do not exist (i.e., $V_i \equiv 0$), P_e and u_i are independent of x , so that the first term of (2.6) vanishes and the second term is independent of x . Actually higher order terms are also independent of x in this case since the

Hamiltonian P does not include terms depending on x . Thus, the Hamilton flow of g is a straight line. This is not obvious because the internal and external motions are not separated in magnetic fields.

- (3) When more than one nucleus exists, these results with the operator G without vector potential term does not seem to hold because each electron is possessed by more than one nucleus at the same time and the electrons around each nucleus does not necessarily offset the influence of magnetic fields.

The next result is concerned with more explicit expression of the solution of the Schrödinger equation for a special initial data. We assume $k = 1$ and set $\tilde{u}_1(x) := \mathcal{V}^* u_1(x)$. Then $\tilde{u}_1(x) \in \text{Ran} \tilde{\Pi}_0(x)$ where

$$\tilde{\Pi}_0(x) := \mathcal{V}^* \Pi_0(x) \mathcal{V},$$

which is the spectral projection of $\tilde{P}_e(x) = \sum_{i=2}^{N+1} \tilde{L}_i(x)^2 + V_0(x, y)$ corresponding to $\sigma_0(x)$.

Let $\alpha_t = (x_t, \xi_t)$ be the solution of

$$\dot{x}_t = \frac{\partial g}{\partial \xi}(x_t, \xi_t), \quad \dot{\xi}_t = -\frac{\partial g}{\partial x}(x_t, \xi_t) \quad (2.7)$$

starting from initial data $\alpha_0 = (x_0, \xi_0)$. Let (η_n, ζ_n) , $n = 1, 2, 3$ be the independent solutions of

$$\begin{pmatrix} \dot{\eta} \\ \dot{\zeta} \end{pmatrix} = J M_t \begin{pmatrix} \eta \\ \zeta \end{pmatrix}, \quad (2.8)$$

with initial data

$$(\eta_n)_j|_{t=0} = \delta_{jn}, \quad (\zeta_n)_j|_{t=0} = i\delta_{jn},$$

where $(\eta_n)_j$ is the j th component of η_n ,

$$J = \begin{pmatrix} 0 & \mathbb{I} \\ -\mathbb{I} & 0 \end{pmatrix},$$

$\mathbb{I}$ being the unit matrix and M_t is the Hessian of g at α_t :

$$(M_t)_{i,j} = \left(\frac{\partial^2 g}{\partial \alpha^2} \right)_{i,j} \Big|_{\alpha=\alpha_t}.$$

By Theorems 2.6 and 2.7 as in Martinez and Sordani [18, Theorem 11.3] (see also Combescure and Robert [3]), we have the following theorem.

Theorem 2.9. *Let $k = 1$ and $\tilde{\varphi}_0 \in L^2(\mathbb{R}^{3(N+1)})$ be as follows*

$$\tilde{\varphi}_0 = (\pi h)^{-3/4} \tilde{\Pi} \chi(\tilde{P}) (e^{ix\xi_0/h - (x-x_0)^2/2h} \tilde{u}_1(x)),$$

where $\tilde{\Pi} = \mathcal{V}^* \Pi \mathcal{V}$, and $\chi = 1$ near $\xi_0^2 + \mu(x_0)$. Then there exists $C > 0$ such that for any integer $J \geq 1$ one has

$$e^{-it\tilde{P}/h}\tilde{\varphi}_0 = (\pi h)^{-3/4}(e^{ix\xi_t/h-(x-x_t)^2/2h}\tilde{u}_1(x)) \\ + h^{1/2}e^{i\delta_t/h}\sum_{\mu=0}^{\ell_J}c_\mu(t;h)\phi_{\ell,t}\tilde{v}_\ell(x) + \mathcal{O}(h^{J/4}), \quad (2.9)$$

where

$$c_\ell(t;h) = \sum_{j=0}^{j_\ell} h^{j/2} c_{\ell,j}(t),$$

$c_{\ell,j}$ are given by the polynomials with respect to $\partial^\gamma g(x_t, \xi_t)$ and $\operatorname{Re}\eta_n$, $\operatorname{Im}\eta_n$, $\operatorname{Re}\zeta_n$, $\operatorname{Im}\zeta_n$, $1 \leq n \leq 3$, $\delta_t := \int_0^t (\dot{x}_s \xi_s - g(x_s, \xi_s)) ds + (x_0 \xi_0 - x_t \xi_t)/2$, $\tilde{v}_\ell \in C^\infty(\mathbb{R}^3; L_y^2(\mathbb{R}^{3N}))$ and $\phi_{\ell,t}$ is a generalized coherent state centered at (x_t, ξ_t) . The estimate is uniform with respect to (t, h) such that $h > 0$ is small enough and $t < C^{-1} \ln \frac{1}{h}$.

- Remark 2.10.* (1) Since in the proof of Theorem 2.9 we apply the stationary phase method for pseudodifferential operators with operator valued symbols (see the proof of [18, Theorem 11.3]), the factor $\tilde{v}_\mu(x)$ consists of the terms like $a(x)\partial_x^\alpha \tilde{u}_1(x)$ with $a(x) \in C^\infty(\mathbb{R}_x^3; \mathcal{L}(L_y^2(\mathbb{R}_y^{3N})))$ which is not in $\operatorname{Ran}\Pi_0$ in general.
- (2) For $k > 1$ the propagation of wave packets through the crossing and avoided crossing of the eigenvalues of the electronic Hamiltonian is also studied by Hagedorn [10] and Hagedorn and Joye [11].

3. Proof of the Main Results

Our method is based on the construction of the almost invariant subspace as in Nenciu and Sordani [21] and Martinez and Sordani [17, 18] but there are difficulties about boundedness of operators which come from the unboundedness of the vector potential and the change of the coordinates. The method of Spohn and Teufel [25] is different from ours and based on rather direct comparison of the evolutions.

The outline of the construction of Π can be illustrated as follows. First we obtain the expression of the operator valued symbol of the resolvent $(P - z)^{-1}$ of P as formal power series $\sum_{j=0}^\infty q_j(x, \xi; z)h^j$ with respect to h . Secondly we integrate each q_j with respect to z along the loop $\Gamma(x, \xi)$ enclosing the set $\{z: z - \xi^2 \in \sigma_0(x)\}$ and denote it by $\hat{\pi}_j(x, \xi)$. Thirdly we quantize $\hat{\pi}_j$ and multiply by the cutoff function $\Phi(P)$ to make each term bounded. Fourthly we make a resummation and symmetrize the operator to obtain a operator $\hat{\Pi}_\Phi$. Finally integrating the resolvent of $\hat{\Pi}_\Phi$ along the loop enclosing the point $z = 1$, we obtain Π .

3.1. Definition of the Formal Symbol of the Resolvent and the Projection Π

We denote by $p(x, \xi; h) = \xi^2 + \sum_{i=2}^{N+1} L_i^2 - 4he \sum_{i=2}^{N+1} A(y_i)\xi + h^2 Q + V(x, y)$ the operator valued symbol of P and by $p_0(x, \xi) := \xi^2 + \sum_{i=2}^{N+1} L_i^2 + V_0(x, y)$ its principal symbol. Let $\gamma(x)$ be the continuous loop as in Remark 2.3. Let us

consider $\Omega := \{(x, \xi, z) \in \mathbb{R}^6 \times \mathbb{C} ; z - \xi^2 \in \gamma(x)\}$. For $(x, \xi, z) \in \Omega$, $p_0(x, \xi) - z$ is invertible and $q_0 := (p_0 - z)^{-1}$ is smooth and bounded. We define the symbol

$$r(x, \xi; h, z) = \sum_{j \geq 1} r^j(x, \xi; z) h^j,$$

as in Sordoni [24] (see also Nenciu and Sordoni [21]) by

$$(p(x, \xi; h) - z) \# q_0(x, \xi; z) = 1 - r(x, \xi; h, z), \quad (3.1)$$

where

$$a(x, \xi) \# b(x, \xi) = \sum_{\alpha, \beta} \frac{h^{|\alpha+\beta|} (-1)^{|\alpha|}}{(2i)^{|\alpha+\beta|} \alpha! \beta!} (\partial_x^\alpha \partial_\xi^\beta a(x, \xi)) (\partial_\xi^\alpha \partial_x^\beta b(x, \xi)),$$

at a formal series level. We define

$$q(x, \xi; h, z) = \sum_{j \geq 1} q^j(x, \xi; z) h^j,$$

by

$$q := q_0 + q_0 \# \sum_{j \geq 1} r^{\#j}.$$

Here $r^{\#j} = r \# \dots \# r$ where the number of $\#$ is j . One can check that, for $j \geq 1$, q_j is given by the sum of terms of the following form

$$C q_0 \prod_{i=1}^m ((\partial_x^{\alpha_i} \partial_\xi^{\beta_i} s_i) q_0), \quad (3.2)$$

where C is a constant, $1 \leq m \leq 2j$, $|\sum_{i=1}^m \alpha_i| = |\sum_{i=1}^m \beta_i| \leq j$ and s_i is one of the following

$$\begin{aligned} \xi^\alpha, \quad |\alpha| = 1, \quad \sum_{i=2}^{N+1} A(y_i) \xi, \quad L_i A(y_i), \quad \left(\sum_{i=2}^{N+1} L_i \right)^2, \quad \left(\sum_{i=2}^{N+1} eA(y_i) \right)^2, \\ \partial_x^\gamma V_1(x) \left(\sum_{i=2}^{N+1} y_i \right)^\gamma + \sum_{i=2}^{N+1} \partial_x^\gamma V_i(x + y_i) \left(\sum_{i=2}^{N+1} y_i \right)^\gamma, \quad \gamma \in \mathbb{N}^3. \end{aligned} \quad (3.3)$$

Since the operators do not commute with each other, the order of the product is specified as $\prod_{i=1}^m ((\partial_x^{\alpha_i} \partial_\xi^{\beta_i} s_i) q_0) = (\partial_x^{\alpha_1} \partial_\xi^{\beta_1} s_1) q_0 \dots (\partial_x^{\alpha_m} \partial_\xi^{\beta_m} s_m) q_0$. Let us set

$$\Gamma(x, \xi) := \{z \in \mathbb{C}; z - \xi^2 \in \gamma(x)\},$$

and let us define

$$\hat{\pi}_j(x, \xi) = \frac{i}{2\pi} \oint_{\Gamma(x, \xi)} q_j(x, \xi; z) dz. \quad (3.4)$$

Then by the change of the variable $z \rightarrow z - \xi^2$, we see $\hat{\pi}_0(x, \xi) = \Pi_0(x)$.

3.2. Quantization of the Symbols and Boundedness Properties

To determine the symbol classes of operators we need the boundedness of $R_e(x; z) = (P_e(x) - z)^{-1}$ multiplied by some operators. We shall also estimate $q'_0(x, \xi; h) := (p'(x, \xi) - i)^{-1}$ where $p'(x, \xi) := p_0(x, \xi) + h^2 Q_1$ which are needed later. Here $Q_1 = \frac{1}{1-Nh^2} \left(\sum_{i=2}^{N+1} L_i \right)^2$ is introduced in Sect. 2. If a potential V in $\mathbb{R}^3$ is Δ -bounded with relative bound ϵ , then $V(y_i)$ and $V(y_i - y_j)$ are relatively bounded with respect to $\sum_{i=2}^{N+1} L_i^2$ with relative bound ϵ (see e.g., [22]). Thus we see easily the following lemma.

Lemma 3.1. *Let $R_e(x; z)$ and $q'_0(x, \xi; h)$ be as above. Then the following conclusions hold:*

(i) *There exists a positive constant C such that*

$$\|(L_i)_k R_e(x; z)\|_{\mathcal{L}(L^2(\mathbb{R}_y^{3N}))} + \|(L_i)_k (L_j)_\ell R_e(x; z)\|_{\mathcal{L}(L^2(\mathbb{R}_y^{3N}))} \leq C,$$

for all $(x, z) \in \mathbb{R}^3 \times \mathbb{C}$ satisfying $z \in \gamma(x)$ where $(L_i)_k$, $k = 1, 2, 3$ denotes the k th component of L_i .

(ii) *There exist positive constants C and h_0 such that*

$$\|(L_i)_k (L_j)_\ell q'_0(x, \xi; h)\|_{\mathcal{L}(L^2(\mathbb{R}_y^{3N}))} \leq C,$$

$$\|(L_i)_k q'_0(x, \xi; h)\|_{\mathcal{L}(L^2(\mathbb{R}_y^{3N}))} \leq C(1 + |\xi|)^{-1},$$

$$\|q'_0(x, \xi; h)\|_{\mathcal{L}(L^2(\mathbb{R}_y^{3N}))} \leq C(1 + |\xi|)^{-2},$$

for all $(x, \xi, h) \in \mathbb{R}^3 \times \mathbb{R}^3 \times (0, h_0]$.

We define for $m \in \mathbb{N}$ and $s \in \mathbb{R}$ the weighted magnetic Sobolev space:

$$H_{A,y}^{m,s} = \left\{ u \in L_{\text{loc}}^2(\mathbb{R}_y^{3N}) : ((1 + |y|^2)^{s/2} L_2^{\alpha_2} \dots L_{N+1}^{\alpha_{N+1}} u \in L^2(\mathbb{R}_y^{3N}), \right. \\ \left. \alpha_i \in \mathbb{N}^3, \sum_{i=2}^{N+1} |\alpha_i| \leq m \right\},$$

(see [6, Appendix A]). To quantize $\hat{\pi}_j$ we need to prove that they satisfy the conditions of the symbols (see the ‘‘Appendix’’).

Lemma 3.2. *For any $j \in \mathbb{N}$ and $r, s \in \mathbb{R}$, $\hat{\pi}_j(x, \xi) \in S^j(\mathbb{R}^3 \times \mathbb{R}^3; \mathcal{L}(L_y^{2,r}, L_y^{2,r+s}))$ and $p\#\hat{\pi}_j \in S^{j+2}(\mathbb{R}^3 \times \mathbb{R}^3; \mathcal{L}(L_y^{2,r}, L_y^{2,r+s}))$ where $L_y^{2,s} := \{u : (1 + |y|^2)^{s/2} u \in L_y^2\}$.*

Proof. By (3.2) q_j can be written as a sum of the terms of the following form

$$C_{q_0} \left(\prod_{i=1}^m (y^{\alpha_i} a_i q_0) \right) \xi^\beta, \quad (3.5)$$

where C is a constant, $\alpha_i \in \mathbb{N}^{3N}$, $\beta \in \mathbb{N}^3$ and $a_i R_e(x; z) \in \mathcal{L}(L^2(\mathbb{R}_y^{3N}))$ uniformly with respect to $(x, z) \in \mathbb{R}^3 \times \mathbb{C}$ satisfying $z \in \gamma(x)$. It is easy to see

$|\beta| \leq j$ from the construction of q_j . By the change of the variable $z \rightarrow z - \xi^2$, $\oint_{\Gamma(x, \xi)} q_0 \prod_{i=1}^m (y^{\alpha_i} a_i q_0) dz$ becomes

$$\oint_{\gamma(x)} R_e(x; z) \prod_{i=1}^m (y^{\alpha_i} a_i R_e(x; z)) dz.$$

We shall prove $\hat{\pi}_j(x, \xi) \in S^j(\mathbb{R}^3 \times \mathbb{R}^3; \mathcal{L}(L_y^{2,r}, H_{A,y}^{2,r+s})) \subset S^j(\mathbb{R}^3 \times \mathbb{R}^3; \mathcal{L}(L_y^{2,r}, L_y^{2,r+s}))$. Thus we need to show for any $\alpha_0 \in \mathbb{N}^{3N}$ there exists a constant $C > 0$ such that for all $x \in \mathbb{R}^3$

$$\begin{aligned} & \left\| y^{\alpha_0} a_0 \oint_{\gamma(x)} R_e(z) \prod_{i=1}^m (y^{\alpha_i} a_i R_e(z)) dz \right\|_{L_y^2} \\ &= \left\| \oint_{\gamma(x)} \prod_{i=0}^m (y^{\alpha_i} a_i R_e(z)) dz \right\|_{L_y^2} \leq C, \end{aligned}$$

where $a_0 = 1$, $(L_i)_k$ or $(L_i)_k(L_j)_\ell$. For this purpose denoting $(y^{\alpha_i} a_i R_e(z))$ by $b_i(z)$, we prove the following by induction:

$$\begin{aligned} & \oint_{\gamma(x)} \prod_{i=0}^m b_i(z) dz \\ &= \sum_{j=0}^{\ell-1} \left(\frac{i}{2\pi} \right)^j \oint_{\gamma(x)} \oint_{\gamma_0(x)} \cdots \oint_{\gamma_{j-1}(x)} \left(\prod_{n=0}^{j-1} \frac{1}{z - z_n} b_n(z_n) \right) \\ & \quad \cdot (b_j(z) \Pi_0) \left(\prod_{i=j+1}^m b_i(z) \right) dz dz_0 \cdots dz_j \\ & \quad + \left(\frac{i}{2\pi} \right)^\ell \oint_{\gamma(x)} \oint_{\gamma_0(x)} \cdots \oint_{\gamma_{\ell-1}(x)} \left(\prod_{n=0}^{\ell-1} \frac{1}{z - z_n} b_n(z_n) \right) \\ & \quad \cdot \left(\prod_{i=\ell}^m b_i(z) \right) dz dz_0 \cdots dz_{\ell-1}, \end{aligned} \quad (3.6)$$

for any $\ell = 0, \dots, m+1$ where $\gamma_j(x)$ is a slightly larger loop enclosing $\gamma(x)$ and z_j is a point on $\gamma_j(x)$. Here we suppose $\sum_{i=0}^{\ell-1} d_i = 0$, $\prod_{i=0}^{\ell-1} d_i = 1$, $\prod_{i=m+1}^m d_i = 1$, $\oint_{\gamma(x)} \oint_{\gamma_0(x)} \cdots \oint_{\gamma_{j-1}(x)} = \oint_{\gamma(x)}$, and $dz dz_0 \cdots dz_{j-1} = dz$. For $\ell = 0$ since the right hand side is $\oint_{\gamma(x)} \prod_{i=0}^m b_i(z) dz$, the equation holds.

We denote $\oint_{\gamma(x)} \oint_{\gamma_0(x)} \cdots \oint_{\gamma_{j-1}(x)}$ by $\oint_j$ and $dz dz_0 \cdots dz_{j-1}$ by dz'_j . Set $B_\ell = \prod_{n=0}^{\ell-1} \frac{1}{z - z_n} b_n(z_n)$ and $D_\ell = \prod_{i=\ell}^m b_i(z)$. Then (3.6) is rewritten as

$$\oint_{\gamma(x)} D_0 dz = \sum_{j=0}^{\ell-1} \left(\frac{i}{2\pi} \right)^j \oint_j B_j(b_j(z) \Pi_0) D_{j+1} dz'_{j+1} + \left(\frac{i}{2\pi} \right)^\ell \oint_\ell B_\ell D_\ell dz'_\ell. \quad (3.7)$$

Supposing (3.7) holds for $\ell = \ell_0 < m + 1$ we prove (3.7) for $\ell = \ell_0 + 1$. We rewrite the last term in (3.7). Using $\Pi_0(x) = \frac{i}{2\pi} \oint_{\gamma_{\ell_0+1}(x)} R_e(z_{\ell_0+1}) dz_{\ell_0+1}$, by the resolvent equation we have

$$\begin{aligned} & \oint_{\ell_0} B_{\ell_0}(y^{\alpha_{\ell_0}} a_{\ell_0} R_e(z) \Pi_0) D_{\ell_0+1} dz'_{\ell_0} \\ &= \oint_{\ell_0} B_{\ell_0} y^{\alpha_{\ell_0}} a_{\ell_0} \frac{i}{2\pi} \oint_{\gamma_{\ell_0+1}(x)} \frac{1}{z - z_{\ell_0+1}} (R_e(z) - R_e(z_{\ell_0+1})) D_{\ell_0+1} dz'_{\ell_0} dz_{\ell_0+1}. \end{aligned}$$

Since $\gamma_{\ell_0+1}(x)$ lies outside $\gamma(x)$, we have

$$\frac{i}{2\pi} \oint_{\gamma_{\ell_0+1}(x)} \frac{1}{z - z_{\ell_0+1}} dz_{\ell_0+1} = 1.$$

Thus we have

$$\begin{aligned} & \oint_{\ell_0} B_{\ell_0}(y^{\alpha_{\ell_0}} a_{\ell_0} R_e(z) \Pi_0) D_{\ell_0+1} dz'_{\ell_0} \\ &= \oint_{\ell_0} B_{\ell_0} D_{\ell_0} dz'_{\ell_0} - \frac{i}{2\pi} \oint_{\ell_0+1} B_{\ell_0+1} D_{\ell_0+1} dz'_{\ell_0+1}. \end{aligned}$$

Transposing the last term we have

$$\begin{aligned} & \oint_{\ell_0} B_{\ell_0} D_{\ell_0} dz'_{\ell_0} \\ &= \oint_{\ell_0} B_{\ell_0}(y^{\alpha_{\ell_0}} a_{\ell_0} R_e(z) \Pi_0) D_{\ell_0+1} dz'_{\ell_0} + \frac{i}{2\pi} \oint_{\ell_0+1} B_{\ell_0+1} D_{\ell_0+1} dz'_{\ell_0+1}. \end{aligned}$$

Substituting this equation into the last term of (3.7) for $\ell = \ell_0$, we obtain (3.7) for $\ell = \ell_0 + 1$.

For $\ell = m + 1$, the last term of (3.6) is written as

$$\left(\frac{i}{2\pi}\right)^{m+1} \oint_m \left(\prod_{n=0}^m \frac{1}{z - z_n}\right) \left(\prod_{n=0}^m b_n(z_n)\right) dz'_m.$$

Since $\gamma_i(x)$ lies outside $\gamma(x)$, the integration with respect to z vanishes.

By the assumption (H3) $y^\beta \Pi_0(x)$ is a bounded operator on $L^2(\mathbb{R}_y^{3N})$ uniformly for x . By

$$[(y_i)_k, R_e(x; z)] = R_e(x; z)[P_e, (y_i)_k] R_e(x; z) = -2i R_e(x; z)(L_i)_k R_e(x; z),$$

and Lemma 3.1 the remaining term of (3.6) are the sum of terms of the following form;

$$\begin{aligned} & C \oint_j \left(\prod_{n=0}^{j-1} \frac{1}{z - z_n} \left(\prod_{i=1}^{i_n} c_{n,i} R_e(x; z_n)\right)\right) \left(\prod_{i=1}^r d_i R_e(x; z_n)\right) \\ & \cdot y^\beta \Pi_0(x) y^\delta \left(\prod_{i=1}^s e_i R_e(x; z)\right) dz'_j, \end{aligned} \quad (3.8)$$

where C is a constant, $y^\beta \Pi_0(x) y^\delta$ is bounded uniformly with respect to x . Here $c_{n,j} R_e(x; z)$, $d_i R_e(x; z)$ and $e_i R_e(x; z)$ are bounded uniformly with respect

to x , $z \in \gamma(x)$ and $(z - z_n)^{-1}$ is uniformly bounded with respect to $z \in \gamma(x)$, $z_n \in \gamma_n(x)$. Therefore (3.8) is bounded uniformly with respect to x . Thus by (3.5), we have $\|y^{\alpha_0} a_0 \hat{\pi}_j(x, \xi)\|_{L_y^2} < C(1 + |\xi|)^j$ uniformly with respect to x , where $C > 0$ is a positive constant. In the same way we can see for all $\alpha \in \mathbb{N}^{3N}$, $\beta, \delta \in \mathbb{N}^3$ and $a = 1$, $(L_i)_k$ or $(L_i)_k(L_j)_\ell$, there exists a constant $C_{\alpha, \beta, \delta}$ such that $\|y^\alpha a \partial_x^\beta \partial_\xi^\delta \hat{\pi}_j(x, \xi)\|_{L_y^2} < C_{\alpha, \beta, \delta}(1 + |\xi|)^{j - |\delta|}$. Thus, $\hat{\pi}_j \in S^j(\mathbb{R}^3 \times \mathbb{R}^3; \mathcal{L}(L_y^{2, r}, H_{A, y}^{2, r+s}))$ for any $r, s \in \mathbb{R}$. Since $p \in S^2(\mathbb{R}^3 \times \mathbb{R}^3; \mathcal{L}(H_{A, y}^{2, r+2}, L_y^r))$ for any $r \in \mathbb{R}$, we have $p \# \hat{\pi}_j \in S^{j+2}(\mathbb{R}^3 \times \mathbb{R}^3; \mathcal{L}(L_y^{2, r}, L_y^{2, r+s}))$ for any $r, s \in \mathbb{R}$. This completes the proof of the lemma. $\square$

Remark 3.3. Since in Martinez and Sordani [18] they use the coordinates where independent variables are positions of the particles, the power of y appears only in the electric Hamiltonian and is not included in q_j . On the other hand in our coordinate the power of y is included in $4h^2 e \sum_{i=2}^{N+1} A(y_i) D_x Q$, and the Taylor expansion of V_i . Thus in our case the power of y in q_j becomes of higher order as j increases, so that the boundedness of $\hat{\pi}_j$ is not obvious.

We define

$$\hat{\Pi}_j = Op_h^w(\hat{\pi}_j(x, \xi)).$$

The operator $\hat{\Pi}_j$ is not bounded but by a localization in energy we obtain a bounded operator. To prove the boundedness of $\Phi(P)\hat{\Pi}_j$ we need the expansion of the resolvent $(P - i)^{-1}$ by the pseudodifferential operators with operator valued symbols as in the construction of q above. However, since $\left(\sum_{i=2}^{N+1} L_i^2\right)^2 q_0(x, \xi) \in S^0(\mathbb{R}^3 \times \mathbb{R}^3; \mathcal{L}(L_y^2, L_y^2))$ is not in the class $S^{-1}(\mathbb{R}^3 \times \mathbb{R}^3; \mathcal{L}(L_y^2, L_y^2))$, the decay with respect to ξ of the symbols of the higher order terms of the expansion is not obtained. To obtain the decay we use $q'_0(x, \xi; h)$ defined before Lemma 3.1 instead of q_0 as first term of the expansion. By Lemma 3.1 it is easy to see that the operator valued symbol q'_0 is in the following symbol classes.

Proposition 3.4. *The operator valued symbol q'_0 is in the symbol classes $S^{-2}(\mathbb{R}^3 \times \mathbb{R}^3; \mathcal{L}(L_y^2, L_y^2))$, $S^{-1}(\mathbb{R}^3 \times \mathbb{R}^3; \mathcal{L}(L_y^2, H_{A, y}^1))$ and $S^0(\mathbb{R}^3 \times \mathbb{R}^3; \mathcal{L}(L_y^2, H_{A, y}^2))$.*

Proof. Since

$$\partial_{x_k} q'_0(x, \xi; h) = -q'_0(x, \xi; h) \partial_{x_k} V_0 q'_0(x, \xi; h),$$

and

$$\partial_{\xi_k} q'_0(x, \xi; h) = -\xi_k (q'_0(x, \xi; h))^2,$$

the proposition follows from Lemma 3.1. $\square$

Lemma 3.5. *There exists a constant $h_0 > 0$ such that for any $\Phi \in C_0^\infty(\mathbb{R})$, $\Phi(P)\hat{\Pi}_j$ and $\Phi(P)\hat{\Pi}_j P$ are bounded in $L^2(\mathbb{R}^{3(N+1)})$ uniformly with respect to $h \in (0, h_0]$.*

Proof. We can write

$$\Phi(P)\hat{\Pi}_j = \Phi(P)(P-i)^j(P-i)^{-j}\hat{\Pi}_j, \quad (3.9)$$

where $\Phi(P)(P-i)^j$ is bounded uniformly with respect to h . By Proposition 3.4 we have

$$(p(x, \xi; h) - i) \# q'_0(x, \xi; h) = 1 - r'(x, \xi; h),$$

where $r'(x, \xi; h) \in S^{-1}(\mathbb{R}^3 \times \mathbb{R}^3; \mathcal{L}(L_y^{2,s}, L_y^{2,s-2}))$ for any $s \in \mathbb{R}$. Since

$$(1 - r') \# \left(\sum_{i=0}^{j-1} r'^{\#i} \right) = 1 - r'^{\#j},$$

we have

$$(P-i)Op_h^w \left(q'_0(x, \xi) \# \left(\sum_{i=0}^{j-1} r'^{\#i} \right) \right) = 1 - Op_h^w(r'^{\#j}).$$

Multiplying the both sides of the equation by $(P-i)^{-1}$ from left, we obtain

$$\begin{aligned} (P-i)^{-1} &= (P-i)^{-1}Op_h^w(r'^{\#j}) + Op_h^w \left(q'_0(x, \xi) \# \left(\sum_{i=0}^{j-1} r'^{\#i} \right) \right) \\ &= (P-i)^{-1}Op_h^w(a) + Op_h^w(b), \end{aligned}$$

where $a \in S^{-j}(\mathbb{R}^3 \times \mathbb{R}^3; \mathcal{L}(L_y^{2,r}, L_y^{2,r-2j}))$ and $b \in S^{-2}(\mathbb{R}^3 \times \mathbb{R}^3; \mathcal{L}(L_y^{2,r}, L_y^{2,r-2(j-1)}))$ for any $r \in \mathbb{R}$. Hence we have

$$(P-i)^{-j}\hat{\Pi}_j = \sum_{k=0}^{j-1} (P-i)^{-j+k}Op_h^w(a)(Op_h^w(b))^k\hat{\Pi}_j + (Op_h^w(b))^j\hat{\Pi}_j. \quad (3.10)$$

Since for any $r \in \mathbb{R}$

$$a \# b^{\#k} \in S^{-j-2k}(\mathbb{R}^3 \times \mathbb{R}^3; \mathcal{L}(L_y^{2,r}, L_y^{2,r-2j-2k(j-1)})),$$

and

$$b^{\#j} \in S^{-2j}(\mathbb{R}^3 \times \mathbb{R}^3; \mathcal{L}(L_y^{2,r}, L_y^{2,r-2j(j-1)})),$$

by Lemma 3.2 we have

$$a \# b^{\#k} \# \hat{\Pi}_j \in S^0(\mathbb{R}^3 \times \mathbb{R}^3; \mathcal{L}(L_y^2)),$$

and

$$b^{\#j} \# \hat{\Pi}_j \in S^0(\mathbb{R}^3 \times \mathbb{R}^3; \mathcal{L}(L_y^2)).$$

Thus by Theorem A.4 it follows that

$$\begin{aligned} Op_h^w(a)(Op_h^w(b))^k\hat{\Pi}_j &\in \mathcal{L}(L^2(\mathbb{R}_x^3; L^2(\mathbb{R}_y^{3N}))) \\ &= \mathcal{L}(L^2(\mathbb{R}^{3(N+1)})), \end{aligned}$$

and

$$(Op_h^w(b))^k\hat{\Pi}_j \in \mathcal{L}(L^2(\mathbb{R}^{3(N+1)})). \quad (3.11)$$

By (3.9)–(3.11), we have $\Phi(P)\hat{\Pi}_j \in \mathcal{L}(L^2(\mathbb{R}^{3(N+1)}))$

To prove the boundedness of $\Phi(P)\hat{\Pi}_jP$ we note that the symbol of $\hat{\Pi}_jP$ is $(p\#\hat{\pi}_j)^*$ where a^* is the adjoint of an operator a on y variables. Since by Lemma 3.2 we have $(p\#\hat{\pi}_j)^* \in S^{j+2}(\mathbb{R}^3 \times \mathbb{R}^3; \mathcal{L}(L_y^{2,r}, L_y^{2,r+s}))$ for any $r, s \in \mathbb{R}$, the boundedness of $\Phi(P)\hat{\Pi}_jP$ follows from the same argument as $\Phi(P)\hat{\Pi}_j$ replacing j in (3.9) with $j+1$. $\square$

Remark 3.6. (1) In Martinez and Sordani [18], they considered only localized initial data, so that unboundedness of the vector potential term is not important. On the other hand, in our case we do not restrict the initial data and therefore, because of the existence of the vector potential terms in $\hat{P}$ in (1.1), D_x is not P -bounded. Thus, to prove $(P-i)^{-j}\hat{\Pi}_j$ is bounded, we need to expand the resolvent at the expense of the presence of the power of y . Since $\hat{\pi}_j$ includes $\Pi_0(x)$, the power of y does not prevent $(P-i)^{-j}\hat{\Pi}_j$ from being bounded.

(2) If we use q_0 instead of q'_0 as the first term of the symbol of the resolvent as in (3.1), r includes the term

$$h^2Q_1q_0 = h^2 \frac{1}{1-Nh^2} \left(\sum_{i=2}^{N+1} L_i \right)^2 q_0,$$

which is not in the symbol class $S^{-1}(\mathbb{R}^3 \times \mathbb{R}^3; \mathcal{L}(L_y^2, L_y^2))$. However, when we use q'_0 , since $(p_0 + h^2Q_1 - i)q'_0 = 1$, such a term without decay with respect to ξ does not appear.

3.3. Definition of Π and Proof of Theorem 2.5

To construct Π we make a resummation. Let $\rho \in C_0^\infty(\mathbb{R})$ be a function such that $0 \leq \rho \leq 1$, $\text{supp } \rho \subset [-2, 2]$ and $\rho = 1$ on $[-1, 1]$. It is easy to see that $P\hat{\Pi}_j\Phi(P)$ and $\Phi(P)\hat{\Pi}_jP$ are bounded in $L^2(\mathbb{R}^{3(N+1)})$ and there exists a decreasing sequence of positive numbers $(\epsilon_j)_{j \in \mathbb{N}}$ converging to zero such that for any $j \in \mathbb{N}$

$$\left(1 - \rho\left(\frac{\epsilon_j}{h}\right)\right) (\|\hat{\Pi}_j\Phi(P)\| + \|\Phi(P)\hat{\Pi}_j\| + \|P\hat{\Pi}_j\Phi(P)\| + \|\Phi(P)\hat{\Pi}_jP\|) \leq h^{-1}.$$

Let us define

$$\begin{aligned} \hat{\Pi}\Phi(P) &:= \Pi_0\Phi(P) + \sum_{j \geq 1} \left(1 - \rho\left(\frac{\epsilon_j}{h}\right)\right) \hat{\Pi}_j\Phi(P)h^j, \\ \Phi(P)\hat{\Pi} &:= \Phi(P)\Pi_0 + \sum_{j \geq 1} \left(1 - \rho\left(\frac{\epsilon_j}{h}\right)\right) \Phi(P)\hat{\Pi}_jh^j. \end{aligned}$$

Then it is easy to prove that for any $K \in \mathbb{N}$ there exists $C_N > 0$ such that

$$\left\| \hat{\Pi}\Phi(P) - \sum_{j=0}^K h^j \hat{\Pi}_j\Phi(P) \right\| + \left\| \Phi(P)\hat{\Pi} - \sum_{j=0}^K h^j \Phi(P)\hat{\Pi}_j \right\| \leq C_K h^{K+1}. \quad (3.12)$$

and

$$\left\| P\hat{\Pi}\Phi(P) - \sum_{j=0}^K h^j P\hat{\Pi}_j\Phi(P) \right\| + \left\| \Phi(P)\hat{\Pi}P - \sum_{j=0}^K h^j \Phi(P)\hat{\Pi}_jP \right\| \leq C_K h^{K+1}, \quad (3.13)$$

where we understand $\hat{\Pi}_0 = \Pi_0$. We define

$$\hat{\Pi}_\Phi := \Phi(P)\hat{\Pi} + (1 - \Phi(P))\hat{\Pi}\Phi(P) + (1 - \Phi(P))\Pi_0(1 - \Phi(P)).$$

Since $\sum_{j \geq 0} q_j h^j$ is formally the symbol of $(P - z)^{-1}$, by (3.12) and (3.13) we have the following lemma as in Sordoni [24].

Lemma 3.7. *Let $\chi \in C_0^\infty(\mathbb{R})$ satisfy $\chi\Phi = \chi$. Then for all $K \in \mathbb{N}$*

$$\begin{aligned} \|\chi(P)[\hat{\Pi}_\Phi, P]\| + \|[\hat{\Pi}_\Phi, P]\chi(P)\| &= \mathcal{O}(h^K), \\ \|\chi(P)(\hat{\Pi}_\Phi^2 - \hat{\Pi}_\Phi)\| + \|(\hat{\Pi}_\Phi^2 - \hat{\Pi}_\Phi)\chi(P)\| &= \mathcal{O}(h^K). \end{aligned}$$

Since $\hat{\Pi}_\Phi - \Pi_0 = \mathcal{O}(h)$, for sufficiently small h the spectrum of $\hat{\Pi}_\Phi$ is concentrated near 0 and 1, so that the set $\{z \in \mathbb{C}; |z - 1| = 1/2\}$ is in the resolvent set of $\hat{\Pi}_\Phi$ for sufficiently small h . We define

$$\Pi := \oint_{|z-1|=1/2} (\hat{\Pi}_\Phi - z)^{-1} dz.$$

Then as in Sordoni [24] we have

$$\Pi - \hat{\Pi}_\Phi = \frac{i}{2\pi} (\hat{\Pi}_\Phi^2 - \hat{\Pi}_\Phi) \oint_{|z-1|=1/2} (\hat{\Pi}_\Phi - z)^{-1} (2\hat{\Pi}_\Phi - 1) (1 - \hat{\Pi}_\Phi - z)^{-1} (1 - z)^{-1} dz.$$

Theorem 2.5 follows from this formula and Lemma 3.7.

3.4. Proof of Theorems 2.6 and 2.7

Proof of Theorem 2.6. First, we prove that $\Pi P \Pi$ and $(1 - \Pi)P(1 - \Pi)$ are self-adjoint on a domain containing $D(P)$. Since it can be proved that $P\Pi(P - i)^{-1}$ is bounded as in the proof of Lemma 3.5, we have $\Pi D(P) \in D(P)$. Hence $\Pi P \Pi$ and $(1 - \Pi)P(1 - \Pi)$ are defined on $D(P)$. Since $\hat{P}$ is lower semibounded, P is also lower semibounded. Thus $\Pi P \Pi$ and $(1 - \Pi)P(1 - \Pi)$ are lower semibounded so that there exist Friedrichs extensions of $\Pi P \Pi$ and $(1 - \Pi)P(1 - \Pi)$ with their domains containing $D(P)$.

Multiplying both side of $ih\partial_t\varphi = P\varphi$ by Π we have

$$ih\partial_t\Pi\varphi = P_1\Pi\varphi + R_1\varphi,$$

where $R_1 = \Pi[\Pi, P]\chi(P) = \mathcal{O}(h^\infty)$. This equation can be re-written as,

$$ih\partial_t(e^{itP_1/h}\Pi\varphi) = \mathcal{O}(h^\infty\|\varphi_0\|).$$

and thus integrating from 0 to t we obtain,

$$\Pi\varphi = e^{-itP_1/h}\Pi\varphi_0 + \mathcal{O}(|t|h^\infty\|\varphi_0\|).$$

In the same way we obtain

$$(1 - \Pi)\varphi = e^{-itP_1/h}(1 - \Pi)\varphi_0 + \mathcal{O}(|t|h^\infty \|\varphi_0\|).$$

Combining these two equations we obtain (2.4). $\square$

Proof of Theorem 2.7. Since $\Pi - \Pi_0 = \mathcal{O}(h)$, for h small enough the operator

$$\mathcal{U} := (\Pi_0\Pi + (1 - \Pi_0)(1 - \Pi))(1 - (\Pi_0 - \Pi)^2)^{-1/2} \quad (3.14)$$

can be defined. $\mathcal{U}$ is a unitary operator and it maps $\text{Ran}\Pi$ onto $\text{Ran}\Pi_0$ (see Kato [15, Chap. I.4]).

We can write $\Phi(P)\hat{\Pi}_j = \Phi(P)(P - i)^j(P - i)^{-j}\hat{\Pi}_j$. By the proof of Lemma 3.5 $(P - i)^{-j}\hat{\Pi}_j$ is a h -admissible operator with a symbol in $S^0(\mathbb{R}^3 \times \mathbb{R}^3; \mathcal{L}(L_y^2, L_y^{2,s}))$ for any $s \in \mathbb{R}$. Writing $\Phi'(x) := \Phi(x)(x - i)^j$ we have

$$\Phi'(P) = \frac{i}{2\pi} \int_{\mathbb{C}} \bar{\partial}_z \tilde{\Phi}'(z)(P - z)^{-1} dz d\bar{z}, \quad (3.15)$$

where $\tilde{\Phi}'(z)$ is an almost analytic expansion (see, e.g., Martinez [16]). Expanding $(P - z)^{-1}$ in (3.15) as in the proof of Lemma 3.5 we can see that $\Phi'(P)$ is h -admissible, so that $\hat{\Pi}_\Phi$ is a h -admissible operator on $L^2(\mathbb{R}_x^3; L_y^2)$. By the equation

$$\begin{aligned} \Pi &= \oint_{|z-1|=1/2} (z - \hat{\Pi}_\Phi)^{-1} dz \\ &= \oint_{|z-1|=1/2} (z - \Pi_0)^{-1} \sum_{k=0}^{\infty} ((\hat{\Pi}_\Phi - \Pi_0)(z - \Pi_0)^{-1})^k dz, \end{aligned} \quad (3.16)$$

we see that $\mathcal{U}$ is a h -admissible operator on $L^2(\mathbb{R}_x^3; L_y^2)$ and its principal symbol is 1.

We define W by

$$W\psi = (\mathcal{U}\psi, u_1(x, y))_{L_y^2(\mathbb{R}^{3N})} \oplus \cdots \oplus (\mathcal{U}\psi, u_k(x, y))_{L_y^2(\mathbb{R}^{3N})},$$

where $(u_1, \dots, u_k)$ is the orthonormal basis of $\text{Ran}\Pi_0$ in (H3). Since $W^*(\alpha_1 \oplus \cdots \oplus \alpha_k) = \mathcal{U}^*(\alpha_1 u_1 + \cdots + \alpha_k u_k)$ for $\alpha_i \in L^2(\mathbb{R}^3)$ we obtain

$$W^*W = \mathcal{U}^*\Pi_0\mathcal{U} = \Pi, \quad WW^* = 1,$$

which implies the unitarity of the restriction U of W to $\text{Ran}\Pi$.

Defining $G := UP_1U^*$, it is easy to see that G is a h -admissible operator on $H^2(\mathbb{R}_x^3; L_y^2)$. To calculate the symbol of G we need the following lemma.

Lemma 3.8. $\Pi_0(x)\hat{\Pi}_1\Pi_0(x)\Phi(P)$ is a h -admissible operator and the zeroth order term of its symbol with respect to h is 0.

Proof. Since $\Pi_0(x)\hat{\Pi}_1\Pi_0(x)$ is pseudodifferential operator and its symbol is in the class $S^1(\mathbb{R}^3 \times \mathbb{R}^3; \mathcal{L}(L^{2,r}, L^{2,r+s}))$, as in the proof of Theorem 2.7 we see that $\Pi_0(x)\hat{\Pi}_1\Pi_0(x)\Phi(P)$ is a h -admissible operator. The symbol of $\Pi_0(x)\hat{\Pi}_1\Pi_0(x)$

is $\Pi_0(x)\#\hat{\pi}_1\#\Pi_0(x)$ and its zeroth order term is $\Pi_0(x)\hat{\pi}_1\Pi_0(x)$. Thus it suffices to show $\Pi_0(x)\hat{\pi}_1\Pi_0(x) = 0$. By the direct calculation $\hat{\pi}_1$ is written as

$$\oint_{\gamma(x)} R_e(x; z) \left(4e \sum_{j=2}^{N+1} A(y_j) \xi + i(\partial_x V_0 \xi) R_e(x; z) - i R_e(x; z) (\partial_x V_0 \xi) \right) \cdot R_e(x; z) dz. \quad (3.17)$$

Let $u'_1(x), \dots, u'_k(x)$ be the eigenfunction with eigenvalues $E_1(x), \dots, E_k(x)$ and the orthonormal basis of $\text{Ran} \Pi_0(x)$. Then we have

$$\Pi_0(x)w(x) = \sum_{j=1}^k (w(x), u'_j(x))_{L_y^2} u'_j(x),$$

for $w(x) \in L^2(\mathbb{R}^3; L_y^2)$. Thus for an operator $S(x, \xi)$ on L_y^2 depending on x and ξ we have

$$\begin{aligned} & \Pi_0(x) \oint_{\gamma(x)} R_e(x; z)^n S(x, \xi) R_e(x; z)^m dz \Pi_0(x) w(x) \\ &= \oint_{\gamma(x)} \sum_{j=1}^k \sum_{\ell=1}^k (E_j - z)^{-n} (E_\ell - z)^{-m} dz \\ & \quad \cdot (S(x, \xi) u'_\ell(x), u'_j(x))_{L_y^2} (w(x), u'_\ell(x))_{L_y^2} u'_j(x). \end{aligned}$$

For $n, m \in \mathbb{N}$, $n + m \neq 1$ by the direct calculation we have $\oint_{\gamma(x)} \sum_{j=1}^k \sum_{\ell=1}^k (E_j - z)^{-n} (E_\ell - z)^{-m} dz = 0$. Therefore, by (3.17) we have $\Pi_0(x)\hat{\pi}_1\Pi_0(x) = 0$. $\square$

Set $\alpha = (\alpha_1, \dots, \alpha_k) \in L^2(\mathbb{R}_x^3)^{\oplus k}$, $\hat{\Pi}'_1 = \Phi(P)\hat{\Pi}_1 + (1 - \Phi(P))\hat{\Pi}_1\Phi(P)$, $P_0 = Op_h^w(p_0)$ and $P_1 = -4h^2e \sum_{i=2}^{N+1} A(y_i)D_x$. Using (3.14) and (3.16) we have

$$\begin{aligned} (G\alpha)_i &= \left(u_i, P_0 \sum_{j=1}^k \alpha_j u_j \right)_{L_y^2} + \left(u_i, P_1 \sum_{j=1}^k \alpha_j u_j \right)_{L_y^2} \\ &+ h \left(u_i, \hat{\Pi}'_1 P_0 \sum_{j=1}^k \alpha_j u_j \right)_{L_y^2} + h \left(u_i, P_0 \hat{\Pi}'_1 \sum_{j=1}^k \alpha_j u_j \right)_{L_y^2} \\ &+ h \left(u_i, (2\Pi_0(x)\hat{\Pi}'_1 - \hat{\Pi}'_1) P_0 \sum_{j=1}^k \alpha_j u_j \right)_{L_y^2} \\ &+ h \left(u_i, P_0 (2\Pi_0(x)\hat{\Pi}'_1 - \hat{\Pi}'_1) \sum_{j=1}^k \alpha_j u_j \right)_{L_y^2} + \mathcal{O}(h^2 \|\alpha\|_{(L_x^2)^{\oplus k}}). \end{aligned}$$

By $\Pi_0(x)u_i(x) = u_i(x)$, $[\Pi_0(x), \Phi(P)] = \mathcal{O}(h)$ and Lemma 3.8 we have

$$(G\alpha)_i = \left(u_i, P_0 \sum_{j=1}^k \alpha_j u_j \right)_{L_y^2} + \left(u_i, P_1 \sum_{j=1}^k \alpha_j u_j \right)_{L_y^2} + \mathcal{O}(h^2 \|\alpha\|_{(L_x^2)^{\oplus k}}).$$

From this (2.5) and (2.6) follow. For higher order terms, it is easy to see $g_j \in S^0(\mathbb{R}^3 \times \mathbb{R}^3; L^2(\mathbb{R}_y^{3N}))$, $j \geq 2$.

We shall prove G is self-adjoint with the domain $U(\text{Ran}\Pi \cap D(P_1))$. For this purpose we only need to prove P_1 is self-adjoint with the domain $\text{Ran}\Pi \cap D(P_1)$ as an operator on $\text{Ran}\Pi$ since U is unitary on $\text{Ran}\Pi$. Let $u, w \in \text{Ran}\Pi$ and suppose $(u, P_1 v) = (w, v)$ for all $v \in \text{Ran}\Pi \cap D(P_1)$. Then for any $\tilde{v} \in D(P_1)$, $(u, P_1 \Pi \tilde{v}) = (w, \Pi \tilde{v})$. Moreover since $P_1 = \Pi P \Pi$ and $w \in \text{Ran}\Pi$, we have $(u, P_1(1 - \Pi)\tilde{v}) = 0$ and $(w, (1 - \Pi)\tilde{v}) = 0$. Thus we have $(u, P_1 \tilde{v}) = (w, \tilde{v})$ for any $\tilde{v} \in D(P_1)$. Since P_1 is self-adjoint, we see $u \in D(P_1)$ and therefore $u \in \text{Ran}\Pi \cap D(P_1)$. Thus P_1 is self-adjoint with the domain $\text{Ran}\Pi \cap D(P_1)$ as an operator on $\text{Ran}\Pi$.

Since G is self-adjoint we can define e^{-itG} and obtain $U^* e^{-itG} U \Pi = e^{-itP_1} \Pi$. This establishes the theorem. $\square$

3.5. Proof of Theorem 2.9

Set $\varphi_0 = (\pi h)^{-3/4} \Pi \chi(P)(e^{ix\xi_0/h - (x-x_0)^2/2h} u_1(x))$. Then by Theorem 2.6 and Theorem 2.7 we have

$$e^{-itP/h} \varphi_0 = U^* e^{-itG/h} U \varphi_0 + \mathcal{O}(|t|h^\infty).$$

Let $\mathcal{U}$ be as in the proof of Theorem 2.7. Then $\mathcal{U} \Pi \chi(P)$ is a h -admissible operator with a symbol b . Applying the stationary phase method to $Op_h^w(b) \varphi_0$ as in Martinez and Sordani [18, Theorem 11.3] we obtain

$$\mathcal{U} \varphi_0 \sim (\pi h)^{-3/4} e^{ix\xi_0/h - (x-x_0)^2/2h} (u_1(x) + h^{1/2} w_0(x)) + \sum_{\ell=1}^{\infty} h^{\ell/2} \phi_\ell w_\ell(x),$$

where ϕ_ℓ is a coherent state centered at (x_0, ξ_0) and $w_\ell(x)$ is a resummation of $w_\ell(x) \sim \sum_{j=0}^{\infty} h^{j/2} w_{\ell,j}$. Here $w_{\ell,j}$ is a sum of terms like $a(x) \partial_x^\alpha u_1(x)$ with $a(x) \in C^\infty(\mathbb{R}_x^3; \mathcal{L}(L^2(\mathbb{R}_y^{3N})))$. Taking the inner product with $u_1(x)$ we have

$$\begin{aligned} U \varphi_0 &\sim (\pi h)^{-3/4} e^{ix\xi_0/h - (x-x_0)^2/2h} (1 + h^{1/2} (u_1(x), w_0(x))_{L_y^2}) \\ &\quad + \sum_{\ell=1}^{\infty} h^{\ell/2} \phi_\ell (u_1(x), w_\ell(x))_{L_y^2}. \end{aligned}$$

Applying the Theorem 3.1 of Combes and Robert [3] we have

$$\begin{aligned} e^{-it\tilde{G}/h} U \varphi_0 &= (\pi h)^{-3/4} e^{ix\xi_t/h - (x-x_t)^2/2h} \\ &\quad + h^{1/2} e^{i\delta_t/h} \sum_{\ell=0}^{\ell_J} c_\ell(t; h) \phi_{\ell,t} + \mathcal{O}(e^{CJt} h^{J/2}), \end{aligned}$$

where δ_t and $c_\ell(t; h)$ are as in the statement of Theorem 2.9 and $C > 0$ is a constant. Multiplying $u_1(x)$ and applying the stationary phase method to $\mathcal{U}^*$ we have

$$\begin{aligned} U^* e^{-itG/h} U \varphi_0 &= (\pi h)^{-3/4} e^{ix\xi_t/h - (x-x_t)^2/2h} u_1(x) \\ &\quad + h^{1/2} e^{i\delta_t/h} \sum_{\ell=0}^{\ell_J} c_\ell(t; h) \phi_{\ell,t} v_\ell(x) + \mathcal{O}(h^{J/4}), \end{aligned} \quad (3.18)$$

for $t < C^{-1} \ln \frac{1}{h}$, where $C > 0$ is a constant, $\delta_t, c_\ell(t; h)$ and $\phi_{\ell,t}$ are satisfying the conditions in Theorem 2.9 and $v_\ell(x) = a(x) \partial_x^\alpha u_1(x)$ with $a(x) \in C^\infty(\mathbb{R}_x^3; \mathcal{L}(L^2(\mathbb{R}_y^{3N})))$.

Since $\mathcal{V}^* \varphi_0 = \tilde{\varphi}_0$, multiplying both sides of (3.18) by $\mathcal{V}^*$ we have (2.9) with $\tilde{v}_\ell(x) = \mathcal{V}^* v_\ell(x)$. We shall prove $\tilde{v}_\ell \in C^\infty(\mathbb{R}_x^3; L_y^2(\mathbb{R}^{3N}))$. Since $a(x) \in C^\infty(\mathbb{R}_x^3; \mathcal{L}(L^2(\mathbb{R}_y^{3N})))$ in $v_\ell(x) = a(x) \partial_x^\alpha u_1(x)$ is the derivative of the symbol of $\mathcal{U}^*$ we have $[y, a(x)] \in C^\infty(\mathbb{R}_x^3; \mathcal{L}(L^2(\mathbb{R}_y^{3N})))$. By the equation

$$\partial_{x_j} (\mathcal{V}^* v_\ell(x)) = -ie \left(A \sum_{i=2}^{N+1} y_i \right)_j \mathcal{V}^* v_\ell(x) + \mathcal{V}^* \partial_{x_j} v_\ell(x) \in C^0(\mathbb{R}_x^3; L_y^2),$$

it is sufficient to prove $y^\alpha \partial_x^\beta u_1(x) \in C^0(\mathbb{R}_x^3; L_y^2)$ for any $\alpha \in \mathbb{N}^{3N}$ and $\beta \in \mathbb{N}^3$. Since

$$\begin{aligned} \partial_{x_j} u_1(x) &= \partial_{x_j} (\Pi_0(x)^2 u_1(x)) \\ &= (\partial_{x_j} \Pi_0(x)) u_1(x) + \Pi_0(x) (\partial_{x_j} \Pi_0(x)) u_1(x) + \Pi_0(x) \partial_{x_j} u_1(x), \end{aligned}$$

we have $y^\alpha \partial_{x_j} u_1(x) \in C^0(\mathbb{R}_x^3; L_y^2)$ for any $\alpha \in \mathbb{N}^{3N}$. In the same way we have $y^\alpha \partial_x^\beta u_1(x) \in C^0(\mathbb{R}_x^3; L_y^2)$ for any $\alpha \in \mathbb{N}^{3N}$ and $\beta \in \mathbb{N}^3$. Thus $\mathcal{V}^* v_\ell(x)$ is differentiable any number of times which completes the proof.

Acknowledgements

The author would like to express his great appreciation to Professor Yoshio Tsutsumi for his helpful advices and encouragements. The author also shows his deep gratitude to Professor Tadayoshi Adachi for his helpful discussions and the referees for their helpful comments about this paper.

Appendix A. Pseudodifferential Operators with Operator Valued Symbols

We introduce the classes of operator valued symbols we use.

Definition A.1. Let $\mathcal{H}_1$ and $\mathcal{H}_2$ be Hilbert spaces and h_0 a positive constant. A function $a(x, \xi; h) \in C^\infty(\mathbb{R}^n \times \mathbb{R}^\ell \times (0, h_0]; \mathcal{L}(\mathcal{H}_1, \mathcal{H}_2))$ is said to be in $S^m(\mathbb{R}^n \times \mathbb{R}^\ell; \mathcal{L}(\mathcal{H}_1, \mathcal{H}_2))$ if for any $\alpha \in \mathbb{N}^n$ and $\beta \in \mathbb{N}^\ell$ one has

$$\sup_{(x, \xi, h) \in \mathbb{R}^{n+\ell} \times (0, h_0]} \|\partial_x^\alpha \partial_\xi^\beta a(x, \xi; h)\|_{\mathcal{L}(\mathcal{H}_1, \mathcal{H}_2)} (1 + |\xi|)^{-m+|\beta|} < \infty. \quad (\text{A.1})$$

Definition A.2. For $a(x, y, \xi) \in S^m(\mathbb{R}^n \times \mathbb{R}^n; \mathcal{L}(\mathcal{H}_1, \mathcal{H}_2))$ and $u \in \mathcal{S}(\mathbb{R}^n; \mathcal{H}_1)$ we set

$$Op_h^w(a)u(x; h) = \frac{1}{(2\pi h)^n} \int \left(\int e^{i(x-y)\xi/h} a\left(\frac{x+y}{2}, \xi\right) u(y) dy \right) d\xi.$$

$Op_h^w(a)$ is called the semiclassical pseudodifferential operator with operator valued symbol.

We can see that $Op_h^w(a)$ is continuous $\mathcal{S}(\mathbb{R}^n; \mathcal{H}_1) \rightarrow \mathcal{S}(\mathbb{R}^n; \mathcal{H}_2)$ as in Martinez [16]. Since the formal adjoint of $Op_h^w(a)$ is $Op_h^w(a^*)$, we can extend it uniquely to a linear continuous operator $\mathcal{S}'(\mathbb{R}^n; \mathcal{H}_1) \rightarrow \mathcal{S}'(\mathbb{R}^n; \mathcal{H}_2)$.

For the composition of pseudodifferential operators we have the following theorem as in the scalar case (see Hörmander [14]).

Theorem A.3. If $a_1 \in S^{m_1}(\mathbb{R}^n \times \mathbb{R}^n; \mathcal{L}(\mathcal{H}_1, \mathcal{H}_2))$, $a_2 \in S^{m_2}(\mathbb{R}^n \times \mathbb{R}^n; \mathcal{L}(\mathcal{H}_2, \mathcal{H}_3))$, then as operators on $\mathcal{S}(\mathbb{R}^n; \mathcal{H}_1)$ or $\mathcal{S}'(\mathbb{R}^n; \mathcal{H}_1)$

$$Op_h^w(a_2)Op_h^w(a_1) = Op_h^w(b),$$

where $b \in S^{m_1+m_2}(\mathbb{R}^n \times \mathbb{R}^n; \mathcal{L}(\mathcal{H}_1, \mathcal{H}_3))$ is given by

$$b(x, \xi) = e^{ih[D_\eta D_x - D_y D_\xi]/2} a_2(y, \eta) a_1(x, \xi) \Big|_{\eta=\xi}^{y=x} =: a_1 \# a_2,$$

and has the expansion

$$b(x, \xi) = \sum_{|\alpha+\beta| \leq N} \frac{h^{|\alpha+\beta|} (-1)^{|\alpha|}}{(2i)^{|\alpha+\beta|} \alpha! \beta!} (\partial_x^\alpha \partial_\xi^\beta a_2(x, \xi)) (\partial_\xi^\alpha \partial_x^\beta a_1(x, \xi)) + h^{N+1} r_N(x, \xi),$$

where $r_N \in S^{m_1+m_2-N-1}(\mathbb{R}^n \times \mathbb{R}^n; \mathcal{L}(\mathcal{H}_1, \mathcal{H}_3))$

The L^2 -boundedness can be established as in the scalar case.

Theorem A.4. Let $a \in S^0(\mathbb{R}^n \times \mathbb{R}^n; \mathcal{L}(\mathcal{H}))$. Then $Op_h^w(a)$ is continuous on $L^2(\mathbb{R}^n; \mathcal{H})$ and

$$\|Op_h^w(a)\|_{\mathcal{L}(L^2(\mathbb{R}^n; \mathcal{H}))} \leq C_n \left(\sum_{|\alpha+\beta| \leq M_n} \|\partial_x^\alpha \partial_\xi^\beta a\|_{L^\infty(\mathbb{R}^n \times \mathbb{R}^n; \mathcal{L}(\mathcal{H}))} \right),$$

where the positive constants C_n and M_n depend only on n .

We also use the useful notion of h -admissible operators (see Martinez and Sordani [18, Appendix]).

Definition A.5. Let $m \in \mathbb{R}$ and let $\mathcal{H}_1$ and $\mathcal{H}_2$ be Hilbert spaces. An operator $A = A(h) : H^m(\mathbb{R}^n; \mathcal{H}_1) \rightarrow L^2(\mathbb{R}^n; \mathcal{H}_2)$ with $h \in (0, h_0]$ is called h -admissible (of degree m) if, for any $N \geq 1$

$$A(h) = \sum_{j=0}^N h^j Op_h^w(a_j(x, \xi; h)) + h^N R_N(h),$$

where R_N is uniformly bounded from $H^m(\mathbb{R}^n; \mathcal{H}_1)$ to $L^2(\mathbb{R}^n; \mathcal{H}_2)$ for $h \in (0, h_0]$, and, for all $h > 0$ small enough, $a_j \in S^m(\mathbb{R}^{2n}; \mathcal{L}(\mathcal{H}_1, \mathcal{H}_2))$.

References

- [1] Agmon, S.: Lectures on exponential decay of solutions of second order elliptic equations. In: *Mathematical Notes*, vol. 29, Princeton Univ. Press, Princeton (1982)
- [2] Combescur, M.: The squeezed state approach of semiclassical limit of the time dependent Schrödinger equation. *J. Math. Phys.* **33**, 3870–3880 (1992)
- [3] Combescur, M., Robert, D.: Semiclassical spreading of quantum wave packets and applications near unstable fixed points of the classical flow. *Asymptot. Anal.* **14**, 377–404 (1997)
- [4] Dereziński, J., Gérard, C.: Scattering theory of classical and quantum N -particle systems. In: *Texts and Monographs in Physics*. Springer, Berlin (1997)
- [5] Gérard, C., Laba, I.: Multiparticle quantum scattering in constant magnetic fields. In: *Math. Surveys and Monographs*, vol. 90. AMS, Providence (2002)
- [6] Gérard, C., Martinez, A., Sjöstrand, J.: A mathematical approach to the effective Hamiltonian in perturbed periodic problems. *Commun. Math. Phys.* **142**, 217–244 (1991)
- [7] Hagedorn, G.A.: A time-dependent Born–Oppenheimer approximation. *Commun. Math. Phys.* **77**, 1–19 (1980)
- [8] Hagedorn, G.A.: High order corrections to the time-dependent Born–Oppenheimer approximation I: smooth potentials. *Ann. Math.* **124**, 571–590 (1986). [Erratum: *Ann. Math.* **126**, 219 (1987)]
- [9] Hagedorn, G.A.: High order corrections to the time-dependent Born–Oppenheimer approximation II: Coulomb systems. *Commun. Math. Phys.* **117**, 387–403 (1988)
- [10] Hagedorn, G.A.: Molecular propagation through electron energy level crossings. *Mem. AMS* **536**, 1–130 (1994)
- [11] Hagedorn, G.A., Joye, A.: Landou-Zener transitions through small electronic eigenvalue gaps in the Born–Oppenheimer approximation. *Ann. Inst. H. Poincaré. Phys. Théor.* **68**, 85–134 (1998)
- [12] Hagedorn, G.A., Joye, A.: A time-dependent Born–Oppenheimer approximation with exponentially small error estimates. *Commun. Math. Phys.* **233**, 583–626 (2001)
- [13] Helffer, B., Sjöstrand, J.: Analyse semi-classique pour l’équation de Harper II. *Mem. Soc. Math. Fr.* **40**, 1–139 (1990)
- [14] Hörmander, L.: *The Analysis of Linear Partial Differential Operators I, III*. Springer, Berlin (1990, 1985)
- [15] Kato, T.: Perturbation theory for linear operators. In: *Classics in Mathematics*, 2nd edn. Springer, Berlin (1976)
- [16] Martinez, A.: An introduction to semiclassical and microlocal analysis. In: *Universitext*. Springer, New York (2002)
- [17] Martinez, A., Sordoni, V.: A general reduction scheme for the time-dependent Born–Oppenheimer approximation. *C. R. Acad. Sci. Paris* **334**, 185–188 (2002)
- [18] Martinez, A., Sordoni, V.: Twisted pseudodifferential calculus and application to the quantum evolution of molecules. *Mem. AMS* **936**, 1–82 (2009)
- [19] Nenciu, G.: Linear adiabatic theory. Exponential estimates. *Commun. Math. Phys.* **152**, 479–496 (1993)

- [20] Nenciu, G.: On asymptotic perturbation theory for quantum mechanics: almost invariant subspaces and gauge invariant magnetic perturbation theory. *J. Math. Phys.* **43**, 1273–1298 (2002)
- [21] Nenciu, G., Sordoni, V.: Semiclassical limit for multistate Klein–Gordon systems: almost invariant subspaces, and scattering theory. *J. Math. Phys.* **45**, 3676–3696 (2004)
- [22] Reed, M., Simon, B.: *Methods of Modern Mathematical Physics. I–IV*. Academic Press, New York (1972, 1975, 1979, 1978)
- [23] Sjöstrand, J.: Projecteurs adiabatiques du point de vue pseudodifférentiel. *C. R. Acad. Sci. Paris* **317**, 217–220 (1993)
- [24] Sordoni, V.: Reduction scheme for semiclassical operator-valued Schrödinger type equation and application to scattering. *Commun. Partial Differ. Equ.* **28**, 1221–1236 (2003)
- [25] Spohn, H., Teufel, S.: Adiabatic decoupling and time-dependent Born–Oppenheimer theory. *Commun. Math. Phys.* **224**, 113–132 (2001)
- [26] Teufel, S.: Adiabatic perturbation theory in quantum dynamics. In: *Lecture Notes in Mathematics*, vol. 1821. Springer, Berlin (2003)

Sohei Ashida
Department of Mathematics
Graduate School of Science
Kyoto University
Kyoto 606, Japan
e-mail: ashida@math.kyoto-u.ac.jp

Communicated by Jan Dereziński.

Received: April 17, 2015.

Accepted: November 12, 2015.