



Good Inducing Schemes for Uniformly Hyperbolic Flows, and Applications to Exponential Decay of Correlations

Ian Melbourne and Paulo Varandas

Abstract. Given an Axiom A attractor for a $C^{1+\alpha}$ flow ($\alpha > 0$), we construct a countable Markov extension with exponential return times in such a way that the inducing set is a smoothly embedded unstable disk. This avoids technical issues concerning irregularity of boundaries of Markov partition elements and enables an elementary approach to certain questions involving exponential decay of correlations for SRB measures.

1. Introduction

Statistical properties of Anosov and Axiom A diffeomorphisms [3, 33] were developed extensively in the 1970s. Key tools were the construction of finite Markov partitions [10, 32] and the spectral properties of transfer operators [28]. In particular, ergodic invariant probability measures were constructed corresponding to any Hölder potential [12, 29, 31]; moreover, it was shown that hyperbolic basic sets for Axiom A diffeomorphisms are always exponentially mixing up to a finite cycle for such measures, see for example [12, 22, 29].

Still in the 1970s, finite Markov partitions were constructed [11, 26] for Anosov and Axiom A flows. This allows us to model each hyperbolic basic set as a suspension flow over a subshift of finite type, enabling the study of thermodynamic formalism (see e.g. [14]) and statistical properties (see e.g. [17, 27]). By the Anosov alternative [3, 23], a transitive Anosov flow is mixing if and only if it is a constant suspension over an Axiom A diffeomorphism.

However, rates of mixing for Axiom A flows are still poorly understood. By [24, 30], mixing Axiom A flows can mix arbitrarily slowly. Although there has been important progress starting with [16, 18, 20], it remains an open question whether mixing Anosov flows have exponential decay of correlations. Very

recently, this question was answered positively [35] in the case of C^∞ three-dimensional flows.

It turns out that using finite Markov partitions for flows raises technical issues due to the irregularity of their boundaries [5, 15, 34]. Even in the discrete-time setting, it is known that the boundaries of elements of a finite Markov partition need not be smooth [13]. In this paper, we propose using the approach of [36] to circumvent such issues at least in the case of SRB measures. In particular, we show that

Any attractor¹ for an Axiom A flow can be modeled by a suspension flow over a full branch countable Markov extension where the inducing set is a smoothly embedded unstable disk. The roof function, though unbounded, has exponential tails.

A precise statement is given in Theorem 2.1 below.

Remark 1.1. The approach of Young towers [36] has proved to be highly effective for studying discrete-time examples like planar dispersing billiards and Hénon-like attractors where suitable Markov partitions are not available. However, as shown in the current paper, there can be advantages (at least in continuous time) to working with countable Markov extensions even when there is a well-developed theory of finite Markov partitions. The extra flexibility of Markov extensions can be used not only to construct the extension but to ensure good regularity properties of the partition elements.

As a consequence of Theorem 2.1, we obtain an elementary proof of the following result:

Theorem 1.2. *Suppose that Λ is an Axiom A attractor with SRB measure μ for a C^{1+} flow ϕ_t with C^{1+} stable holonomies² and such that the stable and unstable bundles are not jointly integrable. Then for all Hölder observables $v, w : \Lambda \rightarrow \mathbb{R}$, there exist constants $c, C > 0$ such that*

$$\left| \int_{\Lambda} v w \circ \phi_t d\mu - \int_{\Lambda} v d\mu \int_{\Lambda} w d\mu \right| \leq C e^{-ct} \quad \text{for all } t > 0.$$

Remark 1.3. Joint nonintegrability holds for an open and dense set of Axiom A flows and their attractors, see [19] and references therein. It implies mixing and is equivalent to mixing for codimension one Anosov flows. It is conjectured to be equivalent to mixing for Anosov flows [23].

Remark 1.4. (a) In the case when the unstable direction is one-dimensional and the stable holonomies are C^2 , this result is due to [4, 5, 8, 9]. In particular, using the fact that stable bunching is a robust sufficient condition for smoothness of stable holonomies together with the robustness of joint nonintegrability, [4] constructed the first robust examples of Axiom A flows with exponential decay of correlations. The smoothness condition on stable holonomies was relaxed from C^2 to C^{1+} in [6] extending the

¹Here, an attractor is an attracting hyperbolic basic set, and so is topologically transitive with an open basin of attraction.

² C^{1+} means C^r for some $r > 1$.

class of examples in [4]. This class of examples is extended further by Theorem 1.2 with the removal of the one-dimensionality restriction on unstable manifolds.

- (b) There is no restriction on the dimension of unstable manifolds in [8], and it is not surprising that the smoothness assumption on stable holonomies can also be relaxed as in [6]. However, there is a crucial hypothesis in [8] on the regularity of the inducing set in the unstable direction which is nontrivial in higher dimensions.

Theorem 1.2 is stated in the special case of Anosov flows in [15]. In [15, Appendix] it is argued that at least in the Anosov case the Markov partitions of [26] are sufficiently regular that the methods in [8] can be pushed through. In [5], a sketch is given of how to prove Theorem 1.2 also in the Axiom A case, but the details are not fully worked out.

As mentioned, our approach in this paper completely bypasses such issues since our inducing set is a smoothly embedded unstable disk. Moreover, our method works equally well for Anosov flows and Axiom A attractors. As a consequence, we recover the examples in [15], in particular that codimension one volume-preserving mixing C^{1+} Anosov flows are exponentially mixing in dimension four and higher.

The remainder of the paper is organized as follows. In Sect. 2, we state precisely and prove our result on good inducing for attractors of Axiom A flows. In Sect. 3, we prove a result on exponential mixing for a class of skew product Axiom A flows, extending/combining the results in [6, 8]. In Sect. 4, we complete the proof of Theorem 1.2.

2. Good Inducing for Attractors of Axiom A Flows

Let $\phi_t : M \rightarrow M$ be a C^{1+} flow defined on a compact Riemannian manifold (M, d_M) , and let $\Lambda \subset M$ be a closed ϕ_t -invariant subset. We assume that Λ is an attracting transitive uniformly hyperbolic set with adapted norm and that Λ is not a single trajectory. In particular, there is a continuous $D\phi_t$ -invariant splitting $T_\Lambda M = E^s \oplus E^c \oplus E^u$ where E^c is the one-dimensional central direction tangent to the flow, and there exists $\lambda \in (0, 1)$ such that $|D\phi_t v| \leq \lambda^t |v|$ for all $v \in E^s$, $t \geq 1$; $|D\phi_{-t} v| \leq \lambda^t |v|$ for all $v \in E^u$, $t \geq 1$. Since the time- s map $\phi_s : \Lambda \rightarrow \Lambda$ is ergodic for all but countably many choices of $s \in \mathbb{R}$ [25], we can scale time by a constant close to one if necessary so that $\phi_{-1} : \Lambda \rightarrow \Lambda$ is transitive. Then there exists $p \in \Lambda$ such that $\bigcup_{i \geq 1} \phi_{-i} p$ is dense in Λ .

We can define (local) stable disks $W_\delta^s(y) = \{z \in W^s(y) : d_M(y, z) < \delta\}$ for $\delta > 0$ sufficiently small for all $y \in \Lambda$. Define local center-stable disks $W_\delta^{cs}(y) = \bigcup_{|t| < \delta} \phi_t W_\delta^s(y)$. Let Leb and d denote induced Lebesgue measure and induced distance on local unstable manifolds. It is convenient to define local unstable disks $W_\delta^u(y) = \{z \in W^u(y) : d(y, z) < \delta\}$ using the induced distance.

For δ_0 small, define $\mathcal{D} = W_{\delta_0}^u(p)$ and $\widehat{\mathcal{D}} = \bigcup_{x \in \mathcal{D}} W_{\delta_0}^{cs}(x)$. Define $\pi : \widehat{\mathcal{D}} \rightarrow \mathcal{D}$ such that $\pi|_{W_{\delta_0}^{cs}(x)} \equiv x$. Whenever $\phi_n y \in \widehat{\mathcal{D}}$, we set $g_n y = \pi(\phi_n y)$.

We are now in a position to give a precise description of our inducing scheme.

Theorem 2.1. *There exists an open unstable disk $Y = W_{\delta}^u(p) \subset \mathcal{D}$ (for some $\delta \in (0, \delta_0)$) and a discrete return time function $R : Y \rightarrow \mathbb{Z}^+ \cup \{\infty\}$ such that*

- (i) $\text{Leb}(R > n) = O(\gamma^n)$ for some $\gamma \in (0, 1)$;
- (ii) *Each connected component of $\{R = n\}$ is mapped by ϕ_n into $\widehat{\mathcal{D}}$ and mapped homeomorphically by g_n onto Y .*

Remark 2.2. Let \mathcal{P} be the partition of Y consisting of connected components of $\{R = n\}$ for $n \geq 1$. (It follows from Theorem 2.1(i) that \mathcal{P} is a partition of $Y \bmod 0$.) Define $F : Y \rightarrow Y$, $F = g_R = \pi \circ \phi_R$. Note that F is locally the composition of a time- R map ϕ_R (where R is constant on each partition element) with a center-stable holonomy. Since center-stable holonomies are Hölder continuous, it follows that F maps partition elements $U \in \mathcal{P}$ homeomorphically onto Y and that $F|_U : U \rightarrow Y$ is a bi-Hölder bijection. If moreover, the center-stable holonomies are C^1 , then the partition elements are diffeomorphic to disks (in contrast to the situation for finite Markov partitions of Λ [13]).

In the remainder of this section, we prove Theorem 2.1. Our proof is essentially the same as in [36, Section 6] for Axiom A diffeomorphisms, but we closely follow the treatment in [2] which provides many of the details of arguments sketched in [36].

Choice of constants. We can choose $\delta_0 > 0$ such that the following bounded distortion property holds³: there exists $C_1 \geq 1$ so that

$$\frac{|\det D\phi_n(x)|E^u|}{|\det D\phi_n(y)|E^u|} \leq C_1 \quad (2.1)$$

for every $n \geq 1$ and all $x, y \in \Lambda$ with $\phi_n x, \phi_n y$ in the same unstable disk such that $d(\phi_j x, \phi_j y) < 4\delta_0$ for all $0 \leq j \leq n$.

By standard results about stable holonomies, π is absolutely continuous and C^α for some $\alpha \in (0, 1)$ when restricted to unstable disks in $\widehat{\mathcal{D}}$. For δ_0 sufficiently small, there exists $C_2, C_3 \geq 1$ such that

$$C_2^{-1} \leq \frac{\text{Leb}(\pi(E))}{\text{Leb}(E)} \leq C_2 \quad (2.2)$$

for all Lebesgue-measurable subset $E \subset W_{\delta_0}^u(y) \cap \widehat{\mathcal{D}}$ and all $y \in \Lambda$, and

$$d(\pi x, \pi y) \leq C_3 d(x, y)^\alpha \quad (2.3)$$

for all $x, y \in \widehat{\mathcal{D}}$ with x, y in the same unstable disk such that $d(x, y) < 4\delta_0$.

³ The function $\psi = \log |\det D\phi_1|E^u|$ is Hölder (since the flow is C^{1+} and the bundle E^u is Hölder). Hence $\sum_{j=0}^{n-1} \{\psi(\phi_j x) - \psi(\phi_j y)\}$ is bounded for the specified n, x, y . Estimate (2.1) follows.

Let $d_u = \dim E^u$ and fix $L > 3$ so that

$$C_1 C_2^2 \frac{2^{d_u} - 1}{(L - 1)^{d_u}} < \frac{1}{4}. \quad (2.4)$$

By the local product structure, there exists $\delta_1 \in (0, \delta_0)$ such that $W_{\delta_0}^{cs}(x) \cap W_{\delta_0}^u(y)$ consists of precisely one point for all $x, y \in \Lambda$ with $d_M(x, y) < 4\delta_1$. Similarly, there exists $\delta \in (0, \delta_1)$ such that $W_{\delta_1}^{cs}(x) \cap W_{\delta_1}^u(y)$ consists of precisely one point for all $x, y \in \Lambda$ with $d_M(x, y) < (L + 1)\delta$. Since local center-stable/unstable manifolds lie in the corresponding cones, and the center-stable/unstable cones are uniformly transverse, the intersection point $z \in W_{\delta_1}^{cs}(x) \cap W_{\delta_1}^u(y)$ satisfies $d(z, y) \leq C_4 d_M(x, y)$ where $C_4 \geq 1$ is a constant. Shrink $\delta > 0$ if necessary so that $C_3(3\delta)^\alpha < \frac{1}{2}\delta_0$ and $C_4(L + 1)\delta < \delta_0$. Choose $N_1 \geq 1$ such that $\bigcup_{i=1}^{N_1} \phi_{-i}p$ is δ -dense in Λ .

Construction of the partition. We consider various small neighborhoods $\mathcal{D}_c = W_{c\delta}^u(p)$ with $c \in \{1, 2, L - 1, L\}$. Define $\tilde{\mathcal{D}}_c = \bigcup_{x \in \mathcal{D}_c} W_{\delta_1}^{cs}(x)$.

Take $Y = \mathcal{D}_1$. Define a partition $\{I_k : k \geq 1\}$ of $\mathcal{D}_2 \setminus \mathcal{D}_1$,

$$I_k = \{y \in \mathcal{D}_2 : \delta(1 + \lambda^{\alpha k}) \leq d(y, p) < \delta(1 + \lambda^{\alpha(k-1)})\}.$$

Fix $\varepsilon > 0$ small (as stipulated in Propositions 2.4 and 2.5 and Lemma 2.9 below). We define sets Y_n and functions $t_n : Y_n \rightarrow \mathbb{N}$, and $R : Y \rightarrow \mathbb{Z}^+$ inductively, with $Y_n = \{R > n\}$. Define $Y_0 = Y$ and $t_0 \equiv 0$. Inductively, suppose that $Y_{n-1} = Y \setminus \{R < n\}$ and that $t_{n-1} : Y_{n-1} \rightarrow \mathbb{N}$ is given. Write $Y_{n-1} = A_{n-1} \dot{\cup} B_{n-1}$ where

$$A_{n-1} = \{t_{n-1} = 0\}, \quad B_{n-1} = \{t_{n-1} \geq 1\}.$$

Consider the neighborhood

$$A_{n-1}^{(\varepsilon)} = \{y \in Y_{n-1} : d(\phi_n y, \phi_n A_{n-1}) < \varepsilon\}$$

of the set A_{n-1} . Define U_{nj}^L , $j \geq 1$, to be the connected components of $A_{n-1}^{(\varepsilon)} \cap \phi_{-n} \tilde{\mathcal{D}}_L$ that are mapped inside $\tilde{\mathcal{D}}_L$ by ϕ_n and mapped homeomorphically onto \mathcal{D}_L by g_n . Let

$$U_{nj}^c = U_{nj}^L \cap g_n^{-1} \mathcal{D}_c \quad \text{for } c = 1, 2, L - 1.$$

Define $R|_{U_{nj}^1} = n$ for each U_{nj}^1 and take $Y_n = Y_{n-1} \setminus \bigcup_j U_{nj}^1$. Finally, define $t_n : Y_n \rightarrow \mathbb{N}$ as

$$t_n(y) = \begin{cases} k, & y \in \bigcup_j U_{nj}^2 \text{ and } g_n y \in I_k \text{ for some } k \geq 1 \\ 0, & y \in A_{n-1} \setminus \bigcup_j U_{nj}^2 \\ t_{n-1}(y) - 1, & y \in B_{n-1} \setminus \bigcup_j U_{nj}^2 \end{cases}$$

and take $A_n = \{t_n = 0\}$, $B_n = \{t_n \geq 1\}$ and $Y_n = A_n \dot{\cup} B_n$.

Remark 2.3. By construction, property (ii) of Theorem 2.1 is satisfied. It remains to verify that $\text{Leb}(R > n)$ decays exponentially.




FIGURE 1. Visualisation of Y after 7 generations where there is one return at time 2 and one return at time 7. The pink region B_7 consists of collars around the sets $\{R = 2\}$ and $\{R = 7\}$ that have made a return. The two outermost shells $\{t_7 = 1\}$ and $\{t_7 = 2\}$ of each collar are shown. The collars in B_7 are diffeomorphic by g_2 and g_7 , respectively, to an annulus; in reality the collar around $\{R = 2\}$ should be slightly distorted and the collar around $\{R = 7\}$ more so (and smaller)

Visualisation of B_n . The set B_n is a disjoint union $B_n = \bigcup_{m=1}^n C_n(m)$ where $C_n(m)$ is a disjoint union of *collars* around each component of $\{R = m\}$. Each collar in $C_n(m)$ is homeomorphic under g_m to $\bigcup_{k \geq n-m+1} I_k$ with outer ring homeomorphic under g_m to I_{n-m+1} , and the union of outer rings is the set $\{t_n = 1\}$. This picture presupposes Proposition 2.4 below which guarantees that each new generation of collars $C_n(n)$ does not intersect the set $\bigcup_{1 \leq m \leq n-1} C_{n-1}(m)$ of collars in the previous generation. A sample visualization after 7 generations is shown in Fig. 1.

We now turn to the proof of Proposition 2.4. A cautionary diagram is shown in Fig. 2.

Proposition 2.4. *Choose $\varepsilon < (C_3^{-1}\delta)^{1/\alpha}$ sufficiently small that $W_\varepsilon^u(x) \subset \widehat{\mathcal{D}}$ for all $x \in \widetilde{\mathcal{D}}_L$. Then $\bigcup_j U_{n_j}^{L-1} \subset A_{n-1}$ for all $n \geq 1$.*

Proof. We argue by contradiction. There is nothing to prove for $n = 1$. Let $n \geq 2$ be least such that the result fails and choose j such that $U_{n_j}^{L-1}$ intersects B_{n-1} . Then either (i) $U_{n_j}^{L-1} \subset B_{n-1}$, or (ii) $U_{n_j}^{L-1}$ intersects ∂A_{n-1} .

In case (i), choose $x \in U_{n_j}^{L-1}$ (so in particular $\phi_n x \in \widehat{\mathcal{D}}$) with $g_n x = p$. Since $U_{n_j}^{L-1} \subset U_{n_j}^L \subset A_{n-1}^{(\varepsilon)}$, there exists $y \in A_{n-1}$ with $d(\phi_n x, \phi_n y) < \varepsilon$. In particular, $\phi_n y \in \widehat{\mathcal{D}}$ so $g_n y$ is well-defined. Note that $x \in U_{n_j}^{L-1}$ and $y \notin U_{n_j}^{L-1}$ since $U_{n_j}^{L-1} \subset B_{n-1}$. Hence the geodesic ℓ in \mathcal{D} joining $g_n x$ and $g_n y$ intersects $g_n \partial U_{n_j}^{L-1}$. Choose $z \in \partial U_{n_j}^{L-1} \cap g_n^{-1} \ell$. Since $g_n = \pi \circ \phi_n$, it follows from (2.3)




FIGURE 2. This diagram indicates some subtleties in the proof of Proposition 2.4. The solid black curve denotes part of the boundary of A_{n-1} formed by a much earlier return. The dashed black curve denotes the corresponding part of the boundary of $A_{n-1}^{(\epsilon)}$. The red curve denotes the boundary of a U_{nj}^L . Even though U_{nj}^L is contained in $A_{n-1}^{(\epsilon)}$ it is not clear that U_{nj}^{L-1} is contained in A_{n-1}

that

$$\delta < (L-1)\delta = d(g_n x, g_n z) < d(g_n x, g_n y) \leq C_3 d(\phi_n x, \phi_n y)^\alpha < C_3 \varepsilon^\alpha < \delta$$

which is a contradiction. This rules out case (i).

In case (ii), choose $x \in U_{nj}^{L-1} \cap \partial A_{n-1}$. We show below that there exists $y \in \partial A_{n-1}^{(\epsilon)}$ such that $d(\phi_n x, \phi_n y) \leq \varepsilon$. In particular, $g_n x$ and $g_n y$ are well-defined and $d(g_n x, g_n y) \leq C_3 \varepsilon^\alpha < \delta$. Since $U_{nj}^L \subset A_{n-1}^{(\epsilon)}$, we have that $y \notin U_{nj}^L$. It follows that $g_n x \in \mathcal{D}_{L-1}$ while $g_n y \notin \mathcal{D}_L$. Hence $d(g_n x, g_n y) \geq \delta$ which is the desired contradiction.

It remains to verify that there exists $y \in \partial A_{n-1}^{(\epsilon)}$ such that $d(\phi_n x, \phi_n y) \leq \varepsilon$. Since n is least, B_{n-1} is a disjoint union of collars as described in the visualization above. Hence there exists a collar $Q \subset C_{n-1}(n-k)$ intersected by U_{nj}^{L-1} for some $1 \leq k < n$ such that x lies in the outer boundary $\partial_o Q$ of Q . Note that $\partial_o Q = \partial A_{n-1} \cap Q$. Let D denote the disk enclosed by $\partial_o Q$ and let

$$S = D \cap \partial(\phi_{-n} B_\varepsilon(\phi_n \partial D)).$$

We claim that $S \neq \emptyset$ and $S \subset Q$. Then S is a $(\dim Y - 1)$ -dimensional sphere contained in $\partial A_{n-1}^{(\epsilon)}$ and there exists $y \in S$ with the desired properties, see Fig. 3. (The point of the claim is that S lies entirely in Y_{n-1} .)

Note that g_{n-k} maps Q homeomorphically onto the set $J = \bigcup_{i \geq k} I_i$ which is an annulus of radial thickness $\delta \lambda^{\alpha k}$. By (2.3), ϕ_{n-k} maps Q homeomorphically onto a set $\tilde{J} = \pi^{-1} J$ of radial thickness at least $(C_3^{-1} \delta \lambda^{\alpha k})^{1/\alpha} = (C_3^{-1} \delta)^{1/\alpha} \lambda^k$.

Moreover, $\phi_k(\tilde{J} \cap \phi_{n-k} A_{n-1}^{(\epsilon)}) \subset \phi_n A_{n-1}^{(\epsilon)}$ is contained in the set of points within d -distance ε of $\phi_n \partial A_{n-1}^{(\epsilon)}$, so by definition of λ we have that $\tilde{J} \cap \phi_{n-k} A_{n-1}^{(\epsilon)}$ is contained in the set of points within d -distance $\varepsilon \lambda^k$ of the outer




FIGURE 3. Schematic of the claim in the proof of Proposition 2.4. The pink region is a collar Q . The blue curve denotes the boundary of a $U_{n_j}^{L-1}$. The claim is that the relevant part S of $\partial A_{n-1}^{(\varepsilon)}$ lies inside Q enabling the choice of a point y close to, but not too close to, x

boundary of \tilde{J} . Since $\varepsilon < (C_3^{-1}\delta)^{1/\alpha}$, we obtain that $\tilde{J} \cap \phi_{n-k}\partial A_{n-1}^{(\varepsilon)}$ is homeomorphic to a $(\dim Y - 1)$ -dimensional sphere contained entirely inside \tilde{J} . Hence $S = Q \cap \partial A_{n-1}^{(\varepsilon)}$ is homeomorphic to a $(\dim Y - 1)$ -dimensional sphere contained entirely inside Q , as required. \square

Proposition 2.5. *Choose $\varepsilon < \{C_3^{-1}\delta(\lambda^{-\alpha} - 1)\}^{1/\alpha}$. Then for all $n \geq 1$,*

- (a) $A_{n-1}^{(\varepsilon)} \subset \{y \in Y_{n-1} : t_{n-1}(y) \leq 1\}$ for all $n \geq 1$.
- (b) $\phi_{-n}W_\varepsilon^u(\phi_n x) \subset A_{n-1}^{(\varepsilon)}$ for all $x \in A_{n-1}$.

Proof. (a) Suppose that $t_{n-1}(y) > 1$. Then there exists a collar in $C_{n-1}(n-k)$ containing y . Let Q denote the outer ring of the collar with outer boundary Q_1 and inner boundary Q_2 . Then $t_{n-1}|_Q \equiv 1$ and $t_{n-1}(y) > 1$, so y lies inside the region bounded by Q_2 .

Suppose for contradiction that $y \in A_{n-1}^{(\varepsilon)}$. Then we can choose $x \in A_{n-1}$ with $d(\phi_n x, \phi_n y) < \varepsilon$. Let ℓ be the geodesic in $W_\varepsilon^u(\phi_n x)$ connecting $\phi_n x$ to $\phi_n y$ and define $q_j \in Q_j \cap \phi_{-n}\ell$ for $j = 1, 2$.

Recall that Q is homeomorphic under g_{n-k} to I_k . Moreover, $g_{n-k}q_j$ lie in distinct components of the boundary of I_k , so

$$d(g_{n-k}q_1, g_{n-k}q_2) \geq \delta(\lambda^{\alpha(k-1)} - \lambda^{\alpha k}) = \delta(\lambda^{-\alpha} - 1)\lambda^{\alpha k}.$$

Hence

$$\begin{aligned} d(\phi_n q_1, \phi_n q_2) &\geq \lambda^{-k} d(\phi_{n-k} q_1, \phi_{n-k} q_2) \\ &\geq \lambda^{-k} \{C_3^{-1} d(g_{n-k} q_1, g_{n-k} q_2)\}^{1/\alpha} \geq \{C_3^{-1} \delta(\lambda^{-\alpha} - 1)\}^{1/\alpha} > \varepsilon. \end{aligned}$$

But $d(\phi_n q_1, \phi_n q_2) \leq d(\phi_n y, \phi_n x) < \varepsilon$ so we obtain the desired contradiction.

(b) Let $x \in A_{n-1}$ and $y \in \phi_{-n}W_\varepsilon^u(\phi_n x)$. Note that $y \in A_{n-1}^{(\varepsilon)}$ if and only if $y \in Y_{n-1}$. Hence we must show that $y \in Y_{n-1}$. If not, then there exists $k \geq 1$

such that $y \in \{R = n - k\}$. Define $Q \subset C_{n-1}(n - k)$ to be the outer ring of the corresponding collar. Choosing q_1 and q_2 as in part (a) we again obtain a contradiction. \square

Lemma 2.6. *There exists $a_1 > 0$ such that for all $n \geq 1$,*

- (a) $\text{Leb}(B_{n-1} \cap A_n) \geq a_1 \text{Leb}(B_{n-1})$.
- (b) $\text{Leb}(A_{n-1} \cap B_n) \leq \frac{1}{4} \text{Leb}(A_{n-1})$.
- (c) $\text{Leb}(A_{n-1} \cap \{R = n\}) \leq \frac{1}{4} \text{Leb}(A_{n-1})$.

Proof. (a) Let $y \in B_{n-1}$. By Proposition 2.4, $y \notin \bigcup_j U_{n_j}^{L-1}$ so in particular $y \in Y_n$. Note that $t_n(y) = 0$ if and only if $t_{n-1}(y) = 1$. Hence $B_{n-1} \cap A_n = \{t_{n-1} = 1\}$.

Now let $Q \subset C_{n-1}(n - k) \subset B_{n-1}$ be a collar ($1 \leq k \leq n$) with outer ring $Q \cap A_n = Q \cap \{t_{n-1} = 1\}$. Then $g_{n-k} = \pi \circ \phi_{n-k}$ maps Q homeomorphically onto $\bigcup_{i \geq k} I_i$ and $Q \cap \{t_{n-1} = 1\}$ homeomorphically onto I_k . Let $d_u = \dim E^u$. By (2.1) and (2.2),

$$\begin{aligned} \frac{\text{Leb}(Q)}{\text{Leb}(Q \cap A_n)} &= \frac{\text{Leb}(Q)}{\text{Leb}(Q \cap \{t_{n-1} = 1\})} \leq C_1 \frac{\text{Leb}(\phi_{n-k}Q)}{\text{Leb}(\phi_{n-k}(Q \cap \{t_{n-1} = 1\}))} \\ &\leq C_1 C_2^2 \frac{\text{Leb}(\bigcup_{i \geq k} I_i)}{\text{Leb}(I_k)} = C_1 C_2^2 D(d_u, \lambda^\alpha, k) \end{aligned}$$

where $D(d_u, \lambda, k) = \frac{(1 + \lambda^{k-1})^{d_u} - 1}{(1 + \lambda^{k-1})^{d_u} - (1 + \lambda^k)^{d_u}}$. Since $\lim_{k \rightarrow \infty} D(d_u, \lambda, k) = (1 - \lambda)^{-1}$, we obtain that $\text{Leb}(Q) \leq C_1 C_2^2 D \text{Leb}(Q \cap A_n)$ where $D = \sup_{k \geq 1} D(d_u, \lambda^\alpha, k)$ is a constant depending only on d_u and λ^α . Summing over collars Q , it follows that $\text{Leb}(B_{n-1}) \leq C_1 C_2^2 D \text{Leb}(B_{n-1} \cap A_n)$.

(b) By Proposition 2.4, $U_{n_j}^2 \subset U_{n_j}^{L-1} \subset A_{n-1}$ for each j . It follows that $A_{n-1} \cap B_n = \bigcup_j U_{n_j}^2 \setminus U_{n_j}^1$. By (2.1), (2.2) and (2.4),

$$\frac{\text{Leb}(U_{n_j}^2 \setminus U_{n_j}^1)}{\text{Leb}(U_{n_j}^{L-1})} \leq C_1 C_2^2 \frac{\text{Leb}(\mathcal{D}_2 \setminus \mathcal{D}_1)}{\text{Leb}(\mathcal{D}_{L-1})} = C_1 C_2^2 \frac{2^{d_u} - 1}{(L - 1)^{d_u}} < \frac{1}{4}.$$

Hence

$$\frac{\text{Leb}(A_{n-1} \cap B_n)}{\text{Leb}(A_{n-1})} \leq \frac{\sum_j \text{Leb}(U_{n_j}^2 \setminus U_{n_j}^1)}{\sum_j \text{Leb}(U_{n_j}^{L-1})} < \frac{1}{4}.$$

(c) Proceeding as in part (b) with $U_{n_j}^2 \setminus U_{n_j}^1$ replaced by $U_{n_j}^1$, leads to the estimate

$$\frac{\text{Leb}(A_{n-1} \cap \{R = n\})}{\text{Leb}(A_{n-1})} \leq \frac{\sum_j \text{Leb}(U_{n_j}^1)}{\sum_j \text{Leb}(U_{n_j}^{L-1})} \leq \frac{C_1 C_2^2}{(L - 1)^{d_u}} < \frac{1}{4}.$$

\square

Corollary 2.7. *For all $n \geq 1$,*

- (a) $\text{Leb}(A_{n-1} \cap A_n) \geq \frac{1}{2} \text{Leb}(A_{n-1})$.
- (b) $\text{Leb}(B_{n-1} \cap B_n) \leq (1 - a_1) \text{Leb}(B_{n-1})$.
- (c) $\text{Leb}(B_n) \leq \frac{1}{4} \text{Leb}(A_{n-1}) + (1 - a_1) \text{Leb}(B_{n-1})$.

$$(d) \text{ Leb}(A_n) \geq \frac{1}{2} \text{ Leb}(A_{n-1}) + a_1 \text{ Leb}(B_{n-1}).$$

Proof. Recall that $A_{n-1} \subset Y_{n-1} = Y_n \dot{\cup} \{R = n\} = A_n \dot{\cup} B_n \dot{\cup} \{R = n\}$. Hence by Lemma 2.6(b,c),

$$\begin{aligned} \text{Leb}(A_{n-1}) &= \text{Leb}(A_{n-1} \cap A_n) + \text{Leb}(A_{n-1} \cap B_n) + \text{Leb}(A_{n-1} \cap \{R = n\}) \\ &\leq \text{Leb}(A_{n-1} \cap A_n) + \frac{1}{2} \text{Leb}(A_{n-1}), \end{aligned}$$

proving (a). Similarly, by Lemma 2.6(a),

$$\begin{aligned} \text{Leb}(B_{n-1}) &= \text{Leb}(B_{n-1} \cap A_n) + \text{Leb}(B_{n-1} \cap B_n) + \text{Leb}(B_{n-1} \cap \{R = n\}) \\ &\geq a_1 \text{Leb}(B_{n-1}) + \text{Leb}(B_{n-1} \cap B_n), \end{aligned}$$

proving (b).

Next, recall that $B_n = B_n \cap Y_{n-1} = B_n \cap (A_{n-1} \dot{\cup} B_{n-1})$. Hence part (c) follows from Lemma 2.6(b) and part (b). Similarly, $A_n = A_n \cap (A_{n-1} \dot{\cup} B_{n-1})$ and part (d) follows from Lemma 2.6(a) and part (a). \square

Corollary 2.8. *There exists $a_0 > 0$ such that $\text{Leb}(B_n) \leq a_0 \text{Leb}(A_n)$ for all $n \geq 0$.*

Proof. Let $a_0 = \frac{2 + a_1}{2a_1}$. We prove the result by induction. The case $n = 0$ is trivial since $B_0 = \emptyset$. For the induction step from $n - 1$ to n , we consider separately the cases $\text{Leb}(B_{n-1}) > \frac{1}{2a_1} \text{Leb}(A_{n-1})$ and $\text{Leb}(B_{n-1}) \leq \frac{1}{2a_1} \text{Leb}(A_{n-1})$.

Suppose first that $\text{Leb}(B_{n-1}) > \frac{1}{2a_1} \text{Leb}(A_{n-1})$. By Corollary 2.7(c), $\text{Leb}(B_n) < \{\frac{1}{2}a_1 + (1 - a_1)\} \text{Leb}(B_{n-1}) = (1 - \frac{1}{2}a_1) \text{Leb}(B_{n-1}) < \text{Leb}(B_{n-1})$. By Corollary 2.7(d),

$$\text{Leb}(A_n) > (\frac{1}{2} + a_1 \frac{1}{2a_1}) \text{Leb}(A_{n-1}) = \text{Leb}(A_{n-1}).$$

Hence by the induction hypothesis,

$$\text{Leb}(B_n) < \text{Leb}(B_{n-1}) \leq a_0 \text{Leb}(A_{n-1}) < a_0 \text{Leb}(A_n),$$

establishing the result at time n .

Finally, suppose that $\text{Leb}(B_{n-1}) \leq \frac{1}{2a_1} \text{Leb}(A_{n-1})$. By Corollary 2.7(a,c),

$$\begin{aligned} \text{Leb}(B_n) &\leq \frac{1}{4} \text{Leb}(A_{n-1}) + \text{Leb}(B_{n-1}) \leq (\frac{1}{4} + \frac{1}{2a_1}) \text{Leb}(A_{n-1}) \\ &\leq (\frac{1}{2} + \frac{1}{a_1}) \text{Leb}(A_n) = a_0 \text{Leb}(A_n), \end{aligned}$$

completing the proof. \square

Lemma 2.9. *Let $\varepsilon \in (0, \frac{1}{2}\delta_0)$ be small as in Propositions 2.4 and 2.5. There exist $c_1 > 0$ and $N \geq 1$ such that*

$$\text{Leb} \left(\bigcup_{i=0}^N \{R = n + i\} \right) \geq c_1 \text{Leb}(A_{n-1}) \quad \text{for all } n \geq 1.$$

Proof. Fix $\lambda \in (0, 1)$, $L > 1$, $0 < \delta < \delta_1 < \delta_0$ and $N_1 \geq 1$ as defined from the outset. Recall that $C_3(3\delta)^\alpha < \frac{1}{2}\delta_0$ and $C_4(L+1)\delta < \delta_0$. Choose $N_2 \geq 1$ such that $\lambda^{N_2} < \varepsilon/\delta_0$ and take $N = N_1 + N_2$.

We claim that

- (*) For all $z \in \Lambda$, there exists $i \in \{1, \dots, N_1\}$ such that $\pi(\phi_{i+N_2}W_\varepsilon^u(z) \cap \tilde{\mathcal{D}}_L) \supset \mathcal{D}_L$.

Fix $z \in \Lambda$. By the definition of N_1 , there exists $1 \leq i \leq N_1$ such that $d_M(\phi_{-i}p, \phi_{N_2}z) < \delta$. Let $y \in \mathcal{D}_L$. Then

$$\begin{aligned} d_M(\phi_{-i}y, \phi_{N_2}z) &\leq d(\phi_{-i}y, \phi_{-i}p) + d_M(\phi_{-i}p, \phi_{N_2}z) \\ &\leq d(y, p) + d_M(\phi_{-i}p, \phi_{N_2}z) < (L+1)\delta. \end{aligned}$$

Using the local product structure and choice of δ , we can define $x \in W_{\delta_1}^{cs}(\phi_{-i}y) \cap W_{\delta_1}^u(\phi_{N_2}z)$. Then $\phi_i x \in W_{\delta_1}^{cs}(y) \subset \tilde{\mathcal{D}}_L$ and $g_i x = \pi \phi_i x = y$. Also,

$$d(x, \phi_{N_2}z) \leq C_4 d_M(\phi_{-i}y, \phi_{N_2}z) < C_4(L+1)\delta < \delta_0.$$

By the definition of N_2 ,

$$\phi_i x \in \phi_i W_{\delta_0}^u(\phi_{N_2}z) \subset \phi_{i+N_2}W_\varepsilon^u(z).$$

Hence we obtain that $y = \pi \phi_i x \in \pi(\phi_{i+N_2}W_\varepsilon^u(z) \cap \tilde{\mathcal{D}}_L)$ proving (*).

Next, we claim that

- (**) For all $z \in \phi_n A_{n-1}$, $n \geq 1$, there exist $i \in \{0, \dots, N\}$ and j such that $U_{n+i,j}^1 \subset \phi_{-n}W_{\delta_0}^u(z)$.

To prove (**), define $V_\varepsilon = \phi_{-n}W_\varepsilon^u(z)$. By Proposition 2.5(b), $V_\varepsilon \subset A_{n-1}^{(\varepsilon)}$. We now consider two possible cases.

Suppose first that $V_\varepsilon \subset A_{n+i}$ for all $0 \leq i \leq N$. By claim (*), there exists $1 \leq i \leq N = N_1 + N_2$ such that

$$\pi(\phi_{n+i}V_\varepsilon \cap \tilde{\mathcal{D}}_L) = \pi(\phi_i W_\varepsilon^u(z) \cap \tilde{\mathcal{D}}_L) \supset \mathcal{D}_L,$$

while $V_\varepsilon \subset A_{n+i-1}$ by assumption. This means that $V_\varepsilon \supset U_{n+i,j}^L$ for some j . Hence

$$U_{n+i,j}^1 \subset U_{n+i,j}^L \subset V_\varepsilon \subset \phi_{-n}W_{\delta_0}^u(z),$$

and we are done.

In this way, we reduce to the second case where there exists $0 \leq i \leq N$ least such that $V_\varepsilon \not\subset A_{n+i}$. Since i is least, $V_\varepsilon \subset A_{n+i-1}^{(\varepsilon)}$. (The ε is required in case $i = 0$.) By Proposition 2.5(a), $V_\varepsilon \subset \{t_{n+i-1} \leq 1\}$. Hence

$$\begin{aligned} V_\varepsilon \setminus A_{n+i} &= (V_\varepsilon \cap B_{n+i}) \cup (V_\varepsilon \cap \{R = n+i\}) \\ &\subset \{t_{n+i-1} \leq 1, t_{n+i} \geq 1\} \cup \{R = n+i\} \subset \bigcup_j U_{n+i,j}^2. \end{aligned}$$

Since $V_\varepsilon \setminus A_{n+i} \neq \emptyset$, this means that there exists j so that V_ε intersects $U_{n+i,j}^2$. Hence we can choose $a_2 \in W_\varepsilon^u(z) \cap \phi_n U_{n+i,j}^2$.

Recall that $\phi_{n+i}U_{n+i,j}^m \subset \tilde{\mathcal{D}}_m$ and $g_{n+i}U_{n+i,j}^m = \mathcal{D}_m$ for $m = 1, 2$. In particular, $b_2 = \phi_i a_2 \in \tilde{\mathcal{D}}_2$ and $c_2 = g_i a_2 \in \mathcal{D}_2$.

Let $c_1 \in \mathcal{D}_1$. Then $d_M(c_1, b_2) \leq d_M(c_1, c_2) + d_M(c_2, b_2) < 3\delta + \delta_1 < 4\delta_1$. Hence, using the local product structure and definition of δ_1 , we can define $b_1 \in W_{\delta_0}^{cs}(c_1) \cap W_{\delta_0}^u(b_2)$ and $a_1 = \phi_{-i}b_1$. Note that

$$\phi_i a_r = b_r, \quad \pi b_r = c_r, \quad r = 1, 2.$$

Hence

$$d(a_1, a_2) \leq d(b_1, b_2) \leq C_3 d(c_1, c_2)^\alpha < C_3 (3\delta)^\alpha < \frac{1}{2} \delta_0,$$

and so $d(a_1, z) \leq d(a_1, a_2) + d(a_2, z) < \frac{1}{2} \delta_0 + \varepsilon < \delta_0$. It follows that $a_1 \in W_{\delta_0}^u(z)$ and thereby that $c_1 \in g_i(W_{\delta_0}^u(z) \cap \phi_{-i}\widehat{\mathcal{D}}_1)$. This proves that $\mathcal{D}_1 \subset g_i(W_{\delta_0}^u(z) \cap \phi_{-i}\widehat{\mathcal{D}}_1)$. Hence $U_{n+i,j}^1 \subset g_{-(n+i)}\mathcal{D}_1 \subset \phi_{-n}W_{\delta_0}^u(z)$ verifying claim (**).

We are now in a position to complete the proof of the lemma. Let $n \geq 1$, and let $Z \subset \phi_n A_{n-1}$ be a maximal set of points such that the balls $W_{\delta_0/2}^u(z)$ are disjoint for $z \in Z$. If $x \in \phi_n A_{n-1}$, then $W_{\delta_0/2}^u(x)$ intersects at least one $W_{\delta_0/2}^u(z)$, $z \in Z$, by maximality of the set Z . Hence $\phi_n A_{n-1} \subset \bigcup_{z \in Z} W_{\delta_0}^u(z)$. It follows that

$$A_{n-1} \subset \bigcup_{z \in Z} \phi_{-n} W_{\delta_0}^u(z).$$

Let $z \in Z$ and let $U_z = U_{n+i,j}^1$ be as in claim (**). In particular, $g_{n+i}U_z = \mathcal{D}_1 = W_\delta^u(p)$. Also, $\text{Leb}(\phi_{n+i}U_z) \leq |D\phi_1|E^u|_\infty^{im} \text{Leb}(\phi_n U_z)$ where $m = \dim E^u$. Hence, by (2.2),

$$\frac{1}{\text{Leb}(\phi_n U_z)} \leq |D\phi_1|E^u|_\infty^{Nm} \frac{1}{\text{Leb}(\phi_{n+i}U_z)} \leq C_3 |D\phi_1|E^u|_\infty^{Nm} \frac{1}{\text{Leb}(W_\delta^u(p))}.$$

By (2.1),

$$\frac{\text{Leb}(\phi_{-n}W_{\delta_0}^u(z))}{\text{Leb}(U_z)} \leq C_1 \frac{\text{Leb}(W_{\delta_0}^u(z))}{\text{Leb}(\phi_n U_z)} \leq K,$$

where $K = C_1 C_3 |D\phi_1|E^u|_\infty^{Nm} \frac{\sup_{y \in Y} \text{Leb}(W_{\delta_0}^u(y))}{\text{Leb}(W_\delta^u(p))}$.

Finally, the sets U_z are connected components of $\bigcup_{0 \leq i \leq N} \{R = n + i\}$ lying in distinct disjoint sets $\phi_{-n}W_{\delta_0}^u(z)$. Hence

$$\begin{aligned} \text{Leb}(A_{n-1}) &\leq \sum_{z \in Z} \text{Leb}(\phi_{-n}W_{\delta_0}^u(z)) \\ &\leq K \sum_{z \in Z} \text{Leb}(U_z) \leq K \text{Leb} \left(\bigcup_{0 \leq i \leq N} \{R = n + i\} \right), \end{aligned}$$

as required. \square

We can now complete the proof of Theorem 2.1.

Corollary 2.10. $\text{Leb}(R > n) = O(\gamma^n)$ for some $\gamma \in (0, 1)$.

Proof. By Corollary 2.8 and Lemma 2.9,

$$\begin{aligned} \text{Leb}(R \geq n) &= \text{Leb}(A_{n-1}) + \text{Leb}(B_{n-1}) \\ &\leq (1 + a_0) \text{Leb}(A_{n-1}) \leq d_2 \text{Leb} \left(\bigcup_{i=0}^N \{R = n + i\} \right) \end{aligned}$$

where $d_2 = c_1^{-1}(1 + a_0)$. It follows that

$$\begin{aligned} d_2^{-1} \text{Leb}(R \geq n) &\leq \text{Leb}(R = n) + \cdots + \text{Leb}(R = n + N) \\ &= \text{Leb}(R \geq n) - \text{Leb}(R > n + N). \end{aligned}$$

Hence

$$\text{Leb}(R > n + N) \leq (1 - d_2^{-1}) \text{Leb}(R \geq n).$$

In particular, $\text{Leb}(R > kN) \leq \gamma^{kN}$ with $\gamma = (1 - d_2^{-1})^{1/N}$ and the result follows. \square

3. Exponential Decay of Correlations for Flows

In this section, we consider exponential decay of correlations for a class of uniformly hyperbolic skew product flows satisfying a uniform nonintegrability condition, generalising from C^2 flows as treated in [8] to $C^{1+\alpha}$ flows. In doing so, we remove the restriction in [6, 9] that unstable manifolds are one-dimensional.

The arguments are a straightforward combination of those in [6, 8]. We follow closely the presentation in [6], with the focus on incorporating the ideas from [8] where required.

Quotienting by stable leaves leads to a class of semiflows considered in Subsection 3.1. The flows are considered in Subsection 3.2.

The current section is completely independent from Sect. 2, so overlaps in notation will not cause any confusion.

3.1. $C^{1+\alpha}$ Uniformly Expanding Semiflows

Fix $\alpha \in (0, 1)$. Let $Y \subset \mathbb{R}^m$ be an open ball⁴ in Euclidean space with Euclidean distance d . We suppose that $\text{diam } Y = 1$. Let Leb denote Lebesgue measure on Y . Let \mathcal{P} be a countable partition mod 0 of Y consisting of open sets.

Suppose that $F : \bigcup_{U \in \mathcal{P}} U \rightarrow Y$ is $C^{1+\alpha}$ on each $U \in \mathcal{P}$ and maps U diffeomorphically onto Y . Let $\mathcal{H} = \{h : Y \rightarrow U : U \in \mathcal{P}\}$ denote the family of inverse branches, and let \mathcal{H}_n denote the inverse branches for F^n . We say that F is a $C^{1+\alpha}$ *uniformly expanding map* if there exist constants $C_1 \geq 1$, $\rho_0 \in (0, 1)$ such that

- (i) $\|Dh\|_\infty \leq C_1 \rho_0^n$ for all $h \in \mathcal{H}_n$, $n \geq 1$;
- (ii) $|\log |\det Dh||_\alpha \leq C_1$ for all $h \in \mathcal{H}$;

⁴More generally, we could consider a John domain as in [8] but the current setting suffices for our purposes.

where $|\psi|_\alpha = \sup_{y \neq y'} |\psi(y) - \psi(y')|/d(y, y')^\alpha$. Under these assumptions, it is standard [1] that there exists a unique F -invariant absolutely continuous measure μ . The density $d\mu/d\text{Leb}$ is C^α , bounded above and below, and μ is ergodic and mixing.

We consider roof functions $r : \bigcup_{U \in \mathcal{P}} U \rightarrow \mathbb{R}^+$ that are C^1 on partition elements U with $\inf r > 0$. Define the suspension $Y^r = \{(y, u) \in Y \times \mathbb{R} : 0 \leq u \leq r(y)\} / \sim$ where $(y, r(y)) \sim (Fy, 0)$. The suspension semiflow $F_t : Y^r \rightarrow Y^r$ is given by $F_t(y, u) = (y, u + t)$ computed modulo identifications, with ergodic invariant probability measure $\mu^r = (\mu \times \text{Lebesgue})/\bar{r}$ where $\bar{r} = \int_Y r d\mu$. We say that F_t is a $C^{1+\alpha}$ *uniformly expanding semiflow* if F is a $C^{1+\alpha}$ uniformly expanding map and we can choose C_1 from condition (i) and $\varepsilon > 0$ such that

- (iii) $|D(r \circ h)|_\infty \leq C_1$ for all $h \in \mathcal{H}$;
- (iv) $\sum_{h \in \mathcal{H}} e^{\varepsilon|r \circ h|_\infty} |\det Dh|_\infty < \infty$.

Let $r_n = \sum_{j=0}^{n-1} r \circ F^j$ and define

$$\psi_{h_1, h_2} = r_n \circ h_1 - r_n \circ h_2 : Y \rightarrow \mathbb{R},$$

for $h_1, h_2 \in \mathcal{H}_n$. We require the following *uniform nonintegrability* condition [8, Equation (6.6)]:

- (UNI) There exists $E > 0$ and $h_1, h_2 \in \mathcal{H}_{n_0}$, for some sufficiently large $n_0 \geq 1$, with the following property: There exists a continuous unit vector field $\ell : \mathbb{R}^m \rightarrow \mathbb{R}^m$ such that $|D\psi_{h_1, h_2}(y) \cdot \ell(y)| \geq E$ for all $y \in Y$.

(The requirement “sufficiently large” can be made explicit as in [6, Equations (2.1) to (2.3)].) From now on, n_0, h_1 and h_2 are fixed.

Define $F_\alpha(Y^r)$ to consist of L^∞ functions $v : Y^r \rightarrow \mathbb{R}$ such that $\|v\|_\alpha = |v|_\infty + |v|_\alpha < \infty$ where

$$|v|_\alpha = \sup_{(y, u) \neq (y', u)} \frac{|v(y, u) - v(y', u)|}{d(y, y')^\alpha}.$$

Define $F_{\alpha, k}(Y^r)$ to consist of functions with $\|v\|_{\alpha, k} = \sum_{j=0}^k \|\partial_t^j v\|_\alpha < \infty$ where ∂_t denotes differentiation along the semiflow direction.

We can now state the main result in this section. Given $v \in L^1(Y^r)$, $w \in L^\infty(Y^r)$, define the correlation function

$$\rho_{v, w}(t) = \int v w \circ F_t d\mu^r - \int v d\mu^r \int w d\mu^r.$$

Theorem 3.1. *Suppose that $F_t : Y^r \rightarrow Y^r$ is a $C^{1+\alpha}$ uniformly expanding semiflow satisfying (UNI). Then there exist constants $c, C > 0$ such that*

$$|\rho_{v, w}(t)| \leq C e^{-ct} \|v\|_{\alpha, 1} \|w\|_{\alpha, 1} \quad \text{for all } t > 0 \text{ and all } v, w \in F_{\alpha, 1}(Y^r),$$

(alternatively, all $v \in F_{\alpha, 2}(Y^r)$, $w \in L^\infty(Y^r)$).

In the remainder of this subsection, we prove Theorem 3.1.

For $s \in \mathbb{C}$, let P_s denote the (non-normalized) transfer operator

$$P_s = \sum_{h \in \mathcal{H}} A_{s, h}, \quad A_{s, h} v = e^{-s r \circ h} |\det Dh| v \circ h.$$

For $v : Y \rightarrow \mathbb{C}$, define $\|v\|_\alpha = \max\{|v|_\infty, |v|_\alpha\}$ where $|v|_\alpha = \sup_{y \neq y'} \frac{|v(y) - v(y')|}{d(y, y')^\alpha}$. Let $C^\alpha(Y)$ denote the space of functions $v : Y \rightarrow \mathbb{C}$ with $\|v\|_\alpha < \infty$. We introduce the family of equivalent norms

$$\|v\|_b = \max\{|v|_\infty, |v|_\alpha/(1 + |b|^\alpha)\}, \quad b \in \mathbb{R}.$$

Proposition 3.2. *Write $s = \sigma + ib$. There exists $\varepsilon \in (0, 1)$ such that the family $s \mapsto P_s$ of operators on $C^\alpha(Y)$ is continuous on $\{\sigma > -\varepsilon\}$. Moreover, $\sup_{|\sigma| < \varepsilon} \|P_s\|_b < \infty$.*

Proof. The first five lines of the proof of [6, Proposition 2.5] should be changed to the following:

Using the inequality $1 - t \leq -\log t$ valid for $t > 0$, we obtain for $a > b > 0$ that $a - b = a(1 - \frac{b}{a}) \leq -a \log \frac{b}{a} = a(\log a - \log b)$. Hence $|\log |\det Dh(x)| - \log |\det Dh(y)|| \leq |\det Dh|_\infty (\log |\det Dh(x)| - \log |\det Dh(y)|)$ and so by (ii),

$$|\log |\det Dh(x)| - \log |\det Dh(y)|| \leq C_1 |\det Dh|_\infty d(x, y)^\alpha \quad \text{for all } h \in \mathcal{H}, x, y \in \mathfrak{X}1$$

The proof now proceeds exactly as for [6, Proposition 2.5] (with R, h' and $|x - y|$ changed to $r, \det Dh$ and $d(x, y)$). \square

The unperturbed operator P_0 has a simple leading eigenvalue $\lambda_0 = 1$ with strictly positive C^α eigenfunction f_0 . By Proposition 3.2, there exists $\varepsilon \in (0, 1)$ such that P_σ has a continuous family of simple eigenvalues λ_σ for $|\sigma| < \varepsilon$ with associated C^α eigenfunctions f_σ . For $s = \sigma + ib$ with $|\sigma| \leq \varepsilon$, we define the normalized transfer operators

$$L_s v = (\lambda_\sigma f_\sigma)^{-1} P_s(f_\sigma v) = (\lambda_\sigma f_\sigma)^{-1} \sum_{h \in \mathcal{H}} A_{s,h}(f_\sigma v).$$

In particular, $L_\sigma 1 = 1$ and $|L_s|_\infty \leq 1$.

Set $C_2 = C_1^2/(1 - \rho)$, $\rho = \rho_0^\alpha$. Then

- (ii₁) $|\log |\det Dh||_\alpha \leq C_2$ for all $h \in \mathcal{H}_n, n \geq 1$,
- (iii₁) $|D(r_n \circ h)|_\infty \leq C_2$ for all $h \in \mathcal{H}_n, n \geq 1$.

Write

$$L_s^n v = \lambda_\sigma^{-n} f_\sigma^{-1} \sum_{h \in \mathcal{H}_n} A_{s,h,n}(f_\sigma v), \quad A_{s,h,n} v = e^{-sr_n \circ h} |\det Dh| v \circ h.$$

Lemma 3.3 (Lasota-Yorke inequality). *There is a constant $C_3 > 1$ such that*

$$|L_s^n v|_\alpha \leq C_3(1 + |b|^\alpha)|v|_\infty + C_3 \rho^n |v|_\alpha \leq C_3(1 + |b|^\alpha)\{|v|_\infty + \rho^n \|v\|_b\},$$

for all $s = \sigma + ib$, $|\sigma| < \varepsilon$, and all $n \geq 1$, $v \in C^\alpha(Y)$.

Proof. It follows from (ii₁) that

$$|\log |\det Dh(x)| - \log |\det Dh(y)|| \leq C_2 |\det Dh|_\infty d(x, y)^\alpha \leq C_2 e^{C_2} |\det Dh(z)| d(x, y)^\alpha$$

for all $h \in \mathcal{H}_n, n \geq 1, x, y, z \in Y$. The proof now proceeds exactly as for [6, Lemma 2.7]. \square

Corollary 3.4. $\|L_s^n\|_b \leq 2C_3$ for all $s = \sigma + ib$, $|\sigma| < \varepsilon$, and all $n \geq 1$.

Proof. This is unchanged from [6, Corollary 2.8]. \square

Given $b \in \mathbb{R}$, we define the cone

$$C_b = \left\{ (u, v) : u, v \in C^\alpha(Y), u > 0, 0 \leq |v| \leq u, |\log u|_\alpha \leq C_4|b|^\alpha, \right. \\ \left. |v(x) - v(y)| \leq C_4|b|^\alpha u(y)d(x, y)^\alpha \text{ for all } x, y \in Y \right\}.$$

(The constant C_4 is specified in Lemma 3.8.)

Throughout $B_\delta(y) = \{x \in \mathbb{R}^m : d(x, y) < \delta\}$.

Lemma 3.5 (Cancellation Lemma). *Assume that the (UNI) condition is satisfied (with associated constants $E > 0$ and $n_0 \geq 1$). Let $h_1, h_2 \in \mathcal{H}_{n_0}$ be the branches from (UNI).*

There exists $0 < \delta < \Delta = 4\pi/E$ such that for all $s = \sigma + ib$, $|\sigma| < \varepsilon$, $|b| \geq 1$, and all $(u, v) \in C_b$ we have the following:

For every $y' \in Y$ with $B_{(\delta+\Delta)/|b|}(y') \subset Y$, there exists $y'' \in B_{\Delta/|b|}(y')$ such that one of the following inequalities holds on $B_{\delta/|b|}(y'')$:

$$\text{Case } h_1: |A_{s, h_1, n_0}(f_\sigma v) + A_{s, h_2, n_0}(f_\sigma v)| \leq \frac{3}{4}A_{\sigma, h_1, n_0}(f_\sigma u) + A_{\sigma, h_2, n_0}(f_\sigma u),$$

$$\text{Case } h_2: |A_{s, h_1, n_0}(f_\sigma v) + A_{s, h_2, n_0}(f_\sigma v)| \leq A_{\sigma, h_1, n_0}(f_\sigma u) + \frac{3}{4}A_{\sigma, h_2, n_0}(f_\sigma u).$$

Proof. Let $\theta = V - b\psi_{h_1, h_2}$ where $\psi_{h_1, h_2} = r_{n_0} \circ h_1 - r_{n_0} \circ h_2$ and $V = \arg(v \circ h_1) - \arg(v \circ h_2)$.

We follow the following steps from [6, Lemma 2.9]:

- (1) Reduce to the situation where $|v(h_m y')| > \frac{1}{2}u(h_m y')$ for both $m = 1$ and $m = 2$.
- (2) Establish the estimate $|V(y) - V(y')| \leq \pi/6$ for all $y \in B_{(\delta+\Delta)/|b|}(y')$.
- (3) Construct $y'' \in B_{\Delta/|b|}(y')$ such that

$$b(\psi_{h_1, h_2}(y'') - \psi_{h_1, h_2}(y')) = \theta(y'') - \pi \pmod{2\pi}.$$

- (4) Deduce that $|\theta(y) - \pi| \leq 2\pi/3$ for all $y \in B_{(\delta+\Delta)/|b|}(y')$.
- (5) Conclude the desired result.

Only step (3) requires any change from the argument in [6, Lemma 2.9]. We provide here the modified argument. Approximate the continuous unit vector field $\ell : \mathbb{R}^m \rightarrow \mathbb{R}^m$ in (UNI) by a smooth vector field $\ell : \mathbb{R}^m \rightarrow \mathbb{R}^m$ with $|\ell(x)| \leq 1$ for all $x \in \mathbb{R}^m$. By condition (iii)₁, the approximation can be chosen close enough that

$$|D\psi_{h_1, h_2}(y) \cdot \ell(y)| \geq \frac{1}{2}E \quad \text{for all } y \in Y. \quad (3.2)$$

Let $g : [0, \Delta/|b|] \rightarrow \mathbb{R}^m$ be the solution to the initial value problem

$$\frac{dg}{dt} = \ell \circ g, \quad g(0) = y'$$

and set $y_t = g(t)$. Note that $d(y_t, y') \leq \int_0^t |\ell(g(s))| ds \leq \Delta/|b|$, so $y_t \in B_{\Delta/|b|}(y')$ for all $t \in [0, \Delta/|b|]$. By the mean value theorem applied to $\psi_{h_1, h_2} \circ g : [0, \Delta/|b|] \rightarrow \mathbb{R}$ and (3.2),

$$|\psi_{h_1, h_2}(y_t) - \psi_{h_1, h_2}(y')| \geq t \inf_{s \in [0, \Delta/|b|]} |D\psi_{h_1, h_2}(y_s) \cdot \ell(y_s)| \geq \frac{1}{2}Et = (2\pi/\Delta)t$$

for all $t \in [0, \Delta/|b|]$. It follows that $b(\psi_{h_1, h_2}(y_t) - \psi_{h_1, h_2}(y'))$ fills out an interval around 0 of length at least 2π as t varies in $[0, \Delta/|b|]$. In particular, we can choose $y'' \in B_{\Delta/|b|}(y')$ such that (3) holds. \square

Let $\{y'_1, \dots, y'_k\} \subset Y$ be a maximal set of points such that the open balls $B_{(\delta+\Delta)/|b|}(y'_i)$ are disjoint and contained in Y .

Let $(u, v) \in \mathcal{C}_b$. For each $i = 1, \dots, k$, there exists a ball $B_i = B_{\delta/|b|}(y''_i)$ on which the conclusion of Lemma 3.5 holds. Write $\text{type}(B_i) = h_m$ if we are in case h_m . Let $\widehat{B}_i = B_{\frac{1}{2}\delta/|b|}(y''_i)$

There exists a universal constant $C > 0$ and a C^1 function $\omega_i : Y \rightarrow [0, 1]$ such that $\omega_i \equiv 1$ on \widehat{B}_i , $\omega_i \equiv 0$ on $Y \setminus B_i$, and $\|\omega_i\|_{C^1} \leq C|b|/\delta$. Define $\omega : Y \rightarrow [0, 1]$,

$$\omega(y) = \begin{cases} \sum_{\text{type}(B_i)=h_m} \omega_i(F^{n_0}y), & y \in \text{range } h_m, m = 1, 2 \\ 0 & \text{otherwise.} \end{cases}$$

Note that $\|\omega\|_{C^1} \leq C'|b|$ where $C' = C\delta$ is independent of $(u, v) \in \mathcal{C}_b$ and $s \in \mathbb{C}$, and we can assume that $C' > 4$. Then $\chi = 1 - \omega/C' : Y \rightarrow [\frac{3}{4}, 1]$ satisfies $|D\chi| \leq |b|$. Moreover, if $\text{type}(B_i) = h_m$ then $\chi \equiv \eta$ on $\text{range } h_m$ where $\eta = 1 - 1/C' \in (0, 1)$.

Corollary 3.6. *Let δ, Δ be as in Lemma 3.5. Let $|b| \geq 1$, $(u, v) \in \mathcal{C}_b$. Let $\chi = \chi(b, u, v)$ be the C^1 function described above (using the branches $h_1, h_2 \in \mathcal{H}_{n_0}$ from (UNI)). Then $|L_s^{n_0}v| \leq L_\sigma^{n_0}(\chi u)$ for all $s = \sigma + ib$, $|\sigma| < \varepsilon$.*

Proof. This is immediate from Lemma 3.5 and the definition of χ . \square

Define the disjoint union $\widehat{B} = \bigcup \widehat{B}_i$.

Proposition 3.7. *Let $K > 0$. There exists $c_1 > 0$ such that $\int_{\widehat{B}} w d\mu \geq c_1 \int_Y w d\mu$ for all C^α function $w : Y \rightarrow (0, \infty)$ with $|\log w|_\alpha \leq K|b|^\alpha$, for all $|b| \geq 16\pi/E$.*

Proof. Let $y \in Y$. Since $(\delta + \Delta)/|b| \leq 2\Delta/|b| = 8\pi/(E|b|) \leq \frac{1}{2}$, there exists $z \in Y$ with $B_{(\delta+\Delta)/|b|}(z) \subset Y$ such that $d(z, y) < (\delta + \Delta)/|b|$. By maximality of the set of points $\{y'_1, \dots, y'_k\}$, there exists y'_i such that $B_{(\delta+\Delta)/|b|}(z)$ intersects $B_{(\delta+\Delta)/|b|}(y'_i)$. Hence $Y \subset \bigcup_{i=1}^k B_i^*$ where $B_i^* = B_{3(\delta+\Delta)/|b|}(y'_i)$. Since the density $d\mu/d\text{Leb}$ is bounded above and below, there is a constant $c_2 > 0$ such that $\mu(\widehat{B}_i) \geq c_2\mu(B_i^*)$ for each i .

Let $x \in \widehat{B}_i, y \in B_i^*$. Then $d(x, y) \leq 4(\delta + \Delta)/|b|$ and so $|w(x)/w(y)| \leq e^{K'}$ where $K' = \{4(\delta + \Delta)\}^\alpha K$. It follows that

$$\int_{\widehat{B}_i} w d\mu \geq \mu(\widehat{B}_i) \inf_{\widehat{B}_i} w \geq c_2 e^{-K'} \mu(B_i^*) \sup_{B_i^*} w \geq c_1 \int_{B_i^*} w d\mu,$$

where $c_1 = c_2 e^{-K'}$. Since the sets $\widehat{B}_i \subset Y$ are disjoint,

$$\int_{\widehat{B}} w d\mu = \sum_i \int_{\widehat{B}_i} w d\mu \geq c_1 \sum_i \int_{B_i^*} w d\mu \geq c_1 \int_Y w d\mu$$

as required. \square

Lemma 3.8 (Invariance of cone). *There is a constant C_4 depending only on $C_1, C_2, |f_0^{-1}|_\infty$ and $|f_0|_\alpha$ such that the following holds:*

For all $(u, v) \in \mathcal{C}_b$, we have that

$$(L_\sigma^{n_0}(\chi u), L_s^{n_0}v) \in \mathcal{C}_b,$$

for all $s = \sigma + ib$, $|\sigma| < \varepsilon$, $|b| \geq 1$. (Here, $\chi = \chi(b, u, v)$ is from Corollary 3.6.)

Proof. This is unchanged from [6, Lemma 2.12]. □

Lemma 3.9 (L^2 contraction). *There exist $\varepsilon, \beta \in (0, 1)$ such that*

$$\int_Y |L_s^{mn_0}v|^2 d\mu \leq \beta^m |v|_\infty^2$$

for all $m \geq 1$, $s = \sigma + ib$, $|\sigma| < \varepsilon$, $|b| \geq \max\{16\pi/E, 1\}$, and all $v \in C^\alpha(Y)$ satisfying $|v|_\alpha \leq C_4|b|^\alpha|v|_\infty$.

Proof. Define $u_0 \equiv 1, v_0 = v/|v|_\infty$ and inductively,

$$u_{m+1} = L_\sigma^{n_0}(\chi_m u_m), \quad v_{m+1} = L_s^{n_0}(v_m),$$

where $\chi_m = \chi(b, u_m, v_m)$. It is immediate from the definitions that $(u_0, v_0) \in \mathcal{C}_b$, and it follows from Lemma 3.8 that $(u_m, v_m) \in \mathcal{C}_b$ for all m . Hence inductively the χ_m are well-defined as in Corollary 3.6.

We proceed as in [6, Lemma 2.13] in the following steps.

- (1) It suffices to show that there exists $\beta \in (0, 1)$ such that $\int_Y u_{m+1}^2 d\mu \leq \beta \int u_m^2 d\mu$ for all m .
- (2) Define $w = L_0^{n_0}(u_m^2)$. Then

$$u_{m+1}^2(y) \leq \begin{cases} \xi(\sigma)\eta_1 w(y) & y \in \widehat{B} \\ \xi(\sigma)w(y) & y \in Y \setminus \widehat{B} \end{cases}$$

where $\xi(\sigma)$ can be made as close to 1 as desired by shrinking ε . Here, $\eta_1 \in (0, 1)$ is a constant independent of v, m, s, y .

- (3) The function $w : Y \rightarrow \mathbb{R}$ satisfies the hypotheses of Proposition 3.7; consequently $\int_{\widehat{B}} w d\mu \geq c_1 \int_{Y \setminus \widehat{B}} w d\mu$. This leads to the desired conclusion. □

Lemma 3.10 (C^α contraction). *Let $E' = \max\{16\pi/E, 2\}$. There exists $\varepsilon \in (0, 1)$, $\gamma \in (0, 1)$ and $A > 0$ such that $\|P_s^n\|_b \leq \gamma^n$ for all $s = \sigma + ib$, $|\sigma| < \varepsilon$, $|b| \geq E'$, $n \geq A \log |b|$.*

Proof. This is unchanged from [6, Proposition 2.14, Corollary 2.15 and Theorem 2.16]. □

Proof of Theorem 3.1. This is identical to [6, Section 2.7]. We note that there is a typo in the statement of [6, Lemma 2.23] where $|b| \leq D'$ should be $|b| \geq D'$

(twice). Also, for the second statement of [6, Proposition 2.18] it would be more natural to argue that

$$\begin{aligned} \int_Y e^{\varepsilon r} d\text{Leb} &= \sum_{h \in \mathcal{H}} \int_{h(Y)} e^{\varepsilon r} d\text{Leb} \\ &= \sum_{h \in \mathcal{H}} \int_Y e^{\varepsilon r \circ h} |\det Dh| d\text{Leb} \leq \text{Leb}(Y) \sum_{h \in \mathcal{H}} e^{\varepsilon |r \circ h|_\infty} |\det Dh|_\infty \end{aligned}$$

which is finite by condition (iv). Hence $\int_Y e^{\varepsilon r} d\mu < \infty$ by boundedness of $d\mu/d\text{Leb}$.

3.2. $C^{1+\alpha}$ Uniformly Hyperbolic Skew Product Flows

Let $X = Y \times Z$ where Y is an open ball of diameter 1 with Euclidean metric d_Y and (Z, d_Z) is a compact Riemannian manifold. Define the metric $d((y, z), (y', z')) = d_Y(y, y') + d_Z(z, z')$ on X . Let $f(y, z) = (Fy, G(y, z))$ where $F : Y \rightarrow Y$, $G : X \rightarrow Z$ are $C^{1+\alpha}$.

We say that $f : X \rightarrow X$ is a $C^{1+\alpha}$ *uniformly hyperbolic skew product* if $F : Y \rightarrow Y$ is a $C^{1+\alpha}$ uniformly expanding map satisfying conditions (i) and (ii) as in Sect. 3.1, with absolutely continuous invariant probability measure μ , and moreover

- (v) There exist constants $C > 0$, $\gamma_0 \in (0, 1)$ such that $d(f^n(y, z), f^n(y, z')) \leq C\gamma_0^n d(z, z')$ for all $y \in Y$, $z, z' \in Z$.

Let $\pi^s : X \rightarrow Y$ be the projection $\pi^s(y, z) = y$. This defines a semiconjugacy between f and F and there is a unique f -invariant ergodic probability measure μ_X on X such that $\pi_*^s \mu_X = \mu$.

Suppose that $r : \bigcup_{U \in \mathcal{P}} U \rightarrow \mathbb{R}^+$ is C^1 on partition elements U with $\inf r > 0$. Define $r : X \rightarrow \mathbb{R}^+$ by setting $r(y, z) = r(y)$. Define the suspension $X^r = \{(x, u) \in X \times \mathbb{R} : 0 \leq u \leq r(x)\} / \sim$ where $(x, r(x)) \sim (fx, 0)$. The suspension flow $f_t : X^r \rightarrow X^r$ is given by $f_t(x, u) = (x, u + t)$ computed modulo identifications, with ergodic invariant probability measure $\mu_X^r = (\mu_X \times \text{Lebesgue}) / \bar{r}$.

We say that f_t is a $C^{1+\alpha}$ *uniformly hyperbolic skew product flow* provided $f : X \rightarrow X$ is a $C^{1+\alpha}$ uniformly hyperbolic skew product as above, and $r : Y \rightarrow \mathbb{R}^+$ satisfies conditions (iii) and (iv) as in Sect. 3.1. If $F : Y \rightarrow Y$ and $r : Y \rightarrow \mathbb{R}^+$ satisfy condition (UNI) from Sect. 3.1, then we say that the skew product flow f_t satisfies (UNI).

Define $F_\alpha(X^r)$ to consist of L^∞ functions $v : X^r \rightarrow \mathbb{R}$ such that $\|v\|_\alpha = |v|_\infty + |v|_\alpha < \infty$ where

$$|v|_\alpha = \sup_{(y, z, u) \neq (y', z', u)} \frac{|v(y, z, u) - v(y', z', u)|}{d((y, z), (y', z'))^\alpha}.$$

Define $F_{\alpha, k}(X^r)$ to consist of functions with $\|v\|_{\alpha, k} = \sum_{j=0}^k \|\partial_t^j v\|_\alpha < \infty$ where ∂_t denotes differentiation along the flow direction.

We can now state the main result in this section. Given $v \in L^1(X^r)$, $w \in L^\infty(X^r)$, define the correlation function

$$\rho_{v,w}(t) = \int v w \circ f_t d\mu_X^r - \int v d\mu_X^r \int w d\mu_X^r.$$

Theorem 3.11. *Assume that $f_t : X \rightarrow X$ is a $C^{1+\alpha}$ hyperbolic skew product flow satisfying the (UNI) condition. Then there exist constants $c, C > 0$ such that*

$$|\rho_{v,w}(t)| \leq C e^{-ct} \|v\|_{\alpha,1} \|w\|_{\alpha,1},$$

for all $t > 0$ and all $v, w \in F_{\alpha,1}(X^r)$ (alternatively all $v \in F_{\alpha,2}(X^r)$, $w \in F_\alpha(X^r)$).

Proof. This is unchanged from [6, Section 4]. \square

4. Proof of Theorem 1.2

We return to the situation of Sect. 2, so $\Lambda \subset M$ is a uniformly hyperbolic attractor for a $C^{1+\alpha}$ flow, $\alpha \in (0, 1)$, defined on a compact Riemannian manifold. Define the open unstable disk $Y = W_\delta^u(p)$ with discrete return time $R : Y \rightarrow \mathbb{Z}^+$ and induced map $F = \pi \circ \phi_R : Y \rightarrow Y$ as in Theorem 2.1.

Under smoothness assumptions on holonomies, we verify the conditions on the suspension flow f_t in Sect. 3 and obtain Theorem 1.2 as an easy consequence.

Proposition 4.1. *Suppose that the center-stable holonomies are $C^{1+\alpha}$. (In particular, $\pi : \widehat{\mathcal{D}} \rightarrow \mathcal{D}$ is $C^{1+\alpha}$.) Then (after shrinking δ_0 in Sect. 2 if necessary) F is a $C^{1+\alpha}$ uniformly expanding map.*

Proof. As in Remark 2.2, it is immediate that $F|_U : U \rightarrow Y$ is a $C^{1+\alpha}$ diffeomorphism for all $U \in \mathcal{P}$. Let $h : Y \rightarrow U$ be an inverse branch with $R|_U = n$, and define $\pi_U = \pi|_{\phi_n(U)} : \phi_n(U) \rightarrow \mathcal{D}$. Then

$$\lambda^{-1}|v| \leq \lambda^{-n}|v| \leq |D\phi_n(x)v| \leq |(D\pi_U)^{-1}|_\infty |DF(x)v|$$

for all $x \in U$, $v \in T_x Y$. Hence $|Dh|_\infty \leq \rho_0$ where $\rho_0 = \lambda \sup_U |(D\pi_U)^{-1}|_\infty$. Shrinking δ_0 , we can ensure that $\rho_0 < 1$. In particular, condition (i) in Sect. 3.1 holds (with $C_1 = 1$). Condition (ii) is the standard distortion estimate. \square

In the remainder of this section, we suppose moreover that the stable holonomies are $C^{1+\alpha}$. Shrink $\delta_0 \in (0, 1)$ as in Proposition 4.1 and shrink $\delta_1 \in (0, \delta_0)$ so that $\phi_t(W_{\delta_1}^s(y)) \subset W_{\delta_0}^s(\phi_t y)$ for all $t > 0$, $y \in \Lambda$. Recall that $\mathcal{D} = W_{\delta_0}^u(p)$ and

$$\widehat{\mathcal{D}} = \bigcup_{y \in \mathcal{D}} W_{\delta_0}^{cs}(y) = \bigcup_{|t| < \delta_0} \phi_t \left(\bigcup_{y \in \mathcal{D}} W_{\delta_0}^s(y) \right).$$

The projection $\pi^s : \bigcup_{y \in \mathcal{D}} W_{\delta_0}^s(y) \rightarrow \mathcal{D}$ given by $\pi^s|_{W_{\delta_0}^s(y)} \equiv y$ is $C^{1+\alpha}$. Moreover, $\pi = \pi^s \circ \phi_{r_0}$ where $\phi_{r_0} : \widehat{\mathcal{D}} \rightarrow \bigcup_{y \in \mathcal{D}} W_{\delta_0}^s(y)$ and $r_0 : \widehat{\mathcal{D}} \rightarrow (-\delta_0, \delta_0)$ is

$C^{1+\alpha}$. Define $r = R + r_0$ on Y . The choice $\delta_0 < 1$ ensures that $\inf r \geq 1 - \delta_0 > 0$. Define the corresponding semiflow $F_t : Y^r \rightarrow Y^r$.

Proposition 4.2. *$F_t : Y^r \rightarrow Y^r$ is a $C^{1+\alpha}$ uniformly expanding semiflow.*

Proof. By Proposition 4.1, F is a $C^{1+\alpha}$ uniformly expanding map. In particular, conditions (i) and (ii) are satisfied.

Notice that $F = \pi^s \circ \phi_r$ where $r = R + r_0 : Y \rightarrow \mathbb{R}^+$ is $C^{1+\alpha}$ on partition elements $U \in \mathcal{P}$. Since $Dr = Dr_0$ on partition elements, it is immediate that $\sup_{h \in \mathcal{H}} |D(r \circ h)|_\infty \leq |Dr_0|_\infty \sup_{h \in \mathcal{H}} |Dh|_\infty \leq \rho_0 |Dr_0|_\infty < \infty$ verifying condition (iii) on r . Recall that $\text{Leb}(R > n) = O(\gamma^n)$ for some $\gamma \in (0, 1)$, so we can choose $\varepsilon > 0$ such that $\int_Y e^{\varepsilon R} d\text{Leb} < \infty$. Condition (ii) ensures that $|\det Dh|_\infty \leq (\text{Leb } Y)^{-1} e^{C_1} \text{Leb}(\text{range } h)$ for all $h \in \mathcal{H}$. Hence $\sum_{h \in \mathcal{H}} e^{\varepsilon |r \circ h|_\infty} |\det Dh|_\infty \ll \sum_{h \in \mathcal{H}} e^{\varepsilon |R \circ h|_\infty} \text{Leb}(\text{range } h) = \int_Y e^{\varepsilon R} d\text{Leb} < \infty$ verifying condition (iv) on r . \square

We now make a $C^{1+\alpha}$ change of coordinates so that $\widehat{\mathcal{D}}$ is identified with $\mathcal{D} \times W_{\delta_0}^s(p) \times (-\delta_0, \delta_0)$ where $\{y\} \times W_{\delta_0}^s(p)$ is identified with $W_{\delta_0}^s(y)$ for all $y \in \mathcal{D}$ and $(-\delta_0, \delta_0)$ is the flow direction. Let $X = Y \times Z$ where $Z = W_{\delta_0}^s(p)$ and define $r : X \rightarrow (0, \infty)$ by $r(y, z) = r(y)$. Also, define $f = \phi_r : X \rightarrow X$ and the corresponding suspension flow $f_t : X^r \rightarrow X^r$

Proposition 4.3. *$f_t : X^r \rightarrow X^r$ is a $C^{1+\alpha}$ uniformly hyperbolic skew product flow.*

Proof. Note that $\pi^s(X) = Y$ and $\pi^s(y, z) = y$. Also, $f(y, z) = (Fy, G(y, z))$ where $G : X \rightarrow Z$ is $C^{1+\alpha}$. Since Z corresponds to the exponential contracting stable foliation, condition (v) in Sect. 3.2 is satisfied. Hence $f : X \rightarrow X$ is a $C^{1+\alpha}$ uniformly hyperbolic skew product and the corresponding suspension flow $f_t : X^r \rightarrow X^r$ is a $C^{1+\alpha}$ uniformly hyperbolic skew product flow. \square

Next we recall the standard argument that joint nonintegrability implies (UNI) in the current situation. (Similar arguments are given for instance in [7, Section 3] and [21, Section 5.3].)

Joint nonintegrability is defined in terms of the temporal distortion function. To define this intrinsically (independently of the inducing scheme) we have to introduce the first return time $\tau : X \rightarrow \mathbb{R}^+$ and the Poincaré map $g : X \rightarrow X$ given by

$$\tau(x) = \inf\{t > 0 : \phi_t(x) \in X\}, \quad g(x) = \phi_{\tau(x)}(x).$$

Note that τ is constant along stable leaves by the choice of X .

For $x_1, x_2 \in X$, define the local product $[x_1, x_2]$ to be the unique intersection point of $W^u(x_1) \cap W^s(x_2)$. The *temporal distortion function* D is defined to be

$$D(x_1, x_2) = \sum_{j=-\infty}^{\infty} \{\tau(g^j x_1) - \tau(g^j [x_1, x_2]) - \tau(g^j [x_2, x_1]) + \tau(g^j x_2)\}$$

at points $x_1, x_2 \in X$. The stable and unstable bundles are jointly integrable if and only if $D \equiv 0$.

Lemma 4.4. *Joint nonintegrability of the stable and unstable bundles implies (UNI).*

Proof. For points $x, x' \in X$ with $x' \in W^u(x)$, we define

$$D_0(x, x') = \sum_{j=1}^{\infty} \{\tau(g^{-j}x) - \tau(g^{-j}x')\}.$$

Since τ is constant along stable leaves,

$$\begin{aligned} D(x_1, x_2) &= \sum_{j=1}^{\infty} \{\tau(g^{-j}x_1) - \tau(g^{-j}[x_1, x_2]) - \tau(g^{-j}[x_2, x_1]) + \tau(g^{-j}x_2)\} \\ &= D_0(x_1, [x_1, x_2]) + D_0(x_2, [x_2, x_1]). \end{aligned}$$

Next, we find a more convenient expression for D_0 in terms of r and f . Note that for any $x \in X$, there exists $N(x) \in \mathbb{Z}^+$ (the number of returns to X up to time $r(x)$) such that

$$r(x) = \sum_{\ell=0}^{N(x)-1} \tau(g^\ell x), \quad f(x) = g^{N(x)}x.$$

Corresponding to the partition \mathcal{P} of Y , we define the collection $\tilde{\mathcal{P}} = \{\bar{U} \times \bar{Z} : U \in \mathcal{P}\}$ of closed subsets of X . Suppose that $x, x' \in V_0$, $V_0 \in \tilde{\mathcal{P}}$, with $x' \in W^u(x)$. The induced map $f : X \rightarrow X$ need not be invertible since it is not the first return to X . However, we may construct suitable inverse branches z_j, z'_j of x, x' as follows. Set $z_0 = x, z'_0 = x'$. Since f is transitive and continuous on closures of partition elements, there exists $V_1 \in \tilde{\mathcal{P}}$ and $z_1 \in V_1$ such that $fz_1 = z_0$. Since F is full-branch, $f(W^u(z_1) \cap V_1) \supset W^u(z_0)$, so there exists $z'_1 \in W^u(z_1) \cap V_1$ such that $fz'_1 = z'_0$. Inductively, we obtain $V_n \in \tilde{\mathcal{P}}$ and $z_j, z'_j \in V_n$ with $z'_j \in W^u(z_j)$ such that $fz_j = z_{j-1}$ and $fz'_j = z'_{j-1}$.

By construction, $z_{j-1} = fz_j = g^{N(z_j)}z_j$. Hence $z_j = g^{-(N(z_1)+\dots+N(z_j))}x$ and

$$r(z_j) = \sum_{\ell=0}^{N(z_j)-1} \tau(g^\ell g^{-(N(z_1)+\dots+N(z_j))}x) = \sum_{\ell=N(z_1)+\dots+N(z_{j-1})+1}^{N(z_1)+\dots+N(z_j)} \tau(g^{-\ell}x).$$

A similar expression holds for $r(z'_j)$. Hence

$$D_0(x, x') = \sum_{j=1}^{\infty} \{r(z_j) - r(z'_j)\}.$$

We are now in a position to complete the proof of the lemma, showing that if (UNI) fails, then $D \equiv 0$. To do this, we make use of [8, Proposition 7.4] (specifically the equivalence of their conditions 1 and 3). Namely, the failure of the (UNI) condition in Sect. 3.1 means that we can write $r = \xi \circ F - \xi + \zeta$ on Y where $\xi : Y \rightarrow \mathbb{R}$ is continuous (even C^1) and ζ is constant on partition elements $U \in \mathcal{P}$. Extending ξ and ζ trivially to $X = Y \times Z$, we obtain that

$r = \xi \circ f - \xi + \zeta$ on X where $\xi : X \rightarrow \mathbb{R}$ is continuous and constant on stable leaves, and ζ is constant on elements $V \in \tilde{\mathcal{P}}$. In particular,

$$\sum_{j=1}^n r(z_j) = \sum_{j=1}^n \{\xi(z_{j-1}) - \xi(z_j) + \zeta(z_j)\} = \xi(x) - \xi(z_n) + \sum_{j=1}^n \zeta(z_j).$$

For $x, x' \in V_0$, $V_0 \in \tilde{\mathcal{P}}$, with $x' \in W^u(x)$, it follows that

$$\sum_{j=1}^n \{r(z_j) - r(z'_j)\} = \xi(x) - \xi(x') - \xi(z_n) + \xi(z'_n).$$

Taking the limit as $n \rightarrow \infty$, we obtain that $D_0(x, x') = \xi(x) - \xi(x')$. Hence $D(x_1, x_2) = \xi(x_1) - \xi([x_1, x_2]) - \xi([x_2, x_1]) + \xi(x_2)$. Since ξ is constant on stable leaves, $D(x_1, x_2) = 0$ as required. \square

Proof of Theorem 1.2. By Proposition 4.3 and Lemma 4.4, f_t is a $C^{1+\alpha}$ uniformly hyperbolic flow satisfying (UNI). The result for $C^{1+\alpha}$ observables follows from Theorem 3.11. As in [18], the result follows from a standard interpolation argument (see also [6, Corollary 2.3]).

Acknowledgements

This research was supported in part by CMUP, which is financed by national funds through FCT - Fundação para a Ciência e a Tecnologia, I.P., under the project with reference UIDB/00144/2020. PV also acknowledges financial support from the project PTDC/MAT-PUR/4048/2021 and from the grant CEECIND/03721/2017 of the Stimulus of Scientific Employment, Individual Support 2017 Call, awarded by FCT. The authors are also grateful to Fundação Oriente for the financial support during the Pedro Nunes Meeting in Convento da Arrábida, where part of this work was done, and to the referee for helpful comments and suggestions.

Open Access. This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

References

- [1] Aaronson, J.: An Introduction to Infinite Ergodic Theory. Math. Surveys and Monographs, vol. 50. Amer. Math. Soc. (1997)
- [2] Alves, J.F., Luzzatto, S., Pinheiro, V.: Markov structures and decay of correlations for non-uniformly expanding dynamical systems. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **22**, 817–839 (2005)
- [3] Anosov, D.V.: Geodesic flows on closed Riemannian manifolds of negative curvature. *Trudy Mat. Inst. Steklov.* **90**, 209 (1967)
- [4] Araújo, V., Butterley, O., Varandas, P.: Open sets of Axiom A flows with exponentially mixing attractors. *Proc. Amer. Math. Soc.* **144**, 2971–2984 (2016)
- [5] Araújo, V., Butterley, O., Varandas, P.: Erratum to “Open sets of Axiom A flows with exponentially mixing attractors”. *Proc. Amer. Math. Soc.* **146**, 5013–5014 (2018)
- [6] Araújo, V., Melbourne, I.: Exponential decay of correlations for nonuniformly hyperbolic flows with a $C^{1+\alpha}$ stable foliation, including the classical Lorenz attractor. *Ann. Henri Poincaré* **17**, 2975–3004 (2016)
- [7] Araújo, V., Melbourne, I., Varandas, P.: Rapid mixing for the Lorenz attractor and statistical limit laws for their time-1 maps. *Comm. Math. Phys.* **340**, 901–938 (2015)
- [8] Avila, A., Gouëzel, S., Yoccoz, J.: Exponential mixing for the Teichmüller flow. *Publ. Math. Inst. Hautes Études Sci.* **104**, 143–211 (2006)
- [9] Baladi, V., Vallée, B.: Exponential decay of correlations for surface semi-flows without finite Markov partitions. *Proc. Amer. Math. Soc.* **133**, 865–874 (2005)
- [10] Bowen, R.: Markov partitions for Axiom A diffeomorphisms. *Amer. J. Math.* **92**, 725–747 (1970)
- [11] Bowen, R.: Symbolic dynamics for hyperbolic flows. *Amer. J. Math.* **95**, 429–460 (1973)
- [12] Bowen, R.: Equilibrium States and the Ergodic Theory of Anosov Diffeomorphisms. *Lecture Notes in Math*, vol. 470. Springer, Berlin (1975)
- [13] Bowen, R.: Markov partitions are not smooth. *Proc. Amer. Math. Soc.* **71**, 130–132 (1978)
- [14] Bowen, R., Ruelle, D.: The ergodic theory of Axiom A flows. *Invent. Math.* **29**, 181–202 (1975)
- [15] Butterley, O., War, K.: Open sets of exponentially mixing Anosov flows. *J. Eur. Math. Soc. (JEMS)* **22**, 2253–2285 (2020)
- [16] Chernov, N.I.: Markov approximations and decay of correlations for Anosov flows. *Ann. of Math.* **147**, 269–324 (1998)
- [17] Denker, M., Philipp, W.: Approximation by Brownian motion for Gibbs measures and flows under a function. *Ergodic Theory Dynam. Systems* **4**, 541–552 (1984)
- [18] Dolgopyat, D.: On the decay of correlations in Anosov flows. *Ann. of Math.* **147**, 357–390 (1998)
- [19] Field, M.J., Melbourne, I., Török, A.: Stability of mixing and rapid mixing for hyperbolic flows. *Ann. of Math.* **166**, 269–291 (2007)
- [20] Liverani, C.: On contact Anosov flows. *Ann. of Math.* **159**, 1275–1312 (2004)

- [21] Melbourne, I.: Superpolynomial and polynomial mixing for semiflows and flows. *Nonlinearity* **31**, R268–R316 (2018)
- [22] Parry, W., Pollicott, M.: *Zeta Functions and the Periodic Orbit Structure of Hyperbolic Dynamics*. Astérisque **187–188**, Société Mathématique de France, Montrouge (1990)
- [23] Plante, J.F.: Anosov flows. *Amer. J. Math.* **94**, 729–754 (1972)
- [24] Pollicott, M.: A complex Ruelle-Perron-Frobenius theorem and two counterexamples. *Ergodic Theory Dynam. Systems* **4**, 135–146 (1984)
- [25] Pugh, C., Shub, M.: Ergodic elements of ergodic actions. *Compositio Math.* **23**, 115–122 (1971)
- [26] Ratner, M.: Markov partitions for Anosov flows on n -dimensional manifolds. *Israel J. Math.* **15**, 92–114 (1973)
- [27] Ratner, M.: The central limit theorem for geodesic flows on n -dimensional manifolds of negative curvature. *Israel J. Math.* **16**, 181–197 (1973)
- [28] Ruelle, D.: Statistical mechanics of a one-dimensional lattice gas. *Comm. Math. Phys.* **9**, 267–278 (1968)
- [29] Ruelle, D.: *Thermodynamic Formalism*. Encyclopedia of Math. and its Applications, vol. 5. Addison Wesley, Massachusetts (1978)
- [30] Ruelle, D.: Flows which do not exponentially mix. *C. R. Acad. Sci. Paris* **296**, 191–194 (1983)
- [31] Sinai, Y.G.: Gibbs measures in ergodic theory. *Russ. Math. Surv.* **27**, 21–70 (1972)
- [32] Sinai, Ja.G.: Construction of Markov partitionings. *Funkcional. Anal. i Priložen.* **2**, 70–80 (1968)
- [33] Smale, S.: Differentiable dynamical systems. *Bull. Amer. Math. Soc.* **73**, 747–817 (1967)
- [34] Stoyanov, L.: Spectra of Ruelle transfer operators for axiom A flows. *Nonlinearity* **24**, 1089–1120 (2011)
- [35] Tsujii, M., Zhang, Z.: Smooth mixing Anosov flows in dimension three are exponentially mixing. *Ann. of Math. (2)* **197**, 65–158 (2023)
- [36] Young, L.-S.: Statistical properties of dynamical systems with some hyperbolicity. *Ann. of Math.* **147**, 585–650 (1998)

Ian Melbourne
Mathematics Institute
University of Warwick
Coventry CV4 7AL
UK
e-mail: I.Melbourne@warwick.ac.uk

Paulo Varandas
CMUP and Departamento de Matemática
Universidade Federal da Bahia
Salvador 40170-110
Brazil

Communicated by Claude-Alain Pillet.

Received: May 17, 2023.

Accepted: April 5, 2024.