On General Position Sets in Cartesian Products

Sandi Klavžar, Balázs Patkós , Gregor Rus, and Ismael G. Yero

Abstract. The general position number $gp(G)$ of a connected graph G is the cardinality of a largest set S of vertices such that no three distinct vertices from S lie on a common geodesic; such sets are referred to as gp-sets of G . The general position number of cylinders $P_r \square C_s$ is deduced. It is proved that $gp(C_r \square C_s) \in \{6, 7\}$ whenever $r \geq s \geq 3$, $s \neq 4$, and $r \geq 6$. A probabilistic lower bound on the general position number of Cartesian graph powers is achieved. Along the way a formula for the number of gp-sets in $P_r \square P_s$, where $r, s \geq 2$, is also determined.

Mathematics Subject Classifications. 05C12, 05C76, 05C30, 05D40.

Keywords. General position problem, Cartesian product of graphs, paths and cycles, probabilistic constructions, exact enumeration.

1. Introduction

Points in general position in the plane are basic objects in classical geometry as well as in computational geometry. The concept naturally extends to arbitrary metric spaces and asking for the maximum number of points in general position appears as a natural problem. The problem to find the maximum number of points that can be placed in the $n \times n$ grid so that no three points lie on a line is known as the *no-three-in-line problem* and was posed back in 1917 by Dudeney [5]. After a century the problem remains open, cf. [12, 15, 20]. Closely related problems are investigated in discrete geometry, see [3, 19].

Recently, the general position problem has been studied for metric spaces generated by graphs. The problem was independently introduced in [13, 25], the present terminology and formalism are from [13]. If $G = (V(G), E(G))$ is a graph, then $S \subseteq V(G)$ is a *general position set* if $d_G(u, v) \neq d_G(u, w) +$

$d_G(w, v)$ holds for every $\{u, v, w\} \in \binom{S}{3}$, where $d_G(x, y)$ denotes the shortest-path distance in G , and $\binom{S}{3}$ the set of all 3-subsets of S . Equivalently, no three vertices lie on a common geodesic. We also say that the vertices from S lie in *general position*. The *general position problem* is to find a largest general position set of G , the order of such a set is the *general position number* $\text{gp}(G)$ of G . A general position set of G of order $\text{gp}(G)$ will be called a *gp-set*.

Following the seminal papers, the general position problem has been investigated in a sequence of papers [1, 6, 10, 14, 16, 18, 22, 26]. As it happens, in the special case of hypercubes, the general position problem was studied back in 1995 by Körner [11] related to some coding theory problems. In this paper, asymptotic lower and upper bounds were proved on the gp-number of hypercubes, and several closely related problems (cf. Sect. 5) were considered. The lower bound from [11] was improved in [17].

The results from [14] on the general position problem in interconnection networks with the emphasis on grid graphs were a starting motivation for the present study. One of the main results of [14] asserts that if P_∞ denotes the two-way infinite path, then $\text{gp}(P_\infty \square P_\infty) = 4$, and consequently $\text{gp}(P_r \square P_s) = 4$ for $r, s \geq 3$. The non-trivial part of this result (that $\text{gp}(P_r \square P_s) \leq 4$ holds) was proved using the so-called Monotone Geodesic Lemma which was in turn derived from the celebrated Erdős-Szekeres theorem, cf. [2, Theorem 1.1]. The result $\text{gp}(P_\infty \square P_\infty) = 4$ was recently generalized in two directions. In [9], it is proved that $\text{gp}(P_\infty^{\square, k}) = 2^{2^{k-1}}$, while in [24], it is demonstrated that if T_1 and T_2 are arbitrary trees, then $\text{gp}(T_1 \square T_2) = \text{gp}(T_1) + \text{gp}(T_2)$. The general position number of several other Cartesian product graphs is studied in [23].

In the rest of this section we prepare material needed later on. In the next section we have a closer look at the structure of gp-sets in grid graphs. We need to do it for the proof of the subsequent theorem on cylinder graphs (Theorem 3.2). As a side result we determine the number of gp-sets in $P_r \square P_s$ for every $r, s \geq 2$, a result that could be of independent interest. In Sect. 3 we then determine $\text{gp}(P_r \square C_s)$ for every $r \geq 2$ and $s \geq 3$. In the subsequent section we prove that if $3 \leq s \leq r$, $4 \neq s$, and $r \geq 6$, then $\text{gp}(C_r \square C_s) \in \{6, 7\}$. We also discuss exact values and in particular prove that $\text{gp}(C_3 \square C_s) = 6$ holds for $s \geq 6$. Motivated by the results of [11], we consider in Sect. 5 how to apply the probabilistic method to obtain asymptotic lower bounds on the gp-number of Cartesian powers of graphs.

1.1. Preliminaries

For a positive integer k we will use the notation $[k] = \{1, \dots, k\}$ and $[k]_0 = \{0, \dots, k-1\}$. The *Cartesian product* $G \square H$ of graphs G and H has the vertex set $V(G \square H) = V(G) \times V(H)$, vertices (g, h) and (g', h') are adjacent if either $gg' \in E(G)$ and $h = h'$, or $g = g'$ and $hh' \in E(H)$. Given a vertex $h \in V(H)$, the subgraph of $G \square H$ induced by the set $\{(g, h) : g \in V(G)\}$, is a *G-layer* and is denoted by G^h . *H-layers* gH are defined analogously. Each *G-layer* and

H -layer is isomorphic to G and H , respectively. If $X \subseteq V(G \square H)$, then the *projection* $p_G(X)$ of X to G is the set $\{g \in V(G) : (g, h) \in X \text{ for some } h \in V(H)\}$. The projection $p_H(X)$ of X to H is defined analogously. The k -tuple Cartesian product of a graph G by itself, alias Cartesian power of G , is denoted by $G^{\square, k}$. This is well-defined since the Cartesian product operation is associative. For more on the Cartesian product see [7]. As stated above, the following result was the primary motivation for the present paper.

Theorem 1.1 [14]. *If $r \geq 3$ and $s \geq 3$, then $\text{gp}(P_r \square P_s) = 4$.*

A subgraph H of a graph G is *isometric* if $d_H(u, v) = d_G(u, v)$ holds for all $u, v \in V(H)$. A set of subgraphs $\{H_1, \dots, H_k\}$ of a graph G is an *isometric cover* of G if each H_i , $i \in [k]$, is isometric in G and $\bigcup_{i=1}^k V(H_i) = V(G)$.

Theorem 1.2 [13, Theorem 3.1]. *If $\{H_1, \dots, H_k\}$ is an isometric cover of G , then*

$$\text{gp}(G) \leq \sum_{i=1}^k \text{gp}(H_i).$$

If G is a connected graph, $S \subseteq V(G)$, and $\mathcal{P} = \{S_1, \dots, S_p\}$ is a partition of S , then $\mathcal{P}$ is *distance-constant* (alias “distance-regular” [8, p. 331]) if for any $i, j \in [p]$, $i \neq j$, the distance $d_G(u, v)$, where $u \in S_i$ and $v \in S_j$, is independent of the selection of u and v . This distance is then the distance $d_G(S_i, S_j)$ between the parts S_i and S_j . A distance-constant partition $\mathcal{P}$ is *intransitive* if $d_G(S_i, S_k) \neq d_G(S_i, S_j) + d_G(S_j, S_k)$ holds for distinct indices $i, j, k \in [p]$.

Theorem 1.3 [1, Theorem 3.1]. *Let G be a connected graph. Then $S \subseteq V(G)$ is a general position set if and only if the components of the subgraph induced by S are complete subgraphs, the vertices of which form an intransitive, distance-constant partition of S .*

Suppose that G is a connected bipartite graph and a general position set S contains two adjacent vertices x and y . Then Theorem 1.3 implies that $|S| = 2$, because no other vertex of G can be at the same distance to x and y . We state this observation for later use.

Corollary 1.4. *If G is a bipartite graph with $\text{gp}(G) \geq 3$, then every gp-set of G is an independent set.*

2. Enumeration of gp-Sets in Grids

In this section we take a closer look at the structure of gp-sets in grids. We have two reasons to do it, the first is that this insight will be used in the proof of Theorem 3.2, the second is that we are able to enumerate the gp-sets in grids as follows.

Theorem 2.1 *If $2 \leq r \leq s$, then*

$$\# \text{gp}(P_r \square P_s) = \begin{cases} 6; & r = s = 2, \\ \frac{s(s-1)(s-2)}{3}; & r = 2, s \geq 3, \\ \frac{rs(r-1)(r-2)(s-1)(s-2)(r(s-3)-s+7)}{144}; & r, s \geq 3. \end{cases}$$

Proof. Set $V(P_n) = [n]$. If $r = s = 2$, then the assertion is clear since $P_2 \square P_2 = C_4$. Let next $r = 2$ and $s \geq 3$. It is straightforward to see that $\text{gp}(P_2 \square P_s) = 3$. Moreover, if X is a gp-set of $P_2 \square P_s$, then X has one vertex in one of the two P_s -layers and two vertices in the other P_s -layer, say $X = \{(1, i), (2, j), (2, k)\}$, where $j < k$. Since X is a gp-set we infer that $j < i < k$. Hence, for a given vertex $(1, i)$, there are $i - 1$ possibilities to select the vertex $(2, j)$, and $s - i$ possibilities for the vertex $(2, k)$. The same holds if X has two vertices in 1P_s and one vertex in 2P_s . From this it follows that

$$\# \text{gp}(P_2 \square P_s) = 2 \cdot \sum_{i=1}^s (i-1)(s-i) = \frac{s(s-1)(s-2)}{3}.$$

Suppose in the rest that $r, s \geq 3$, so that $\text{gp}(P_r \square P_s) = 4$ by Theorem 1.1. Hence by Corollary 1.4, every gp-set is an independent set (of cardinality 4). Let X be an arbitrary such set and assume first that $|p_{P_s}(X)| = 2$. Then, clearly, X has two vertices in one P_r -layer and two vertices in another P_r -layer. Let $(i, j) \in X$ be a vertex that has the smallest first coordinate among the vertices of X . Then (i, j) and the two vertices of X from the P_r -layer not containing (i, j) lie on a common geodesic. Analogously, X cannot be a general position set if $|p_{P_r}(X)| = 2$. Since also $|p_{P_r}(X)| = 1$ or $|p_{P_s}(X)| = 1$ are not possible, we only need to distinguish the following two cases.

Case 1: $|p_{P_r}(X)| = 4$ and $|p_{P_s}(X)| = 4$.

Let $p_{P_r}(X) = \{a, b, c, d\}$, where $a < b < c < d$, and let $p_{P_s}(X) = \{a', b', c', d'\}$, where $a' < b' < c' < d'$. Then in the set $p_{P_r}(X) \times p_{P_s}(X)$ there are $4!$ different 4-sets of vertices that project onto both $p_{P_r}(X)$ and $p_{P_s}(X)$. They can be described with permutations π of $p_{P_s}(X)$. That is, if $\pi : p_{P_s}(X) \rightarrow p_{P_r}(X)$ is a bijection, then the corresponding gp-set of vertices of $P_r \square P_s$ is $S_\pi = \{(a, \pi(a')), (b, \pi(b')), (c, \pi(c')), (d, \pi(d'))\}$. Now, by the metric structure of $P_r \square P_s$ (cf. [14]), S_π is a general position set if and only if the sequence $(\pi(a'), \pi(b'), \pi(c'), \pi(d'))$ contains no monotone subsequence of length 3. By a direct inspection we find that if $\pi(a') = a'$ or if $\pi(a') = d'$, then we get no general position sets. If $\pi(a') = b'$, then exactly the sequences (b', a', d', c') and (b', d', a', c') yield general position sets. Symmetrically, if $\pi(a') = c'$, then exactly the sequences (c', a', d', b') and (c', d', a', b') yield general position sets. Hence, if $|p_{P_r}(X)| = 4$ and $|p_{P_s}(X)| = 4$, then there are exactly $4 \binom{r}{4} \binom{s}{4}$ gp-sets.

Case 2: $|p_{P_r}(X)| = 3$ (and $|p_{P_s}(X)| = 3$ or $|p_{P_s}(X)| = 4$).

Let $p_{P_r}(X) = \{a, b, c\}$, where two vertices from X project to a , say (a, a') , $(a, b') \in X$, where $a' < b'$. Let (x, x') be a vertex of $P_r \square P_s$, where $x' \leq a'$ and $(x, x') \neq (a, a')$. Then $d((x, x'), (a, b')) = d((x, x'), (a, a')) + d((a, a'), (a, b'))$, which means that $(x, x') \notin X$. Similarly, if (x, x') is a vertex of $P_r \square P_s$ with $x' \geq b'$ and $(x, x') \neq (a, b')$, then also $(x, x') \notin X$. We have thus shown that

$$X \cap ([r] \times \{1, \dots, a'\}) = \{(a, a')\} \text{ and } X \cap ([r] \times \{b', \dots, s\}) = \{(a, b')\}.$$

Let $X = \{(a, a'), (a, b'), (x, x'), (y, y')\}$, where $x \neq y$. By a similar argument as above we see that, without loss of generality,

$$\begin{aligned} (x, x') &\in \{1, \dots, a-1\} \times \{a'+1, \dots, b'-1\} \text{ and} \\ (y, y') &\in \{a+1, \dots, r\} \times \{a'+1, \dots, b'-1\}. \end{aligned}$$

Since the vertices (x, x') and (y, y') are arbitrary vertices from $\{1, \dots, a-1\} \times \{a'+1, \dots, b'-1\}$ and $\{a+1, \dots, r\} \times \{a'+1, \dots, b'-1\}$, respectively, for fixed a, a', b' we obtain precisely

$$[(b' - a' - 1)(a - 1)] \cdot [(b' - a' - 1)(r - a)] = (b' - a' - 1)^2(a - 1)(r - a)$$

gp-sets. To obtain the number of all gp-sets in this case, we need to sum up over all coordinates a of the factor P_r , onto which two vertices from X project. Consequently, the number of gp-sets in Case 2 is

$$\sum_{a=1}^r \sum_{a'=1}^s \sum_{b'=a'+1}^s [(b' - a' - 1)^2(a - 1)(r - a)] = \frac{rs(r^2 - 3r + 2)(s^3 - 4s^2 + 5s - 2)}{72}.$$

By the above two cases, if $r, s \geq 3$, then

$$\begin{aligned} \#gp(P_r \square P_s) &= 4 \binom{r}{4} \binom{s}{4} + \frac{rs(r^2 - 3r + 2)(s^3 - 4s^2 + 5s - 2)}{72} \\ &= \frac{rs(r-1)(r-2)(s-1)(s-2)(r(s-3) - s + 7)}{144}, \end{aligned}$$

which is the claimed expression. $\square$

If $r = 3$ and $s \geq 3$, then Theorem 2.1 yields

$$\#gp(P_3 \square P_s) = \frac{s(s-2)(s-1)^2}{12},$$

which, after substituting s with $s+1$ gives the sequence A002415 from OEIS [21]. In addition, the case $r = 2$ and $s \geq 3$ yields the sequence A007290.

3. Cylinders

In this section we determine the general position number of cylinders. For this task, the following function will be useful. If G is a connected graph and $X \subseteq V(G)$ is a general position set, then

$F(X) = \{u \in V(G) - X : X \cup \{u\} \text{ is not a general position set}\}.$

If $X = \{x, y\}$, we will simplify the notation $F(\{x, y\})$ to $F(x, y)$.

Set $V(P_r) = [r]_0$ and $V(C_s) = [s]_0$. From now on, operations with the integers in $V(C_s)$ are done modulo s .

Lemma 3.1 *Let $r \geq 2$, $s \geq 3$, and let S be a general position set of the cylinder graph $P_r \square C_s$. Then the following assertions hold.*

- (i) *If $|S| \geq 5$, then S is an independent set.*
- (ii) *If $|S| \geq 4$, then $|S \cap V(iC_s)| \leq 2$ for every $i \in [r]_0$.*
- (iii) *If $|S \cap V(iC_s)| = 2$ for some $i \in [r]_0$, then $|S| \leq 4$.*
- (iv) *If $r \geq 6$, $|S| = 5$, and $|S \cap V(iC_s)| \leq 1$ for every $i \in [r]_0$, then $\text{gp}(P_5 \square C_s) \geq 5$.*

Proof. (i) Suppose $|S| \geq 5$ and S is not independent. If $(i, k), (i+1, k) \in S$, then we observe that $F((i, k), (i+1, k)) = V(P_r \square C_s) - \{(i, k), (i+1, k)\}$, which means that $S = \{(i, k), (i+1, k)\}$, a contradiction. On the other hand, if $(i, k), (i, k+1) \in S$, then either $F((i, k), (i, k+1)) = V(P_r \square C_s) - \{(i, k), (i, k+1)\}$ (when s is even), or $F((i, k), (i, k+1)) = V(P_r \square C_s) - (\{(i, k), (i, k+1)\} \cup (V(P_r) \times \{j\}))$ (when s is odd), where j is the vertex which is on C_s opposite to k and $k+1$. The first possibility directly leads to a contradiction. For the second one, since every P_r -layer, being an isometric subgraph, contributes at most two vertices to a general position set of $P_r \square C_s$, it follows that $|S| \leq 4$, which is again a contradiction. Consequently S must be an independent set.

(ii) The result follows directly from the following fact. If $\{(i, k_1), (i, k_2), (i, k_3)\} \subseteq S$, then (since k_1, k_2 , and k_3 are distinct) $F(\{(i, k_1), (i, k_2), (i, k_3)\}) = V(P_r \square C_s) - \{(i, k_1), (i, k_2), (i, k_3)\}$, which means that $|S| = 3$.

(iii) Let $i \in [r]_0$ be such that $|S \cap V(iC_s)| = 2$. We may assume without loss of generality that $S \cap V(iC_s) = \{(i, 0), (i, j)\}$, where $j \leq \lfloor s/2 \rfloor$. Then

$$\begin{aligned} F((i, 0), (i, j)) &= [r]_0 \times (\{j - \lfloor s/2 \rfloor, \dots, 0\} \cup \{j, \dots, \lfloor s/2 \rfloor\}) \\ &\cup (\{i\} \times I) - \{(i, 0), (i, j)\}, \end{aligned}$$

where $I = [s]_0$ if s is even and $j = s/2$; or $I = \{0, \dots, j\}$ otherwise.

If $j = 1$, then $F((i, 0), (i, j))$ equals $V(P_r \square C_s) - \{(i, 0), (i, j)\}$ when s is even, or $F((i, 0), (i, j))$ equals $V(P_r \square C_s) - (([r]_0 \times \{\lfloor s/2 \rfloor\}) \cup \{(i, 0), (i, j)\})$ when s is odd. In the first situation we clearly have $|S| \leq 2$. In the latter one, the set of vertices $[r]_0 \times \{\lfloor s/2 \rfloor\}$ could contain at most two vertices of S , since it induces an isometric path in $P_r \square C_s$, and so $|S| \leq 4$. We may assume in the rest that $j \geq 2$. Note that $V(P_r \square C_s) - (F((i, 0), (i, j)) \cup \{(i, 0), (i, j)\}) = Y_1 \cup Y_2 \cup Y_3 \cup Y_4$, where

$$\begin{aligned} Y_1 &= [i]_0 \times \{1, \dots, j-1\} \\ Y_2 &= [i]_0 \times \{\lfloor s/2 \rfloor + 1, \dots, j - \lfloor s/2 \rfloor - 1\} \\ Y_3 &= ([r]_0 - [i+1]_0) \times \{1, \dots, j-1\} \\ Y_4 &= ([r]_0 - [i+1]_0) \times \{\lfloor s/2 \rfloor + 1, \dots, j - \lfloor s/2 \rfloor - 1\}. \end{aligned}$$

Consider two vertices $(i', j'), (i'', j'') \in S - \{(i, 0), (i, j)\}$. (If there are no two such vertices, then clearly $|S| \leq 3$.) If either $(i', j'), (i'', j'') \in Y_1$; or $(i', j'), (i'', j'') \in Y_2$; or $(i', j'), (i'', j'') \in Y_3$; or $(i', j'), (i'', j'') \in Y_4$, then it happens that $(i, 0) \in F((i', j'), (i'', j''))$ or $(i, j) \in F((i', j'), (i'', j''))$, which is not possible. As a consequence, each of the sets Y_1, Y_2, Y_3 , and Y_4 must contain at most one vertex of S .

Assume now there is a vertex $(i', j') \in Y_1 \cap S$. Then the set $F((i', j'), (i, 0)) \cup F((i', j'), (i, j))$ contains the whole set Y_4 . By symmetry, if there is a vertex $(i', j') \in Y_4 \cap S$, then $Y_1 \cap S = \emptyset$. Hence, $|(Y_1 \cup Y_4) \cap S| \leq 1$. Similarly we obtain that $|(Y_2 \cup Y_3) \cap S| \leq 1$. Therefore, the sets Y_1, Y_2, Y_3 , and Y_4 can contain at most two vertices of S , which gives $|S| \leq 4$.

(iv) Let $S = \{(i_k, j_k) : k \in [5]_0\}$. Since $|S \cap V(iC_s)| \leq 1$, the coordinates i_k are pairwise different, hence we may assume without loss of generality that $i_0 < i_1 < i_2 < i_3 < i_4$. Set $S' = \{(k, j_k) : k \in [5]_0\}$. We claim that S' is a general position set of $G_5 = P_5 \square C_s$. Assume on the contrary that

$$d_{G_5}((p, j_p), (r, j_r)) = d_{G_5}((p, j_p), (q, j_q)) + d_{G_5}((q, j_q), (r, j_r))$$

for some $p, q, r \in [5]_0$, $p < q < r$. Since the distance function in Cartesian products is additive, we get that

$$d_{P_5}(p, r) + d_{C_s}(j_p, j_r) = d_{P_5}(p, q) + d_{C_s}(j_p, j_q) + d_{P_5}(q, r) + d_{C_s}(j_q, j_r).$$

Since $d_{P_5}(p, r) = d_{P_5}(p, q) + d_{P_5}(q, r)$, we thus have

$$d_{C_s}(j_p, j_r) = d_{C_s}(j_p, j_q) + d_{C_s}(j_q, j_r).$$

From this we get that in $G_r = P_r \square C_s$,

$$\begin{aligned} d_{G_r}((i_p, j_p), (i_r, j_r)) &= d_{P_r}(i_p, i_r) + d_{C_s}(j_p, j_r) \\ &= [d_{P_r}(i_p, i_q) + d_{P_r}(i_q, i_r)] + [d_{C_s}(j_p, j_q) + d_{C_s}(j_q, j_r)] \\ &= [d_{P_r}(i_p, i_q) + d_{C_s}(j_p, j_q)] + [d_{P_r}(i_q, i_r) + d_{C_s}(j_q, j_r)] \\ &= d_{G_r}((i_p, j_p), (i_q, j_q)) + d_{G_r}((i_q, j_q), (i_r, j_r)). \end{aligned}$$

This contradiction proves that S' is a general position set of $P_5 \square C_s$. We conclude that $\text{gp}(P_5 \square C_s) \geq 5$. $\square$

Note that Lemma 3.1(iv) allows us to map a general position set of cardinality 5 in long cylinders to a general position set of the same cardinality in cylinders over P_5 .

Theorem 3.2 *If $r \geq 2$ and $s \geq 3$, then*

$$\text{gp}(P_r \square C_s) = \begin{cases} 3; & r = 2, s = 3, \\ 5; & r \geq 5, \text{ and } s = 7 \text{ or } s \geq 9, \\ 4; & \text{otherwise.} \end{cases}$$

Proof. First, it is easy to verify that $\text{gp}(P_2 \square C_3) = 3$.

Assume next that $r \leq 4$ and suppose that there exists a general position set S with $|S| \geq 5$. If $r = 2$, this is not possible by Lemma 3.1(ii). Let next

$r \in \{3, 4\}$. Then there exists a C_s -layer iC_s with $|V({}^iC_s) \cap S| \geq 2$. The case $|V({}^iC_s) \cap S| > 2$ is not possible by Lemma 3.1(ii), while the case $|V({}^iC_s) \cap S| = 2$ is excluded by Lemma 3.1(iii). Hence $\text{gp}(P_r \square C_s) \leq 4$ for $r \in \{2, 3, 4\}$. It is straightforward to see that the set $\{(0, 0), (1, 1), (0, \lfloor s/2 \rfloor), (1, \lfloor s/2 \rfloor + 1)\}$ is a general position set of $P_r \square C_s$ for $r \geq 2$ and $s \geq 4$. Moreover, if $s = 3$, then the set $\{(0, 1), (1, 0), (1, 2), (2, 1)\}$ is a general position set of $P_r \square C_3$ for $r \geq 3$. Hence $\text{gp}(P_r \square C_s) \geq 4$ for $r \in \{2, 3, 4\}$ and so $\text{gp}(P_r \square C_s) = 4$ for $r \in \{2, 3, 4\}$.

The general position set $\{(0, 0), (1, 2), (2, 4), (3, 6), (4, 1)\}$ of $P_5 \square C_7$ demonstrates that $\text{gp}(P_5 \square C_7) \geq 5$.

Suppose next that for some $r \geq 6$ the cylinder $P_r \square C_8$ contains a general position set S with $|S| = 5$. From Lemma 3.1(iii) it follows that $|S \cap V({}^iC_8)| \leq 1$ for every $i \in [r]_0$. Hence the assumptions of Lemma 3.1(iv) are fulfilled, and we can map the general position set S of cardinality 5 in the cylinder $P_r \square C_8$, $r \geq 6$, to a general position set of the same cardinality (5) in the cylinder over $P_5 \square C_8$, which indeed implies that $\text{gp}(P_5 \square C_8) \geq 5$. However, we have checked by computer that $\text{gp}(P_5 \square C_8) = 4$. Thus, we have obtained a contradiction. Therefore, $\text{gp}(P_r \square C_8) \leq 4$ for $r \geq 5$ (the case $r = 5$ was computationally made). Since clearly $\text{gp}(P_r \square C_8) \geq 4$, we conclude that $\text{gp}(P_r \square C_8) = 4$ for $r \geq 5$.

Suppose now that $r = 5$, $s \geq 9$, and consider the set

$$S = \{u_0 = (0, 1), u_1 = (1, 4), u_2 = (2, \lfloor s/2 \rfloor + 2), u_3 = (3, 0), u_4 = (4, 3)\}.$$

We claim that S is a general position set. Note first that the vertices u_0, u_1, u_3, u_4 lie in a general position. Further,

$$\begin{aligned} d(u_2, u_0) &= (s - \lfloor s/2 \rfloor - 1) + 2 = s - \lfloor s/2 \rfloor + 1, \\ d(u_2, u_1) &= \lfloor s/2 \rfloor - 1, \\ d(u_2, u_3) &= s - \lfloor s/2 \rfloor - 1, \\ d(u_2, u_4) &= \lfloor s/2 \rfloor + 1. \end{aligned}$$

Then $d(u_0, u_2) = s - \lfloor s/2 \rfloor + 1 < \lfloor s/2 \rfloor + 3 = d(u_0, u_1) + d(u_1, u_2)$. Similarly we see that u_2 is not on a geodesic containing three vertices of S . Hence, S is a general position set and thus $\text{gp}(P_5 \square C_s) \geq 5$ for $s \geq 9$.

Note finally that the general position set for $P_5 \square C_7$ and the general position set for $P_5 \square C_s$, $s \geq 9$, are also general position sets for $P_r \square C_7$, $r \geq 6$, and for $P_r \square C_s$, $r \geq 6$, respectively. We conclude that $\text{gp}(P_r \square C_s) \geq 5$ for $r \geq 5$ and $s \geq 7$, $s \neq 8$.

It remains to prove that the above constructed general position sets of cardinality 5 are gp-sets. Hence let S be a gp-set of $P_r \square C_s$, where $|S| \geq 5$. Note that S is an independent set, by Lemma 3.1(i). Then make a partition of $V(P_r \square C_s)$ into two sets A_1 and A_2 inducing two grids that are isometric subgraphs of $P_r \square C_s$. Without loss of generality, we may assume $A_1 = V(P_r) \times [\lfloor s/2 \rfloor]_0$ and $A_2 = V(P_r) \times ([s]_0 - [\lfloor s/2 \rfloor]_0)$. Now, let $S_1 = S \cap A_1$ and $S_2 =$

FIGURE 1. Two possible configurations of the set S_1 (edges of the grid have not been drawn)

$S \cap A_2$. Since $\text{gp}(P_r \square C_s) \geq 5$, it follows $|S_1| \geq 3$ or $|S_2| \geq 3$. Moreover, since A_1 and A_2 induce isometric grid graphs, Theorem 1.1 implies that $|S_1| \leq 4$ and $|S_2| \leq 4$. To simplify the notation, we write $s_d = \lfloor s/2 \rfloor$. Note that here we do not need to change floor to ceil. We consider the following cases.

Case 1: $|S_1| = 4$.

Let $S_1 = \{(a', a), (b', b), (c', c), (d', d)\}$. Note that S_1 is then a gp-set of the grid induced by A_1 . Since the structure of every gp-set of a grid graph is known from the proof of Theorem 2.1, we can assume without loss of generality the following facts: $a < b < d$, $a < c < d$, and either $(c' < a' < b'$ and $c' < d' < b')$ or $(b' < a' < c'$ and $b' < d' < c')$. Note that there is neither relationship between a, b nor between a', b' , namely, it can happen $a \leq b$ or $b \leq a$ and $a' \leq b'$ or $b' \leq a'$. Examples of such sets are shown in Fig. 1. For presentation purposes, in this and the subsequent figures an orientation is selected such that C_s -layers are drawn horizontally and P_r -layers vertically.

We now consider the set $F(S_1)$ in $V(P_r \square C_s)$. Figure 2 shows an example of the forbidden area generated by only two vertices of S_1 ((c', c) and (d', d) in this case). Since it is not necessary for our purposes, we do not look at the whole set S_1 , only a significant part.

We detail now the case $a < b \leq c < d$ and $c' < a' \leq d' < b'$, see Fig. 3. Observe that:

$$\begin{aligned}
 \{a', \dots, r-1\} \times \{a, a-1, \dots, c-s_d\} &\subset F((c', c), (a', a)), \\
 \{0, \dots, a'\} \times \{a, a-1, \dots, b-s_d\} &\subset F((b', b), (a', a)), \\
 \{d', \dots, r-1\} \times \{d, d+1, \dots, c+s_d\} &\subset F((c', c), (d', d)), \\
 \{0, \dots, d'\} \times \{d, d+1, \dots, b+s_d\} &\subset F((b', b), (d', d)).
 \end{aligned}$$

FIGURE 2. The forbidden area $F((c', c), (d', d))$ appears surrounded by the dashed rectangles. Vertices b_1 and b_2 of C_s are diametral with b while c_1 and c_2 are diametral with c . Here $b_1 = b + s_d$, $b_2 = b - s_d$, $c_1 = c + s_d$, and $c_2 = c - s_d$. Similar convention will be used in the subsequent figures. Notice that if the cycle would have an even order, then $b_1 = b_2$ and $c_1 = c_2$

Let us define a subset A of A_2 by $A = A_2 - F(S_1)$. Observe that A is empty if and only if one of the following situations occur: $a' = d'$, or $a' = d' - 1$, or $b = c$, or $(b = c - 1$ and s is even). More precisely, $A = \{a' + 1, \dots, d' - 1\} \times \{b_2, \dots, c_1\}$. Figure 3 shows an example of this where the set A is not empty.

As a consequence, it must happen that $S_2 \subseteq A$, since otherwise we get a contradiction with S being a gp-set. If $|S_2| \geq 2$, then let $(x', x), (y', y) \in S_2$. It is then not difficult to observe that $(c', c), (x', x), (y', y)$ or $(d', d), (x', x), (y', y)$ lie in a geodesic of $P_r \square C_s$, as well as $(a', a), (x', x), (y', y)$ or $(b', b), (x', x), (y', y)$, which is not possible. Thus $|S_2| \leq 1$.

By using a similar reasoning, we deduce the same conclusion for any other relationship of a, b, c, d and a', b', c', d' (from those ones that allow to obtain a gp-set of the grid induced by A_1 , according to the proof of Theorem 2.1). As a consequence of the whole deduction of this case, we obtain that $\text{gp}(P_r \square C_s) = |S| = |S_1| + |S_2| \leq 5$.

Case 2: $|S_2| = 4$.

The situation is similar to Case 1, although if s is odd, then the set A_2 is smaller than A_1 by a difference of one P_r -layer. Notwithstanding, this fact does not influence the arguments considered in Case 1.

FIGURE 3. Part of the forbidden area of the bolded set of vertices appears in dashed rectangles. The two gray vertices of the thick rectangle (denoted by A) do not belong to the forbidden area of the bolded vertices

FIGURE 4. An example of a configuration for the set S_1

Case 3: $|S_1| = 3$ or $|S_2| = 3$.

Assume first that $|S_1| = 3$, and let $S_1 = \{(a', a), (b', b), (c', c)\}$. Clearly, the three elements of S_1 cannot lie simultaneously in the same iC_s -layer, or in the same $P_r{}^j$ -layer. Moreover, it cannot happen that $a' \leq b' \leq c'$ and $a \leq b \leq c$ at the same time, or any other similar double monotone sequence. This means that, for instance, if $a' \leq b' \leq c'$, then either $(b < a \text{ and } b < c)$ or $(b > a \text{ and } b > c)$.

We may assume now that $a' \leq b' \leq c'$, $b < a$ and $b < c$. Figure 4 shows an example of this.

FIGURE 5. A significant part of the set $F(S_1)$ appears surrounded by dashed rectangles. For $x \in \{a, b, c\}$, the vertices x_1 and x_2 from C_s ($x_1 = x + s_d$ and $x_2 = x - s_d$) are diametral vertices with x . Note that if C_s is an even cycle, then $x_1 = x_2$

We now consider the set $F(S_1)$ in $V(P_r \square C_s)$, and observe the following. Recalling that $s_d = \lfloor s/2 \rfloor$ we have:

$$\begin{aligned} \{c', \dots, r-1\} \times \{c, c+1, \dots, a+s_d\} &\subset F((c', c), (a', a)), \\ \{0, \dots, a'\} \times \{a, a-1, \dots, c-s_d\} &\subset F((c', c), (a', a)), \\ \{b', \dots, r-1\} \times \{b, b-1, \dots, a-s_d\} &\subset F((b', b), (a', a)), \\ \{0, \dots, a'\} \times \{a, a+1, \dots, b+s_d\} &\subset F((b', b), (a', a)), \\ \{c', \dots, r-1\} \times \{c, c+1, \dots, b+s_d\} &\subset F((c', c), (b', b)), \\ \{0, \dots, b'\} \times \{b, b-1, \dots, c-s_d\} &\subset F((c', c), (b', b)). \end{aligned}$$

See Fig. 5 for an example of the situations above.

Observe now that there are four sets, say B_1 , B_2 , B_3 , and B_4 , such that $B_1 \cup B_2 \cup B_3 \cup B_4 = A_2 - F(S_1)$, and satisfying the following. If B_1 , B_2 , B_3 , and B_4 are not empty, then

$$\begin{aligned} B_1 &= \{a'+1, \dots, c'-1\} \times \{s_d+1, \dots, b+s_d\}, \\ B_2 &= \{a'+1, \dots, c'-1\} \times \{b+s_d+1, \dots, a+s_d\}, \\ B_3 &= \{0, \dots, a'\} \times \{b+s_d+1, \dots, a+s_d\}, \\ B_4 &= \{0, \dots, a'\} \times \{a+s_d+1, \dots, c+s_d\}. \end{aligned}$$

Note that some of these sets could be empty, or could have non-empty intersection, depending on the parity of s and on the structure of the set S_1 .

If $|S_2 \cap B_i| \geq 2$ for some $i \in [4]$, then we shall find an isometric subgraph of $P_r \square C_s$ isomorphic to a grid graph such that it contains four vertices of the set

S . Hence, we can change the partition given by A_1 and A_2 from the beginning, to a new one, and proceed as in Case 1, to prove that $\text{gp}(P_r \square C_s) \leq 5$. That is, if $|S_2 \cap B_i| \geq 2$ for some $i \in [3]$, then we can use the partition $A'_1 = [r]_0 \times \{a, a+1, \dots, a+s_d\}$ and $A'_2 = V(P_r \square C_s) - A'_1$, and if $|S_2 \cap B_4| \geq 2$, then we can use the partition $A'_1 = [r]_0 \times \{a+s_d+1, a+s_d+2, \dots, a\}$ (note that $a+s_d+1 = a-s_d$) and $A'_2 = V(P_r \square C_s) - A'_1$. In concordance, we may assume that $|S_2 \cap B_i| \leq 1$ for every $i \in [4]$.

We consider now the three sets B_1 , B_2 and B_3 . If at least two of them contain one element from S_2 , then, as above, we can find a different partition of $V(P_r \square C_s)$ and proceed like in Case 1. Thus, $|(B_1 \cup B_2 \cup B_3) \cap S_1| \leq 1$.

Finally, we deduce that $\text{gp}(P_r \square C_s) = |S| = |S_1| + |S_2| = |S_1| + |(B_1 \cup B_2 \cup B_3) \cap S_1| + |B_4 \cap S_1| \leq 5$. By using similar arguments, we can again obtain a similar conclusion for any other possible relationship between a, b, c and a', b', c' .

Now, if $|S_2| = 3$ and s is even, then an identical procedure as above leads to the same conclusion. Finally, if $|S_2| = 3$ and s is odd, then the arguments above do not apply since A_2 has only $\lfloor s/2 \rfloor$ P_r -layers. However, it must happen $|S_1| \leq 2$, since otherwise, we proceed as in the previous situation (when $|S_1| = 3$). Consequently, we have that $\text{gp}(P_r \square C_s) = |S| = |S_1| + |S_2| \leq 5$. This completes the proof of this case, and therefore, of the whole theorem. $\square$

4. Torus Graphs

Knowing $\text{gp}(P_r \square P_s)$ and $\text{gp}(P_r \square C_s)$, the next task is to consider the *torus graphs* $C_r \square C_s$, $r, s \geq 3$, where we keep the convention that $V(C_n) = [n]_0$. In contrast to the former two cases, for the torus graphs we are not able to give an exact result, but we will prove a theorem which reduces the general position number of the torus graphs to only two cases. The following lemma and short remarks will be useful for our proof.

Lemma 4.1 *If S is a general position set in $P_r \square P_s$ and there exists $x \in S$, with $\deg(x) = 2$, then $|S| \leq 3$.*

Proof. Suppose on the contrary that $|S| = 4$. Assume without loss of generality that $(0, 0) \in S$. Let $(i, i'), (j, j'), (k, k')$ be the other three vertices of S where $i \leq j \leq k$. If $i' \leq j'$, then $(0, 0), (i, i'), (j, j')$ lie on a common geodesic. So $i' > j'$. Similarly, $j' > k'$. But now $(i, i'), (j, j'), (k, k')$ lie on a common geodesic. $\square$

Remark 4.2. If $S = \{(0, 0), (i, i'), (j, j')\}$ is a general position set of $P_r \square P_s$, then $i \neq j$ and $i' \neq j'$.

Remark 4.3. The graph P_r is an isometric subgraph of C_s if $r \leq \lfloor \frac{s}{2} \rfloor + 1$.

We now bound $\text{gp}(C_r \square C_s)$ from the above as follows.

Theorem 4.4 *If $r \geq 3$ and $s \geq 3$, then $\text{gp}(C_r \square C_s) \leq 7$.*

Proof. Let $S \subseteq V(C_r \square C_s)$ be a general position set and let $w \in S$. Since $C_r \square C_s$ is a vertex-transitive graph (meaning that for each pair of vertices of $C_r \square C_s$ there is an automorphism which maps the first vertex into the second), we can without loss of generality set $w = (\lfloor \frac{r}{2} \rfloor, \lfloor \frac{s}{2} \rfloor)$. Consider the following four subgraphs of $C_r \square C_s$ (where addition is done modulo r or s):

- $X_1 = \{(u_i, v_j) : u_i \in [\lfloor \frac{r}{2} \rfloor + 1]_0, v_j \in [\lfloor \frac{s}{2} \rfloor + 1]_0\}$,
- $X_2 = \{(u_i, v_j) : u_i \in \{\lfloor \frac{r}{2} \rfloor, \lfloor \frac{r}{2} \rfloor + 1, \dots, \lfloor \frac{r}{2} \rfloor + \lfloor \frac{r}{2} \rfloor\}, v_j \in [\lfloor \frac{s}{2} \rfloor + 1]_0\}$,
- $X_3 = \{(u_i, v_j) : u_i \in \{\lfloor \frac{r}{2} \rfloor, \lfloor \frac{r}{2} \rfloor + 1, \dots, \lfloor \frac{r}{2} \rfloor + \lfloor \frac{r}{2} \rfloor\}, v_j \in \{\lfloor \frac{s}{2} \rfloor, \lfloor \frac{s}{2} \rfloor + 1, \dots, \lfloor \frac{s}{2} \rfloor + \lfloor \frac{s}{2} \rfloor\}$,
- $X_4 = \{(u_i, v_j) : u_i \in [\lfloor \frac{r}{2} \rfloor + 1]_0, v_j \in \{\lfloor \frac{s}{2} \rfloor, \lfloor \frac{s}{2} \rfloor + 1, \dots, \lfloor \frac{s}{2} \rfloor + \lfloor \frac{s}{2} \rfloor\}$.

For each $i \in [4]$, the graph X_i is isomorphic to $P_{\lfloor \frac{r}{2} \rfloor + 1} \square P_{\lfloor \frac{s}{2} \rfloor + 1}$. Hence, by Remark 4.3, every X_i is an isometric subgraph of $C_r \square C_s$. Since $w \in X_i$, $i \in [4]$, it is clear that $\text{gp}(C_r \square C_s) \leq 9$ (by Lemma 4.1). This bound can be further improved using the following:

Claim: If $|(X_1 \cup X_3) \cap S| = 5$, then $|(X_2 \cup X_4) \cap S| \leq 2$.

Clearly, if the claim is proved, then by symmetry we also infer that if $|(X_2 \cup X_4) \cap S| = 5$, then $|(X_1 \cup X_3) \cap S| \leq 2$. From these two facts the assertion of the theorem will follow. It thus remains to prove the claim.

Let $u_1 = (x_1, y_1)$, $u_2 = (x_2, y_2)$, $u_3 = (x_3, y_3)$, and $u_4 = (x_4, y_4)$ be vertices such that $X_1 \cap S = \{u_1, u_2, w\}$ and $X_3 \cap S = \{u_3, u_4, w\}$. Suppose that there are vertices $u_5 = (x_5, y_5)$ and $u_6 = (x_6, y_6)$ such that $\{u_5, u_6\} \subseteq X_2 \cap S$. As S is a general position, Remark 4.2 implies that we may without loss of generality assume that $x_1 < x_2 < x_3 < x_4$ and $y_2 < y_1 < y_4 < y_3$. Further, we may also assume that $x_5 < x_6$ and $y_5 < y_6$.

If $x_3 - x_1 \leq \frac{r}{2}$, then either w lies on a u_1, u_3 -geodesic (if $y_3 - y_1 \leq \frac{s}{2}$) or u_2 lies on such a geodesic (when $y_3 - y_1 \geq \frac{s}{2}$). It follows that $x_3 - x_1 > \frac{r}{2}$. With a similar reasoning we also get that $y_3 - y_1 > \frac{s}{2}$, since otherwise u_4 is on a u_1, u_3 -geodesic.

Applying the same reasoning the following is obtained: $x_4 - x_2 > \frac{r}{2}$ and $y_4 - y_2 > \frac{s}{2}$. We now distinguish the following cases.

Case 1: $(y_3 - y_6 \geq \frac{s}{2})$.

Subcase 1a: $x_3 < x_6$.

First, $x_3 > x_5$, since otherwise u_5 lies on a u_3, u_6 -geodesic. If $y_4 - y_5 \geq \frac{s}{2}$, then u_3 lies on a u_4, u_5 -geodesic. So, $y_4 - y_5 < \frac{s}{2}$. If $x_4 > x_6$, then u_6 is on a u_4, u_5 -geodesic, so $x_6 > x_4$. If $y_1 \leq y_6$, then u_6 is on a u_1, u_4 -geodesic, so $y_1 > y_6$. But if $y_1 > y_6$, then u_6 is on a u_1, u_3 -geodesic.

Subcase 1b: $x_3 > x_6$.

FIGURE 6. General position set with seven vertices

If $y_4 - y_6 \geq \frac{s}{2}$, then u_3 is on a u_4, u_6 -geodesic. So, $y_4 - y_6 < \frac{s}{2}$. But now u_6 is on a u_4, u_5 -geodesic.

Case 2: $(y_3 - y_6 < \frac{s}{2})$.

Subcase 2a: $x_3 < x_6$.

If $x_6 > x_4$, then u_4 is on a u_3, u_6 -geodesic. So, $x_4 > x_6$. Also, $y_4 - y_5 > \frac{s}{2}$, since otherwise u_6 is on a u_4, u_5 -geodesic. If $x_3 > x_5$, then u_3 is on a u_4, u_5 -geodesic, so $x_3 < x_5$.

We observe that $y_1 < y_6$, for otherwise u_6 is on a u_1, u_5 -geodesic. If $x_6 - x_2 > \frac{r}{2}$, then u_1 is on a u_2, u_6 -geodesic. Therefore, $x_6 - x_2 < \frac{r}{2}$. This implies that $y_2 > y_5$, for otherwise u_5 lies on a u_2, u_6 -geodesic. But now, u_5 lies on a u_1, u_3 -geodesic.

Subcase 2b ($x_3 > x_6$).

If $y_4 - y_5 \geq \frac{s}{2}$, then u_3 is on a u_4, u_5 -geodesic, and if $y_4 - y_5 < \frac{s}{2}$, then u_6 is on a u_4, u_5 -geodesic. $\square$

A general position set S of $C_7 \square C_7$ with $|S| = 7$ is shown on Fig. 6. It can be easily checked that the minimum distance between the pairs of vertices from S is 3, and that the maximal distance is 5. Therefore, S is indeed a general position set.

The construction from Fig. 6 cannot be extended to arbitrary $r, s \geq 7$. We have checked by computer that $\text{gp}(C_8 \square C_7) \leq 6$. In fact, 6 is also the exact lower bound in this case as follows from the following, main result of this section.

Theorem 4.5 *If $r \geq s \geq 3$, $s \neq 4$, and $r \geq 6$, then $\text{gp}(C_r \square C_s) \in \{6, 7\}$.*

FIGURE 7. The set S in $C_6 \square C_3$ appears in bold

Proof. From Theorem 4.4 we know that $\text{gp}(C_r \square C_s) \leq 7$. It thus remains to prove that $\text{gp}(C_r \square C_s) \geq 6$.

The condition $s \neq 4$ assures that $S_s = \{0, \lfloor \frac{s}{3} \rfloor, \lfloor \frac{2s}{3} \rfloor\}$ is a gp-set of C_s . The condition that $r \geq 6$ assures that $\lfloor r/6 \rfloor \geq 1$. Consider now the set

$$S = \{(0, 0), (\lfloor r/2 \rfloor, 0), (\lfloor r/6 \rfloor, \lfloor s/3 \rfloor), (\lfloor r/6 \rfloor + \lfloor r/2 \rfloor, \lfloor s/3 \rfloor), \\ (\lfloor (2r)/6 \rfloor, \lfloor (2s)/3 \rfloor), (\lfloor (2r)/6 \rfloor + \lfloor r/2 \rfloor, \lfloor (2s)/3 \rfloor)\}.$$

In Fig. 7 the set S is shown for the case $C_6 \square C_3$.

We claim that S is a general position set. Since C_p -layers are isometric subgraphs, no other vertex is on a geodesic between the pair of vertices with the same second coordinate. Hence we only need to consider the triples of vertices from S with pairwise different second coordinates. We do this for the vertices $x_1 = (0, 0)$, $x_2 = (\lfloor r/6 \rfloor, \lfloor s/3 \rfloor)$, and $x_3 = (\lfloor (2r)/6 \rfloor, \lfloor (2s)/3 \rfloor)$, the other cases are treated similarly. Since x_1, x_2, x_3 lie in a subgraph of $C_r \square C_s$ isomorphic to $P_{\lfloor r/3 \rfloor + 1} \square C_s$ which is an isometric subgraph, it suffices to show that $d(x_1, x_3) < d(x_1, x_2) + d(x_2, x_3)$. This can be verified using the facts $d(x_1, x_3) = \lfloor r/3 \rfloor + (s - \lfloor (2s)/3 \rfloor)$, $d(x_1, x_2) = \lfloor r/6 \rfloor + \lfloor s/3 \rfloor$, and $d(x_2, x_3) \geq \lfloor r/6 \rfloor + \lfloor s/3 \rfloor$. $\square$

Theorems 4.4 and 4.5 yield the following problem, for which we believe that if $\min\{r, s\} \leq 6$, then $\text{gp}(C_r \square C_s) = 6$.

Problem 4.6 Determine for $r \geq 3, r \neq 4, s \geq 6$, whether $\text{gp}(C_r \square C_s) = 6$ or whether $\text{gp}(C_r \square C_s) = 6$.

5. Cartesian Powers

In this section, we consider the general position number of Cartesian powers and obtain asymptotically exponential lower bounds using a probabilistic approach.

The n -dimensional hypercube Q_n is defined as $K_2^{\square, n}$. In particular, $Q_1 = K_2$, $Q_2 = C_4$, and Q_3 is the graph of the 3-D cube. Cartesian products of complete graphs, known as Hamming graphs, form a natural generalization of hypercubes. In [6], it was proved that if $k \geq 2$ and $n_1, \dots, n_k \geq 2$, then

$$\text{gp}(K_{n_1} \square \dots \square K_{n_k}) \geq n_1 + \dots + n_k - k. \quad (1)$$

Moreover, this lower bound is sharp on products of two complete graphs, that is, $\text{gp}(K_{n_1} \square K_{n_2}) = n_1 + n_2 - 2$.

The situation above changes dramatically as k grows. Körner [11] obtained a probabilistic construction of general position sets in Q_n of size $\frac{1}{2} \frac{2^n}{\sqrt{3^n}}$. He also pointed out that the problem of finding the size of the largest point set in general position in Q_n is equivalent to finding the largest size of what is called a $(2, 1)$ -separating system in coding theory. (For more on separating systems, see [4].) Körner was interested in

$$\alpha = \limsup_{n \rightarrow \infty} \frac{\log_2 \text{gp}(Q_n)}{n}.$$

His probabilistic lower bound gives $\alpha \geq 1 - \frac{1}{2} \log_2 3$ and he also proved $\alpha \leq 1/2$. Later, Randriambololona [17] improved the lower bound to $\alpha \geq \frac{3}{50} \log_2 11$ with an explicit construction.

The first moment method can be applied in a general setting to obtain large general position sets. For any graph G , let $p(G)$ denote the probability that if one picks a triple $(x, y, z) \in V(G)^3$ uniformly at random, then $d_G(y, z) = d_G(y, x) + d_G(x, z)$ holds. Let us call such triples *bad*. Note that this is never the case if $x \neq y$ and $y = z$, so $p(G) \leq 1 - \frac{|V(G)|-1}{|V(G)|^2} < 1$. Let $H = H_1 \square \cdots \square H_k$. Observe that the triple $\mathbf{x} = (x_1, \dots, x_k), \mathbf{y} = (y_1, \dots, y_k), \mathbf{z} = (z_1, \dots, z_k) \in V(H)$ is bad in H if and only if the triples x_i, y_i, z_i are bad in H_i for all $i \in [k]$. So if we pick M vertices uniformly at random with repetition from $V(H)$, then the expected value $\mathbb{E}(X)$ of the number $X = X(M)$ of unordered triples on a geodesic will be $3 \binom{M}{3} \prod_{i=1}^k p(H_i)$. If $X \leq M/2$, then removing one vertex from every bad triple will leave us a general position set of size at least $M/2$. As there is always an instance for which $X \leq \mathbb{E}(X)$ holds, we obtain a general position set of size $M/2$ provided $3 \binom{M}{3} \prod_{i=1}^k p(H_i) \leq M/2$ holds. Therefore, it seems to be interesting to examine

$$\text{gp}_{\square}(G) := \limsup_{n \rightarrow \infty} \frac{\log_{|V(G)|} \text{gp}(G^{\square, n})}{n}.$$

Clearly, we have $\text{gp}_{\square}(G) \leq 1$ and the above reasoning yields the following theorem.

For a graph G one can consider its Cartesian power $G^{\square, n}$. Then the required inequality is $3 \binom{M}{3} p(G)^n \leq M/2$ which is equivalent to $(M-1)(M-2) \leq p(G)^{-n}$. Thus there exists a general position set in $G^{\square, n}$ of size $\frac{1}{2} p(G)^{-n/2}$. This and the inequality $p(G) \leq 1 - \frac{|V(G)|-1}{|V(G)|^2}$ yields the following statement.

Theorem 5.1 *If G is a graph, then*

$$\text{gp}_{\square}(G) \geq \log_{|V(G)|} p(G)^{-1/2} \geq 1 - \log_{|V(G)|} (|V(G)|^2 - |V(G)| + 1).$$

Let us calculate $p(G)$ for some graphs. First of all, $p(K_n) = \frac{2n-1}{n^2}$ as in K_n the equality $d(y, z) = d(y, x) + d(x, z)$ holds if and only if $x = y$ or $x = z$. (The

case $p(K_2) = \frac{3}{4}$ in Theorem 5.1 is just Körner's result.) For even cycles we have $p(C_{2k}) = \frac{k(k+3)-1}{4k^2}$. If the vertices are $\{-(k-1), -(k-2), \dots, 0, \dots, k-1, k\}$ in this cyclic order, then by symmetry we can assume $x = 0$. There are $4k-1$ triples with $x = y$ or $x = z$ that form bad triples. If $y = k$ or $z = k$, then there are no other bad triples, otherwise for any y , there are $k - |y|$ ways to choose z to obtain a bad triple. Similarly, one can verify $p(C_{2k+1}) = \frac{k(k+3)+1}{(2k+1)^2}$. Finally, consider the star S_k with k leaves. Then conditioning on whether x is the center or not one obtains $p(S_k) = \frac{1}{k+1} + \frac{k}{k+1} \frac{2k+1}{(k+1)^2}$. Observe that if one picks uniformly at random only among the leaves of S_k , then the probability of picking a bad triple is $p'(S_k) = \frac{2k-1}{k^2}$ which for large enough k is roughly $2/3$ of $p(S_k)$, so in this way one obtains the better bound $\text{gp}_{\square}(S_k) \geq \log_2 p'(S_k)^{-1/2}$.

Concerning $\text{gp}_{\square}(G)$ we wonder whether one can write limit instead of limit superior in the definition of $\text{gp}_{\square}(G)$. Moreover, by the above we have $\lim_{k \rightarrow \infty} p(C_k) = \frac{1}{4}$. We also pose:

Problem 5.2 *Decide whether $\liminf_{k \rightarrow \infty} \frac{\text{gp}_{\square}(C_k)}{\log_k 2} > 1$ holds.*

Acknowledgements

We acknowledge the financial support from the Slovenian Research Agency (research core funding No. P1-0297 and projects J1-9109, J1-1693, N1-0095, N1-0108). This research was done while the last author was visiting the University of Ljubljana, Slovenia, supported by “Ministerio de Educación, Cultura y Deporte”, Spain, under the “José Castillejo” program for young researchers (reference number: CAS18/00030). The research of the second author was supported by the National Research, Development and Innovation Office - NKFIH under the grants SNN 129364 and FK 132060.

Funding Open access funding provided by ELKH Alfréd Rényi Institute of Mathematics

Declarations

Conflict of Interest The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's

Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- [1] Anand, B.S., Ullas Chandran, S.V., Changat, M., Klavžar, S., Thomas, E.J.: Characterization of general position sets and its applications to cographs and bipartite graphs. *Appl. Math. Comput.* **359**, 84–89 (2019)
- [2] Bukh, B., Matoušek, J.: Erdős–Szekeres-type statements: Ramsey function and decidability in dimension 1. *Duke Math. J.* **163**, 2243–2270 (2014)
- [3] Cardinal, J., Toth, C.D., Wood, D.R.: General position subsets and independent hyperplanes in d -space. *J. Geometry* **108**, 33–43 (2017)
- [4] Cohen, G.D., Schaathun, H.G.: Asymptotic overview on separating codes, Tech. Report 248, Department of Informatics, University of Bergen, Bergen, Norway (2003)
- [5] Dudeney, H.E.: *Amusements in Mathematics*. Nelson, Edinburgh (1917)
- [6] Ghorbani, M., Klavžar, S., Maimani, H.R., Momeni, M., Rahimi-Mahid, F., Rus, G.: The general position problem on Kneser graphs and on some graph operations. *Discuss. Math. Graph Theory* (2019). <https://doi.org/10.7151/dmgt.2269>
- [7] Imrich, W., Klavžar, S., Rall, D.F.: *Topics in Graph Theory: Graphs and Their Cartesian Product*. A K Peters, Wellesley, MA (2008)
- [8] Kanté, M.M., Sampaio, R.M., dos Santos, V.F., Szwarcfiter, J.L.: On the geodetic rank of a graph. *J. Combin.* **8**, 323–340 (2017)
- [9] Klavžar, S., Rus, G.: The general position number of integer lattices. *Appl. Math. Comput.* **390**, Paper 125664 (2021)
- [10] Klavžar, S., Yero, I.G.: The general position problem and strong resolving graphs. *Open Math.* **17**, 1126–1135 (2019)
- [11] Körner, J.: On the extremal combinatorics of the Hamming space. *J. Combin. Theory Ser. A* **71**, 112–126 (1995)
- [12] Ku, C.Y., Wong, K.B.: On no-three-in-line problem on m -dimensional torus. *Graphs Combin.* **34**, 355–364 (2018)
- [13] Manuel, P., Klavžar, S.: A general position problem in graph theory. *Bull. Aust. Math. Soc.* **98**, 177–187 (2018)
- [14] Manuel, P., Klavžar, S.: The graph theory general position problem on some interconnection networks. *Fund. Inf.* **163**, 339–350 (2018)
- [15] Misiak, A., Stepień, Z., Szymaszkiewicz, A., Szymaszkiewicz, L., Zwierzchowski, M.: A note on the no-three-in-line problem on a torus. *Discrete Math.* **339**, 217–221 (2016)
- [16] Neethu, P.K., Ullas Chandran, S.V., Changat, M., Klavžar, S.: On the general position number of complementary prisms. *Fund. Inf.* **178**, 267–281 (2021)

- [17] Randriambololona, H.: $(2, 1)$ -separating systems beyond the probabilistic bound. *Israel J. Math.* **195**, 171–186 (2013)
- [18] Patkós, B.: On the general position problem on Kneser graphs. *ARS Math. Contemp.* **18**, 273–280 (2020)
- [19] Payne, M., Wood, D.R.: On the general position subset selection problem. *SIAM J. Discrete Math.* **27**, 1727–1733 (2013)
- [20] Por, A., Wood, D.R.: No-three-in-line-in-3D. *Algorithmica* **47**, 481–488 (2007)
- [21] Sloane, N.J.A. (ed.): The On-Line Encyclopedia of Integer Sequences (2019). published electronically at <https://oeis.org>
- [22] Thomas, E.J., Ullas Chandran, S.V.: Characterization of classes of graphs with large general position number. *AKCE Int. J. Graphs Combin.* **17**, 935–939 (2020)
- [23] Tian, J., Xu, K.: The general position number of Cartesian products involving a factor with small diameter. *Appl. Math. Comput.* **403**, 126206 (2021)
- [24] Tian, J., Xu, K., Klavžar, S.: The general position number of Cartesian product of two trees. *Bull. Aust. Math. Soc.* (2020). <https://doi.org/10.1017/S0004972720001276>
- [25] Ullas Chandran, S.V., Parthasarathy, G.Jaya: The geodesic irredundant sets in graphs. *Int. J. Math. Combin.* **4**, 135–143 (2016)
- [26] Yao, Y., He, M., Ji, S., Li, G.: On the general position set of two classes of graphs (2020). [arXiv:2002.03108](https://arxiv.org/abs/2002.03108)

Sandi Klavžar

Faculty of Mathematics and Physics
University of Ljubljana
Ljubljana
Slovenia

e-mail: sandi.klavzar@fmf.uni-lj.si

and

Faculty of Natural Sciences and Mathematics
University of Maribor
Maribor
Slovenia

Sandi Klavžar and Gregor Rus

Institute of Mathematics, Physics and Mechanics
Ljubljana
Slovenia

e-mail: gregor.rus4um.si

Balázs Patkós

Alfréd Rényi Institute of Mathematics
Hungarian Academy of Sciences
Budapest
Hungary

e-mail: patkos.balazs@renyi.hu

Gregor Rus
Faculty of Organizational Sciences
University of Maribor
Maribor
Slovenia

Ismael G. Yero
Departamento de Matemáticas
Universidad de Cádiz
Algeciras
Spain
e-mail: ismael.gonzalez@uca.es

Received: November 4, 2020.

Accepted: May 17, 2021.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.