



Norm Attaining Elements of the Ball Algebra $H^\infty(B_N)$

Richard M. Aron, José Bonet , and Manuel Maestre

Abstract. Let B_N be the Euclidean ball of \mathbb{C}^N . The space $H^\infty(B_N)$ of bounded holomorphic functions on B_N is known to have a predual, denoted by $G^\infty(B_N)$. We study the functions in $H^\infty(B_N)$ that attain their norm as elements of the dual of $G^\infty(B_N)$. We also examine similar questions for the polydisc algebra $H^\infty(\mathbb{D}^N)$ and for the space of Dirichlet series $\mathcal{D}^\infty(\mathbb{C}_+)$.

Mathematics Subject Classification. Primary 46E15; Secondary 46B04, 46B10, 46B20.

Keywords. Norm attaining, predual, ball algebra, polydisc algebra, Dirichlet series.

1. Introduction

Ando [1] proved that the Banach space $H^\infty(\mathbb{D})$ of bounded holomorphic functions on the unit disc \mathbb{D} has a unique isometric predual. Let us denote it by $G^\infty(\mathbb{D})$. By the Bishop-Phelps theorem, the set $NA(G^\infty(\mathbb{D}))$ of functions $f \in H^\infty(\mathbb{D})$ which attain their norm as elements of the dual of $G^\infty(\mathbb{D})$ is a norm-dense subset of $H^\infty(\mathbb{D})$. Fisher [6] showed that $f \in H^\infty(\mathbb{D})$, $\|f\| = 1$, attains its norm as an element of the dual of $G^\infty(\mathbb{D})$ if and only if the radial limits $f^*(w)$ of f in the torus \mathbb{T} satisfy that the set $\{w \in \mathbb{T} : |f^*(w)| = 1\}$ has positive Lebesgue measure on \mathbb{T} . The aim of this article is to investigate versions of Fisher's result for the Banach space of bounded holomorphic functions on the N -dimensional ball and the N -dimensional polydisc. Our main results are Theorems 5 and 8 and Propositions 6 and 7 in the case of the ball.

Richard M. Aron, José Bonet and Manuel Maestre have contributed equally to this work.

The case of the polydisc is treated in Sect. 3. The final section deals with the Banach space of bounded Dirichlet series.

Let X be a complex Banach space. Its open unit ball is denoted by B_X and its closed unit ball by U_X . The space of all holomorphic functions on B_X (i.e. the \mathbb{C} -Fréchet differentiable functions $f : B_X \rightarrow \mathbb{C}$) will be denoted $H(B_X)$. The Banach space $H^\infty(B_X)$ of all bounded holomorphic functions f in $H(B_X)$ is endowed with the supremum norm $\|f\|_\infty = \sup_{x \in B_X} |f(x)|$. We denote by τ_0 the compact-open topology on $H^\infty(B_X)$, that is, the topology of uniform convergence on compact subsets of B_X . Recall that τ_0 is Hausdorff and coarser than the norm topology. Let $U_{H^\infty(B_X)}$ denote the closed unit ball of $H^\infty(B_X)$. The vector space $G^\infty(B_X)$, given by

$$G^\infty(B_X) := \{\varphi \in H^\infty(B_X)^* : \varphi|_{U_{H^\infty(B_X)}} \text{ is } \tau_0\text{-continuous}\}$$

is a Banach space when endowed with the dual norm. By using the Ng-Dixmier Theorem [12], Mujica [11], proved that the topological dual of $G^\infty(B_X)$ is isometrically isomorphic to $H^\infty(B_X)$. We abbreviate this fact by

$$G^\infty(B_X)^* \stackrel{1}{=} H^\infty(B_X).$$

For each $x \in B_X$ we denote by $\delta_x : H^\infty(B_X) \rightarrow \mathbb{C}$ the evaluation $\delta_x(f) := f(x)$ at the point x . Clearly δ_x is τ_0 continuous. Moreover, the vector space $\text{span}\{\delta_x : x \in B_X\}$ is a norm-dense subset in $G^\infty(B_X)$. Indeed, $\{\delta_x : x \in B_X\}$ separates points of $H^\infty(B_X)$. Hence $\text{span}\{\delta_x : x \in B_X\}$ is a subspace of $G^\infty(B_X)$ that is $w(G^\infty(B_X), H^\infty(B_X))$ -dense in $G^\infty(B_X)$. Thus it is also norm-dense subset of $G^\infty(B_X)$. We collect the following consequence for reference later in the paper.

Lemma 1. *If \mathcal{F} is a closed subspace of $G^\infty(B_X)$ containing $\{\delta_x : x \in B_X\}$, then $\mathcal{F} = G^\infty(B_X)$.*

Let Y be a Banach space. The set of norm attaining functionals is defined to be the following subset of Y^* :

$$NA(Y) := \{y^* \in Y^* : \text{there exists } y \in Y, \|y\| = 1 \text{ such that } \|y^*\| = y^*(y)\}$$

The Bishop–Phelps theorem (see, e.g., Theorem 8.11 in [2]) ensures that the set $NA(Y)$ of norm attaining functionals is a norm-dense subset of Y^* . As a consequence, for each non-trivial, complex Banach space X , there exists a norm-dense subset $NA(G^\infty(B_X))$ of $H^\infty(B_X)$, such that for every $f \in NA(G^\infty(B_X))$, there exists an element $\varphi \in G^\infty(B_X)$ with $\|\varphi\| = 1$ such that

$$\|f\|_\infty = \varphi(f).$$

The aim of this paper is to study those functions $f \in H^\infty(B_X)$ that attain their norm as elements of the dual of $G^\infty(B_X)$, that is, those $f \in NA(G^\infty(B_X))$. We mainly concentrate on the case $X = (\mathbb{C}^N, \|\cdot\|_2)$ and hence, B_X is the N -dimensional Euclidean ball which henceforth will be denoted B_N .

In the one dimensional case, $B_N = \mathbb{D}$ and its boundary is the torus $\mathbb{T} = \{z \in \mathbb{C} : |z| = 1\}$. In this case, by a result by Fatou, there is an isometric isomorphism between $H^\infty(\mathbb{D})$ and

$$H^\infty(\mathbb{T}) := \left\{ g \in L^\infty(\mathbb{T}) : \hat{g}(k) = \int_{\mathbb{T}} w^{-k} g(w) dm_1(w) = 0, k = -1, -2, \dots \right\}.$$

The isometric isomorphism $H^\infty(\mathbb{D}) \rightarrow H^\infty(\mathbb{T})$ is given by

$$\begin{aligned} H^\infty(\mathbb{D}) &\longrightarrow H^\infty(\mathbb{T}) \\ f &\longrightarrow f^* \end{aligned}$$

where the radial limit

$$f^*(w) := \lim_{r \rightarrow 1^-} f(rw),$$

exists almost everywhere on \mathbb{T} (with respect to the Lebesgue normalized measure on \mathbb{T} , denoted by $dm_1(w) = \frac{dt}{2\pi}$, where $w = e^{it}$.) From this point of view $H^\infty(\mathbb{D}) \stackrel{1}{=} H^\infty(\mathbb{T})$ is a closed subspace of $L^\infty(\mathbb{T})$, and hence it is a dual space. In fact, if $H_0^1(\mathbb{T})$ is the closed subspace of $L^1(\mathbb{T})$ given by

$$H_0^1(\mathbb{T}) = \left\{ f \in L_1(\mathbb{T}) : \hat{f}(-n) = \int_{\mathbb{T}} f(w) w^n dm_1(w) = 0, \text{ for all } n = 0, 1, 2, \dots \right\},$$

then

$$H^\infty(\mathbb{T}) \stackrel{1}{=} \left(L^1(\mathbb{T}) / H_0^1(\mathbb{T}) \right)^*.$$

Ando in [1] proved that $H^\infty(\mathbb{D})$ has a unique isometric predual. Accordingly, $L^1(\mathbb{T}) / H_0^1(\mathbb{T}) \stackrel{1}{=} G^\infty(\mathbb{D})$. As far as we know, it is an open question for $N \geq 2$ whether there is a unique predual of the corresponding H^∞ -spaces in the case of the N -dimensional ball and the N -polydisc. In this paper, we will introduce another natural predual and show, in Theorems 5 and 10, that it coincides with $G^\infty(B_X)$.

The characterization of norm attaining elements of $f \in H^\infty(\mathbb{D})$ was obtained by S. Fisher in 1969.

Theorem 2 (Fisher [6, Theorem 2]). *Let f be an element of norm one in $H^\infty(\mathbb{D})$. The function f attains its norm as an element of the dual of $L^1(\mathbb{T}) / H_0^1(\mathbb{T}) = G^\infty(\mathbb{D})$ if and only if $f^*(w) = \lim_{r \rightarrow 1^-} f(rw)$ (a.e. in \mathbb{T}) satisfies that*

$$\{w \in \mathbb{T} : |f^*(w)| = 1\}$$

has positive Lebesgue measure on \mathbb{T} .

In this paper, in Sect. 2, we explore several variable versions of Fisher’s result. We also examine, in Sects. 3 and 4, similar questions for the polydisc algebra $H^\infty(\mathbb{D}^N)$ and for the space of Dirichlet series $\mathcal{D}^\infty(\mathbb{C}_+)$.

2. The Case of the Euclidean Ball

Recall that the Euclidean open unit ball in \mathbb{C}^N is:

$$B_N := \left\{ z = (z_1, \dots, z_N) \in \mathbb{C}^N : \|z\|_N := \sqrt[2]{|z_1|^2 + \dots + |z_N|^2} < 1 \right\}.$$

The unit sphere in \mathbb{C}^N is:

$$S_N := \left\{ z = (z_1, \dots, z_N) \in \mathbb{C}^N : \|z\|_N := \sqrt[2]{|z_1|^2 + \dots + |z_N|^2} = 1 \right\}.$$

(Observe that this is not completely standard notation since the usual notation for the N -dimensional real sphere in \mathbb{R}^N is S_{N-1} .)

By σ_N we denote the unique rotation-invariant positive Borel measure on S_N for which

$$\sigma_N(S_N) = 1.$$

In other words, σ_N is the Haar measure of the N -dimensional sphere.

In [15, p.84], the space $H^\infty(B_N)$, is defined as

$$H^\infty(B_N) := \left\{ f \in H(B_N) : \|f\|_\infty := \sup_{z \in B_N} |f(z)| < \infty \right\}.$$

The ball algebra is the Banach subalgebra of $H^\infty(B_N)$ given by

$$A(B_N) := \{ f : \bar{B}_N \rightarrow \mathbb{C} : f \text{ is continuous on } \bar{B}_N \text{ and holomorphic on } B_N \}.$$

Finally, by $A(S_N) = A(B_N) \cap C(S_N)$, we understand the restrictions of the elements of $A(B_N)$ to the sphere S_N , i.e.

$$A(S_N) := \{ f|_{S_N} : f \in A(B_N) \}.$$

By the maximum modulus theorem, the mapping $\pi : A(B_N) \rightarrow A(S_N)$ defined by $\pi(f) := f|_{S_N}$ is an isometry.

Hardy spaces have a dual definition. The Hardy space $H^\infty(S_N)$ is the weak-star closure of $A(S_N)$ in $L^\infty(S_N, \sigma_N)$. i.e.

$$H^\infty(S_N) := \overline{A(S_N)}^{w(L^\infty(S_N), L_1(S_N))}.$$

As the polynomials are dense in $A(B_N)$ we have that $\text{span}\{z^\beta : \beta \in \mathbb{N}_0^N\}$ is a $\|\cdot\|_\infty$ dense subspace of $A(B_N)$. Hence, $\text{span}\{w^\beta : \beta \in \mathbb{N}_0^N\}$ is $\|\cdot\|_\infty$ dense in $A(S_N)$. Thus

$$H^\infty(S_N) = \overline{\text{span}\{w^\beta : \beta \in \mathbb{N}_0^N\}}^{w(L^\infty(S_N), L_1(S_N))}.$$

At this point, we show that $H^\infty(S_N)$ and $H^\infty(B_N)$ are isometrically isomorphic. We need some notation and results that can be found, for example, in the books [15] and [16]. *The invariant Poisson kernel of B_N* is the kernel function $P_N : B_N \times S_N \rightarrow [0, +\infty[$

$$P_N(z, w) := \frac{(1 - |z|^2)^N}{|1 - \langle z, w \rangle|^{2N}}.$$

The *Poisson integral* $P(g)$ of a function g in $L^1(S_N, \sigma_N)$ is defined, for $z \in B_N$, by

$$P_N(g)(z) := \int_{S_N} P(z, w)g(w)d\sigma_N(w).$$

We have that $P_N : H^\infty(S_N) \rightarrow H^\infty(B_N)$ is a linear isometry onto.

To prove that this mapping is onto, the concept of Korányi, or K -limit, of a holomorphic function on B_N is needed. For $\alpha > 1$ and $w \in S_N$ we set

$$D_\alpha(w) := \left\{ z \in \mathbb{C}^N : |w - z| < \frac{\alpha}{2}(1 - |z|^2) \right\}.$$

Clearly $D_\alpha(w) \subset B_N$. We say that a function $F : B_N \rightarrow \mathbb{C}$ has K -limit $\lambda \in \mathbb{C}$ at $w \in S_N$ if the following is true: For every $\alpha > 1$ and for every sequence (z_j) in $D_\alpha(w)$ that converges to a point $w \in S_N$, we have that $F(z_j)$ converges to λ and write

$$(K - \lim F)(w) = \lambda.$$

The following result (see e.g. [15, Section 5.4.]) is important and very useful for our paper.

Theorem 3. *If f is a function in $H^\infty(B_N)$ then f has finite K -limits f^* σ_N -almost everywhere on S_N . Moreover, $f^* \in H^\infty(S_N)$, $\|f^*\|_\infty = \|f\|_\infty$ and*

$$P_N(f^*) = f.$$

In other words, the mapping $f \rightarrow f^$ is a linear isometry from $H^\infty(B_N)$ onto $H^\infty(S_N)$.*

We also need the following well known fact, a proof of which is given for the sake of completeness.

Lemma 4. *Let X be a Banach space and let Y be a weak-star closed subspace of X^* . The subspace*

$$Y_\perp := \{x \in X : y^*(x) = 0, \text{ for all } y^* \in Y\},$$

satisfies

$$Y_\perp^\perp := \{x^* \in X^* : x^*(x) = 0, \text{ for all } x \in Y_\perp\} = Y,$$

and Y is isometrically isomorphic to $(X/Y_\perp)^$.*

Proof. Clearly, by the definition, $Y \subset Y_\perp^\perp$. Assume that the reverse inclusion is not true. Hence there exists $x_0^* \in Y_\perp^\perp \setminus Y$.

Since Y is $w(X^*, X)$ closed and convex we can find $\varphi : X^* \rightarrow \mathbb{C}$, $w(X^*, X)$ -continuous, such that

$$\varphi(x_0^*) = 1 \text{ and } \varphi(y^*) = 0,$$

for all $y^* \in Y$. Since φ is weak-star continuous, there exists $x_0 \in X$ such that

$$\varphi(x^*) = x^*(x_0),$$

for all $x^* \in X^*$. Thus, $x_0^*(x_0) = 1$ and $y^*(x_0) = 0$ for all $y^* \in Y$. Hence x_0 belongs Y_\perp . But, $x_0^* \in Y_\perp^\perp$, which, by definition implies

$$x_0^*(x_0) = 0.$$

This is a contradiction.

Finally, we have $(X/Y_\perp)^* \stackrel{1}{=} Y_\perp^\perp = Y$, as follows from [10, Theorem 1.10.17] for example. □

Now we define

$$H_0^1(S_N) = \left\{ g \in L_1(S_N) : \int_{S_N} g(w)f(w)d\sigma_N(w) = 0 \text{ for all } f \in A(S_N) \right\}.$$

Since

$$H^\infty(S_N) := \overline{A(S_N)}^{w(L_\infty(S_N), L_1(S_N))} = \overline{\text{span}\{w^\beta : \beta \in \mathbb{N}_0^N\}}^{w(L_\infty(S_N), L_1(S_N))},$$

the subspace $H^\infty(S_N) \subset L_\infty(S_N)$ is $w(L_\infty(S_N), L_1(S_N))$ -closed in $L_\infty(S_N)$ and

$$\begin{aligned} H_0^1(S_N) &= \left\{ g \in L_1(S_N) : \int_{S_N} g(w)f(w)d\sigma_N(w) = 0, \text{ for all } f \in H^\infty(S_N) \right\} \\ &= \left\{ g \in L_1(S_N) : \hat{g}(-\beta) := \int_{S_N} g(w)w^\beta d\sigma_N(w) = 0, \text{ for all } \beta \in \mathbb{N}_0^N \right\}. \end{aligned}$$

In the notation of Lemma 4, with $X = L_1(S_N)$, $X^* = L_\infty(S_N)$ and $Y = H^\infty(S_N)$ (which is weak-star closed in X^*), we have

$$\begin{aligned} Y_\perp &= H^\infty(S_N)_\perp = H_0^1(S_N), \\ Y_\perp^\perp &= H_0^1(S_N)^\perp = H^\infty(S_N). \end{aligned}$$

Lemma 4 implies the isometric isomorphism

$$H^\infty(S_N) \stackrel{1}{=} (L_1(S_N)/H_0^1(S_N))^*.$$

Next we show that $G^\infty(B_N)$ and $L^1(S_N)/H_0^1(S_N)$ are isometrically isomorphic. Thus, these two natural preduals of $H^\infty(B_N)$ coincide, and so the extension of Ando’s result on the uniqueness of the predual of $H^\infty(\mathbb{D})$ to several variables is still open.

Theorem 5. *For every $N \in \mathbb{N}$ we have that*

$$L^1(S_N)/H_0^1(S_N) = G^\infty(B_N)$$

isometrically.

Proof. First we prove that $L^1(S_N)/H_0^1(S_N) \subset G^\infty(B_N)$.

Let $[\varphi] \in L^1(S_N)/H_0^1(S_N)$ and $g \in H^\infty(S_N)$. The duality is given by

$$\langle [\varphi], g \rangle = \int_{S_N} \varphi(w)g(w)d\sigma_N(w) = \int_{S_N} (\varphi(w) + \eta(w))g(w)d\sigma_N(w),$$

for every $\varphi \in L_1(S_N)$ and every $\eta \in H_0^1(S_N)$.

We identify $L^1(S_N)/H_0^1(S_N)$ as a subspace of the dual of $H^\infty(S_N)$ in the following natural way. Define $T_{[\varphi]} : H^\infty(B_N) \rightarrow \mathbb{C}$ by

$$T_{[\varphi]}(f) := \langle [\varphi], f^* \rangle = \int_{S_N} \varphi(w) f^*(w) d\sigma_N(w).$$

We check that $T_{[\varphi]}$ belongs to $G^\infty(B_N)$ for every equivalence class $[\varphi] \in L^1(S_N)/H_0^1(S_N)$.

Clearly

$$|T_{[\varphi]}(f)| \leq \int_{S_N} |\varphi(w)| \|f^*\|_\infty d\sigma_N(w) = \|\varphi\|_1 \|f\|_\infty.$$

Hence, $T_{[\varphi]}$ belongs to $H^\infty(B_N)^*$. This fact and the equality $\|T_{[\varphi]}\| = \|[\varphi]\|$ are consequences of the isometric isomorphism $H^\infty(S_N) \stackrel{1}{=} (L_1(S_N)/H_0^1(S_N))^*$ and Theorem 3.

Let us check that $T_{[\varphi]}$ is τ_0 -continuous when restricted to the closed unit ball $U_{H^\infty(B_N)}$ of $H^\infty(B_N)$.

By Theorem 3, we know that if $f \in H^\infty(B_N)$ and $f^* \in H^\infty(S_N)$ is its K -limit that exists a.e. in S_N , then

$$f(z) = \int_{S_N} P_N(z, w) f^*(w) d\sigma_N(w)$$

for all $z \in B_N$. Conversely, if $h \in H^\infty(S_N)$, then $P_N(h) \in H^\infty(B_N)$ and we have

$$P_N(h)^*(w) = h(w)$$

a.e. on S_N .

For each $z \in B_N$ the mapping $P_N(z, \cdot) : S_N \rightarrow]0, +\infty[$ is continuous on S_N . Hence $P_N(z, \cdot) \in L^1(S_N)$.

Given $(f_n) \cup \{f\} \subset U_{H^\infty(B_N)}$ such that (f_n) converges to f with respect to the compact-open topology on B_N , we have $(f_n^*) \cup \{f^*\} \subset U_{H^\infty(S_N)}$. But $U_{H^\infty(S_N)}$ is a weak-star closed subset of $U_{L^\infty(S_N)}$ which, in turn, is a $w(L^\infty(S_N), L^1(S_N))$ -compact set. Since $L^1(S_N)$ is separable, it follows that $U_{H^\infty(S_N)}$ is a metrizable compact set with the weak-star topology. Consider now any subsequence $(f_{n_k}^*)$ that is $w(L^\infty(S_N), L^1(S_N))$ -convergent to some $h \in U_{H^\infty(S_N)}$. We will have

$$\begin{aligned} P_N(h)(z) &= \int_{S_N} P_N(z, w) h(w) d\sigma_N(w) \\ &= \langle P_N(z, \cdot), h \rangle = \lim_{k \rightarrow \infty} \langle P_N(z, \cdot), f_{n_k} \rangle \\ &= \lim_{k \rightarrow \infty} \int_{S_N} P_N(z, w) f_{n_k}^*(w) d\sigma_N(w) \\ &= \lim_{k \rightarrow \infty} f_{n_k}(z) = f(z), \end{aligned}$$

for all $z \in B_N$. Hence,

$$h(w) = P_N(h)^*(w) = f^*(w)$$

a.e in S_N . We have just proved that the only weak-star adherent point of (f_n^*) is f^* . Thus (f_n^*) weak-star converges to f^* . In particular

$$\begin{aligned} T_{[\varphi]}(f) &= \int_{S_N} f^*(w)\varphi(w)d\sigma_N(w) \\ &= \langle [\varphi], f^* \rangle = \lim_{n \rightarrow \infty} \langle [\varphi], f_n^* \rangle \\ &= \lim_{n \rightarrow \infty} T_{[\varphi]}(f_n), \end{aligned}$$

and $T_{[\varphi]}$ is continuous with the compact-open topology when restricted to the closed unit ball of $H^\infty(B_N)$; i.e. $T_{[\varphi]} \in G^\infty(B_N)$.

For the other inclusion observe that

$$\delta_z(f) = P_N(f^*)(z) = \int_{S_N} P_N(z, w)f^*(w)d\sigma_N(w) = T_{[P_N(z, \cdot)]}(f),$$

for every $z \in B_N$ and every $f \in H^\infty(B_N)$. Thus

$$\text{span}\{\delta_z : z \in B_N\} \subset L^1(S_N)/H_0^1(S_N).$$

The conclusion follows from Lemma 1. □

Theorem 5 permits us to get a sufficient condition for a function on $H^\infty(B_N)$ to attain the norm.

Proposition 6. *If f is an element of $H^\infty(B_N)$ of norm one such that the set*

$$E := \{w \in S_N : |f^*(w)| = 1\},$$

has positive σ_N measure in S_N , then f attains its norm as an element of the dual of $L^1(S_N)/H_0^1(S_N) = G^\infty(B_N)$.

Proof. Define $\varphi : S_N \rightarrow \mathbb{C}$ by

$$\varphi(w) = \begin{cases} \frac{|f^*(w)|}{f^*(w)} \frac{1}{\sigma_N(E)}, & \text{if } w \in E \\ 0, & \text{otherwise.} \end{cases}$$

We have that φ is a bounded measurable function on S_N . Thus $\varphi \in L^1(S_N)$ and

$$\int_{S_N} |\varphi(w)|d\sigma_N(w) = \frac{1}{\sigma_N(E)} \int_E d\sigma_N(w) = 1.$$

Define $T_{[\varphi]} : H^\infty(B_N) \rightarrow \mathbb{C}$ by

$$T_{[\varphi]}(g) := \langle [\varphi], g^* \rangle = \int_{S_N} \varphi(w)g^*(w)d\sigma_N(w).$$

By Theorem 5, $T_{[\varphi]} \in L^1(S_N)/H_0^1(S_N) = G^\infty(B_N)$ and

$$|T_{[\varphi]}(g)| \leq \|g^*\|_\infty \|\varphi\|_1 = \|g\|_\infty \|\varphi\|_1 = \|g\|_\infty,$$

for every $g \in H^\infty(B_N)$. Hence

$$\|T_{[\varphi]}\| \leq 1.$$

But

$$T_{[\varphi]}(f) = \int_{S_N} \varphi(w)f^*(w)d\sigma_N(w) = \frac{1}{\sigma_N(E)} \int_E |f^*(w)|d\sigma_N(w) = 1 = \|f\|.$$

and f in the dual of $G^\infty(B_N)$ attains its norm at $T_{[\varphi]}$. □

A partial converse to the above proposition is the following.

Proposition 7. *If f is an element of $H^\infty(B_N)$ of norm one such that there exists $\varphi \in L^1(S_N)$ with $\|\varphi\|_1 = 1$ and $T_{[\varphi]}(f) = 1$, then*

$$\sigma_N(\{w \in S_N : |f^*(w)| = 1\}) > 0.$$

Proof. We denote $E = \{w \in S_N : |f^*(w)| = 1\}$.

Assume that $\sigma_N(E) = 0$.

Let

$$K_n = \left\{ w \in S_N : |f^*(w)| < \frac{n-1}{n} \right\}.$$

Clearly $S_N \setminus E = \cup_{n=1}^\infty K_n$.

We have that $T_{[\varphi]} \in G^\infty(B_N)$ and is of norm one since

$$1 = T_{[\varphi]}(f) \leq \|[\varphi]\| \|f\|_\infty = \|[\varphi]\| \leq \|\varphi\|_1 = 1.$$

For each n , we get

$$\begin{aligned} \int_{S_N \setminus K_n} |\varphi(w)|d\sigma_N(w) + \int_{K_n} |\varphi(w)|d\sigma_N(w) &= 1 = \int_{S_N} f(w)\varphi(w)d\sigma_N(w) \\ &= \int_{S_N \setminus K_n} f(w)\varphi(w)d\sigma_N(w) + \int_{K_n} f(w)\varphi(w)d\sigma_N(w) \\ &\leq \int_{S_N \setminus K_n} |f(w)\varphi(w)|d\sigma_N(w) + \int_{K_n} |f(w)\varphi(w)|d\sigma_N(w) \\ &\leq \int_{S_N \setminus K_n} |\varphi(w)|d\sigma_N(w) + \frac{n-1}{n} \int_{K_n} |\varphi(w)|d\sigma_N(w). \end{aligned}$$

Thus, $\int_{K_n} |\varphi(w)|d\sigma_N(w) = 0$. Since n is arbitrary, we get

$$\int_{S_N \setminus E} |\varphi(w)|d\sigma_N(w) = 0.$$

But, by hypothesis $\sigma_N(E) = 0$ and finally we arrive at the contradiction

$$1 = \int_{S_N} |\varphi(w)|d\sigma_N(w) = 0.$$

□

A subset E of S_N is called a *peak set* if there exists $f \in A(B_N)$ such that $f(z) = 1$ for every $z \in E$ and $|f(z)| < 1$ for every $z \in \overline{B_N} \setminus E$. Every peak set is a null set.

A result by Fatou states that every compact subset of \mathbb{T} of Lebesgue measure zero is a peak set of $A(\mathbb{D})$, a fact which is instrumental in the proof of Fisher’s Theorem 2. On the other hand, there are null sets on S_N (respectively in \mathbb{T}^N), which are not peak sets [15, 10.1.1 and 11.2.5] (respectively [14, Theorem 6.3.4, p. 149-150]). We do not know if the converse of Proposition 6 is true or not. But, if we restrict ourselves to functions in $A(B_N)$ that attain their norm, we get the following characterization in terms of peak sets.

Theorem 8. *Let f be an element of $A(B_N)$ of norm one. The function f attains its norm as an element of $H^\infty(B_N)$ if and only if the set*

$$E(f) = \{w \in S_N : |f(w)| = 1\}$$

is not a peak set.

Before presenting the proof we need some notation and a lemma.

We recall that a complex Borel measure μ on S_N is a *Henkin measure* (See [15, 9.1.5, p. 186]) if

$$\lim_{n \rightarrow \infty} \int_{S_N} f_n(w) d\mu(w) = 0,$$

for every sequence (f_n) contained in the closed unit ball $U_{A(B_N)}$ of $A(B_N)$ that converges uniformly to 0 on the compact subsets of B_N , that is, converges to 0 in the τ_0 topology in B_N . (By the Montel theorem, a sequence (f_n) contained in $U_{A(B_N)}$ converges to 0 in τ_0 if and only if converges to 0 pointwise on B_N).

Lemma 9. (1) *For every Henkin measure μ there is $T \in G^\infty(B_N)$ such that*

$$T(f) = \int_{S_N} f(w) d\mu(w)$$

for each $f \in A(B_N)$, and $\|\mu\| \geq \|T\|$.

(2) *If $T \in G^\infty(B_N)$, then there is a Henkin measure μ on S_N such that*

$$T(f) = \int_{S_N} f(w) d\mu(w)$$

for each $f \in A(B_N)$, and $\|\mu\| = \|T\|$.

Proof. (1) Define $T_1 : A(B_N) \rightarrow \mathbb{C}$ by

$$T_1(g) := \int_{S_N} g(w) d\mu(w).$$

Clearly, T_1 is a continuous linear form on $A(B_N)$ which is τ_0 -continuous on $U_{A(B_N)}$ and

$$\|T_1\| \leq \|\mu\|.$$

Given $f \in H^\infty(B_N)$, the function $f_r(z) := f(rz)$, $0 \leq r < 1$, belongs to $A(B_N)$. In addition, (f_r) converges to f uniformly on the compact subsets of B_N and

$$\|f_r\| \leq \|f\|, \quad \|f\| = \sup_r \|f_r\|. \tag{1}$$

By [15, 11.3.1], since μ is a Henkin measure, the limit

$$\lim_{r \rightarrow 1^-} \int_{S_N} f_r(w) d\mu(w) = \lim_{r \rightarrow 1^-} T_1(f_r) \in \mathbb{C},$$

exists for every $f \in H^\infty(B_N)$.

We define $T : H^\infty(B_N) \rightarrow \mathbb{C}$, by

$$T(f) := \lim_{r \rightarrow 1^-} T_1(f_r).$$

T is linear and $T \in (H^\infty(B_N))^*$, since

$$|T(f)| \leq \sup_r |T_1(f_r)| \leq \|T_1\| \|f\|,$$

for every $f \in H^\infty(B_N)$. Moreover, $\|T\| = \|T_1\|$.

We claim that the restriction of T_1 to $U_{A(B_N)}$ is τ_0 -uniformly continuous. Indeed, given $\varepsilon > 0$ there are a compact subset K of B_N and $\delta > 0$ such that $|T_1(g)| < \varepsilon$ if $g \in U_{A(B_N)}$ and $\sup_{z \in K} |g(z)| < \delta$. Hence, if $g, h \in U_{A(B_N)}$ and $\sup_{z \in K} |g(z) - h(z)| < \delta$, then

$$|T_1(g) - T_1(h)| = |2T_1\left(\frac{g-h}{2}\right)| < 2\varepsilon.$$

Since $U_{A(B_N)}$ is τ_0 -dense in $U_{H^\infty(B_N)}$, there exists a unique $\widetilde{T}_1 : U_{H^\infty(B_N)} \rightarrow \mathbb{C}$ that is τ_0 -continuous and such that

$$\widetilde{T}_1(g) = T_1(g),$$

for all $g \in U_{A(B_N)}$. Given $f \in U_{H^\infty(B_N)}$, then $(f_r) \subset U_{A(B_N)}$, and (f_r) converges to f in τ_0 as $r \rightarrow 1^-$. Thus

$$\widetilde{T}_1(f_r) \rightarrow \widetilde{T}_1(f),$$

in \mathbb{C} . But $\widetilde{T}_1(f_r) = T_1(f_r)$ for each $r \in [0, 1[$ and $T_1(f_r)$ converges to $T(f)$ by definition. This implies $\widetilde{T}_1(f) = T(f)$ for each $f \in U_{H^\infty(B_N)}$. We have obtained that the restriction of T to $U_{H^\infty(B_N)}$ is τ_0 continuous. This, by definition, implies that T belongs to $G^\infty(B_N)$.

- (2) If $T \in G^\infty(B_N)$, the restriction of T to $U_{H^\infty(B_N)}$ is τ_0 continuous. If we define $T_1 : A(B_N) \rightarrow \mathbb{C}$, by $T_1(g) := T(g)$, then T_1 is continuous for the sup-norm, $\|T_1\| \leq \|T\|$ and T_1 is τ_0 -continuous on $U_{A(B_N)}$. By (1), we have

$$|T(f)| = \lim_{r \rightarrow 1^-} |T(f_r)| = \lim_{r \rightarrow 1^-} |T_1(f_r)| \leq \|T_1\| \sup_{r < 1} \|f_r\| = \|T_1\| \|f\|,$$

for every $f \in H^\infty(B_N)$. Thus, $\|T_1\| = \|T\|$. We can consider $T_1 : A(S_N) \rightarrow \mathbb{C}$. By the Riesz theorem, there is a complex Borel measure μ on S_N such that

$$T_1(h) := \int_{S_N} h(w)d\mu(w),$$

for every $h \in A(B_N)$ with $\|T_1\| = |\mu|(S_N) = \|\mu\|$.

The properties of T_1 imply that μ is a Henkin measure. □

Now we give the proof of Theorem 8:

Proof. Assume that $f \in A(B_N)$, $\|f\| = 1$ and that $E(f)$ is not a peak set. By [15, 10.1.1] there exists a Borel measure ρ , such that $\rho(E(f)) \neq 0$ such that

$$h(0) = \int_{S_N} h(w)d\rho(w), \tag{2}$$

for every $h \in A(B_N)$.

Define

$$g(z) = \begin{cases} 0 & \text{if } z \in S_N \setminus E(f) \\ \frac{f(z)}{|\rho|(E(f))} & \text{if } z \in E(f) \end{cases} .$$

Since g is bounded and measurable, $g \in L^1(|\rho|)$. Hence, the measure $g|\rho|$ defined by $g|\rho|(M) = \int_M g d|\rho|$ for Borel measurable sets M is absolutely continuous with respect to ρ . The measure ρ is Henkin (this fact is a direct consequence of (2) and the definition of Henkin measure as given in [15, p. 187]), and so $g|\rho|$ is also a Henkin measure by [15, 9.3.1]. We set $T_1 : A(B_N) \rightarrow \mathbb{C}$.

$$T_1(h) = \int_{S_N} h(w)g(w)d|\rho|(w).$$

We have

$$T_1(f) = \int_{S_N} f(w)g(w)d|\rho|(w) = 1,$$

and

$$|T_1(h)| \leq \int_{S_N} |h(w)||g(w)d|\rho|(w) \leq \frac{|\rho|(E(f))}{|\rho|(E(f))} = 1,$$

for every $h \in A(B_N)$ with $\|h\| \leq 1$, and we have $g|\rho|$ a Henkin measure such that

$$T_1(f) = 1 \quad \text{and} \quad \|T_1\| = 1.$$

By Lemma 9.(1), there is $T \in G^\infty(B_N)$ with $\|T\| = \|T_1\| = 1$ such that

$$T(h) = T_1(h),$$

for every $h \in A(B_N)$.

In particular, $T(f) = 1$ and f attains its norm on $G^\infty(B_N)$.

Suppose now that $f \in A(B_N)$, $\|f\| = 1$, satisfies that $E(f)$ is a peak set and that there is $T \in G^\infty(B_N)$, $\|T\| = 1$ with $T(f) = 1$. To get a contradiction we are going to give an argument that follows closely the one given in Proposition 7:

By Lemma 9.(2) there exists μ a Henkin measure such that

$$T(h) = \int_{S_N} h(w)d\mu(w),$$

for every $h \in A(B_N)$ and

$$\|T\| = \|T|_{A(B_N)}\| = |\mu|(S_N) = 1.$$

By [15, 9.3.1] $|\mu|$ is also a Henkin measure. Hence, by [15, Lemma 11.3.3] (see also [15, Lemma 11.3.1]), $|\mu|(E(f)) = 0$. Let

$$K_n = \left\{ w \in S_N : |f(w)| < \frac{n-1}{n} \right\}.$$

Clearly $S_N \setminus E(f) = \cup_{n=1}^\infty K_n$.

We have that, for each n ,

$$\begin{aligned} |\mu|(S_N \setminus K_n) + |\mu|(K_n) &= |\mu|(S_N) = 1 = T(f) = \int_{S_N} |f(w)|d|\mu|(w) \\ &= \int_{S_N \setminus K_n} |f(w)|d|\mu|(w) + \int_{K_n} |f(w)|d|\mu|(w) \\ &\leq \int_{S_N \setminus K_n} d|\mu|(w) + \frac{n-1}{n} \int_{K_n} d|\mu|(w) \\ &= |\mu|(S_N \setminus K_n) + \frac{n-1}{n}|\mu|(K_n). \end{aligned}$$

This implies $|\mu|(K_n) = 0$ for each n and $|\mu|(S_N \setminus E(f)) = 0$.

Therefore, $1 = |\mu|(S_N) = |\mu|(S_N \setminus E(f)) + |\mu|(E(f)) = 0$, a contradiction. □

3. The Case of the Polydisc

For a fixed $N \in \mathbb{N}$, the N -dimensional Poisson kernel [14, p. 17] $P_N : \mathbb{D}^N \times \mathbb{T}^N \rightarrow (0, \infty)$ is defined as

$$P_N(z, w) := \prod_{j=1}^N P_1(z_j, w_j) = \prod_{j=1}^N \frac{1 - |z_j|^2}{|1 - z_j \bar{w}_j|^2}.$$

It is well known ([14, Theorem 3.3.3, p.45]) that if $f \in H^\infty(\mathbb{D}^N)$ then the limit

$$f^*(w) = \lim_{r \rightarrow 1^-} f(rw)$$

exists almost everywhere in \mathbb{T}^N , and

$$f(z) = \int_{\mathbb{T}^N} P_N(z, w) f^*(w) dm_N(w) \tag{3}$$

for all $z \in \mathbb{D}^N$. As a consequence there exists an isometric isomorphism

$$\begin{aligned} H^\infty(\mathbb{D}^N) &\longrightarrow H^\infty(\mathbb{T}^N) \\ f &\longrightarrow f^* \end{aligned}$$

where $H^\infty(\mathbb{T}^N) := \overline{A(\mathbb{T}^N)}^{w(L^\infty(\mathbb{T}^N), L^1(\mathbb{T}^N))}$,

$$A(\mathbb{D}^N) = \{f : \overline{\mathbb{D}}^N \rightarrow \mathbb{C} : f \text{ is continuous on } \overline{\mathbb{D}}^N \text{ and holomorphic on } \mathbb{D}^N\}$$

and

$$A(\mathbb{T}^N) := \{f|_{\mathbb{T}^N} : f \in A(\mathbb{D}^N)\}.$$

By the maximum modulus theorem $A(\mathbb{D}^N)$ and $A(\mathbb{T}^N)$ are isometrically isomorphic. By Fejer’s theory for the polydisc we have

$$\begin{aligned} H^\infty(\mathbb{T}^N) &:= \left\{ g \in L^\infty(\mathbb{T}) : \hat{g}(\alpha) \right. \\ &= \left. \int_{\mathbb{T}^N} w^{-\alpha} g(w) dm_N(w) = 0, \text{ for all } \alpha \in \mathbb{Z}^N \setminus \mathbb{N}_0^N \right\}. \end{aligned}$$

On the other hand, by applying Lemma 4,

$$H^\infty(\mathbb{T}^N) \stackrel{1}{=} (L^1(\mathbb{T}^N)/H_0^1(\mathbb{T}^N))^*,$$

where

$$\begin{aligned} H_0^1(\mathbb{T}^N) &= \left\{ f \in L^1(\mathbb{T}^N) : \hat{f}(-\beta) = \int_{\mathbb{T}} f(w) w^\beta dm_N(w) = 0, \text{ for all } \beta \in \mathbb{N}_0^N \right\} \\ &= \overline{\text{span}\{w^{-\alpha} : \text{for all } \alpha \in \mathbb{Z}^N \setminus \mathbb{N}_0^N\}}. \end{aligned}$$

Very similar arguments to the ones given for the N -dimensional Euclidean ball can be given for the N -polydisc to obtain the following results.

Theorem 10. *For every $N \in \mathbb{N}$ we have*

$$L^1(\mathbb{T}^N)/H_0^1(\mathbb{T}^N) \stackrel{1}{=} G^\infty(\mathbb{D}^N).$$

Proposition 11. *Let f be an element of $H^\infty(\mathbb{D}^N)$ of norm one such that the set*

$$E := \{w \in \mathbb{T}^N : |f^*(w)| = 1\},$$

has positive Lebesgue measure (in \mathbb{T}^N). Then f attains its norm as an element of the dual of $L^1(\mathbb{T}^N)/H_0^1(\mathbb{T}^N)$.

Proposition 12. *If f is an element of $H^\infty(\mathbb{D}^N)$ of norm one such that there exists $\varphi \in L^1(\mathbb{T}^N)$ with $\|\varphi\|_1 = 1$ and $T_{[\varphi]}(f) = 1$, then the set*

$$\{w \in \mathbb{T}^N : |f^*(w)| = 1\},$$

has positive Lebesgue measure in the polytorus \mathbb{T}^N .

Example 13. The following example, which is inspired by [3, Theorem 3.1], shows that a polydisc (for $N > 1$) version of Theorem 8 does not hold. Let $f : \mathbb{D} \times \mathbb{D} \rightarrow \mathbb{C}$ be the function $f(z, w) := (1/2)(1 + z)$, which belongs to $A(\mathbb{D} \times \mathbb{D})$. This function does not attain its norm on $G^\infty(\mathbb{D} \times \mathbb{D})$. Indeed, if it did, the function $g(z) = (1/2)(1 + z)$, as an element of $H^\infty(\mathbb{D})$, would attain its norm on $G^\infty(\mathbb{D})$, because $H^\infty(\mathbb{D})$ is canonically isometrically contained in $H^\infty(\mathbb{D} \times \mathbb{D})$. But the function g does not attain its norm on $G^\infty(\mathbb{D})$ by Fisher’s Theorem 2, because $\{z \in \mathbb{T} \mid |g(z)| = 1\} = \{1\}$. On the other hand, $E(f) = \{(z, w) \in \mathbb{T} \times \mathbb{T} \mid |f(z, w)| = 1\} = \{1\} \times \mathbb{T}$, as it is easy to check. The set $E(f)$ is not a peak set of $A(\mathbb{D} \times \mathbb{D})$. Otherwise, it would be a zero set; see [14, 6.1.2. Theorem, p.132]. But if a function $h \in A(\mathbb{D} \times \mathbb{D})$ vanishes on $E(f)$, then $h(1, w) \in A(\mathbb{D})$ vanishes on \mathbb{T} . The maximum principle then implies that h vanishes on $\{1\} \times \mathbb{D}$, and therefore, there is no function $h \in A(\mathbb{D} \times \mathbb{D})$ vanishing only in $E(f)$. Observe that $E(f)$ is a null set in $\mathbb{T} \times \mathbb{T}$ which is not a peak set.

4. The Case of the Space of Dirichlet Series $\mathcal{D}^\infty(\mathbb{C}_+)$

Let $\mathcal{D}^\infty(\mathbb{C}_+)$ denote the Banach space of the Dirichlet series $D(s) = \sum_{n=1}^\infty \frac{a_n}{n^s}$ convergent and bounded on the right half plane \mathbb{C}_+ endowed with the supremum norm. We refer the reader to [4] and [13] for detailed information about this space.

The space $\mathcal{D}^\infty(\mathbb{C}_+)$ is a closed subspace of the Banach space $H^\infty(\mathbb{C}_+)$ of all bounded holomorphic functions in the right half plane \mathbb{C}_+ endowed with the supremum norm. Since, by the Montel theorem, the closed unit ball of $H^\infty(\mathbb{C}_+)$ is τ_0 -compact, we can apply the Dixmier-Ng theorem [12] to obtain that

$$G^\infty(\mathbb{C}_+) := \{R \in H^\infty(\mathbb{C}_+)^* : \text{the restriction of } R \text{ to } U_{H^\infty(\mathbb{C}_+)} \text{ is } \tau_0 \text{ continuous}\},$$

is a predual of $H^\infty(\mathbb{C}_+)$.

It is well-known that the spaces $H^\infty(\mathbb{C}_+)$ and $H^\infty(\mathbb{D})$ are isometrically isomorphic. We are going to show that their preduals are also isometrically isomorphic.

Proposition 14. *$H^\infty(\mathbb{C}_+)$ is isometrically isomorphic to $H^\infty(\mathbb{D})$, and $G^\infty(\mathbb{C}_+)$ is isometrically isomorphic to $G^\infty(\mathbb{D})$.*

Proof. It is enough to consider the Cayley transformation $\varphi : \mathbb{C} \setminus \{1\} \rightarrow \mathbb{C} \setminus \{-1\}$ defined by

$$\varphi(z) = \frac{1+z}{1-z}$$

The Cayley transformation is a biholomorphic mapping with inverse

$$\varphi^{-1}(s) = \frac{s-1}{s+1}.$$

Actually it is also biholomorphic from \mathbb{D} onto \mathbb{C}_+ , and it is a homeomorphism from $\mathbb{T} \setminus \{1\}$ onto $\{ti : t \in \mathbb{R}\}$. Clearly the composition operator $T_\varphi : H^\infty(\mathbb{C}_+) \rightarrow H^\infty(\mathbb{D})$ defined by

$$T_\varphi(g) = g \circ \varphi,$$

for $g \in H^\infty(\mathbb{C}_+)$ is an isometry with inverse $(T_\varphi)^{-1} = T_{\varphi^{-1}}$. Its adjoint $T_\varphi^* : H^\infty(\mathbb{D})^* \rightarrow H^\infty(\mathbb{C}_+)^*$ is also an isometric isomorphism with

$$(T_\varphi^*)^{-1} = T_{\varphi^{-1}}^*.$$

It is enough to check that $T_\varphi^*(G^\infty(\mathbb{D})) = G^\infty(\mathbb{C}_+)$ to prove that $G^\infty(\mathbb{D})$ and $G^\infty(\mathbb{C}_+)$ are isometrically isomorphic.

Let $R \in G^\infty(\mathbb{D})$. We have

$$T_\varphi^*(R)(g) = R(T_\varphi(g)) = R(g \circ \varphi)$$

for all $g \in H^\infty(\mathbb{C}_+)$. Let $K \subset \mathbb{D}$ be a compact set. The set $\varphi(K)$ is a compact subset of \mathbb{C}_+ . Take (g_n) and g in the closed unit ball of $H^\infty(\mathbb{C}_+)$ such that (g_n) converges with respect to the compact open topology on \mathbb{C}_+ to g . Since (g_n) converges to g uniformly on $\varphi(K)$, we have $(g_n \circ \varphi)$ converges to $g \circ \varphi$ uniformly on $K = \varphi^{-1}(\varphi(K))$, for every K . Thus, $(g_n \circ \varphi)$ converges to $g \circ \varphi$ with respect to the compact open topology on \mathbb{D} . Hence, $(R(g_n \circ \varphi))$ converges to $R(g \circ \varphi)$ and we get

$$T_\varphi^*(R) \in G^\infty(\mathbb{C}_+).$$

Analogously we obtain $T_{\varphi^{-1}}^*(G^\infty(\mathbb{C}_+)) \subset G^\infty(\mathbb{D})$, from which it follows that

$$G^\infty(\mathbb{C}_+) = T_\varphi^* \circ T_{\varphi^{-1}}^*(G^\infty(\mathbb{C}_+)) \subset T_\varphi^*(G^\infty(\mathbb{D})).$$

□

Remark 15. Recall that for any fixed $\alpha > 1$ and $w \in \mathbb{T}$ the Stolz region is $S(\alpha, w) = \{z \in \mathbb{D} : |z - w| < \alpha(1 - |z|)\}$ ([9, Definition 8.1.9.]). Since w is an accumulation point of $S(\alpha, w)$ it makes sense to speak about the limit at w of any function $f : S(\alpha, w) \rightarrow \mathbb{C}$. Actually, in [9, Theorem 8.1.11], it is proved that if $f \in H^\infty(\mathbb{D})$ the following equality holds on \mathbb{T}

$$f^*(w) = \lim_{z \in S(\alpha, w) \rightarrow w} f(z),$$

almost everywhere with respect to the Lebesgue measure.

In [4, p. 286 and 287] it is observed that if $g \in H^\infty(\mathbb{C}_+)$, then there exists a Lebesgue null set $A \subset \mathbb{R}$ such that the limit

$$\lim_{r \rightarrow 0^+} g(r + it) := g^*(it)$$

exists for every $t \in \mathbb{R} \setminus A$ and actually that

$$g^*(it) = \lim_{z \in S(\alpha, \varphi^{-1}(it)) \rightarrow \varphi^{-1}(it)} T_\varphi(g)(z).$$

In other words, the “horizontal” limits of g exist a.e. and coincide with the Fatou radial limits of its associated function $T_\varphi(g)$ belonging to $H^\infty(\mathbb{D})$.

We can now get the following consequence of Ando’s Theorem [1] and Fisher’s Theorem 2.

Corollary 16. *The space $H^\infty(\mathbb{C}_+)$ has a unique predual. Moreover, $g \in H^\infty(\mathbb{C}_+)$ with $\|g\|_{\mathbb{C}_+} = 1$ is norm attaining if and only if the set*

$$E := \{t \in \mathbb{R} : |g^*(it)| = 1\}$$

has positive (including $+\infty$) Lebesgue measure.

Proposition 17. *$\mathcal{D}^\infty(\mathbb{C}_+)$ is a dual space.*

Proof. By a result of F. Bayart (see e.g. [4, Theorem 3.11]), it is known that if (D_n) is a bounded sequence in $\mathcal{D}^\infty(\mathbb{C}_+)$ then there exists a subsequence (D_{n_k}) and a Dirichlet series $D \in \mathcal{D}^\infty(\mathbb{C}_+)$ such that for every $\sigma > 0$ the sequence (D_{n_k}) converges to D uniformly on $\mathbb{C}_\sigma := \{s \in \mathbb{C} ; \text{Res} \geq \sigma\}$. Thus, if we denote by τ_+ the topology of uniform convergence on these half planes \mathbb{C}_σ , Bayart’s result says that the closed unit ball of $\mathcal{D}^\infty(\mathbb{C}_+)$ is a compact set. Now the Dixmier-Ng theorem [12] implies that

$$\mathcal{G}^\infty(\mathbb{C}_+) := \{R \in \mathcal{D}^\infty(\mathbb{C}_+)^* : \text{the restriction of } R \text{ to } U_{\mathcal{D}^\infty(\mathbb{C}_+)} \text{ is } \tau_+ \text{ continuous}\}, \tag{4}$$

endowed with the topology induced by $\mathcal{D}^\infty(\mathbb{C}_+)^*$ is a predual of $\mathcal{D}^\infty(\mathbb{C}_+)$. □

We can now get a positive result about norm attaining elements of $\mathcal{D}^\infty(\mathbb{C}_+)$ with respect to that predual.

Proposition 18. *Consider the space $\mathcal{D}^\infty(\mathbb{C}_+)$ as the dual of $\mathcal{G}^\infty(\mathbb{C}_+)$. Given $D \in \mathcal{D}^\infty(\mathbb{C}_+)$ of norm one, if the set*

$$E := \{t \in \mathbb{R} : |D^*(it)| = 1\}$$

has positive (including $+\infty$) Lebesgue measure, then D is norm attaining.

Proof. As $\mathcal{D}^\infty(\mathbb{C}_+)$ is a closed subspace of $H^\infty(\mathbb{C}_+)$, we can consider $D \in H^\infty(\mathbb{C}_+)$. By Corollary 16, we know that there exists $R \in \mathcal{G}^\infty(\mathbb{C}_+)$ such that

$$\|R\| = 1 = R(D).$$

Recall that $R \in H^\infty(\mathbb{C}_+)^*$ and satisfies that the restriction of R to $U_{H^\infty(\mathbb{C}_+)}$ is τ_0 continuous. We denote by S the restriction of R to $\mathcal{D}^\infty(\mathbb{C}_+)$. Since $U_{\mathcal{D}^\infty(\mathbb{C}_+)} \subset U_{H^\infty(\mathbb{C}_+)}$ we have that S is τ_0 continuous when restricted to $U_{\mathcal{D}^\infty(\mathbb{C}_+)}$. The theorem of Bayart [4, Theorem 3.11] implies that $U_{\mathcal{D}^\infty(\mathbb{C}_+)}$ is a compact set with respect to τ_+ . The compact open topology τ_0 on \mathbb{C}_+ is Hausdorff and weaker than τ_+ on that ball. Hence both topologies coincide on $U_{\mathcal{D}^\infty(\mathbb{C}_+)}$ and $S \in \mathcal{G}^\infty(\mathbb{C}_+)$. Moreover

$$1 = \|R\| \geq \|S\| \geq |S(D)| = S(D) = R(D) = 1,$$

and D attains its norm. □

It is natural to ask whether the converse of Proposition 18 holds. Actually, by the Hahn-Banach theorem one can extend R in $\mathcal{G}^\infty(\mathbb{C}_+)$ to an element T belonging to $H^\infty(\mathbb{C}_+)^*$ with the same norm. But we don't know if it is possible to choose an extension T in $G^\infty(\mathbb{C}_+)$.

Acknowledgements

The authors are very grateful to the referee for the careful reading of our manuscript and for the suggestions which improved our article. The research of R. Aron was partially supported by the project PID2021-122126NB-C33/MCIN/AEI/ 10.13039/ 501100011033 (FEDER). The research of J. Bonet was partially supported by the project PID2020-119457GB-100 funded by MCIN/AEI/ 10.13039/501100011033 and by “ERFD A way of making Europe” and by the project GV AICO/2021/170. The research of M. Maestre was partially supported by the project PID2021-122126NB-C33/MCIN/AEI/10.13039/501100011033 (FEDER) and the project GV PROMETEU/2021/070.

Author contributions All authors contributed equally to the study conception and design. All authors read and approved the final manuscript.

Funding Open Access funding provided thanks to the CRUE-CSIC agreement with Springer Nature. The research of R. Aron was partially supported by the project PID2021-122126NB-C33/MCIN/AEI/ 10.13039/ 501100011033 and by “ERFD A way of making Europe”. The research of J. Bonet was partially supported by the project PID2020-119457GB-100 funded by MCIN/AEI/10.13039/501100011033 and by “ERFD A way of making Europe” and by the project GV AICO/2021/170. The research of M. Maestre was partially supported by the project PID2021-122126NB-C33/MCIN/AEI/ 10.13039/ 501100011033 and by “ERFD A way of making Europe” and the project GV PROMETEU/2021/070.

Declarations

Conflict of interest The authors have no relevant financial or non-financial interests to disclose.

Open Access. This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- [1] Ando, T.: On the predual of H^∞ . *Commentationes Mathematicae: Tomus Specialis in Honorem Ladislai Orlicz* **1**, 33–40 (1978)
- [2] Bollobás, B.: *Linear analysis*, 2nd edn. An introductory course, Cambridge Mathematical Textbooks, Cambridge University Press (1999)
- [3] Boos, L.J.: Totally null sets for $A(X)$. *Bull. Aust. Math. Soc.* **87**, 108–114 (2013)
- [4] Defant, A., García, D., Maestre, M., Sevilla-Peris, P.: *Dirichlet Series and Holomorphic Functions in High Dimensions*, New Mathematical Monographs, vol. 37, Cambridge University Press (2019)
- [5] Fatou, P.: Séries trigonométriques et séries de Taylor. *Acta Math.* **30**, 335–400 (1906)
- [6] Fisher, S.: Exposed points in spaces of bounded analytic functions. *Duke Math. J.* **36**, 479–484 (1969)
- [7] Garnett, J.B.: *Bounded Analytic Functions*, Graduate Texts in Mathematics, vol. 236, Springer (2007)
- [8] Koosis, P.: *Introduction to H^p spaces*, 2nd edition, Cambridge Tracts in Mathematics 115. Cambridge University Press, Cambridge (1998)
- [9] Krantz, S.G.: *Function theory of several complex variables*, 2nd edn. AMS Chelsea Publishing, American Mathematical Society, Providence (2001)
- [10] Megginson, R.E.: *An Introduction to Banach Space Theory*, Graduate Texts in Mathematics, vol. 183. Springer, New York (1998)
- [11] Mujica, J.: Linearization of bounded holomorphic mappings on Banach spaces. *Trans. Am. Math. Soc.* **324–2**, 867–887 (1991)
- [12] Ng, K.-F.: On a theorem of Dixmier. *Math. Scand.* **29**, 279–280 (1971)
- [13] Queffelec, H., Queffelec, M.: *Diophantine Approximation and Dirichlet Series*. Springer, Singapore (2020)
- [14] Rudin, W.: *Function Theory in Polydiscs*, Mathematics Lecture Note Series. W. A. Benjamin Inc, New York-Amsterdam (1969)
- [15] Rudin, W.: *Function Theory in the Unit ball of \mathbb{C}^n* , Springer, reprint of the 1980 edition (2008)

- [16] Zhu, K.: Spaces of Holomorphic Functions in the Unit Ball, Graduate Texts in Mathematics, vol. 226. Springer, New York (2006)

Richard M. Aron
Department of Mathematical Sciences
Kent State University
Kent OH 44242
USA
e-mail: aron@math.kent.edu

José Bonet
Instituto Universitario de Matemática Pura y Aplicada IUMPA
Universitat Politècnica de València
46071 Valencia
Spain
e-mail: jbonet@mat.upv.es

Manuel Maestre
Departamento de Análisis Matemático
Universidad de Valencia
Doctor Moliner 50
46100 Burjassot, Valencia
Spain
e-mail: manuel.maestre@uv.es

Received: June 14, 2023.

Accepted: December 19, 2023.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.