



# Correction to: Experimental desiccation indicates high moisture content maintains hyporheic biofilm processes during drought in temperate intermittent streams

Laura E. Coulson<sup>1,2</sup> · Jakob Schelker<sup>1,3</sup> · Katrin Attermeyer<sup>1,3</sup> · Christian Griebler<sup>3</sup> · Thomas Hein<sup>1,2</sup> · Gabriele Weigelhofer<sup>1,2</sup>

Published online: 30 June 2021  
© Springer Nature Switzerland AG 2021

**Correction to:** *Aquatic Sciences* (2021) 83:46  
<https://doi.org/10.1007/s00027-021-00799-3>

In the original publication of the article, on page No 3, The colonization flow rate was reported as 0.9 mL per hour but the actual flow rate was 62 mL per second.

This has been corrected in this article.

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

---

The original article can be found online at <https://doi.org/10.1007/s00027-021-00799-3>.

---

✉ Laura E. Coulson  
laura.coulson@wcl.ac.at

- <sup>1</sup> WasserCluster Lunz–Biologische Station, Dr. Carl Kupelwieser Promenade 5, 3293 Lunz am See, Austria
- <sup>2</sup> Institute of Hydrobiology and Aquatic Ecosystem Management, University of Natural Resources and Life Sciences (BOKU), Gregor-Mendel-Str. 33/DG, 1180 Vienna, Austria
- <sup>3</sup> Division of Limnology, Department of Functional and Evolutionary Ecology, University of Vienna, Althanstr. 14, 1090 Vienna, Austria