



Toric G -solid Fano threefolds

Ivan Cheltsov¹ · Adrien Dubouloz² · Takashi Kishimoto³

Accepted: 27 September 2022 / Published online: 2 February 2023
© The Author(s) 2023

Abstract

We study toric G -solid Fano threefolds that have at most terminal singularities, where G is an algebraic subgroup of the normalizer of a maximal torus in their automorphism groups. All varieties are assumed to be projective and defined over the field of complex numbers.

Mathematics Subject Classification 14J50 · 14M25 · 14N05

1 Introduction

Fano varieties with many symmetries appear naturally in several geometric problems. A special role among them is played by the so-called G -Fano varieties [27], which naturally occur as the end product of the equivariant Minimal Model Program for rationally connected varieties. Recall from [27] that a G -Fano variety is a pair (X, G) consisting of a Fano variety X and an algebraic subgroup G in $\text{Aut}(X)$ such that

- (1) The singularities of X are terminal (mild);
- (2) The G -invariant part $\text{Cl}(X)^G$ of the class group of X has rank 1 (G -minimal).

In dimension two, we know the complete list of G -Fano varieties [18], which are traditionally called G -del Pezzo surfaces. In [26, 27], Prokhorov obtained many deep results about G -Fano threefolds for G finite. A complete classification is still lacking.

✉ Ivan Cheltsov
I.Cheltsov@ed.ac.uk

Adrien Dubouloz
adrien.dubouloz@u-bourgogne.fr

Takashi Kishimoto
kisimoto.takasi@gmail.com

¹ University of Edinburgh, Edinburgh, Scotland

² Université de Bourgogne, Dijon, France

³ Saitama University, Saitama, Japan

In higher dimensions, our knowledge of G -Fano varieties is limited to some sporadic examples.

Since by definition a G -Fano variety X is a G -Mori fibre space (see [10, Definition 1.1.5]), to describe its G -equivariant birational geometry, it is enough to classify all G -birational maps from X to other G -Mori fibre spaces. By [12, 20], each such birational map can be decomposed into a sequence of elementary links, which are known as G -Sarkisov links. Then, following [3, 10, 11, 13], we say that a G -Fano variety X is:

- G -birationally super-rigid if no G -Sarkisov link starts at X ;
- G -birationally rigid if every G -Sarkisov link that starts at X also ends at X ;
- G -solid if X is not G -birational to a G -Mori fibre space with positive dimensional base.

If X is G -solid, then all G -Mori fibre spaces that are G -birational to X are terminal Fano threefolds — they form a set $\mathcal{P}_G(X)$, which we call the G -pliability of X [14]. For instance, if X is G -solid, then $\mathcal{P}_G(X) = \{X\}$ if and only if X is G -birationally rigid.

In this paper, we consider toric G -Fano varieties in the case where G is an algebraic subgroup in $\text{Aut}(X)$ that normalizes a maximal torus $\mathbb{T} \cong \mathbb{G}_m^n$, where $n = \dim(X)$. In this case, letting G_X be the normalizer of the torus \mathbb{T} in $\text{Aut}(X)$, we have a split exact sequence of groups

$$1 \longrightarrow \mathbb{T} \longrightarrow G_X \xrightarrow{\nu_X} \mathbb{W}_X \longrightarrow 1,$$

where \mathbb{W}_X is a finite subgroup of $\text{GL}_n(\mathbb{Z})$, known as the Weyl group. It is a natural problem to determine for such groups G which G -Fano varieties are G -solid, and to characterize which of these are G -birationally rigid or super-rigid. For surfaces, it is easy to give a satisfactory answer to these questions.

Exercise 1.1 (cf. [18, 21, 30, 33]) *Let X be a smooth toric del Pezzo surface and let G be a subgroup of G_X . If X is G -minimal and G -solid, then one of the following cases holds:*

- (i) $X = \mathbb{P}^2$, $\mathbb{W}_X \cong \mathfrak{S}_3$ and $\nu_X(G)$ contains the subgroup isomorphic to μ_3 ;
- (ii) $X = \mathbb{P}^1 \times \mathbb{P}^1$, $\mathbb{W}_X \cong \text{D}_8$ and $\nu_X(G)$ contains the subgroup isomorphic to μ_4 ;
- (iii) X is the del Pezzo surface of degree 6, $\mathbb{W}_X \cong \mathfrak{S}_3 \times \mu_2$, and either $\nu_X(G)$ contains the subgroup isomorphic to μ_6 or it contains the subgroup isomorphic to \mathfrak{S}_3 that acts transitively on the (-1) -curves in X .

In each of these three cases, X is G -minimal and G -birationally rigid provided that $|G| \geq 72$.

In this paper, we obtain a similar answer for three-dimensional toric Fano varieties. To state it, let $V_6 = \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$, let V_4 be the toric complete intersection in \mathbb{P}^5 given by

$$xu - yw = xu - zt = 0,$$

let Y_{24} be the toric divisor of degree $(1, 1, 1, 1)$ in $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ given by

$$x_1x_2x_3x_4 - y_1y_2y_3y_4 = 0,$$

and let X_{24} be the toric Fano threefold No 47 in [7] (see Sect. 4.3 for its construction). The Weyl groups \mathbb{W}_X of these toric Fano threefolds are all isomorphic to the group $\mathfrak{S}_4 \times \mu_2$, and we have the following result:

Theorem 1.2 *Let X be a toric Fano threefold that have at most terminal singularities, let \mathbb{T} be a maximal torus in $\text{Aut}(X)$ and let G_X be its normalizer in $\text{Aut}(X)$. Then the following two conditions are equivalent:*


- (i) X is G_X -minimal and G_X -solid;
- (ii) X is one of the threefolds V_6, V_4, X_{24}, Y_{24} and \mathbb{P}^3 .

Let G be an algebraic subgroup in G_X and let $\nu_X: G_X \rightarrow \mathbb{W}_X = G_X/\mathbb{T}$ be the quotient homomorphism. If X is one of the toric Fano threefolds V_6, V_4, X_{24}, Y_{24} and \mathbb{P}^3 , then the following assertions hold:

- (1) if X is G -minimal and G -solid, then $\nu_X(G)$ contains a subgroup isomorphic to \mathfrak{A}_4 ;
- (2) if $\nu_X(G)$ contains a subgroup isomorphic to \mathfrak{A}_4 , then X is G -minimal unless
 - (a) $X = V_4, \nu_X(G) \cong \mathfrak{S}_4$ and G acts intransitively on \mathbb{T} -invariant surfaces;
 - (b) $X = V_4$ and $\nu_X(G) \cong \mathfrak{A}_4$.
- (3) if X is G -minimal, $\nu_X(G)$ contains a subgroup isomorphic to \mathfrak{A}_4 and $|G| \geq 32 \cdot 24^4$, then X is G -solid.

□

If $X = V_4$ and $\nu_X(G)$ contains a subgroup isomorphic to \mathfrak{A}_4 , then X is not G -minimal if and only if there exists the following G -commutative:



where β is the blow-up of the four \mathbb{T} -invariant points, α is the contraction of the proper transforms of the six \mathbb{T} -invariant lines, and the dashed arrow is the birational map that is given by the linear system of quadric surfaces that pass through the four \mathbb{T} -invariant points.

Remark 1.3 If X is one of the toric Fano threefolds V_4, X_{24} or Y_{24} , then $G_X = \text{Aut}(X)$.

If X is one of the toric Fano threefolds $V_6, V_4, X_{24}, Y_{24}, \mathbb{P}^3$, and G is an algebraic subgroup in $\text{Aut}(X)$ such that the threefold X is G -minimal, $\nu_X(G)$ contains a subgroup isomorphic to \mathfrak{A}_4 , and $|G| \geq 32 \cdot 24^4$, then the threefold X is G -solid by Theorem 1.2. In this case, we describe all (possible) G -birational maps between these Fano threefolds. We summarize this description in the table presented in Appendix A. It gives

Corollary 1.4 (cf. [4, 11]) *Let X be one of the toric Fano threefolds V_6 , V_4 , X_{24} , Y_{24} , \mathbb{P}^3 , let \mathbb{T} be a maximal torus in $\text{Aut}(X)$ and let G_X be its normalizer in $\text{Aut}(X)$. Then the following three conditions are equivalent:*

- (i) X is G_X -minimal and G_X -birationally rigid;
- (ii) X is G_X -minimal and G_X -birationally super-rigid;
- (iii) X is isomorphic to either V_6 or Y_{24} .

Let G be an algebraic subgroup in G_X and let $\nu_X: G_X \rightarrow \mathbb{W}_X = G_X/\mathbb{T}$ be the quotient homomorphism. Assume that $\nu_X(G)$ contains a subgroup isomorphic to \mathfrak{A}_4 . Then the following assertions hold:

- (1) if X is G -minimal and G -birationally rigid, then $X = V_6$ or $X = Y_{24}$;
- (2) if $X = V_6$ or $X = Y_{24}$, and $|G| \geq 32 \cdot 24^4$, then X is G -birationally super-rigid.

□

In Sect. 3, we provide a criterion for a G -minimal toric Fano variety X to be G -solid in the case where G is an algebraic subgroup of G_X that contains the maximal torus \mathbb{T} . Unfortunately, we do not know how to generalize this criterion for finite subgroups in G_X . Nevertheless, Exercise 1.1 and Theorem 1.2 suggest the following conjecture.

Conjecture 1.5 *Let X be a toric Fano variety with at most terminal singularities and let G be a subgroup in G_X that contains \mathbb{T} such that X is G -minimal and G -solid. Then there exists a constant $c_X > 0$ such that for every finite subgroup $H \subset G$ such that $\nu_X(H) = \nu_X(G)$, the Fano variety X is H -solid provided that $|H| \geq c_X$.*

The structure of the article is the following. In Sect. 2, we present results that will be used in the proof of Theorem 1.2. In Sects. 3 and 4, we prove Theorem 1.2 for infinite algebraic groups using toric geometry. In Sect. 5, we explicitly describe two (known) equivariant toric Sarkisov links that start at X_{24} . In Sect. 6, we give an alternative proof of Theorem 1.2(3) for infinite algebraic groups, which can be generalized for large finite groups. In Sect. 7, we complete the proof of Theorem 1.2 by proving its part (3) for finite groups (the remaining assertions of Theorem 1.2 for finite groups follows from the results proven in Sects. 3 and 4).

2 Preliminary results

In this section, we review results and notions that will be used in the proof of Theorem 1.2.

2.1 Varieties with regular group actions

Let G and G' be two algebraic groups, and let X and X' be two algebraic varieties endowed with regular actions $m_X: G \times X \rightarrow X$ and $m_{X'}: G' \times X' \rightarrow X'$ of the groups G and G' respectively.

Definition 2.1 An equivariant rational map between the varieties X and X' is a pair consisting of morphism of algebraic groups $\varphi: G \rightarrow G'$ and a rational map $\Phi: X \dashrightarrow X'$ such that the following diagram of rational maps commutes

$$\begin{array}{ccc}
 G \times X & \xrightarrow{m_X} & X \\
 \downarrow \varphi \times \Phi & & \downarrow \Phi \\
 G' \times X' & \xrightarrow{m_{X'}} & X'
 \end{array}$$

We say that the rational map Φ is φ -equivariant.

If the morphism φ in Definition 2.1 is an isomorphism and the map Φ is birational, then letting $\rho: G \rightarrow \text{Aut}(X)$ and $\rho': G' \rightarrow \text{Aut}(X')$ be the group homomorphisms determined by m_X and $m_{X'}$, respectively, the commutativity of the diagram in Definition 2.1 is equivalent to the property that

$$\rho' \circ \varphi(g) = \Phi \rho(g) \Phi^{-1}$$

for every $g \in G$. In this paper, we are mostly interested in the case when $G \cong G'$. Because of this, we will assume in the following that $G = G'$, so that both varieties X and X' are endowed with regular actions of the same group G .


Definition 2.2 A G -equivariant rational map $X \dashrightarrow X'$ is an id_G -equivariant rational map $\Phi: X \dashrightarrow X'$. A rational G -map $X \dashrightarrow X'$ is a rational map $\Phi: X \dashrightarrow X'$ which is φ -equivariant for some automorphism φ of G .

We denote by $\text{Bir}^G(X, X')$ the set of birational G -maps between X and X' , and we denote by $\text{Bir}_G(X, X')$ its subset consisting of G -equivariant birational maps $X \dashrightarrow X'$. If $X = X'$ then these sets are groups (with respect to composition of birational maps), which we denote by $\text{Bir}^G(X)$ and $\text{Bir}_G(X)$, respectively.

As an illustration, we describe an equivariant version of a birational map of threefolds which appeared in [11, Proposition 2.1]. Its nature is local, but we will use global language for simplicity of exposition.

Example 2.3 Let X be a smooth threefold, let P be a point in X , let G be an algebraic group that acts faithfully on X , and let C be a G -irreducible curve in X consisting of three irreducible components C_1, C_2 and C_3 meeting at a unique point P and such that the curves C_1, C_2 and C_3 are smooth at P , and their tangent directions at P generate the tangent space $T_P(X)$. Let $\alpha: \tilde{X} \rightarrow X$ be the blow-up of the point P , and let E be its exceptional surface. Denote by \tilde{C}_i the proper transform of the curve C_i on the threefold \tilde{X} . Let L_{ij} be the line in $E \cong \mathbb{P}^2$ that pass through the points $\tilde{C}_i \cap E$ and

$\tilde{C}_j \cap E$. Then there exists the following commutative diagram of birational G -maps:



where β is the blow-up of the curves \tilde{C}_1, \tilde{C}_2 and \tilde{C}_3 , the map γ is a composition of Atiyah flops of the proper transforms on \bar{X} of the curves L_{12}, L_{13} and L_{23} , and δ is the birational contraction of the proper transform of the surface E to a singular point of type $\frac{1}{2}(1, 1, 1)$. The morphism π is a G -equivariant extremal divisorial contraction.

We will use the following result, which is a consequence of Luna’s étale slice theorem (see e. g. [1, Section 2.1] and [8, p. 98]).

Lemma 2.4 *Let G be a reductive group acting faithfully on a variety X and let $P \in X$ be a smooth point which is fixed by the action of G . Then the induced linear action of G on the Zariski tangent space $T_P(X)$ is faithful.* □

In the case of curves, we have the following more precise consequence:

Corollary 2.5 *A finite group of automorphisms of a curve fixing a smooth point is a cyclic group.* □

Corollary 2.6 *Let X be an algebraic variety with a faithful action of the group $G = \mu_n^2$ fixing a smooth point $P \in X$. Let C be a G -invariant curve in X containing P and assume that the stabilizer in G of every irreducible component of C passing through P acts on this component faithfully. Then $\text{mult}_P(C) \geq n$.* □

Proof Let $f: \tilde{C} \rightarrow C$ be the normalization of the curve C . The action of the group G on C lifts to an action on \tilde{C} preserving the preimage $F = f^{-1}(P)$ of the point P . Let Q be a point in F , and let G_Q be its stabilizer in G . Note that Q is contained in a unique irreducible component of the smooth curve \tilde{C} , which then must be G_Q -invariant. Since the group G_Q acts faithfully on this component, we conclude that G_Q is a cyclic subgroup of the group $G \cong \mu_n^2$ by Corollary 2.5. Then $|G_Q| \leq n$, so that

$$\text{mult}_P(C) \geq |F| \geq \frac{|G|}{|G_Q|} \geq n$$

as required. □

2.2 Singularities of log-pairs

Let X be a threefold with at most terminal singularities, let \mathcal{M}_X be a non-empty mobile linear system on X that consists of \mathbb{Q} -Cartier divisors and let λ be a positive rational number.

Lemma 2.7 ([15, Exercise 6.18]) *Let C be an irreducible curve in X . Assume that*

$$\text{mult}_C(\mathcal{M}_X) \leq \frac{1}{\lambda}.$$

Then C is not a center of non-canonical singularities of the log pair $(X, \lambda\mathcal{M}_X)$. \square

The following result is due to Alessio Corti.

Lemma 2.8 ([13, Theorem 3.1]) *Let C be an irreducible curve in X . Assume that C is a center of non-log canonical singularities of the log pair $(X, \lambda\mathcal{M}_X)$. Then*

$$\text{mult}_C(M_1 \cdot M_2) > \frac{4}{\lambda^2}$$

for any two general surfaces M_1 and M_2 in the linear system \mathcal{M}_X . \square

The following result is due to Alexander Pukhlikov.

Lemma 2.9 ([29], see also [13, Corollary 3.4]) *Let P be a smooth point of X . Assume that P is a center of non-canonical singularities of the log pair $(X, \lambda\mathcal{M}_X)$. Then*

$$\text{mult}_P(M_1 \cdot M_2) > \frac{4}{\lambda^2}$$

for any two general surfaces M_1 and M_2 in the linear system \mathcal{M}_X . \square

We will also need the following two results due to Kawamata [23] and Corti, respectively.

Lemma 2.10 *Let P be a singular point of X of type $\frac{1}{2}(1, 1, 1)$, let $\pi : V \rightarrow X$ be the Kawamata blow-up of P , let E be the exceptional surface, let \mathcal{M}_V be the proper transform of the linear system \mathcal{M}_X via π , and let $m \in \mathbb{Q}$ be such that*

$$\mathcal{M}_V \sim_{\mathbb{Q}} \pi^*(\mathcal{M}_X) - mE.$$

If $(X, \lambda\mathcal{M}_X)$ is not canonical at P then $m > \frac{1}{2\lambda}$. \square

Proof Suppose that $m \leq \frac{1}{2\lambda}$. Let us seek for a contradiction. Since

$$K_V + \lambda\mathcal{M}_V + \left(\lambda m - \frac{1}{2}\right)E \sim_{\mathbb{Q}} \pi^*(K_X + \lambda\mathcal{M}_X),$$

the pair $(V, \lambda\mathcal{M}_V)$ is not canonical at some point $O \in E$. Then $\text{mult}_O(\mathcal{M}_V) > \frac{1}{\lambda}$, so that

$$\text{mult}_O(\mathcal{M}_{V|E}) > \frac{1}{\lambda},$$

which is impossible, since $\mathcal{M}_{V|E} \sim_{\mathbb{Q}} 2mL$, where L is a line in $E \cong \mathbb{P}^2$. □

Lemma 2.11 *Let P be an ordinary double point of X , let $\pi : V \rightarrow X$ be the blow-up of P , let E be the exceptional surface of π , let \mathcal{M}_V be the proper transform of the linear system \mathcal{M}_X via π , and let $m \in \mathbb{Q}$ be such that*

$$\mathcal{M}_V \sim_{\mathbb{Q}} \pi^*(\mathcal{M}_X) - mE.$$

If P is a center of non-canonical singularities of the log pair $(X, \lambda\mathcal{M}_X)$ then $m > \frac{1}{\lambda}$. □

Proof This is [9, Theorem 1.7.20], which is equivalent to [13, Theorem 3.10]. □

Finally, we will need one local result proved in [2]. To state it, we suppose that the threefold X is endowed with an action of an algebraic group G , and \mathcal{M}_X is G -invariant.

Lemma 2.12 ([2, Lemma 2.4]) *Suppose that the group G fixes a smooth point $P \in X$ and that its induced linear action on the Zariski tangent space $T_P X$ is an irreducible representation. If P is a non-canonical center of the log pair $(X, \lambda\mathcal{M}_X)$ then $\text{mult}_P(\mathcal{M}_X) > \frac{2}{\lambda}$.* □

2.3 Finite groups acting on toric varieties

Let \mathbb{T} be a torus of dimension $d \geq 2$, and let Γ_n be a subgroup of \mathbb{T} isomorphic to μ_n^d , where n is a positive integer (note that \mathbb{T} contains such a subgroup for every n). Let X be a projective toric \mathbb{T} -variety of dimension d .

Lemma 2.13 *Let C be a Γ_n -invariant Γ_n -irreducible curve in X , and let H be a very ample divisor on X . If $n > H \cdot C$, then C is \mathbb{T} -invariant.* □

Proof Suppose that C is not \mathbb{T} -invariant. By replacing X by a \mathbb{T} -invariant toric closed subvariety if necessary, we can assume that the curve C is not contained in any proper \mathbb{T} -invariant subvariety of X so that Γ_n acts faithfully on C . The curve C is \mathbb{T} -invariant if and only if each of its irreducible components is \mathbb{T} -invariant.

Let k be the number of irreducible components of the curve C , let Z be an irreducible component of C and let Γ_Z be the stabilizer of the curve Z in the group Γ_n . Then Γ_Z is an index k subgroup of μ_n^d , equal to the product of d cyclic subgroups μ_{m_i} for some positive integers m_i which divide n , say $n = m_i k_i$, $i = 1, \dots, d$. Let $m = \text{gcd}\{m_i\}_{i=1, \dots, d}$ and write $m_i = m r_i$ where $r_i \geq 1$. Then Γ_Z contains a subgroup isomorphic to μ_m^d . By construction, we have $mn^{d-1} \geq \prod_{i=1}^d m_i$ so that

$$k = \prod_{i=1}^d k_i = \prod_{i=1}^d \frac{n}{m_i} = \frac{n^d}{\prod_{i=1}^d m_i} \geq \frac{n^d}{mn^{d-1}} = \frac{n}{m}.$$

Thus $m \geq n/k$ and since by hypothesis $n > H \cdot C$, it follows that $m > H \cdot Z$.

Replacing C by Z and n by m , we assume from now on that C is irreducible. Let us show that $n \leq H \cdot C$. Let $f: \tilde{C} \rightarrow C$ be the normalization of C . Then the action of the group Γ_n lifts to a faithful action on \tilde{C} . Let D be a \mathbb{T} -invariant effective divisor such that $D \sim H$. Then $C \not\subset \text{Supp}(D)$ by assumption on C not being \mathbb{T} -invariant. Let $\Sigma = C \cap \text{Supp}(D)$, and let $\tilde{\Sigma}$ be its preimage in \tilde{C} . Then

$$|\tilde{\Sigma}| \leq \deg(f^*(D|_C)) = \deg(f^*(H|_C)) = H \cdot C.$$

Let P be a point in $\tilde{\Sigma}$, and let G_P be its stabilizer in Γ_n . Then G_P is cyclic by Lemma 2.5. On the other hand, we have

$$|G_P| \geq \frac{|\Gamma_n|}{|\tilde{\Sigma}|} \geq \frac{|\Gamma_n|}{H \cdot C} \geq \frac{n^d}{H \cdot C} \geq \frac{n^2}{H \cdot C}.$$

Therefore, if $n > H \cdot C$, then the order of the cyclic group G_P is strictly larger than n , which is impossible, since G_P is a subgroup of the group $\Gamma_n \cong \mu_n^d$. \square

3 Lattices and toric geometry

Let $\mathbb{T} \cong \mathbb{G}_m^n$ be an algebraic torus of dimension n . We identify \mathbb{T} with the spectrum of the group algebra $\mathbb{C}[M]$ of its character lattice $M = \text{Hom}(\mathbb{T}, \mathbb{G}_m) \cong \mathbb{Z}^n$. The action of the torus \mathbb{T} on itself by translations determines an injective group homomorphism $\mathbb{T} \rightarrow \text{Aut}(\mathbb{T})$ and we have split exact sequence

$$1 \longrightarrow \mathbb{T} \longrightarrow \text{Aut}(\mathbb{T}) \longrightarrow \text{GL}(M) \longrightarrow 1.$$

The splitting is given by mapping every $A \in \text{GL}(M) \cong \text{GL}_n(\mathbb{Z})$ to the algebraic group automorphism of \mathbb{T} associated to the group algebra automorphism $\mathbb{C}[M] \rightarrow \mathbb{C}[M]$ given by

$$\chi^u \mapsto \chi^{A(u)}.$$

We henceforth identify $\text{Aut}(\mathbb{T}) = \mathbb{T} \rtimes \text{GL}(M)$ and we denote its subgroup $\mathbb{T} \times \{\text{id}_M\}$ simply by \mathbb{T} . Every algebraic subgroup $G \subset \text{Aut}(\mathbb{T})$ containing \mathbb{T} is then of the form $G = \mathbb{T} \rtimes \mathbb{W}$ for some finite subgroup \mathbb{W} of $\text{GL}(M)$.

Let \mathbb{W}_1 and \mathbb{W}_2 be finite subgroups of $\text{GL}(M)$, let $G_1 = \mathbb{T} \rtimes \mathbb{W}_1$ and $G_2 = \mathbb{T} \rtimes \mathbb{W}_2$ be the corresponding algebraic subgroups of $\text{Aut}(\mathbb{T})$ that contain the torus \mathbb{T} , let $m_1: G_1 \times \mathbb{T} \rightarrow \mathbb{T}$ and $m_2: G_2 \times \mathbb{T} \rightarrow \mathbb{T}$ be the algebraic actions they determine.

Lemma 3.1 *The following conditions are equivalent:*

- (a) *There exist an isomorphism $\varphi: G_1 \rightarrow G_2$ and a φ -equivariant biregular map $\Phi: \mathbb{T} \rightarrow \mathbb{T}$;*
- (b) *The groups G_1 and G_2 are conjugate in $\text{Aut}(\mathbb{T})$;*

(c) The groups \mathbb{W}_1 and \mathbb{W}_2 are conjugate in $\text{GL}(M)$.

□

Proof Assume (a). Then we have a group automorphism $c_\Phi : \text{Aut}(\mathbb{T}) \rightarrow \text{Aut}(\mathbb{T})$ given by $\alpha \mapsto \Phi \circ \alpha \circ \Phi^{-1}$ and the hypothesis that Φ is φ -equivariant implies that the diagram

$$\begin{array}{ccc} G_1 & \hookrightarrow & \text{Aut}(\mathbb{T}) \\ \varphi \downarrow & & \downarrow c_\Phi \\ G_2 & \hookrightarrow & \text{Aut}(\mathbb{T}) \end{array}$$

must be commutative, so that the algebraic groups G_1 and G_2 are conjugate in $\text{Aut}(\mathbb{T})$. This shows that (a) implies (b).

Now we assume (b). Then $G_2 = \Phi G_1 \Phi^{-1}$ for some $\Phi = (\lambda, A)$ in $\text{Aut}(\mathbb{T})$. Then $\mathbb{W}_2 = A \mathbb{W}_1 A^{-1}$ in $\text{GL}(M)$. This shows that (b) implies (c).

Assume (c). Then $\mathbb{W}_2 = A \mathbb{W}_1 A^{-1}$ for some $A \in \text{GL}(M)$. Let $\Phi = (1, A) \in \text{Aut}(\mathbb{T})$, and let $\varphi : G_1 \rightarrow G_2$ be the homomorphism defined by $g_1 \mapsto \Phi g_1 \Phi^{-1}$. Then φ is an isomorphism for which we have the same commutative diagram as above. It follows in turn that the pair $(\varphi : G_1 \rightarrow G_2, \Phi : \mathbb{T} \rightarrow \mathbb{T})$ is an equivariant isomorphism, which proves that (c) implies (a). □

Now we fix a finite subgroup \mathbb{W} in $\text{GL}(M)$ and we let $G = \mathbb{T} \rtimes \mathbb{W}$ be the corresponding algebraic subgroup in $\text{Aut}(\mathbb{T})$ that contains \mathbb{T} . The group \mathbb{W} acts naturally on the vector space

$$N_{\mathbb{Q}} = \text{Hom}(M, \mathbb{Z}) \otimes \mathbb{Q}.$$

By [16, Chapter 2], the choice of a \mathbb{W} -invariant convex lattice polytope in $N_{\mathbb{Q}}$ determines a projective toric variety X with an open \mathbb{T} -orbit $\mathbb{T}_X \cong \mathbb{T}$ such that the G -action on the torus \mathbb{T}_X extends to a faithful regular G -action $m_X : G \times X \rightarrow X$. Thus, we can identify G with its image in the group $\text{Aut}(X)$ by the injective group homomorphism $\rho_X : G \rightarrow \text{Aut}(X)$ given by $g \mapsto m_X(g, \cdot)$. Then $\text{Aut}(X)$ is an affine algebraic group having \mathbb{T} as a maximal torus [17].

Let G_X be the normalizer of the torus \mathbb{T} in the group $\text{Aut}(X)$. Then G_X is an algebraic group that contains G . Moreover, the torus \mathbb{T}_X is G_X -invariant, and the induced effective action of the group G_X on the torus \mathbb{T}_X corresponds to an injective group homomorphism

$$G_X \rightarrow \text{Aut}(\mathbb{T}_X) \cong \text{Aut}(\mathbb{T}),$$

whose image is equal to $\mathbb{T} \rtimes \mathbb{W}_X$ for a finite subgroup $\mathbb{W}_X \subset GL(M)$ that contains \mathbb{W} . Thus, we have the following commutative diagram of exact sequences:

$$\begin{array}{ccccccccc}
 1 & \longrightarrow & \mathbb{T} & \longrightarrow & G = \mathbb{T} \rtimes \mathbb{W} & \longrightarrow & \mathbb{W} & \longrightarrow & 1 \\
 & & \parallel & & \downarrow & & \downarrow & & \\
 1 & \longrightarrow & \mathbb{T} & \longrightarrow & G_X & \xrightarrow{\nu_X} & \mathbb{W}_X & \longrightarrow & 1.
 \end{array}$$

The group \mathbb{W}_X is usually called the *Weyl group* of the toric variety X .

Corollary 3.2 *If \mathbb{W} is a maximal finite subgroup of $GL(M)$ then $G_X \cong \mathbb{T} \rtimes \mathbb{W}$.* □

As a consequence of Lemma 3.1, we obtain the following two assertions:

Corollary 3.3 *There exists a functorial one-to-one correspondence between finite subgroups $\mathbb{W} \subset GL(M)$ up to conjugacy and projective toric \mathbb{T} -varieties X whose Weyl groups contain a subgroup isomorphic to \mathbb{W} up to $\mathbb{T} \rtimes \mathbb{W}$ -equivariant birational equivalence.* □

Corollary 3.4 *Let \mathbb{W} be a finite subgroup in $GL(M)$ and let X be a projective toric variety whose Weyl group \mathbb{W}_X contains \mathbb{W} . Then*

$$\text{Bir}^{\mathbb{T} \rtimes \mathbb{W}}(X) \cong \mathbb{T} \rtimes \widehat{\mathbb{W}}$$

where $\widehat{\mathbb{W}}$ is the normalizer of the group \mathbb{W} in $GL(M)$. □

Given a subgroup $\mathbb{W} \subset GL(M)$, we say that the lattice $M \cong \mathbb{Z}^n$ is \mathbb{W} -irreducible (or an irreducible \mathbb{W} -module) if M does not contain any proper \mathbb{W} -invariant sublattice M' such that M/M' is torsion free.

Corollary 3.5 *Let \mathbb{W} be a maximal finite subgroup in $GL(M)$ and let X be a projective toric variety whose Weyl group is \mathbb{W} . Suppose that M is \mathbb{W} -irreducible. Then*

$$\text{Bir}^{\mathbb{T} \rtimes \mathbb{W}}(X) \cong \mathbb{T} \rtimes \mathbb{W}.$$

□

Proof Since M is \mathbb{W} -irreducible, $M \otimes \mathbb{Q}$ is an irreducible \mathbb{Q} -representation of the group \mathbb{W} . Applying Maschke’s theorem, we conclude that the centralizer of \mathbb{W} in $GL(M)$ is finite. Since \mathbb{W} is finite, the normalizer $\widehat{\mathbb{W}}$ of the group \mathbb{W} in $GL(M)$ is also finite, and hence, $\widehat{\mathbb{W}} = \mathbb{W}$ because \mathbb{W} is a maximal finite subgroup. The assertion then follows from Corollary 3.4. □

The choice of the n -dimensional toric variety X whose Weyl group contains \mathbb{W} is not unique. In particular, taking a G -equivariant toric resolution of singularities and then applying the G -equivariant toric Minimal Model Program, we can assume that:

- The toric variety X has terminal singularities,

Thus, if X is a G -minimal toric Fano variety, we have a purely group theoretical criterion for its G -solidity. Similarly, we can obtain a criterion for G -birational rigidity.

Proposition 3.8 *Let X be a G -minimal toric Fano variety with Weyl group \mathbb{W}_X . Assume that the character lattice M is \mathbb{W}_X -irreducible. Then the following two conditions are equivalent:*

- (a) X is G -birationally rigid;
- (b) X is the only toric Fano variety with terminal singularities that is G -minimal.

Proof This follows from Proposition 3.7 and definition of G -birational rigidity. □

Finally, using Corollary 3.4, we can obtain a criterion for G -birational super-rigidity.

Proposition 3.9 *Let X be a G -minimal toric Fano variety with Weyl group \mathbb{W} . Assume that the character lattice M is \mathbb{W} -irreducible. Then X is G -birationally super-rigid if and only if the following two conditions are satisfied:*

- (a) X is the only toric Fano variety with terminal singularities that is G -minimal;
- (b) \mathbb{W} is not a proper normal subgroup of any finite subgroup in $GL(M)$.

Proof The assertion follows from Proposition 3.8 and the proof of Corollary 3.5. □

The condition (b) in Proposition 3.8 is combinatorial. A priori, it can be checked using computer, since there are finitely many toric Fano varieties with terminal singularities [6]. For example, there are 634 toric Fano threefolds with terminal singularities [22].

3.1 Toric terminal Fano threefolds

Now let us assume that \mathbb{T} is three-dimensional and that X is a G -minimal toric Fano threefold with terminal singularities. All such threefolds are described in [31]. They are listed in the following table:

Toric Fano threefold	Weyl group	Number in [7]
Divisor Y_{24} of type $(1, 1, 1, 1)$ in $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$	$\mathfrak{S}_4 \times \mu_2$	No 625
$V_6 = \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$	$\mathfrak{S}_3 \times \mu_2^3 \cong \mathfrak{S}_4 \times \mu_2$	No 62
Toric Fano–Enriques threefold X_{24}	$\mathfrak{S}_4 \times \mu_2$	No 47
Toric complete intersection $V_4 \subset \mathbb{P}^3$ of two quadrics	$\mathfrak{S}_4 \times \mu_2$	No 297
Three-dimensional projective space \mathbb{P}^3	\mathfrak{S}_4	No 4
Quadric cone in \mathbb{P}^4 with one singular point	D_8	No 32
Terminal toric Fano threefold X with $-K_X^3 = \frac{81}{2}$	\mathfrak{S}_3	No 92
Weighted projective space $\mathbb{P}(1, 1, 1, 2)$	\mathfrak{S}_3	No 7
Quotient of the space \mathbb{P}^3 by μ_5 -action fixing 5 points	μ_2^2	No 1
Weighted projective space $\mathbb{P}(1, 1, 2, 3)$	μ_2	No 8

where α is Kawamata blow-up of these two singular points, ι is a composition of Francia antiflips of the proper transforms of all \mathbb{T} -invariant curves that contain one of these points, and β is a \mathbb{P}^1 -bundle over a del Pezzo surface of degree 6. The threefolds $X_{24}, \overline{X}_{24}, \widehat{X}_{24}$ are quotients by involutions of the threefolds $V_6, \overline{V}_6, \widehat{V}_6$ from Example 3.12.

Moreover, if X is one of the threefolds $Y_{24}, V_6, X_{24}, V_4, \mathbb{P}^3$ and \mathbb{W} contains a subgroup isomorphic to \mathfrak{A}_4 , then X is G -minimal except in the following two cases:

- (1) $X = V_4, \mathbb{W} \cong \mathfrak{S}_4$ and G acts intransitively on the set of \mathbb{T} -invariant surfaces,
- (2) $X = V_4$ and $\mathbb{W} \cong \mathfrak{A}_4$.

We show this in Corollaries 4.5 and 4.9 and Lemma 5.1 below. Summing up, we get

Corollary 3.14 *The assertion of Theorem 1.2 holds in the case when G is infinite. \square*

If \mathbb{W} contains a subgroup isomorphic to \mathfrak{A}_4 , then \mathbb{W} is conjugate to one of 15 finite subgroups in $GL(M)$ that are described in [32]. Using [24] and notation from [32], we can present these 15 subgroups and the corresponding G -minimal toric Fano threefolds with terminal singularities in the following table.

	$\mathfrak{S}_4 \times \mu_2$	\mathfrak{S}_4	$\mathfrak{A}_4 \times \mu_2$	\mathfrak{A}_4
$\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$	W_1	W_6 or W_7	W_1	W_9
V_4	W_3	W_{10}	W_3	
X_{24}	W_3	W_{10} or W_{11}	W_3	W_{11}
Y_{24}	W_2	W_8 or W_9	W_2	W_{10}
\mathbb{P}^3		W_{11}		W_{11}

Now using Propositions 3.8 and 3.9, we obtain

Corollary 3.15 *The assertion of Corollary 1.4 holds in the case when G is infinite. \square*

In the rest of this paper, we will give another proof of Theorem 1.2(3) in the case when the group G is infinite that is independent on the classification of toric Fano threefolds with terminal singularities, and which also applies to the case of finite groups as well. We also believe that this approach can be used in higher-dimensions.

4 Toric Fano threefolds and lattices of rank three

Among the 73 conjugacy classes of finite subgroups \mathbb{W} of $GL_3(\mathbb{Z})$ classified in [32], there are 4 maximal ones, and only three of them give rise to an irreducible action on \mathbb{Z}^3 . In each of these three cases, one has $\mathbb{W} \cong \mathfrak{S}_4 \times \mu_2$. Let us describe these three conjugacy classes in terms of the actions of the group $\mathfrak{S}_4 \times \mu_2$ on certain lattices.

Let $L = \mathbb{Z}^4$ endowed with the faithful transitive \mathfrak{S}_4 -action given by permutations of the basis vectors h_1, h_2, h_3 and h_4 . Let σ be the involution of the lattice L such that $h_i \mapsto -h_i$ for each $i \in \{1, 2, 3, 4\}$. Then σ commutes with the \mathfrak{S}_4 -action. This

defines a faithful action of the group $\mathfrak{S}_4 \times \mu_2$ on the lattice L , which leaves invariant the sublattice spanned by the element $h_1 + h_2 + h_3 + h_4$. Let

$$M_1 = L / \langle h_1 + h_2 + h_3 + h_4 \rangle.$$

Then the $\mathfrak{S}_4 \times \mu_2$ -action on L induces an action of $\mathfrak{S}_4 \times \mu_2$ on the quotient lattice M_1 . Let e_1, e_2 and e_3 be the basis of M_1 given by the classes of h_1, h_2 and h_3 , respectively. In this basis, we have $\sigma(e_i) = -e_i$ for every $i \in \{1, 2, 3\}$, and for every $g \in \mathfrak{S}_4$, we have

$$g(e_i) = \begin{cases} e_{g(i)} & \text{if } g(i) \neq 4 \\ -e_1 - e_2 - e_3 & \text{otherwise.} \end{cases}$$

We denote by $\mathbb{W}_1 \cong \mathfrak{S}_4 \times \mu_2$ the corresponding subgroup of $\text{GL}_3(\mathbb{Z})$.

Let M_3 be the dual lattice to M_1 , and let e_1^\vee, e_2^\vee and e_3^\vee be the basis of M_3 that is dual to the previously fixed basis of M_1 . Then $\sigma(e_i^\vee) = -e_i^\vee$ for each $i \in \{1, 2, 3\}$. For every $g \in \mathfrak{S}_4$, we have

$$g(e_i^\vee) = \sum_{j=1}^3 e_j^\vee (g^{-1}(e_j)) e_j^\vee.$$

We denote by $\mathbb{W}_3 \cong \mathfrak{S}_4 \times \mu_2$ the corresponding subgroup of $\text{GL}_3(\mathbb{Z})$.

Finally, let M_2 be the lattice \mathbb{Z}^3 , let \mathfrak{S}_3 be the subgroup in $\text{GL}_3(\mathbb{Z})$ consisting of six permutation matrices, let

$$\tau_1 = \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \tau_2 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \tau_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix},$$

and let $\mathbb{W}_2 \cong \mathfrak{S}_3 \times \mu_2^3$ be the subgroup in $\text{GL}_3(\mathbb{Z})$ that is generated by \mathfrak{S}_3 and involutions τ_1, τ_2 and τ_3 . Note that the subgroup generated by $\mathfrak{S}_3, \tau_1 \tau_2$ and that $\tau_1 \tau_3$ is isomorphic to \mathfrak{S}_4 , and $\tau_1 \tau_2 \tau_3$ generates the center of the subgroup \mathbb{W}_2 . Thus, \mathbb{W}_2 is isomorphic to the group $\mathfrak{S}_4 \times \mu_2$.

Proposition 4.1 ([32]) *Let \mathbb{W} be a maximal finite subgroup of the group $\text{GL}_3(\mathbb{Z})$ such that \mathbb{Z}^3 is \mathbb{W} -irreducible. Then \mathbb{W} is conjugate to one of the subgroups $\mathbb{W}_1, \mathbb{W}_2$ or \mathbb{W}_3 . Moreover, the subgroups $\mathbb{W}_1, \mathbb{W}_2, \mathbb{W}_3$ are pairwise non-conjugate in $\text{GL}_3(\mathbb{Z})$.*

Notation 4.2 The center of each of the three finite subgroups \mathbb{W}_i in $\text{GL}_3(\mathbb{Z}), i = 1, 2, 3$ is isomorphic to μ_2 , generated by the involution

$$\sigma = \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}.$$

For every $i = 1, 2, 3$, the image of the subgroup $\mathfrak{A}_4 \times \{1\}$ of $\mathfrak{S}_4 \times \mu_2$ by the isomorphism $GL(M_i) \cong GL_3(\mathbb{Z})$ given by our choice of bases is the unique subgroup of \mathbb{W}_i isomorphic to \mathfrak{A}_4 . We denote this subgroup by $\mathbb{W}_i^{\mathfrak{A}_4}$. In the notation of [32, Proposition 7], these groups correspond respectively to the subgroups

$$\begin{aligned} \mathbb{W}_1^{\mathfrak{A}_4} = W_{10} &= \left\{ \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & -1 & 1 \\ 0 & -1 & 0 \\ 1 & -1 & 0 \end{pmatrix} \right\} \\ \mathbb{W}_2^{\mathfrak{A}_4} = W_9 &= \left\{ \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \right\} \\ \mathbb{W}_3^{\mathfrak{A}_4} = W_{11} &= \left\{ \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} -1 & -1 & -1 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \right\} \end{aligned}$$

of $SL_3(\mathbb{Z}) \subset GL_3(\mathbb{Z})$.

On the other hand, each of the subgroups \mathbb{W}_i contains two different subgroups isomorphic to \mathfrak{S}_4 (see [32, Proposition 9]):

(1) One is the image of the subgroup $\mathfrak{S}_4 \times \{1\}$ of $\mathfrak{S}_4 \times \mu_2$ by the isomorphism $GL(M_i) \cong GL_3(\mathbb{Z})$ given by our choice of bases. We denote it by $\overline{\mathbb{W}}_i^{\mathfrak{S}_4}$. It is easily seen that this subgroup is not contained in $SL_3(\mathbb{Z})$.

(2) The second one is the intersection $\mathbb{W}_i \cap SL_3(\mathbb{Z})$. It is generated by the images under the isomorphism $GL(M_i) \cong GL_3(\mathbb{Z})$ of the transpositions in the subgroup $\mathfrak{S}_4 \times \{1\}$ multiplied by the element $\sigma \in GL_3(\mathbb{Z})$. We denote it by $\mathbb{W}_i^{\mathfrak{S}_4}$.

The lattice M_3 can be seen as the root lattice of the root system A_3 endowed with the natural action of the group $Aut(A_3) \cong \mathfrak{S}_4 \times \mu_2$. Similarly, one can show that M_1 is the weight lattice of this root lattice, so that there is an inclusion $M_3 \hookrightarrow M_1$ as a sublattice of index 4. With our choice of bases, it is given by the matrix

$$\begin{pmatrix} 2 & 1 & 1 \\ 1 & 2 & 1 \\ 1 & 1 & 2 \end{pmatrix}.$$

Likewise, the lattice M_2 is the root lattice of the root system B_3 endowed with the natural action of the group $Aut(B_3) \cong \mathfrak{S}_3 \times \mu_2^3 \cong \mathfrak{S}_4 \times \mu_2$. The inclusion $M_3 \hookrightarrow M_1$ factors as the composition of two inclusions $M_3 \hookrightarrow M_2$ and $M_2 \hookrightarrow M_1$ as sublattices of index two. With our choice of bases, they are given by the matrices

$$\begin{pmatrix} 1 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 1 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 1 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 1 \end{pmatrix},$$

respectively. All the inclusions $M_3 \hookrightarrow M_2 \hookrightarrow M_1$ are $\mathfrak{S}_4 \times \mu_2$ -equivariant.

Let $\mathbb{T}_1 = \text{Spec}(\mathbb{C}[M_1])$, $\mathbb{T}_2 = \text{Spec}(\mathbb{C}[M_2])$ and $\mathbb{T}_3 = \text{Spec}(\mathbb{C}[M_3])$ be the three-dimensional tori that correspond to the lattices M_1 , M_2 and M_3 , respectively. We

write

$$\mathbb{C}[M_1] = \mathbb{C}[\hat{t}_1^{\pm 1}, \hat{t}_2^{\pm 1}, \hat{t}_3^{\pm 1}]$$

and identify (using the inclusions $M_3 \hookrightarrow M_2 \hookrightarrow M_1$) the algebras $\mathbb{C}[M_2]$ and $\mathbb{C}[M_3]$ with the subalgebras of the algebra $\mathbb{C}[M_1]$ as follows:

$$\mathbb{C}[M_2] = \mathbb{C}[\bar{t}_1^{\pm 1}, \bar{t}_2^{\pm 1}, \bar{t}_3^{\pm 1}] = \mathbb{C}[(\hat{t}_1 \hat{t}_2)^{\pm 1}, (\hat{t}_1 \hat{t}_3)^{\pm 1}, (\hat{t}_2 \hat{t}_3)^{\pm 1}]$$

and

$$\begin{aligned} \mathbb{C}[M_3] &= \mathbb{C}[t_1^{\pm 1}, t_2^{\pm 1}, t_3^{\pm 1}] = \mathbb{C}[(\bar{t}_1 \bar{t}_2)^{\pm 1}, (\bar{t}_1 \bar{t}_3)^{\pm 1}, (\bar{t}_2 \bar{t}_3)^{\pm 1}] \\ &= \mathbb{C}[(\hat{t}_1^2 \hat{t}_2 \hat{t}_3)^{\pm 1}, (\hat{t}_1 \hat{t}_2^2 \hat{t}_3)^{\pm 1}, (\hat{t}_1 \hat{t}_2 \hat{t}_3^2)^{\pm 1}]. \end{aligned}$$

This gives us morphisms $q_{12}: \mathbb{T}_1 \rightarrow \mathbb{T}_2$ and $q_{23}: \mathbb{T}_2 \rightarrow \mathbb{T}_3$, which are quotients by the involutions $(\hat{t}_1, \hat{t}_2, \hat{t}_3) \mapsto (-\hat{t}_1, -\hat{t}_2, -\hat{t}_3)$ and $(\bar{t}_1, \bar{t}_2, \bar{t}_3) \mapsto (-\bar{t}_1, -\bar{t}_2, -\bar{t}_3)$, respectively.

In the next three subsections, we present three toric Fano varieties with terminal singularities that are natural equivariant compactifications of the tori $\mathbb{T}_1, \mathbb{T}_2, \mathbb{T}_3$ following the scheme described in Sect. 3. Before doing this, let us first fix some notation.

Notation 4.3 Let $([u_1 : v_1], \dots, [u_n : v_n])$ be homogeneous coordinates on $(\mathbb{P}^1)^n$. We equip $(\mathbb{P}^1)^n$ with its standard structure of a toric variety with open orbit $\mathbb{T}_{(\mathbb{P}^1)^n}$ given by

$$\prod_{i=1}^n u_i v_i \neq 0.$$

We view the collection of ratios $(\frac{u_1}{v_1}, \dots, \frac{u_n}{v_n})$ as natural “toric coordinates” on $(\mathbb{P}^1)^n$. We use these to identify each torus-invariant irreducible closed subvariety of $(\mathbb{P}^1)^n$ with the toric coordinates of its general point. For example, for $n = 4$, this yields:

- $(0, 1, 1, 1)$ is the torus-invariant divisor $u_1 = 0$;
- $(0, 1, \infty, 1)$ is the torus-invariant surface given by $u_1 = v_3 = 0$;
- $(0, 0, 0, 0)$ is the torus-invariant point $u_1 = u_2 = u_3 = u_4 = 0$.

Finally, we denote by v be the involution of \mathbb{P}^1 given by $[u : v] \mapsto [v : u]$.

4.1 Toric Fano threefold with Weyl group \mathbb{W}_1

The convex hull of the points

$$(0, \pm 1, \pm 1), (\pm 1, 0, \pm 1), (\pm 1, \pm 1, 0)$$

in $\text{Hom}(M_1, \mathbb{Z}) \otimes \mathbb{Q}$ is a \mathbb{W}_1 -invariant convex polytope. One can show that the associated toric Fano threefold is the hypersurface Y_{24} in $(\mathbb{P}^1)^4$ that is given by

$$u_1 u_2 u_3 u_4 - v_1 v_2 v_3 v_4 = 0.$$

The open \mathbb{T}_1 -orbit is the subset $\mathbb{T}_{Y_{24}}$ that is given by

$$u_1 u_2 u_3 u_4 v_1 v_2 v_3 v_4 \neq 0.$$

We have $\mathbb{W}_{Y_{24}} \cong \mathbb{W}_1$, so that we identify $\mathbb{W}_{Y_{24}} = \mathbb{W}_1$, $\mathbb{T}_{Y_{24}} = \mathbb{T}_1$ and $G_{Y_{24}} = \mathbb{T}_1 \times \mathbb{W}_1$.

The \mathbb{W}_1 -action on Y_{24} is given by the permutations of the factors in $(\mathbb{P}^1)^4$ and the involution $v \times v \times v \times v$, which corresponds to the element σ of \mathbb{W}_1 . We also denote this involution by $\sigma_{Y_{24}}$.

The threefold Y_{24} has fourteen \mathbb{T}_1 -fixed points: the six points

$$(0, 0, \infty, \infty), (0, \infty, 0, \infty), (0, \infty, \infty, 0), (\infty, 0, \infty, 0), (\infty, 0, 0, \infty), (\infty, \infty, 0, 0),$$

which are isolated ordinary double points forming the singular locus of Y_{24} , and the eight smooth points

$$(0, \infty, \infty, \infty), (\infty, 0, \infty, \infty), (\infty, \infty, 0, \infty), (\infty, \infty, \infty, 0),$$

$$(\infty, 0, 0, 0), (0, \infty, 0, 0), (0, 0, \infty, 0), (0, 0, 0, \infty).$$

Similarly, it has twenty four irreducible \mathbb{T}_1 -invariant curves

$$(0, 0, 1, \infty), (0, 1, 0, \infty), (1, 0, 0, \infty), (0, 0, \infty, 1), (0, 1, \infty, 0), (1, 0, \infty, 0),$$

$$(0, \infty, 0, 1), (0, \infty, 1, 0), (1, \infty, 0, 0), (\infty, 0, 0, 1), (\infty, 0, 1, 0), (\infty, 1, 0, 0),$$

$$(\infty, \infty, 1, 0), (\infty, 1, \infty, 0), (1, \infty, \infty, 0), (\infty, \infty, 0, 1), (\infty, 1, 0, \infty), (1, \infty, 0, \infty),$$

$$(\infty, 0, \infty, 1), (\infty, 0, 1, \infty), (1, 0, \infty, \infty), (0, \infty, \infty, 1), (0, \infty, 1, \infty), (0, 1, \infty, \infty),$$

and twelve irreducible \mathbb{T}_1 -invariant surfaces

$$(0, 1, 1, \infty), (1, 0, 1, \infty), (1, 1, 0, \infty), (0, 1, \infty, 1), (1, 0, \infty, 1), (1, 1, \infty, 0),$$

$$(0, \infty, 1, 1), (1, \infty, 0, 1), (1, \infty, 1, 0), (\infty, 0, 1, 1), (\infty, 1, 0, 1), (\infty, 1, 1, 0).$$

With this description, the following lemma is straightforward to check.

Lemma 4.4 *Let \mathbb{W} be a subgroup in \mathbb{W}_1 that contains $\mathbb{W}_1^{\mathfrak{A}_1}$. Then the following hold:*

- (1) *The group $\mathbb{W}_1^{\mathfrak{A}_1}$ acts transitively on the set of \mathbb{T}_1 -invariant surfaces and on the set of singular points of Y_{24} .*
- (2) *The groups $\mathbb{W}_1^{\mathfrak{A}_1}$ and $\overline{\mathbb{W}}_1^{\mathfrak{S}}$ act on the set of smooth \mathbb{T}_1 -fixed points and on the set of \mathbb{T}_1 -invariant curves with the same orbits. The action on the set of smooth \mathbb{T}_1 -fixed points has two orbits: one consisting of the points*

$$(0, \infty, \infty, \infty), (\infty, 0, \infty, \infty), (\infty, \infty, 0, \infty), (\infty, \infty, \infty, 0),$$

and another one consisting of the points

$$(\infty, 0, 0, 0), (0, \infty, 0, 0), (0, 0, \infty, 0), (0, 0, 0, \infty),$$

Similarly, the action on the set of irreducible \mathbb{T}_1 -invariant curves has two orbits: one consisting of the curves

$$(0, 0, 1, \infty), (0, 1, 0, \infty), (1, 0, 0, \infty), (0, 0, \infty, 1), (0, 1, \infty, 0), (1, 0, \infty, 0), (0, \infty, 0, 1), (0, \infty, 1, 0), (1, \infty, 0, 0), (\infty, 0, 0, 1), (\infty, 0, 1, 0), (\infty, 1, 0, 0),$$

and the other one consisting of the curves

$$(\infty, \infty, 1, 0), (\infty, 1, \infty, 0), (1, \infty, \infty, 0), (\infty, \infty, 0, 1), (\infty, 1, 0, \infty), (1, \infty, 0, \infty), (\infty, 0, \infty, 1), (\infty, 0, 1, \infty), (1, 0, \infty, \infty), (0, \infty, \infty, 1), (0, \infty, 1, \infty), (0, 1, \infty, \infty).$$

- (3) The group $\mathbb{W}_1^{\mathfrak{S}}$ acts transitively on the set of smooth \mathbb{T}_1 -fixed points and on the set of irreducible \mathbb{T}_1 -invariant curves.
- (4) If $\sigma_{Y_{24}} \in \mathbb{W}$, then \mathbb{W} acts transitively on the set of smooth \mathbb{T}_1 -fixed points and on the set of \mathbb{T}_1 -invariant curves of Y_{24} .

□

Corollary 4.5 *Let G be a subgroup of $G_{Y_{24}}$ that contains $\mathbb{W}_1^{\mathfrak{A}}$. Then $\text{rk}(\text{Cl}(Y_{24})^G) = 1$.* □

4.2 Toric Fano threefold with Weyl group \mathbb{W}_2

The convex hull of the lattice points $(0, 0, \pm 1), (0, \pm 1, 0), (\pm 1, 0, 0)$ in $\text{Hom}(M_2, \mathbb{Z}) \otimes \mathbb{Q}$ is a \mathbb{W}_2 -invariant convex polytope. One can check that the associated toric Fano threefold is $V_6 = \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$. Moreover, one has $\mathbb{W}_{V_6} \cong \mathbb{W}_2$. Therefore, we identify $\mathbb{W}_{V_6} = \mathbb{W}_2, \mathbb{T}_{V_6} = \mathbb{T}_2$ and $G_{V_6} = \mathbb{T}_2 \rtimes \mathbb{W}_2$.

The action of $\mathbb{W}_2 \cong \mathfrak{S}_3 \times \mu_2^2$ on the threefold V_6 is given by the permutations of three factors, the involutions $v \times v \times \text{id}_{\mathbb{P}^1}$ and $v \times \text{id}_{\mathbb{P}^1} \times v$, and the involution $v \times v \times v$, which corresponds to the element σ of \mathbb{W}_2 . We also denote this involution by σ_{V_6} .

Lemma 4.6 *Let \mathbb{W} be a subgroup of \mathbb{W}_2 that contains $\mathbb{W}_2^{\mathfrak{A}}$. Then the following hold:*

- (1) The group $\mathbb{W}_2^{\mathfrak{A}}$ acts transitively on the set of irreducible \mathbb{T}_2 -invariant surfaces and on the set of irreducible \mathbb{T}_2 -invariant curves.
- (2) The groups $\mathbb{W}_2^{\mathfrak{A}}$ and $\overline{\mathbb{W}}_2^{\mathfrak{S}}$ act on the set of \mathbb{T}_2 -fixed points with the same two orbits: one consisting of the points

$$(0, 0, 0), (\infty, \infty, 0), (\infty, 0, \infty), (0, \infty, \infty),$$

and another one consisting of the points

$$(\infty, \infty, \infty), (0, 0, \infty), (0, \infty, 0), (\infty, 0, 0).$$

- (3) The group $\mathbb{W}_2^{\mathfrak{S}}$ acts transitively on the set of \mathbb{T}_2 -fixed points.
- (4) If $\sigma_{V_6} \in \mathbb{W}$, then \mathbb{W} acts transitively on the set of \mathbb{T}_2 -fixed points.

□

Corollary 4.7 *Let G be a subgroup of G_{V_6} that contains $\mathbb{W}_2^{\mathfrak{A}}$. Then $\text{rk}(\text{Cl}(V_6)^G) = 1$. □*

Remark 4.8 Recall that we have the quotient morphism $q_{12} : \mathbb{T}_1 \rightarrow \mathbb{T}_2$, which is given by

$$(\hat{t}_1, \hat{t}_2, \hat{t}_3) \mapsto (\hat{t}_1\hat{t}_2, \hat{t}_1\hat{t}_3, \hat{t}_2\hat{t}_3).$$

By construction, this morphism is equivariant for the actions of the group $\mathfrak{S}_4 \times \mu_2$ given by the subgroups \mathbb{W}_1 and \mathbb{W}_2 , respectively. Moreover, it induces a q_{12} -equivariant rational map $\varphi : Y_{24} \dashrightarrow V_6$ that has generic degree 2. This rational map is equivariant for the actions of the group $\mathfrak{S}_4 \times \mu_2$ on the threefolds Y_{24} and V_6 . With our choice of coordinates, this rational map can be explicitly written as follows. For $i \in \{1, 2, 3, 4\}$, let

$$U_i = \frac{u_i}{v_i} u_1 u_2 u_3 u_4 v_1 v_2 v_3 v_4 \quad \text{and} \quad V_i = \frac{v_i}{u_i} u_1 u_2 u_3 u_4 v_1 v_2 v_3 v_4.$$

Then the image of the rational $\Phi : Y_{24} \dashrightarrow \mathbb{P}^7$ defined by

$$\begin{aligned} & \left([u_1 : v_1], [u_2 : v_2], [u_3 : v_3], [u_4 : v_4] \right) \mapsto \\ & [U_1 : V_1 : U_2 : V_2 : U_3 : V_3 : U_4 : V_4] \end{aligned} \tag{1}$$

is equal to the image of V_6 by the Segre embedding $j : V_6 \hookrightarrow \mathbb{P}^7$ given by

$$\begin{aligned} & \left([u_1 : v_1], [u_2 : v_2], [u_3 : v_3] \right) \mapsto \left[u_1 u_2 v_3 : v_1 v_2 u_3 : u_1 v_2 u_3 : v_1 u_2 v_3 : \right. \\ & \left. v_1 u_2 v_3 : v_1 u_2 u_3 : u_1 v_2 v_3 : v_1 v_2 v_3 : u_1 u_2 u_3 \right] \end{aligned}$$

and $j \circ \varphi = \Phi$. Moreover, there exists $\mathfrak{S}_4 \times \mu_2$ -equivariant commutative diagram

$$\begin{array}{ccc} \tilde{Y}_{24} & \overset{\beta}{\dashrightarrow} & \hat{Y}_{24} \\ \alpha \downarrow & & \downarrow \gamma \\ Y_{24} & \overset{\varphi}{\dashrightarrow} V_6 \longleftarrow \overset{\delta}{\dashrightarrow} & Y_{12} \end{array} \tag{2}$$

where α is the blow-up of all the singular points of Y_{24} , β is the composition of Atiyah flops of the proper transforms of all \mathbb{T}_1 -invariant curves in Y_{24} , γ is the contraction of the proper transforms of all \mathbb{T}_1 -invariant surfaces in Y_{24} , and δ is the double cover branched over the union of all \mathbb{T}_2 -invariant surfaces. The threefold Y_{12} is the canonical toric Fano threefold No 525553 in [7].

4.3 Toric Fano threefold with Weyl group \mathbb{W}_3

The convex hull of the eight points

$$(\pm 1, \pm 1, \pm 1)$$

in $\text{Hom}(M_3, \mathbb{Z}) \otimes \mathbb{Q}$ is a \mathbb{W}_3 -invariant convex polytope. One can check that the associated toric Fano threefold is the toric Fano threefold X_{24} , No 47 in [7]. Following [5, § 6.3.2], we can also view X_{24} as the quotient of the threefold V_6 by the involution $\tau_{V_6} : V_6 \rightarrow V_6$ defined by

$$([u_1 : v_1], [u_2, v_2], [u_3, v_3]) \mapsto ([u_1 : -v_1], [u_2 : -v_2], [u_3 : -v_3]). \tag{3}$$

In this presentation, the threefold X_{24} comes with a closed embedding $X_{24} \hookrightarrow \mathbb{P}^{13}$ which is induced by the rational map $V_6 \dashrightarrow \mathbb{P}^{13}$ defined by

$$([u_1 : v_1], [u_2 : v_2], [u_3 : v_3]) \mapsto \left[\begin{array}{l} u_1^2 u_2^2 u_3^2 : u_1^2 u_2^2 v_3^2 : u_1^2 u_2 v_2 u_3 v_3 : u_1^2 v_2^2 u_3^2 : \\ u_1^2 v_2^2 v_3^2 : u_1 v_1 u_2^2 u_3 v_3 : u_1 v_1 u_2 v_2 u_3^2 : \\ u_1 v_1 u_2 v_2 v_3^2 : u_1 v_1 v_2^2 u_3 v_3 : v_1^2 u_2^2 u_3^2 : \\ v_1^2 u_2^2 v_3^2 : v_1^2 u_2 v_2 u_3 v_3 : v_1^2 v_2^2 u_3^2 : v_1^2 v_2^2 v_3^2 \end{array} \right]. \tag{4}$$

The action of the torus \mathbb{T}_3 on the threefold X_{24} coincides with that induced from the action of the torus \mathbb{T}_2 on the threefold V_6 via the quotient morphism $\pi : V_6 \rightarrow X_{24}$. Namely, the torus \mathbb{T}_3 is the quotient of the torus \mathbb{T}_2 by the involution

$$(\bar{t}_1, \bar{t}_2, \bar{t}_3) \mapsto (-\bar{t}_1, -\bar{t}_2, -\bar{t}_3),$$

and the quotient morphism $\pi : V_6 \rightarrow X_{24} = V_6/\tau_{V_6}$ becomes equivariant with respect to the quotient morphism $q_{23} : \mathbb{T}_2 \rightarrow \mathbb{T}_3$ when we equip the threefold X_{24} with the induced structure of toric variety.

The involution τ_{V_6} commutes with the action of the Weyl group $\mathbb{W}_{V_6} \cong \mathbb{W}_2$, so that we have $\mathbb{W}_{X_{24}} \cong \mathbb{W}_3$. Hence, we identify $\mathbb{W}_{X_{24}} = \mathbb{W}_3$, $\mathbb{T}_{X_{24}} = \mathbb{T}_3$ and $G_{X_{24}} = \mathbb{T}_3 \rtimes \mathbb{W}_3$.

The action of the group $\mathbb{W}_{X_{24}}$ on the threefold X_{24} coincides with the action induced from the action of the group \mathbb{W}_{V_6} on the threefold V_6 via the quotient morphism $\pi : V_6 \rightarrow X_{24}$. We denote by $\sigma_{X_{24}}$ the involution in $\mathbb{W}_{X_{24}}$ induced by $\sigma_{V_6} \in \mathbb{W}_{V_6}$.

The morphism $\pi : V_6 \rightarrow X_{24}$ maps \mathbb{W}_{V_6} -orbits of irreducible \mathbb{T}_2 -invariants closed subvarieties in V_6 to the $\mathbb{W}_{X_{24}}$ -orbits of irreducible \mathbb{T}_3 -invariant closed subvarieties in X_{24} . Because of this, we will denote irreducible \mathbb{T}_3 -invariant closed subvarieties in X_{24} by the same symbols as the corresponding irreducible \mathbb{T}_2 -invariant subvarieties in V_6 .

Observe that the Fano threefold X_{24} has exactly eight \mathbb{T}_3 -fixed points, which are singular points of type $\frac{1}{2}(1, 1, 1)$. They are the images of the fixed points of the involution τ_{V_6} . One can check that the divisor class group of the threefold X_{24} is

isomorphic to $\mathbb{Z}^3 \oplus \mathbb{Z}_2$. It is generated by the images of the toric divisors in the threefold V_6 . We have

$$-2K_{X_{24}} \sim \mathcal{O}_{\mathbb{P}^{13}}(2)|_{X_{24}}$$

and $-K_{X_{24}}$ is not a Cartier divisor. This together with the adjunction formula imply that every smooth hyperplane section of the threefold $X_{24} \subset \mathbb{P}^{13}$ is an Enriques surface.

As a consequence of Corollary 4.7, we obtain.

Corollary 4.9 *Let G be a subgroup in $G_{X_{24}}$ that contains $\mathbb{W}_3^{\mathbb{Q}}$. Then $\text{rk}(\text{Cl}(X_{24})^G) = 1$. □*

5 Two equivariant Sarkisov links

In this section, we present two known toric birational maps between X_{24} and two other terminal toric Fano threefolds (see Lemmas 5.2 and 5.3 below), which will play a central role in the proof of Theorem 1.2.

Let X_8 be the complete intersection of three quadrics in \mathbb{P}^6 with homogeneous coordinates $[y_1 : y_2 : y_3 : y_4 : y_5 : y_6 : y_7]$ given by the equations

$$\begin{cases} y_7^2 - y_1y_6 = 0 \\ y_7^2 - y_2y_5 = 0 \\ y_7^2 - y_3y_4 = 0. \end{cases} \tag{5}$$

We view X_8 as a toric variety for the torus \mathbb{T}_2 , with open orbit \mathbb{T}_{X_8} that is given by

$$y_1y_2y_3y_4y_5y_6y_7 \neq 0.$$

Then X_8 has six \mathbb{T}_2 -fixed points, which are its singular points, it has twelve \mathbb{T}_2 -invariants irreducible curves, which are lines in \mathbb{P}^6 , and it has eight \mathbb{T}_2 -invariants irreducible surfaces, which are planes in \mathbb{P}^6 . The rational map $\mathbb{P}^6 \dashrightarrow V_6$ given by

$$[y_1 : y_2 : y_3 : y_4 : y_5 : y_6 : y_7] \mapsto ([y_1 : y_7], [y_2 : y_7], [y_3 : y_7]) \tag{6}$$

induces a \mathbb{T}_2 -equivariant birational map $\Phi : X_8 \dashrightarrow V_6$ whose inverse is given by

$$([u_1 : v_1], [u_2 : v_2], [u_3 : v_3]) \mapsto \left[\begin{aligned} &u_1^2W_{2,3} : u_2^2W_{1,3} : u_3^2W_{1,2} : v_3^2W_{1,2} : \\ &v_2^2W_{1,3} : v_1^2W_{2,3} : u_1u_2u_3v_1v_2v_3 \end{aligned} \right], \tag{7}$$

where $W_{i,j} = u_i v_i u_j v_j$ for every i and j in $\{1, 2, 3\}$. Moreover, there exists a commutative diagram

$$\begin{array}{ccc}
 \tilde{V}_6 & \dashrightarrow & \tilde{X}_8 \\
 \downarrow \beta & & \downarrow \alpha \\
 V_6 & \xrightarrow{\Phi^{-1}} & X_8
 \end{array} \tag{8}$$

where β is the blow-up of all eight \mathbb{T}_2 -fixed points, the top dashed arrow consists of flips in the proper transforms of the twelve \mathbb{T}_2 -invariant lines in X_8 , and α is the crepant contraction of the proper transforms of the six \mathbb{T}_2 -invariant surfaces in V_6 .

The action of \mathbb{W}_{V_6} on V_6 given in Subsection 4.2, and the formulas (6) and (7) imply that

$$\Phi^{-1} \mathbb{W}_{V_6} \Phi = \mathbb{W}_{X_8} \cong \mathbb{W}_2,$$

so that Φ is a birational $\mathbb{T}_2 \times \mathbb{W}_2$ -map. The diagram (8) is a so-called *bad* $\mathbb{T}_2 \times \mathbb{W}_2$ -Sarkisov link.

The composition $\Phi^{-1} \circ \tau_{V_6} \circ \Phi$ (see (3) for the definition of τ_{V_6}) is the biregular involution τ_{X_8} of the threefold X_8 defined by

$$[y_1 : y_2 : y_3 : y_4 : y_5 : y_6 : y_7] \mapsto [y_1 : y_2 : y_3 : y_4 : y_5 : y_6 : -y_7].$$

The projection $\mathbb{P}^6 \dashrightarrow \mathbb{P}^5$ from the point $[0 : 0 : 0 : 0 : 0 : 0 : 1]$ induces an isomorphism between the quotient X_8/τ_{X_8} and the complete intersection $V_4 \subset \mathbb{P}^5$ given by

$$\begin{cases} y_1 y_6 - y_2 y_5 = 0 \\ y_1 y_6 - y_3 y_4 = 0. \end{cases} \tag{9}$$

We view V_4 as a toric variety for the torus \mathbb{T}_3 (see Sect. 4), with open orbit \mathbb{T}_{V_4} given by

$$y_1 y_2 y_3 y_4 y_5 y_6 \neq 0,$$

so that the quotient morphism $\pi : X_8 \rightarrow V_4$ is equivariant with respect to the quotient morphism $q_{23} : \mathbb{T}_2 \rightarrow \mathbb{T}_3$.

The threefold V_4 has six \mathbb{T}_3 -fixed points, twelve irreducible \mathbb{T}_3 -invariant curves, which are lines in \mathbb{P}^5 and eight irreducible \mathbb{T}_3 -invariant surfaces which are planes in \mathbb{P}^5 . These \mathbb{T}_3 -invariant irreducible subvarieties are the images of the \mathbb{T}_2 -invariant irreducible subvarieties of X_8 by the quotient morphism $\pi : X_8 \rightarrow V_4$.

Since τ_{V_6} commutes with the action of $\mathbb{T}_2 \times \mathbb{W}_2$ on the threefold X_8 , we obtain an induced regular action of $\mathfrak{S}_4 \times \mu_2$ on the threefold V_4 . Moreover, one has $\mathbb{W}_{V_4} \cong \mathbb{W}_3$, and the threefold V_4 endowed with the action of $G_{V_4} = \mathbb{T}_3 \rtimes \mathbb{W}_{V_4}$ is another projective

terminal toric Fano model for the subgroup \mathbb{W}_3 of $GL_3(\mathbb{Z})$. As usual, we identify $\mathbb{W}_{V_4} = \mathbb{W}_3$, we let σ_{V_4} to be the involution in \mathbb{W}_{V_4} defined by

$$[y_1 : y_2 : y_3 : y_4 : y_5 : y_6] \mapsto [y_6 : y_5 : y_4 : y_3 : y_2 : y_1],$$

and we let $\nu : G_{V_4} \rightarrow \mathbb{W}_3$ be the natural homomorphism.

Lemma 5.1 ([4, Theorem 10]) *Let G be a subgroup of G_{V_4} such that $\nu(G)$ contains \mathbb{W}_3^{al} . Then $\text{rk}(\text{Cl}(V_4)^G) = 1$ if and only if $\sigma_{V_4} \in \nu(G)$ or $\nu(G) = \mathbb{W}_3^{\text{S}}$. \square*

Proof By construction, the eight irreducible \mathbb{T}_3 -invariant surfaces in V_4 are the images by $\pi : X_8 \rightarrow V_4$ of the eight irreducible \mathbb{T}_2 -invariant surfaces in X_8 . By the $\mathbb{T}_2 \times \mathbb{W}_2$ -equivariant commutative diagram (8), the latter are the images by $\alpha : \tilde{X}_8 \rightarrow X_8$ of the proper transforms of exceptional divisors of the blow-up $\beta : \tilde{V}_8 \rightarrow V_6$ of the eight \mathbb{T}_2 -fixed points of V_6 . Since the action of \mathbb{W}_3 on the character lattice of \mathbb{T}_3 is induced from that of \mathbb{W}_2 on the character lattice of \mathbb{T}_2 , it follows that $\nu(G)$ acts transitively on the eight irreducible \mathbb{T}_3 -invariant surfaces in V_4 if and only this group acts transitively on the eight \mathbb{T}_2 -fixed points of V_6 . The assertion then follows from Lemma 4.6. \square

The birational $\mathbb{T}_2 \times \mathbb{W}_2$ -map Φ in (8) induces a birational $\mathbb{T}_3 \times \mathbb{W}_3$ -map $\varphi : V_4 \dashrightarrow X_{24}$ defined by

$$[y_1 : y_2 : y_3 : y_4 : y_5 : y_6] \mapsto [y_1 y_2 y_3 : y_1 y_2 y_4 : y_1 y_3 y_4 : y_1 y_3 y_5 : y_1 y_4 y_5 : y_1 y_2 y_6 : y_1 y_3 y_6 : y_1 y_4 y_6 : y_1 y_5 y_6 : y_2 y_3 y_6 : y_2 y_4 y_6 : y_2 y_5 y_6 : y_3 y_5 y_6 : y_4 y_5 y_6], \tag{10}$$

and we eventually obtain the following:

Lemma 5.2 *There exists a $\mathbb{T}_3 \times \mathbb{W}_3$ -Sarkisov link*

$$\begin{array}{ccc}
 \tilde{X}_{20} & \overset{\rho}{\dashrightarrow} & \bar{X}_{20} \\
 \swarrow \gamma & & \searrow \delta \\
 V_4 & \overset{\varphi}{\dashrightarrow} & X_{24}
 \end{array} \tag{11}$$

where γ is the blow-up of the six singular points of the threefold V_4 , the map ρ is a composition of Atiyah flops in the proper transforms of the twelve \mathbb{T}_3 -invariant lines in V_4 , and δ is the composition of Kawamata blow-ups of the eight singular points of X_{24} . \square

Let V_2 be the hypersurface of degree 4 in $\mathbb{P}(1, 1, 1, 1, 2)$ defined by the equation

$$w^2 - x_1 x_2 x_3 x_4 = 0, \tag{12}$$

where x_1, x_2, x_3 and x_4 are coordinates of weight 1, and w is a coordinate of weight 2. We view V_2 as a toric variety for the torus \mathbb{T}_3 with open orbit \mathbb{T}_{V_2} given by

$x_1x_2x_3x_4w \neq 0$. The threefold V_2 has four \mathbb{T}_2 -fixed points, it has six \mathbb{T}_2 -invariants irreducible curves, which are singular curves of the threefold V_2 , and it has four \mathbb{T}_2 -invariants irreducible surfaces.

The rational map $\mathbb{P}(1, 1, 1, 1, 2) \dashrightarrow V_6$ given by

$$[x_1 : x_2 : x_3 : x_4 : w] \mapsto ([x_1x_2 : w], [x_1x_3 : w], [x_2x_3 : w])$$

induces a \mathbb{T}_2 -equivariant birational map $\Psi_\infty : V_2 \dashrightarrow V_6$, whose inverse is given by

$$([u_1 : v_1], [u_2 : v_2], [u_3 : v_3]) \mapsto [u_1u_2v_3 : u_1u_3v_2 : u_2u_3v_1 : v_1v_2v_3 : u_1u_2u_3v_1v_2v_3]. \tag{13}$$

With the Notation 4.3, this birational map Ψ_∞ fits in the following commutative diagram:

$$\begin{array}{ccc}
 & \tilde{V}_2 & \\
 \alpha_\infty \swarrow & & \searrow \beta_\infty \\
 V_2 & \dashrightarrow \Psi_\infty & V_6
 \end{array} \tag{14}$$

where β_∞ is the blow-up of the four points (∞, ∞, ∞) , $(0, 0, \infty)$, $(0, \infty, 0)$ and $(\infty, 0, 0)$, of X_{24} , and α is the contraction of the proper transforms of the six \mathbb{T}_2 -invariant irreducible surfaces in V_6 onto the six singular curves of V_2 .

Arguing as in the construction of Φ in Sect. 5, we see that

$$\Psi_\infty^{-1} \overline{\mathbb{W}}_2^\mathbb{G} \Psi_\infty = \mathbb{W}_{V_2},$$

where we identified $\mathbb{W}_{V_6} = \mathbb{W}_2$. Therefore, Φ_∞ is a birational $\mathbb{T}_2 \times \overline{\mathbb{W}}_2^\mathbb{G}$ -map. The diagram (14) is a *bad* $\mathbb{T}_2 \times \overline{\mathbb{W}}_2^\mathbb{G}$ -Sarkisov link.

The composition $\Psi_\infty^{-1} \circ \tau_{V_6} \circ \Psi_\infty$ is the biregular involution τ_{V_2} of V_2 defined by

$$[x_1 : x_2 : x_3 : x_4 : w] \mapsto [x_1 : x_2 : x_3 : x_4 : -w].$$

Viewing \mathbb{P}^3 as a toric variety for the torus \mathbb{T}_3 with open orbit $\mathbb{T}_{\mathbb{P}^3}$ given by $x_1x_2x_3x_4 \neq 0$, the quotient morphism $V_2 \rightarrow \mathbb{P}^3$ is equivariant for the quotient morphism $q_{23} : \mathbb{T}_2 \rightarrow \mathbb{T}_3$. We can identify $\mathbb{W}_{\mathbb{P}^3} = \mathbb{W}_{V_2} = \overline{\mathbb{W}}_3^\mathbb{G}$, so that $G_{\mathbb{P}^3} = \mathbb{T}_3 \times \overline{\mathbb{W}}_3^\mathbb{G}$.

It follows that the map Ψ_∞ in (14) induces a birational $\mathbb{T}_3 \times \overline{\mathbb{W}}_3^\mathbb{G}$ -map $\psi_\infty : \mathbb{P}^3 \dashrightarrow X_{24}$ given by

$$\begin{aligned}
 [x_1 : x_2 : x_3 : x_4] \mapsto & \left[x_1^2x_2^2x_3^2 : x_1^3x_2x_3x_4 : x_1^2x_2^2x_3x_4 : x_1x_2^3x_3x_4 : x_1^2x_2^2x_4^2 : \right. \\
 & x_1^2x_2x_3^2x_4 : x_1x_2^2x_3^2x_4 : x_1^2x_2x_3x_4^2 : x_1x_2^2x_3x_4^2 : \\
 & \left. x_1x_2x_3^3x_4 : x_1^2x_3^2x_4^2 : x_1x_2x_3^2x_4^2 : x_2^2x_2^2x_4^2 : x_1x_2x_3x_4^3 \right], \tag{15}
 \end{aligned}$$

and we eventually obtain the following:

Lemma 5.3 *There exists a $\mathbb{T}_3 \rtimes \overline{\mathbb{W}}_3^{\mathfrak{S}}$ -Sarkisov link*

$$\begin{array}{ccc}
 & X_{22} & \\
 \gamma_{\infty} \swarrow & & \searrow \delta_{\infty} \\
 \mathbb{P}^3 & \overset{\psi_{\infty}}{\dashrightarrow} & X_{24}
 \end{array} \tag{16}$$

where δ_{∞} is the composition of Kawamata blow-ups of four points in $\text{Sing}(X_{24})$ that form the $\overline{\mathbb{W}}_3^{\mathfrak{S}}$ -orbit of the singular point (∞, ∞, ∞) of X_{24} , and γ_{∞} is the contraction of the proper transforms of the six \mathbb{T}_3 -invariant surfaces in X_{24} to the six \mathbb{T}_3 -invariant lines in \mathbb{P}^3 . \square

Note that the birational map (15) is defined by the linear system consisting of all sextic surfaces that are singular along the six \mathbb{T}_3 -invariant lines in \mathbb{P}^3 . This recovers the original construction of the $\mathbb{T}_3 \rtimes \overline{\mathbb{W}}_3^{\mathfrak{S}}$ -Sarkisov link (16) that is given in [11].

Remark 5.4 Considering \mathbb{P}^3 as a toric variety for the torus \mathbb{T}_3 and considering the action of the group $\mathbb{W}_{\mathbb{P}^3} \cong \mathfrak{S}_4$ on the character lattice M_3 of \mathbb{T}_3 , we see that $\mathbb{W}_{\mathbb{P}^3} = \overline{\mathbb{W}}_3^{\mathfrak{S}}$. Then there exists the following $\mathbb{T}_3 \rtimes \overline{\mathbb{W}}_3^{\mathfrak{S}}$ -equivariant commutative diagram:

$$\begin{array}{ccccc}
 \tilde{V}_4 & \xrightarrow{\alpha} & V_4 & \xrightarrow{\sigma_{V_4}} & V_4 & \xleftarrow{\alpha} & \tilde{V}_4 \\
 \searrow \beta & & & & & & \swarrow \beta \\
 \mathbb{P}^3 & \dashrightarrow & & & & & \mathbb{P}^3
 \end{array}$$

where β is the blow-up of the four \mathbb{T}_3 -invariant points, α is the contraction of the proper transforms of the six \mathbb{T}_3 -invariant lines, and the dashed arrow is the standard Cremona involution.

6 Proof of Theorem 1.2 (infinite groups)

In this section, we give an alternative proof of Theorem 1.2(3) in the case when the group G is infinite. We will treat each of the threefolds Y_{24} , V_6 , X_{24} , V_4 and \mathbb{P}^3 in a separate subsection.

6.1 Singular Fano threefold Y_{24}

We use the notation introduced in Sect. 4.1. Let $G_{Y_{24}} = \mathbb{T}_1 \rtimes \mathbb{W}_1$, let \mathbb{W} be a subgroup in \mathbb{W}_1 that contains \mathbb{W}_1^{21} (see Notation 4.2), and let $G = \mathbb{T}_1 \rtimes \mathbb{W} \subset G_{Y_{24}}$.

Lemma 6.1 *The threefold Y_{24} is G -birationally super-rigid.* \square

Proof Suppose that Y_{24} is not G -birationally super-rigid. Then, see for instance [10, Theorem 3.3.1], there exists a G -invariant mobile linear system \mathcal{M} on the threefold Y_{24} such that the pair $(Y_{24}, \lambda\mathcal{M})$ is not canonical, where λ is a positive rational number defined by

$$\lambda\mathcal{M} \sim_{\mathbb{Q}} -K_{Y_{24}}.$$

Let Z be a G -irreducible center of non-canonical singularities of the log pair $(Y_{24}, \lambda\mathcal{M})$. Then, by Lemma 4.4, we have one of the following possibilities:

- (1) Z is the G -orbit of the singular point $(0, 0, \infty, \infty)$,
- (2) Z is the G -orbit of the smooth point $(0, \infty, \infty, \infty)$ or of the smooth point $(0, 0, 0, \infty)$,
- (3) Z is the G -orbit of the curve $(0, 0, 1, \infty)$ or of the curve $(0, 1, \infty, \infty)$.

Let us show that none of these three cases is actually possible.

Let S be the surface $(0, 1, 1, \infty)$. Then $S \cong \mathbb{P}^1 \times \mathbb{P}^1$, and the restriction $\lambda\mathcal{M}|_S$ is an effective \mathbb{Q} -linear system of bi-degree $(1, 1)$. Then S contains the singular points $(0, 0, \infty, \infty)$ and $(0, \infty, 0, \infty)$, it also contains the smooth points $(0, 0, 0, \infty)$ and $(0, \infty, \infty, \infty)$, and it also contains the curves $(0, 0, 1, \infty)$, $(0, 1, 0, \infty)$, $(0, 1, \infty, \infty)$ and $(0, \infty, 1, \infty)$.

If Z is a curve, then the multiplicity of the restriction $\lambda\mathcal{M}|_S$ at the curve Z is strictly larger than 1 by Lemma 2.7. Clearly, this is impossible, since $\lambda\mathcal{M}|_S$ has bi-degree $(1, 1)$. Thus Z must be zero dimensional.

Suppose that Z is the G -orbit of the smooth point $(0, 0, 0, \infty)$ or of the smooth point $(0, \infty, \infty, \infty)$. Denote this point by P . Then the tangent space $T_P(Y_{24})$ is an irreducible representation of the stabilizer of the point P in the group G . Thus, by Lemma 2.12, we have

$$\text{mult}_P(\mathcal{M}) > \frac{2}{\lambda}.$$

Let C be a general curve in S of bi-degree $(1, 1)$ that passes through P . Such curves span the whole surface S , so that C is not contained in the base locus of the linear system \mathcal{M} . Then, for a general surface $M \in \mathcal{M}$, we have

$$\frac{2}{\lambda} = M \cdot C \geq \text{mult}_P(\mathcal{M}) > \frac{2}{\lambda},$$

which is absurd.

It thus remains to consider the case where Z consists of singular points of the threefold Y_{24} . Let $\alpha: \tilde{Y}_{24} \rightarrow Y_{24}$ be the blow-up of the points $(0, 0, \infty, \infty)$ and $(0, \infty, 0, \infty)$, let \tilde{M} be the proper transform of a general surface in the linear system \mathcal{M} on the threefold \tilde{Y}_{24} , let E_1 and E_2 be the α -exceptional surfaces. Then

$$\lambda\tilde{M} \sim_{\mathbb{Q}} \alpha^*(-K_{Y_{24}}) - m_1E_1 - m_2E_2$$

for some rational numbers m_1 and m_2 . By Lemma 2.11, we have $m_1 > 1$ and $m_2 > 1$. Now let \tilde{C} be the proper transform on \tilde{Y}_{24} of a general curve in S of bi-degree $(1, 1)$ that passes through both points $(0, 0, \infty, \infty)$ and $(0, \infty, 0, \infty)$. Then $\tilde{C} \not\subset \tilde{M}$, so that

$$0 \leq \lambda \tilde{M} \cdot \tilde{C} = \left(\alpha^*(-K_{Y_{24}}) - m_1 E_1 - m_2 E_2 \right) \cdot \tilde{C} = 2 - m_1 - m_2 < 0,$$

which is absurd. This completes the proof of the lemma. □

Remark 6.2 Since the Fano threefold Y_{24} is $G_{Y_{24}}$ -birationally super-rigid, there is no $G_{Y_{24}}$ -Sarkisov link starting at Y_{24} . But there are *bad* $G_{Y_{24}}$ -Sarkisov links that start at Y_{24} , which implicitly appear in the proof of Lemma 6.1. For example, blowing-up all singular points of the threefold Y_{24} , we obtain the *bad* $G_{Y_{24}}$ -Sarkisov link (2).

6.2 Fano threefold $V_6 = \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$

We use the notation introduced in Sect. 4.2. Let $G_{V_6} = \mathbb{T}_2 \rtimes \mathbb{W}_2$, let \mathbb{W} be one of the subgroups in \mathbb{W}_2 that contains the group $\mathbb{W}_2^{\mathfrak{A}_1}$, and let $G = \mathbb{T}_2 \rtimes \mathbb{W} \subset G_{V_6}$.

Lemma 6.3 *The threefold V_6 is G -birationally super-rigid.* □

Proof We may assume that $\mathbb{W} = \mathbb{W}_2^{\mathfrak{A}_1}$. Suppose that V_6 is not G -birationally super-rigid. Then there exists a G -invariant mobile linear system \mathcal{M} on V_6 such that $(V_6, \lambda \mathcal{M})$ is not canonical, where λ is the positive rational number defined by $\lambda \mathcal{M} \sim_{\mathbb{Q}} -K_{V_6}$.

Let Z be a G -irreducible center of non-canonical singularities of the log pair $(V_6, \lambda \mathcal{M})$. If Z is a curve, we can assume that Z is the $\mathbb{W}_2^{\mathfrak{A}_1}$ -orbit of the \mathbb{T}_2 -invariant curve $(0, 0, 1)$. Otherwise, we can assume that Z is the $\mathbb{W}_2^{\mathfrak{A}_1}$ -orbit of the point $(0, 0, 0)$.

Let S be the surface $(0, 1, 1) \subset V_6$. Then $S \cong \mathbb{P}^1 \times \mathbb{P}^1$, and $\lambda \mathcal{M}|_S$ is an effective \mathbb{Q} -linear system of bi-degree $(2, 2)$. If Z is a curve, then

$$S \cap Z = (0, 0, 1) \cup (0, \infty, 1) \cup (0, 1, 0) \cup (0, 1, \infty).$$

We let $C_1 = (0, 0, 1)$ and $C_2 = (0, \infty, 1)$. Note that $C_1 \cap C_2 = \emptyset$. If Z is a point, then

$$S \cap Z = (0, 0, 0) \cup (0, \infty, \infty).$$

We let $P_1 = (0, 0, 0)$ and $P_2 = (0, \infty, \infty)$.

If Z is a curve, then it follows from Lemma 2.7 that

$$\text{mult}_{C_1}(\mathcal{M}) = \text{mult}_{C_2}(\mathcal{M}) > \frac{1}{\lambda}$$

This implies that the coefficient of these curves in the restriction $\lambda \mathcal{M}|_S$ is larger than 1, contradicting the fact that $\lambda \mathcal{M}|_S$ is of bi-degree $(2, 2)$. Thus Z must be zero dimensional.

Since the stabilizer of the point P_1 in the group G contains a subgroup μ_3 that permutes transitively the \mathbb{T}_2 -invariant curves that pass through P_1 , the tangent space $T_{P_1}(X)$ is an irreducible three-dimensional representation of the stabilizer of P_1 . Therefore, it follows from Lemma 2.12 that


$$\text{mult}_{P_1}(\mathcal{M}) = \text{mult}_{P_2}(\mathcal{M}) > \frac{2}{\lambda}.$$

Let C be a general curve in the surface S of bi-degree $(1, 1)$ that passes through P_1 and P_2 . Such curves span the whole surface S , so that the curve C is not contained in the base locus of the linear system \mathcal{M} . Thus, for a general surface $M \in \mathcal{M}$, we have

$$4 = \lambda M \cdot C \geq \lambda \left(\text{mult}_{P_1}(\mathcal{M}) + \text{mult}_{P_2}(\mathcal{M}) \right) > 4,$$

which is absurd. This completes the proof. □


Remark 6.4 Since threefold V_6 is G_{V_6} -birationally super-rigid, there is no G_{V_6} -Sarkisov link that starts at V_6 . However, there exist *bad* G_{V_6} -Sarkisov links that start at V_6 . For instance, blowing-up all \mathbb{T}_2 -invariant points leads to the *bad* G_{V_6} -Sarkisov link (14). Likewise, the G_{V_6} -equivariant symbolic blow-up α of the union of all \mathbb{T}_2 -invariant curves (see Example 2.3) also leads to a *bad* G_{V_6} -Sarkisov link:



where β is the contraction of the proper transforms of the G_{V_6} -invariant surfaces in V_6 . One can show that X_{12} is the canonical toric Fano threefold No 9099 in [7], which can also be obtained as the quotient of the singular Fano threefold Y_{24} , viewed as a toric variety for the action of the torus \mathbb{T}_1 , by an involution that fixes only \mathbb{T}_1 -invariant points.

6.3 Singular Fano threefolds V_4 and X_{24}

We now treat the threefolds V_4 and X_{24} . We use the notation of Sects. 4.3 and 5, and we identify $G_{V_4} = G_{X_{24}} = \mathbb{T}_3 \rtimes \mathbb{W}_3$. By Sect. 5, we have a $\mathbb{T}_3 \rtimes \mathbb{W}_3$ -Sarkisov link



where γ is the blow-up of the eight singular points of V_4 , the map ρ is a composition of Atiyah flops in the proper transforms of the twelve \mathbb{T}_3 -invariant lines in V_4 , and δ is the composition of Kawamata blow-ups of the eight singular points of X_{24} .

Let \mathbb{W} be a subgroup of \mathbb{W}_3 that contains $\mathbb{W}_3^{\mathfrak{A}}$ such that either $\mathbb{W} = \mathbb{W}_3^{\mathfrak{A}}$, or \mathbb{W} contains the involution σ (see Notation 4.2), and let $G = \mathbb{T}_3 \rtimes \mathbb{W}$. The proof of the following lemma is straightforward.

Lemma 6.5 *Let $\mathcal{M}_{X_{24}}$ be a G -invariant mobile linear system on X_{24} , and let \mathcal{M}_{V_4} be its proper transform on V_4 via φ . Then the following assertions hold:*

(1) *There are $k \in \mathbb{Z}_{\geq 0}$ and $m \in \frac{1}{2}\mathbb{Z}_{\geq 0}$ such that $\mathcal{M}_{X_{24}} \sim_{\mathbb{Q}} -kK_{X_{24}}$ and*

$$\delta_*^{-1}(\mathcal{M}_{X_{24}}) \sim_{\mathbb{Q}} \delta^*(-kK_{X_{24}}) - m \sum_{i=1}^8 F_i,$$

where each F_i is a δ -exceptional surface.

(2) *There are $n \in \mathbb{Z}_{\geq 0}$ and $m' \in \mathbb{Z}_{\geq 0}$ such that $\mathcal{M}_{V_4} \sim n\mathcal{H}_{V_4}$ and*

$$\gamma_*^{-1}(\mathcal{M}_{V_4}) \sim_{\mathbb{Q}} \gamma^*(n\mathcal{H}_{V_4}) - m' \sum_{i=1}^6 E_i,$$

where \mathcal{H}_{V_4} is a hyperplane section of V_4 , and each E_i is a γ -exceptional surface.

(3) *One has $n = 3k - 2m$ and $k = n - m'$.*

□

Now we are ready to prove

Proposition 6.6 *One has $\mathcal{P}_G(V_4) = \{V_4, X_{24}\}$.*

Proof Let $\chi : X_{24} \dashrightarrow Y$ be a G -equivariant birational map such that Y is a threefold with terminal singularities, and there exists a G -equivariant morphism $f : Y \rightarrow Z$ that is a G -Mori fiber space. Fix a sufficiently large positive integer $n \gg 0$. Let D_Z be a sufficiently general very ample divisor on Z , let $\mathcal{M}_Y = |-nK_Y + f^*(D_Z)|$, let

$$\mathcal{M}_{X_{24}} = (\chi)_*^{-1}(\mathcal{M}_Y)$$

and let

$$\mathcal{M}_{V_4} = (\chi \circ \varphi)_*^{-1}(\mathcal{M}_Y).$$

Then $\mathcal{M}_{X_{24}}$ and \mathcal{M}_{V_4} are G -invariant mobile linear systems on X_{24} and V_4 , respectively. Let k and n be the non-negative integers such that $\mathcal{M}_{X_{24}} \sim_{\mathbb{Q}} -kK_{X_{24}}$ and $\mathcal{M}_{V_4} \sim n\mathcal{H}_{V_4}$. If the log pair $(X_{24}, \frac{1}{k}\mathcal{M}_{X_{24}})$ has canonical singularities, then it follows from the Noether–Fano inequality that χ is an isomorphism, see [10, Theorems 3.2.1 and 3.2.6] or [12]. Similarly, if the log pair $(V_4, \frac{2}{n}\mathcal{M}_{V_4})$ has canonical singularities, then the birational map $\chi \circ \varphi$ is an isomorphism. Therefore, to prove the

required assertion, it is enough to show that either $(X_{24}, \frac{1}{k}\mathcal{M}_{X_{24}})$ or $(V_4, \frac{2}{n}\mathcal{M}_{V_4})$ has canonical singularities.

Suppose that the singularities of the log pair $(X_{24}, \frac{1}{k}\mathcal{M}_{X_{24}})$ are worse than canonical. Then, using Lemma 2.10 and Lemma 6.5, we obtain the inequality $n < 2k$. Let us show then that the log pair $(V_4, \frac{2}{n}\mathcal{M}_{V_4})$ has canonical singularities.

By construction of φ , if the singularities of the log pair $(V_4, \frac{2}{n}\mathcal{M}_{V_4})$ are not canonical, then the union of its centers of non-canonical singularities is either the union of all singular points of the threefold V_4 , or a \mathbb{W} -orbit of \mathbb{T}_3 -invariant curves, which are lines in \mathbb{P}^5 .

In the first case, using Lemma 2.11 and Lemma 6.5, we get $n > 2k$, which is impossible, since we already proved that $n < 2k$.


The twelve \mathbb{T}_3 -invariant lines in V_4 form a unique \mathbb{W} -irreducible curve. Suppose that all of them are centers of non-canonical singularities of the log pair $(V_4, \frac{2}{n}\mathcal{M}_{V_4})$. Then

$$\text{mult}_L(\mathcal{M}_{V_4}) > \frac{n}{2} \tag{17}$$

for each such line L by Lemma 2.7. On the other hand, each of the eight \mathbb{T}_3 -invariant planes in V_4 contains three \mathbb{T}_3 -invariant lines. Thus, restricting \mathcal{M}_{V_4} on one such plane, we obtain a contradiction to (17). \square

6.4 Fano threefolds \mathbb{P}^3 and X_{24}


Finally, we deal with the Fano threefolds \mathbb{P}^3 and X_{24} . For X_{24} , we use the same notation as in Sect. 6.3. As in Sect. 5, we view \mathbb{P}^3 as a toric variety for the torus \mathbb{T}_3 , and we identify $\mathbb{W}_{\mathbb{P}^3} = \overline{\mathbb{W}}_3^{\mathfrak{S}}$ and $G_{\mathbb{P}^3} = \mathbb{T}_3 \rtimes \overline{\mathbb{W}}_3^{\mathfrak{S}}$. In Sect. 5, we constructed the following $\mathbb{T}_3 \rtimes \overline{\mathbb{W}}_3^{\mathfrak{S}}$ -Sarkisov link:



where δ_∞ is a composition of Kawamata blow-ups of the four singular points of X_{24} that form the $\overline{\mathbb{W}}_3^{\mathfrak{S}}$ -orbit of the point (∞, ∞, ∞) in X_{24} , and γ_∞ is the contraction of the proper transforms of the six \mathbb{T}_3 -invariant surfaces in X_{24} to the six \mathbb{T}_3 -invariant lines in \mathbb{P}^3 .

Recall subsection 4.3 that the regular involution $\sigma_{X_{24}}$ of $X_{24} = V_6/\tau_{V_6}$ induced by the involution $\sigma_{V_6} = \nu \times \nu \times \nu$ (see Notation 4.3) commutes with the action of $\overline{\mathbb{W}}_3^{\mathfrak{S}}$. We thus obtain a second birational $\mathbb{T}_3 \rtimes \overline{\mathbb{W}}_3^{\mathfrak{S}}$ -map $\psi_0 = \sigma_{X_{24}} \circ \psi_\infty : \mathbb{P}^3 \dashrightarrow X_{24}$. Note that $\sigma_{X_{24}}(\infty, \infty, \infty) = (0, 0, 0)$. Consequently, we have a second $\mathbb{T} \rtimes \overline{\mathbb{W}}_3^{\mathfrak{S}}$ -Sarkisov

link




where δ_0 is a composition of Kawamata blow-ups of the four singular points of X_{24} which form the $\overline{\mathbb{W}}_3^G$ -orbit of the point $(0, 0, 0)$, and γ_0 is the contraction of the proper transforms of the six \mathbb{T}_3 -invariant surfaces in X_{24} to the six \mathbb{T}_3 -invariant lines in \mathbb{P}^3 .

Remark 6.7 Using (4) and (15), one can show that $\psi_\infty^{-1} \circ \sigma_{X_{24}} \circ \psi_\infty$ is equal to the standard Cremona involution $\sigma_{\mathbb{P}^3} : \mathbb{P}^3 \dashrightarrow \mathbb{P}^3$ defined by

$$[x_1 : x_2 : x_3 : x_4] \mapsto [x_2x_3x_4 : x_1x_3x_4 : x_1x_2x_4 : x_1x_2x_3]. \tag{18}$$

In other words, we have a commutative diagram of birational $\mathbb{T} \rtimes \overline{\mathbb{W}}_3^G$ -maps



Let E be the sum of all γ -exceptional surfaces, let F_0 be the sum of all δ_0 -exceptional surfaces, let F_∞ be the sum of all δ_∞ -exceptional surfaces, and let \mathbb{W} be either \mathbb{W}_3^{21} or $\overline{\mathbb{W}}_3^G$. We also let $G = \mathbb{T}_3 \rtimes \mathbb{W}$. The next lemma is straightforward.

Lemma 6.8 *Let $\mathcal{M}_{X_{24}}$ be a G -invariant mobile linear system on the Fano threefold X_{24} , let $\mathcal{M}_{X_{22},\infty}$ and $\mathcal{M}_{X_{22},0}$ be its proper transforms on X_{22} via δ_∞ and δ_0 , respectively, and let $\mathcal{M}_{\mathbb{P}^3,\infty}$ and $\mathcal{M}_{\mathbb{P}^3,0}$ be its proper transforms on \mathbb{P}^3 via ψ_∞ and ψ_0 , respectively. Furthermore, let $k \in \frac{1}{2}\mathbb{Z}$ and let n_∞ and n_0 be integers such that*

$$\begin{cases} \mathcal{M}_{X_{24}} \sim_{\mathbb{Q}} -kK_{X_{24}}, \\ \mathcal{M}_{\mathbb{P}^3,\infty} \sim_{\mathbb{Q}} n_\infty H, \\ \mathcal{M}_{\mathbb{P}^3,0} \sim_{\mathbb{Q}} n_0 H. \end{cases}$$

where H is a hyperplane in \mathbb{P}^3 . Then the following assertions hold:

(1) *There are m_0 and m_∞ in $\frac{1}{2}\mathbb{Z}_{\geq 0}$ such that*

$$\begin{cases} \mathcal{M}_{X_{22},0} \sim_{\mathbb{Q}} \delta_0^*(-kK_{X_{24}}) - m_0F_0, \\ \mathcal{M}_{X_{22},\infty} \sim_{\mathbb{Q}} \delta_\infty^*(-kK_{X_{24}}) - m_\inftyF_\infty. \end{cases}$$

(2) There are m'_0 and m'_∞ in $\mathbb{Z}_{\geq 0}$ such that

$$\begin{cases} \mathcal{M}_{X_{22},0} \sim_{\mathbb{Q}} \gamma_0^*(n_0\mathcal{H}) - m'_0E, \\ \mathcal{M}_{X_{22},\infty} \sim_{\mathbb{Q}} \gamma_\infty^*(n_\infty\mathcal{H}) - m'_\infty E. \end{cases}$$

(3) Furthermore, one has

$$\begin{cases} n_0 = 6k - 4m_0, \\ n_\infty = 6k - 4m_\infty, \\ n_0 = 3n_\infty - 4m'_\infty, \\ k = \frac{n_0}{2} - m'_0, \\ k = \frac{n_\infty}{2} - m'_\infty. \end{cases}$$

□

Now we are ready to prove:

Proposition 6.9 *One has $\mathcal{P}_G(\mathbb{P}^3) = \{\mathbb{P}^3, X_{24}\}$.*

Proof Let $\mathcal{M}_{X_{24}}$ be a G -invariant mobile linear system on the threefold X_{24} . With the notation of Lemma 6.8, if the log pair $(X_{24}, \frac{1}{k}\mathcal{M}_{X_{24}})$ does not have canonical singularities, then, combining Lemmas 2.10 and 6.8, we obtain that

- either $(X_{24}, \frac{1}{k}\mathcal{M}_{X_{24}})$ is not canonical at the point $(0, 0, 0)$ and $n_0 < 4k$,
- or $(X_{24}, \frac{1}{k}\mathcal{M}_{X_{24}})$ is not canonical at the point (∞, ∞, ∞) and $n_\infty < 4k$.

If $(\mathbb{P}^3, \frac{4}{n_\infty}\mathcal{M}_{\mathbb{P}^3_\infty})$ does not have canonical singularities, we let Z be its G -irreducible center of non-canonical singularities. In this case, one of the following cases holds:

- Z is the $G_{\mathbb{P}^3}$ -irreducible curve and $k < \frac{n_\infty}{4}$;
- Z is the $G_{\mathbb{P}^3}$ -orbit of length 4 and $n_0 < n_\infty$.

Indeed, if Z is the $G_{\mathbb{P}^3}$ -irreducible curve, then $m'_\infty > \frac{n_\infty}{4}$ by Lemma 2.7, so that

$$k = \frac{n_\infty}{2} - m'_\infty < \frac{n_\infty}{4},$$

by Lemma 6.8. Similarly, if Z is the $G_{\mathbb{P}^3}$ -orbit of length 4, then

$$m'_\infty > \frac{n_\infty}{2}$$

by Lemma 2.12, because the tangent space $T_P(\mathbb{P}^3)$ at a point $P \in Z$ is an irreducible representation of the stabilizer of the point P in the group G . Thus, in this case, we have

$$n_0 = 3n_\infty - 4m'_\infty < n_\infty$$

by Lemma 6.8.

Now, we let q be the smallest number among $\frac{n_\infty}{4}$, $\frac{n_0}{4}$ and k . Without loss of generality, we may assume that

$$q = \min \left\{ \frac{n_\infty}{4}, k \right\}.$$

In view of the above alternatives, we obtain the following:

- if $q = \frac{n_\infty}{4}$, then $(\mathbb{P}^3, \frac{4}{n_\infty} \mathcal{M}_{\mathbb{P}^3, \infty})$ has canonical singularities;
- if $q = k$, then $(X_{24}, \frac{1}{k} \mathcal{M}_{X_{24}})$ has canonical singularities.

Now, arguing as in the proof of Proposition 6.6, we deduce that \mathbb{P}^3 and X_{24} are the only G -Mori fibre spaces G -birational to \mathbb{P}^3 . □

7 Proof of Theorem 1.2 (finite groups)

All assertions of Theorem 1.2 follow from the results of Sects. 3 and 4 except for the part (3), which has been already proved in Sects. 3 and 6 for infinite groups. The aim of this section is to prove Theorem 1.2(3) for finite groups. To do this, we need some results on finite subgroups of the groups $\mathbb{T}_1 \rtimes \mathbb{W}_1$, $\mathbb{T}_2 \rtimes \mathbb{W}_2$ and $\mathbb{T}_3 \rtimes \mathbb{W}_3$.

7.1 Finite subgroups

We use the notation of Sect. 4.

Lemma 7.1 *Let \mathbb{W} be a subgroup of the finite group \mathbb{W}_2 that contains the group \mathbb{W}_2^{2l} , and let G be a \mathbb{W} -invariant finite subgroup of \mathbb{T}_2 . Then there exists $n \in \mathbb{N}$ such that one of the following three possibilities holds:*

- (1) $G \cong \mu_n^3$;
- (2) n is even and $G \cong \mu_n^2 \times \mu_{\frac{n}{2}}$;
- (3) n is even and $G \cong \mu_n \times \mu_{\frac{n}{2}}$.

□

Proof Let n be the maximal order of elements in G , and let h be an element in G that has maximal order. Then the order of every element of G divides n . This implies that $G \subseteq \mu_n^3$. Thus, there are positive integers a, b, c such that $\gcd(a, b, c, n) = 1$ and

$$h = (\epsilon^a, \epsilon^b, \epsilon^c)$$

for some primitive n th root of unity ϵ .

With respect to the basis f_1, f_2, f_3 of the lattice M_2 , the subgroup $\mathbb{W}_2^{2l} \subset \text{GL}_3(\mathbb{Z})$ is generated by permutation matrices and the matrices

$$A = \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad \text{and} \quad B = \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}.$$

Applying cyclic permutations of order 3 to h , we see that

$$(\epsilon^c, \epsilon^a, \epsilon^b) \in G \ni (\epsilon^b, \epsilon^c, \epsilon^a),$$

so that the group G contains an element of the form $(\epsilon, \epsilon^\beta, \epsilon^\gamma)$ for some integers β and γ . Thus, we may assume that $a = 1$. Applying AB to the element h , we see that

$$h' = (\epsilon, \epsilon^{-b}, \epsilon^{-c}) \in G.$$

It follows that $hh' = (\epsilon^2, 1, 1)$ and its transforms $(1, \epsilon^2, 1)$ and $(1, 1, \epsilon^2)$ by permutations matrices in \mathbb{W} are also contained in the group G .

Let $h_2 = (1, \epsilon^2, 1)$ and $h_3 = (1, 1, \epsilon^2)$. If n is odd then we can replace h by

$$hh_2^\beta h_3^\gamma = (\epsilon, 1, 1),$$

where β and γ are integers such that $\beta b \equiv 1 \pmod n$ and $\gamma c \equiv 1 \pmod n$. If n is even, we can replace h by $hh_2^\beta h_3^\gamma \in G$ for $\beta = -\lfloor \frac{b}{2} \rfloor$ and $\gamma = -\lfloor \frac{c}{2} \rfloor$. Therefore, we may assume that one of the following three cases holds: $(b, c) = (0, 0)$, $(b, c) = (1, 0)$ and $(b, c) = (1, 1)$.

If $(b, c) = (0, 0)$, then $h = (\epsilon, 1, 1)$ from which it follows that $G = \mu_n^3$.

If $(b, c) = (1, 0)$, then n is even and $h = (\epsilon, \epsilon, 1)$. In this case, applying permutations, we get that $(1, \epsilon, \epsilon) \in G$ and hence that G contains the subgroup

$$G' = \langle (\epsilon, \epsilon, 1), (1, \epsilon, \epsilon), (1, 1, \epsilon^2) \rangle \cong \mu_n^2 \times \mu_{\frac{n}{2}}.$$

Since $G \subset \mu_n^3$, it follows that either $G = G'$ or $G \cong \mu_n^3$.

Finally, if $(b, c) = (1, 1)$, then G contains the subgroup

$$G' = \langle (\epsilon, \epsilon, \epsilon), (1, \epsilon^2, 1), (1, 1, \epsilon^2) \rangle \cong \mu_n \times \mu_{\frac{n}{2}}^2,$$

and hence either $G = G'$ or $G \cong \mu_n^3$, which completes the proof. □

Corollary 7.2 *Let \mathbb{W} be a subgroup of the finite group \mathbb{W}_1 that contains the group \mathbb{W}_1^{2l} , and let G be a \mathbb{W} -invariant finite subgroup of \mathbb{T}_1 . Then there exists $n \in \mathbb{N}$ such that one of the following five possibilities holds:*

- (1) $G \cong \mu_n^3$;
- (2) n is even and $G \cong \mu_n^2 \times \mu_{\frac{n}{2}}$;
- (3) n is even and $G \cong \mu_n \times \mu_{\frac{n}{2}}^2$;

- (4) n is divisible by 4 and $G \cong \mu_n \times \mu_{\frac{n}{2}} \times \mu_{\frac{n}{4}}$;
- (5) n is divisible by 4 and $G \cong \mu_n \times \mu_{\frac{n}{4}}^2$.

□

Proof Let $q_{12}: \mathbb{T}_1 \rightarrow \mathbb{T}_2$, be the quotient map that corresponds to the inclusion $M_2 \hookrightarrow M_1$ described in Sect. 4. Then q_{12} is given by

$$(\hat{t}_1, \hat{t}_2, \hat{t}_3) \mapsto (\hat{t}_1 \hat{t}_2, \hat{t}_1 \hat{t}_3, \hat{t}_2 \hat{t}_3),$$

and its kernel consists of two elements $\pm(1, 1, 1)$. Let \overline{G} be the image of G by the map q_{12} . Then either $G \cong \overline{G}$ or there exists an exact sequence of groups

$$1 \longrightarrow \mu_2 \longrightarrow G \longrightarrow \overline{G} \longrightarrow 1.$$

Since the \mathbb{W}_2 -action on M_2 is induced by the restriction of the \mathbb{W}_1 -action on M_1 , we see that \overline{G} is normalized by the action of the subgroup \mathbb{W}_2^{al} . By Lemma 7.1, we obtain that

- (1) either $\overline{G} \cong \mu_n^3$,
- (2) or n is even and $\overline{G} \cong \mu_n^2 \times \mu_{\frac{n}{2}}$,
- (3) or n is even and $\overline{G} \cong \mu_n \times \mu_{\frac{n}{2}}^2$.

This immediately implies the result. □

Corollary 7.3 (cf. [19]) *Let \mathbb{W} be a subgroup of the finite group \mathbb{W}_3 that contains \mathbb{W}_3^{al} , and let G be a \mathbb{W} -invariant finite subgroup of \mathbb{T}_3 . Then there exists $n \in \mathbb{N}$ such that one of the following three possibilities holds:*

- (1) $G \cong \mu_n^3$;
- (2) n is even and $G \cong \mu_n^2 \times \mu_{\frac{n}{2}}$;
- (3) n is divisible by 4 and $G \cong \mu_n^2 \times \mu_{\frac{n}{4}}$.

□

Proof Let $q_{23}: \mathbb{T}_2 \rightarrow \mathbb{T}_3$ be the quotient by the involution $(\bar{t}_1, \bar{t}_2, \bar{t}_3) \mapsto (-\bar{t}_1, -\bar{t}_2, -\bar{t}_3)$, which corresponds to the inclusion of lattices $M_3 \hookrightarrow M_2$ described in Sect. 4, and let \widehat{G} be the preimage of the group G via q_{23} . Then \widehat{G} is a finite subgroup in \mathbb{T}_2 , which is normalized by the group \mathbb{W}_2^{al} . Since \widehat{G} contains $\pm(1, 1, 1)$, we see that $|\widehat{G}|$ is even.

It follows from the proof of Lemma 7.1 that there exist an integer $m \in 2\mathbb{N}$ and a primitive m -th root of unity ϵ such that \widehat{G} is one of the following subgroups:

- (1) $\langle (\epsilon, 1, 1), (1, \epsilon, 1), (1, 1, \epsilon) \rangle \cong \mu_m^3$,
- (2) $\langle (\epsilon, \epsilon, 1), (1, \epsilon, \epsilon), (1, 1, \epsilon^2) \rangle \cong \mu_m^2 \times \mu_{\frac{m}{2}}$,
- (3) $\langle (\epsilon, \epsilon, \epsilon), (1, \epsilon^2, 1), (1, 1, \epsilon^2) \rangle \cong \mu_m \times \mu_{\frac{m}{2}}^2$,

Thus, in the first case, we have

$$G = \langle (\epsilon, 1, \epsilon), (\epsilon, \epsilon^{-1}, 1), (1, \epsilon^{-1}, \epsilon) \rangle = \langle (\epsilon, 1, \epsilon), (1, \epsilon, \epsilon), (1, \epsilon^2, 1) \rangle \cong \mu_m^2 \times \mu_{\frac{m}{2}}.$$

Similarly, in the second case, we have

$$G = \langle (\epsilon^2, \epsilon^{-1}, \epsilon), (\epsilon, \epsilon^{-2}, \epsilon), (1, \epsilon^{-2}, \epsilon^2) \rangle \\ = \langle (\epsilon, 1, \epsilon^{-1}), (1, \epsilon, \epsilon), (1, 1, \epsilon^4) \rangle = \mu_m^2 \times \mu_{\frac{m}{4}},$$

where n is divisible by 4, because $(-1, -1, -1) \in \widehat{G}$. Finally, in the third case, we have

$$G = \langle (\epsilon^2, \epsilon^{-2}, \epsilon^2), (\epsilon^2, \epsilon^2, 1), (1, \epsilon^{-2}, \epsilon^2) \rangle = \langle (\epsilon^2, 1, 1), (1, \epsilon^2, 1), (1, 1, \epsilon^2) \rangle = \mu_{\frac{m}{2}}^3.$$

This completes the proof of the corollary. □

7.2 The proof of Theorem 1.2(3) for finite groups

Let X be one of the threefolds $V_6 = \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$, X_{24} , Y_{24} , V_4 or \mathbb{P}^3 , let \mathbb{T} be a maximal torus in $\text{Aut}(X)$, and let G_X be its normalizer in $\text{Aut}(X)$. Using the split exact sequence of groups

$$1 \longrightarrow \mathbb{T} \longrightarrow G_X \xrightarrow{\nu_X} \mathbb{W}_X \longrightarrow 1,$$

we consider the Weyl group \mathbb{W}_X as a subgroup of the group G_X .

Let G be a finite subgroup of the group G_X , let $\mathbb{W} = \nu_X(G)$, and let $\Gamma = \mathbb{T} \cap G$. Suppose that the group \mathbb{W} contains the unique subgroup in \mathbb{W}_X that is isomorphic to \mathcal{A}_4 . Then Lemma 7.1 and Corollaries 7.2 and 7.3 imply the following:

Corollary 7.4 *The group Γ contains a subgroup $\Gamma' \cong \mu_n^3$ such that $|\Gamma' : \Gamma| \leq 16$. □*

We have $G_X = \langle \mathbb{T}, \mathbb{W}_X \rangle \cong \mathbb{T} \rtimes \mathbb{W}_X$ and $G = \langle \Gamma, \mathbb{W} \rangle \cong \Gamma \rtimes \mathbb{W}$. Note that

$$\text{rk}(\text{Cl}(X)^G) = \text{rk}(\text{Cl}(X)^{\mathbb{W}}),$$

so that X is G -minimal if and only if it is \mathbb{W} -minimal. Now we suppose that X is \mathbb{W} -minimal. To prove Theorem 1.2(3), we have to show that X is G -solid if $|G| \geq 32 \cdot 24^4$.

Suppose that $|\Gamma| \geq 16 \cdot 24^3$. Note that this inequality follows from $|G| \geq 32 \cdot 24^4$. By Corollary 7.4, the group Γ contains a subgroup that is isomorphic to μ_n^3 for $n \geq 24$. Let us prove that X is G -solid.

Let $\mathbf{G} = \langle \mathbb{T}_X, \mathbb{W} \rangle \cong \mathbb{T}_X \rtimes \mathbb{W}$. In Sect. 6, we proved that the threefold X is \mathbf{G} -solid. Moreover, this proof implies that X is G -solid provided that the following condition is satisfied:

- (★) For every non-empty G -invariant mobile linear system \mathcal{M} on the threefold X , all non-canonical centers (if any) of the mobile log pair $(X, \lambda\mathcal{M})$ are \mathbb{T}_X -invariant, where λ is a positive rational number such that

$$\lambda\mathcal{M} \sim_{\mathbb{Q}} -K_X.$$

Moreover, if a \mathbb{T}_X -invariant smooth point P of the threefold X is a non-canonical center of the log pair $(X, \lambda\mathcal{M})$, then $\text{mult}_P(\mathcal{M}) > \frac{2}{\lambda}$.

In the remaining part of this section, we will prove that \star holds.

Let \mathcal{M} be a non-empty G -invariant mobile linear system on X , and let λ be a positive rational number such that $\lambda\mathcal{M} \sim_{\mathbb{Q}} -K_X$.

Lemma 7.5 *Let P be a smooth \mathbb{T} -invariant point of the toric Fano threefold X such that P is a non-canonical center of the log pair $(X, \lambda\mathcal{M})$. Then $\text{mult}_P(\mathcal{M}) > \frac{2}{\lambda}$. \square*

Proof Since P is a \mathbb{T} -invariant smooth point of X , we have $X = V_6$, $X = Y_{24}$ or $X = \mathbb{P}^3$. Let G_P be the stabilizer of the point P in G . Then the tangent space $T_P(X)$ is a faithful representation of the group G_P by Lemma 2.4. Moreover, this representation is irreducible, because G_P contains Γ . Thus, the assertion follows from Lemma 2.12. \square

Now we suppose that $(X, \lambda\mathcal{M})$ is not canonical and we let Z be a non-canonical G -center. To complete the proof, we have to show that Z is \mathbb{T} -invariant. In what follows, we denote by I_X the Fano index of the threefold X .

Lemma 7.6 *If Z is a curve, then Z is \mathbb{T} -invariant. \square*

Proof Let H be an ample Cartier divisor on X such that $-K_X \sim I_X H$. Then

$$\frac{H^3 I_X^2}{\lambda^2} = H \cdot M_1 \cdot M_2 \geq (H \cdot Z) \text{mult}_Z^2(\mathcal{M}) > \frac{H \cdot Z}{\lambda^2}$$

for two general surfaces M_1 and M_2 in \mathcal{M} , because $\text{mult}_Z(\mathcal{M}) > \frac{1}{\lambda}$ by Lemma 2.7. Then

$$H \cdot Z < H^3 I_X^2 \leq 24.$$

But H is very ample, and the group Γ contains a subgroup isomorphic to μ_n^3 for $n \geq 24$. Thus, the curve Z must be \mathbb{T} -invariant by Lemma 2.13. \square

Thus, we may assume that Z is the G -orbit of a point in X .

Lemma 7.7 *The G -orbit Z is contained in the union of \mathbb{T} -invariant curves. \square*

Proof Suppose that Z is not contained in the union of \mathbb{T} -invariant curves. Let us seek for a contradiction. Observe that the G -orbit Z is a G -center of non-log canonical singularities of the log pair $(X, 2\lambda\mathcal{M})$. We claim that Z is an isolated center of non-log canonical singularities of this log pair. Indeed, suppose that there is a G -irreducible curve C that is a center of non-log canonical singularities of the log pair $(X, 2\lambda\mathcal{M})$. Let M_1 and M_2 be general surfaces in \mathcal{M} . Then

$$M_1 \cdot M_2 = mC + \Omega,$$

where m is a non-negative integer, and Ω is an effective one-cycle whose support does not contain C . Then $m > \frac{1}{\lambda^2}$ by Lemma 2.8.

Let H be an ample Cartier divisor on X such that $-K_X \sim I_X H$. Then

$$\frac{H^3 I_X^2}{\lambda^2} = H \cdot M_1 \cdot M_2 = mH \cdot C + H \cdot \Omega \geq mH \cdot C > \frac{H \cdot C}{\lambda^2},$$

so that $H \cdot C < H^3 I_X^2 \leq 24$. Thus, the curve C must be \mathbb{T}_X -invariant by Lemma 2.13. Since Z is not contained in the union of \mathbb{T} -invariant curves, we see that Z is an isolated center of non-log canonical singularities of the log pair $(X, 2\lambda\mathcal{M})$.

Let μ be a positive rational number such that $\mu < \lambda$, and Z is an isolated G -irreducible center of log canonical singularities of the log pair $(X, 2\mu\mathcal{M})$. Let \mathcal{I} be the multiplier ideal sheaf of the log pair $(X, 2\mu\mathcal{M})$. Then the ideal \mathcal{I} defines a subscheme Z' in X whose support contains Z . Using Nadel vanishing theorem (see [25, Theorem 9.4.8]), we get

$$h^1(\mathcal{I} \otimes \mathcal{O}_X(-K_X)) = 0.$$

Now using the exact sequence of sheaves

$$0 \rightarrow \mathcal{I} \otimes \mathcal{O}_X(-K_X) \rightarrow \mathcal{O}_X(-K_X) \rightarrow \mathcal{O}_{Z'} \otimes \mathcal{O}_X(-K_X) \rightarrow 0,$$

we see that $|Z| \leq h^0(\mathcal{O}_X(-K_X))$. But on the other hand, since Z is not contained in the union of \mathbb{T} -invariant curves, we have $|Z| \geq n^2 \geq 24^2$, a contradiction. \square

Finally, we prove the following lemma:

Lemma 7.8 *The G -orbit Z is \mathbb{T} -invariant.* \square

Proof We know from Lemma 7.7 that Z is contained in union of \mathbb{T} -invariant curves. Suppose that Z is not \mathbb{T} -invariant. Let us seek for a contradiction.

Let H be an ample Cartier divisor on X such that $-K_X \sim I_X H$. Recall that H is very ample,

$$I_X = \begin{cases} 2 & \text{if } X = V_6, \\ 1 & \text{if } X = X_{24}, \\ 1 & \text{if } X = Y_{24}, \\ 2 & \text{if } X = V_4, \\ 4 & \text{if } X = \mathbb{P}^3. \end{cases}$$

and

$$H^3 = \begin{cases} 6 & \text{if } X = V_6, \\ 24 & \text{if } X = X_{24}, \\ 24 & \text{if } X = Y_{24}, \\ 4 & \text{if } X = V_4, \\ 1 & \text{if } X = \mathbb{P}^3. \end{cases}$$

Let \mathcal{C} the union of all \mathbb{T} -invariant curves. Then \mathcal{C} is G -invariant. Moreover, it follows from Lemmas 4.4 and 4.6 that \mathbb{W} acts transitively on the set of irreducible components of \mathcal{C} except the following two cases:

- (1) $X = Y_{24}$ and $\mathbb{W} = \mathbb{W}_1^{24}$,
- (2) $X = Y_{24}$ and $\mathbb{W} = \overline{\mathbb{W}}_1^{\mathfrak{S}}$.

In these two cases, the curve \mathcal{C} splits as a union of two G -irreducible curves that are swapped by the group $\mathbb{W}_X \cong \mathfrak{S}_4 \times \mu_2$. Thus, we let \mathcal{C}_1 be the G -irreducible component of the curve \mathcal{C} that contains the G -orbit Z . Then, because Z is G -irreducible, the other G -irreducible component of the curve \mathcal{C} (if any) does not contain Z ,

Let $d = H \cdot \mathcal{C}_1$, and let k be the number of irreducible components of the curve \mathcal{C}_1 . Then $k = d$ in each possible case. Moreover, we have

$$d = \begin{cases} 12 & \text{if } X = V_6, \\ 12 & \text{if } X = X_{24}, \\ 24 & \text{if } X = Y_{24}, \mathbb{W} \neq \mathbb{W}_1^{24} \text{ and } \mathbb{W} \neq \overline{\mathbb{W}}_1^{\mathfrak{S}}, \\ 12 & \text{if } X = Y_{24}, \text{ and either } \mathbb{W} = \mathbb{W}_1^{24} \text{ or } \mathbb{W} = \overline{\mathbb{W}}_1^{\mathfrak{S}}, \\ 12 & \text{if } X = V_4, \\ 6 & \text{if } X = \mathbb{P}^3. \end{cases}$$

Recall that $G \cap \mathbb{T}$ contains a subgroup isomorphic to μ_n^3 for $n \geq 24$. Thus, since by assumption Z is not \mathbb{T} -invariant, we have $|Z| \geq kn \geq 24k = 24d$.

Let M_1 and M_2 be general surfaces in \mathcal{M} . If the curve \mathcal{C} is G -irreducible, then $\mathcal{C} = \mathcal{C}_1$ and

$$M_1 \cdot M_2 = m_1 \mathcal{C}_1 + \Delta,$$

where m_1 is a non-negative integer, and Δ is an effective one-cycle whose support does not contain the curve \mathcal{C} . Similarly, if $X = Y_{24}$ and either $\mathbb{W} = \mathbb{W}_1^{24}$ or $\mathbb{W} = \overline{\mathbb{W}}_1^{\mathfrak{S}}$, then $\mathcal{C} = \mathcal{C}_1 + \mathcal{C}_2$, where both \mathcal{C}_1 and \mathcal{C}_2 are G -irreducible curves, so that we have

$$M_1 \cdot M_2 = m_1 \mathcal{C}_1 + m_2 \mathcal{C}_2 + \Delta,$$

where m_1 and m_2 are non-negative integers, and Δ is an effective one-cycle whose support does not contain the curves \mathcal{C}_1 and \mathcal{C}_2 . In both cases, we have

$$\frac{I_X^2 H^3}{\lambda^2} = H \cdot M_1 \cdot M_2 \geq H \cdot (m_1 \mathcal{C}_1 + \Delta) = m_1 d + H \cdot \Delta \geq m_1 d,$$

which implies that $m_1 \leq \frac{I_X^2 H^3}{d \lambda^2} < \frac{4}{\lambda^2}$.

Let \mathcal{B} be the linear subsystem in $|lH|$ consisting of surfaces that contain \mathcal{C} , where

$$l = \begin{cases} 2 & \text{if } X = V_6, \\ 2 & \text{if } X = X_{24}, \\ 2 & \text{if } X = Y_{24}, \\ 3 & \text{if } X = V_4, \\ 3 & \text{if } X = \mathbb{P}^3. \end{cases}$$

Then \mathcal{C} is the base locus of the linear system \mathcal{B} . Indeed, the generators of the linear system \mathcal{B} are contained in the formulas (4), (1), (10), (13) and (18). Looking at them, we see that the curve \mathcal{C} is the base locus of the linear system \mathcal{B} .

Now we use Lemma 2.9 to deduce that

$$\text{mult}_O(M_1 \cdot M_2) > \frac{4}{\lambda^2}$$

for every point $O \in Z \cap \mathcal{C}$. Thus, for every point $O \in Z \cap \mathcal{C}$, we have

$$\text{mult}_O(\Delta) > \frac{4}{\lambda^2} - m_1,$$

because the curve \mathcal{C}_1 is smooth at O , and \mathcal{C}_1 is the only G -irreducible component of the curve \mathcal{C} that contains points in Z . Now let S be a general surface in \mathcal{B} . Then

$$\begin{aligned} \frac{lI_X^2 H^3}{\lambda^2} - m_1 l d &= S \cdot (M_1 \cdot M_2 - m_1 \mathcal{C}_1) \geq S \cdot \Delta \\ &\geq \sum_{O \in Z} \text{mult}_O(\Delta) > |Z| \left(\frac{4}{\lambda^2} - m_1 \right) \geq dn \left(\frac{4}{\lambda^2} - m_1 \right). \end{aligned}$$

This gives

$$\frac{lI_X^2 H^3}{\lambda^2} + m_1 d(n - l) > \frac{4dn}{\lambda^2}.$$

Since $l \leq n$ and $m_1 \leq \frac{l^2 H^3}{d\lambda^2}$, we obtain $I_X^2 H^3 > 4d$, which is absurd. The obtained contradiction completes the proof of the lemma. \square

Acknowledgements This research was initiated during a research stay at Nemuro city in Hokkaido and continued at the occasion of the Kinosaki Algebraic Geometry Symposium 2019. We would like to thank the Nemuro city council for providing us with excellent working conditions. We are very grateful to Yuri Prokhorov, Hendrick Süß, Yuri Tschinkel and Yuri Zarhin for useful discussions and helpful comments. The first author was supported by Leverhulme Trust Grant RPG-2021-229. The second author was partially supported by the French ANR projects ISITE-BFC "Motivic invariants of Algebraic Varieties" ANR-15-IDEX-0008 and "FIBALGA" ANR-18-CE40-0003. The third author was partially funded by Grant-in-Aid for Scientific Research of JSPS No.19K03395.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Appendix A. Table of G -solid toric Fano threefolds

With the notation of Sect. 1, we let X be one of the toric threefolds among V_6 , V_4 , X_{24} , Y_{24} and \mathbb{P}^3 and we let G be a subgroup in the group G_X whose image $\nu_X(G)$ in the Weyl group \mathbb{W}_X of X contains the subgroup \mathfrak{A}_4 . Then X is G -minimal except the following two cases:

- (1) $X = V_4$, $\nu_X(G) \cong \mathfrak{S}_4$ and G acts intransitively on \mathbb{T} -invariant surfaces,
- (2) $X = V_4$ and $\nu_X(G) \cong \mathfrak{A}_4$.

Moreover, if X is G -minimal and $|G| \geq 32 \cdot 24^4$, then X is G -solid by Theorem 1.2. In this case, the following table summarizes additional information on the G -equivariant birational geometry of the threefold X obtained in the proof of Theorem 1.2.

If $X \neq \mathbb{P}^3$, then $\mathbb{W}_X \cong \mathfrak{S}_4 \times \mu_2$ contains two subgroups isomorphic to \mathfrak{S}_4 , which we call the subgroups \mathfrak{S}_4 of type I and II, see Notation 4.2 for the precise definition. If $\nu_{V_4}(G)$ is the subgroup \mathfrak{S}_4 of type II, then G acts transitively on the set of irreducible \mathbb{T} -invariant surfaces, so that V_4 is G -minimal in this case. In contrast, if $\nu_{V_4}(G)$ is the subgroup \mathfrak{S}_4 of type I, then V_4 is not G -minimal.

Similarly, if $\nu_{X_{24}}(G)$ is the subgroup \mathfrak{S}_4 of type II, then G acts transitively on the set of singular points of the threefold X_{24} . On the other hand, if $\nu_{X_{24}}(G)$ is the subgroup \mathfrak{S}_4 of type I, then G does not act transitively on this set.

$\nu_X(G)$	$\mathfrak{S}_4 \times \mu_2$	\mathfrak{S}_4 (type I)	\mathfrak{S}_4 (type II)	$\mathfrak{A}_4 \times \mu_2$	\mathfrak{A}_4
V_6	G -rationally superrigid	G -rationally superrigid	G -rationally superrigid	G -rationally superrigid	G -rationally superrigid
V_4	G -birational to X_{24}	not G -minimal	G -birational to X_{24}	G -birational to X_{24}	not G -minimal
X_{24}	G -birational to V_4	G -birational to \mathbb{P}^3	G -birational to V_4	G -birational to V_4	G -birational to \mathbb{P}^3
Y_{24}	G -rationally superrigid	G -rationally superrigid	G -birationally superrigid	G -rationally superrigid	G -rationally superrigid
\mathbb{P}^3		G -birational to X_{24}		G -birational to X_{24}	G -birational to X_{24}

References

1. Akhiezer, D.: Lie group actions in complex analysis, In: Vieweg, F. and Braunschweig, S. (1995)
2. Ahmadiñezhad, H., Cheltsov, I., Park, J., Shramov, C.: Double Veronese cones with 28 nodes, preprint, [arXiv:1910.10533](https://arxiv.org/abs/1910.10533) (2019)
3. Ahmadiñezhad, H., Okada, T.: Birationally rigid Pfaffian Fano 3-folds. *Algebr. Geom.* **5**, 160–199 (2018)
4. Avilov, A.: Automorphisms of threefolds that can be represented as an intersection of two quadrics. *Sbornik: Math.* **207**, 315–330 (2016)
5. Bayle, L.: Classification des variétés complexes projectives de dimension trois dont une section hyperplane générale est une surface d’Enriques. *J. Reine Angew. Math.* **449**, 9–63 (1994)
6. Borisov, A., Borisov, L.: Singular Toric Fano varieties. *Acad. Sci. USSR Sb. Math.* **75**, 277–283 (1993)
7. Brown, G. Kasprzyk, A.: Graded Ring Database, <http://www.grdb.co.uk>
8. Cartan, H.: Quotient d’un espace analytique par un groupe d’automorphismes, In *Algebraic Geometry and Topology*, pages 90–102. Princeton University Press, Princeton, N. J., 1957. A symposium in honor of S. Lefschetz
9. Cheltsov, I.: Birationally rigid Fano varieties. *Russ. Math. Surv.* **60**, 875–965 (2005)
10. Cheltsov, I., Shramov, C.: Cremona Groups and the Icosahedron. CRC Press, Boca Raton, FL (2016)
11. Cheltsov, I., Shramov, C.: Finite collineation groups and birational rigidity. *Selecta Math (N.S.)* **25**(5), 71 (2019)
12. Corti, A.: Factorizing birational maps of threefolds after Sarkisov. *J. Alg. Geom.* **4**, 223–254 (1995)
13. Corti, A.: Singularities of linear systems and 3-fold birational geometry. *L.M.S. Lecture Note Series* **281**, 259–312 (2000)
14. Corti, A., Mella, M.: Birational geometry of terminal quartic 3-folds, I. *Am. J. Math.* **126**, 739–761 (2004)
15. Corti, A., Kollár, J., Smith, K.: Rational and nearly rational varieties. Cambridge University Press (2003)
16. Cox, D., Little, J., Schenck, H.: Toric varieties, Vol. 124, Graduate Studies in Mathematics. American Mathematical Society, Providence, RI (2011)
17. Demazure, M.: Sous-groupes algébriques de rang maximum du groupe de Cremona. *Ann. Sci. Ecole Norm. Sup.* **3**, 507–588 (1970)
18. Dolgachev, I., Iskovskikh, V.: Finite subgroups of the plane Cremona group, in: *Algebra, Arithmetic, and Geometry: in Honor of Yu. I. Manin, Vol. I*, Progress in Mathematics, Vol. 269, Birkhäuser Boston, Boston, MA, , pp. 443–548 (2009)
19. Flannery, D.: The finite irreducible monomial linear groups of degree 4. *J. Algebr.* **218**, 436–469 (1999)
20. Hacon, C., McKernan, J.: The Sarkisov program. *J. Algebr. Geom.* **22**, 389–405 (2013)
21. Iskovskikh, V.: Two non-conjugate embeddings of $S_3 \times \text{mathbb{Z}}_2$ into the Cremona group II. *Adv. Study Pure Math.* **50**, 251–267 (2008)
22. Kasprzyk, A.: Toric Fano threefolds with terminal singularities. *Tohoku Math. J.* **58**, 101–121 (2006)
23. Kawamata, Y., Divisorial contractions to 3-dimensional terminal quotient singularities, Higher-dimensional complex varieties (Trento; de Gruyter. Berlin 1996, 241–246 (1994)
24. Kunyavskii, B.: Three-dimensional algebraic tori. *Sel. Math. Sov.* **9**, 1–21 (1991)
25. Lazarsfeld, R.: Positivity in Algebraic Geometry II. Springer-Verlag, Berlin (2004)
26. Prokhorov, Yu.: Simple finite subgroups of the Cremona group of rank 3. *J. Algebr. Geom.* **21**, 563–600 (2012)
27. Prokhorov, Yu.: G -Fano threefolds, I, II. *Adv. Geom.* **13**, 389–434 (2013)
28. Prokhorov, Yu., Reid, M.: On \mathbb{Q} -Fano threefolds of Fano index 2. *Adv. Stud. Pure Math.* **70**, 397–420 (2016)
29. Pukhlikov, A.: Birational automorphisms of Fano hypersurfaces. *Invent. Math.* **134**, 401–426 (1998)
30. Sakovics, D.: G -birational rigidity of the projective plane. *Eur. J. Math.* **5**, 1090–1105 (2019)
31. Sarikyan, A.: Toric G -Fano threefolds. *Internat. J. Math* **33**(5), 2250034 (2022)
32. Tahara, K.I.: On the finite subgroups of $GL(3, \mathbb{Z})$. *Nagoya Math. J.* **41**, 169–209 (1971)
33. Walter, J.: Equivariant birational geometry of quintic del Pezzo surface. *Eur. J. Math.* **4**, 1278–1292 (2018)