Geodesically convex energies and confinement of solutions for a multi-component system of nonlocal interaction equations

Jonathan Zinsl

Abstract. We consider a system of n nonlocal interaction evolution equations on $\mathbb{R}^d$ with a differentiable matrix-valued interaction potential W . Under suitable conditions on convexity, symmetry and growth of W , we prove λ -geodesic convexity for some $\lambda \in \mathbb{R}$ of the associated interaction energy with respect to a weighted compound distance of Wasserstein type. In particular, this implies existence and uniqueness of solutions to the evolution system. In one spatial dimension, we further analyse the qualitative properties of this solution in the non-uniformly convex case. We obtain, if the interaction potential is sufficiently convex far away from the origin, that the support of the solution is uniformly bounded. Under a suitable Lipschitz condition for the potential, we can exclude finite-time blow-up and give a partial characterization of the long-time behaviour.

Mathematics Subject Classification. 35R09, 35B30, 35B40.

Keywords. Nonlocal evolution equation, Interaction potential, Geodesic convexity, Wasserstein distance, Gradient flow, Multi-species system.

1. Introduction

1.1. The evolution system and its variational structure

In this work, we analyse the following system of $n \in \mathbb{N}$ nonlocal interaction evolution equations

$$\begin{aligned} \partial_t \mu_1 &= \operatorname{div}[m_1 \mu_1 \nabla(W_{11} * \mu_1 + W_{12} * \mu_2 + \cdots + W_{1n} * \mu_n)], \\ \partial_t \mu_2 &= \operatorname{div}[m_2 \mu_2 \nabla(W_{21} * \mu_1 + W_{22} * \mu_2 + \cdots + W_{2n} * \mu_n)], \\ &\vdots \\ \partial_t \mu_n &= \operatorname{div}[m_n \mu_n \nabla(W_{n1} * \mu_1 + W_{n2} * \mu_2 + \cdots + W_{nn} * \mu_n)]. \end{aligned} \quad (1)$$

The sought-for n -vector-valued solution $\mu(t) = (\mu_1(t), \dots, \mu_n(t))$ describes the distribution or concentration of n different populations or agents on $\mathbb{R}^d$ at time $t \geq 0$, $d \in \mathbb{N}$ denoting the spatial dimension. Apart from the constant *mobility magnitudes* $m_1, \dots, m_n > 0$, system (1) is mainly governed by the matrix-valued *interaction potential* $W : \mathbb{R}^d \rightarrow \mathbb{R}^{n \times n}$ satisfying the following requirements:

- (W1) $W(z)$ is a symmetric matrix for each $z \in \mathbb{R}^d$.
- (W2) $W_{ij} \in C^1(\mathbb{R}^d; \mathbb{R})$ for all $i, j \in \{1, \dots, n\}$.
- (W3) $W(z) = W(-z)$ for all $z \in \mathbb{R}^d$.
- (W4) There exists a matrix $\overline{W} \in \mathbb{R}^{n \times n}$ such that for each $i, j \in \{1, \dots, n\}$ and all $z \in \mathbb{R}^d$:

$$|W_{ij}(z)| \leq \overline{W}_{ij}(1 + |z|^2).$$

- (W5) There exists a symmetric matrix $\kappa \in \mathbb{R}^{n \times n}$ such that, for each $i, j \in \{1, \dots, n\}$, W_{ij} is κ_{ij} -(semi-)convex, i.e. the map $z \mapsto W_{ij}(z) - \frac{1}{2}\kappa_{ij}|z|^2$ is convex.

System (1) possesses a formal gradient flow structure: On the subspace $\mathcal{P}$ of those n -vector Borel measures on $\mathbb{R}^d$ with fixed *total masses* $\mu_j(\mathbb{R}^d) = p_j > 0$, fixed *(joint) weighted first moment*

$$\sum_{j=1}^n \frac{1}{m_j} \int_{\mathbb{R}^d} x \, d\mu_j(x) = E \in \mathbb{R}^d,$$

and finite second moments $\mathbf{m}_2(\mu_j) := \int_{\mathbb{R}^d} |x|^2 \, d\mu_j(x)$, the *multi-component interaction energy* functional

$$\mathcal{W}(\mu) := \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} W_{ij}(x-y) \, d\mu_i(x) \, d\mu_j(y) \quad (2)$$

induces (1) as its gradient flow w.r.t. the following compound metric of Wasserstein-type distances for each of the components of the vector measures $\mu^0, \mu^1 \in \mathcal{P}$:

$$\mathbf{W}_{\mathcal{P}}(\mu^0, \mu^1) = \left[\sum_{j=1}^n \frac{1}{m_j} \inf \left\{ \int_{\mathbb{R}^d \times \mathbb{R}^d} |x-y|^2 \, d\gamma_j(x, y) \mid \gamma_j \in \Gamma(\mu_j^0, \mu_j^1) \right\} \right]^{1/2}, \quad (3)$$

where $\Gamma(\mu_j^0, \mu_j^1)$ is the subset of finite Borel measures on $\mathbb{R}^d \times \mathbb{R}^d$ with marginals μ_j^0 and μ_j^1 . It easily follows from the properties of the usual Wasserstein distance for probability measures with finite second moment that $\mathbf{W}_{\mathcal{P}}$ defines a distance on the (geodesic) space $\mathcal{P}$ (see for instance [2, 56] for more details on optimal transport and gradient flows). Note that (W1)–(W5) imply (at least formally) that $\mathcal{P}$ is a positively invariant set along the flow of the evolution (1). In this work, we give a rigorous proof for these formal arguments.

1.2. Main results

We present our main results and improvements upon the case of a scalar interaction equation ($n = 1$).

1.2.1. Convexity along generalized geodesics and existence of solutions. We obtain in the case of genuine *irreducible* systems (see Definition 2.6 below) a novel sufficient condition on the model parameters such that the interaction energy functional $\mathcal{W}$ becomes λ -convex along *generalized* geodesics on $\mathcal{P}$ with respect to the distance $\mathbf{W}_{\mathcal{P}}$ for some $\lambda \in \mathbb{R}$:

Theorem (Convexity of $\mathcal{W}$ and generation of gradient flow solution) *Let $n > 1$, assume that (W1)–(W5) hold and let (1) be irreducible (see Definition 2.6). Define for $i \in \{1, \dots, n\}$ the quantity $\eta_i := \min_{j \neq i} \kappa_{ij} m_j \in \mathbb{R}$. Then, $\mathcal{W}$ is λ -convex along generalized geodesics on $\mathcal{P}$ w.r.t. $\mathbf{W}_{\mathcal{P}}$ (see Definition 2.1) for*

$$\lambda = \min_{i \in \{1, \dots, n\}} \left[p_i \min(0, m_i \kappa_{ii} - \eta_i) + \frac{1}{2} \sum_{j=1}^n p_j \left(\eta_j + \eta_i \frac{m_i}{m_j} \right) \right]. \quad (4)$$

Furthermore, the system (1), endowed with an initial datum $\mu^0 \in \mathcal{P}$, induces a λ -contractive gradient flow $\mu \in AC_{\text{loc}}^2([0, \infty); (\mathcal{P}, \mathbf{W}_{\mathcal{P}}))$ (see Theorem 2.13).

If $\mathcal{W}$ is uniformly geodesically convex, i.e., λ -convex along generalized geodesics for some $\lambda > 0$, the measure

$$\mu^\infty := (p_1, \dots, p_n)^T \delta_{x^\infty}, \quad \text{with} \quad x^\infty := E \left[\sum_{j=1}^n \frac{p_j}{m_j} \right]^{-1} \in \mathbb{R}^d,$$

is the unique minimizer—the ground state—of $\mathcal{W}$ and the unique stationary state of (1) on $\mathcal{P}$. It is globally asymptotically stable: the gradient flow solution converges exponentially fast in $(\mathcal{P}, \mathbf{W}_{\mathcal{P}})$ with rate λ to μ^∞ .

Comparison to the scalar case Our condition on convexity along generalized geodesics (4) extends the well-known condition by McCann [44] to multiple components: for $n = 1$, if the interaction potential W is κ -convex on $\mathbb{R}^d$, then $\mathcal{W}$ is $m\kappa p$ -convex along generalized geodesics in the space $\mathcal{P}$ defined as above for $n = 1$. The novel condition (4) is a *non-straightforward* generalization of McCann's condition to the case of systems $n > 1$, especially if one seeks a strictly positive modulus of convexity λ . On the one hand, putting $\kappa_{ij} > 0$ for all $i, j \in \{1, \dots, n\}$ (i.e., all cross- and self-interactions are attractive) is sufficient for $\lambda > 0$, but *far from necessary*. On the other hand, the value of λ in (4) does not seem to depend (at least not in an obvious manner) on the eigenvalues of the matrix $\kappa \in \mathbb{R}^{n \times n}$ from (W5). Hence, our new convexity condition uncovers—in a quantitative way—the dynamics of a multi-component network of (e.g. particle) interactions driven by C^1 potentials. In contrast to [19, 30], we restrict ourselves to the symmetric and smooth framework provided by (W1)–(W5), even if existence and uniqueness of solutions possibly could also be proved in more general settings. Let us now illustrate our novel

condition (4) by several examples. Set, for simplicity, $m_j = 1$ and $p_j = 1$ for all $j \in \{1, \dots, n\}$. Then, (4) simplifies to

$$\lambda = \min_{i \in \{1, \dots, n\}} \left[\min(0, \kappa_{ii} - \eta_i) + \frac{1}{2} \sum_{j=1}^n \eta_j + \frac{n}{2} \eta_i \right] \quad (n > 1),$$

where here $\eta_i = \min_{j \neq i} \kappa_{ij}$. In the even more specific setting of two species ($n = 2$), one has $\eta_1 = \kappa_{12} = \eta_2$ and

$$\lambda = \min\{\kappa_{11}, \kappa_{12}, \kappa_{22}\} + \kappa_{12} \quad (n = 2).$$

Clearly, if all interactions are attractive, e.g. if

$$\kappa = \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix},$$

then $\mathcal{W}$ is uniformly convex (here, the modulus is equal to 2) along generalized geodesics. One can also allow for repulsive self-interaction if overall attraction dominates repulsion, e.g. if

$$\kappa = \begin{pmatrix} -1 & 2 \\ 2 & -1 \end{pmatrix}.$$

Formula (4) yields in this case the 1-convexity of $\mathcal{W}$. Note that κ is an indefinite matrix. In contrast to that, repulsive cross-interactions, as in the case

$$\kappa = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix},$$

may lead to non-uniform convexity $\lambda \leq 0$; here $\lambda = -2$. Remarkably, κ is positive definite.

1.2.2. Qualitative behaviour for non-uniformly convex energy. If $\mathcal{W}$ is λ -geodesically convex with only $\lambda \leq 0$, the dynamics of system (1) are more involved. There, we restrict to one spatial dimension ($d = 1$) and rewrite the system in terms of *inverse distribution functions*: given the (scaled) cumulative distribution functions

$$F_i(t, x) = \int_{-\infty}^x \frac{1}{p_i} d\mu_i(t, y) \quad \in [0, 1], \quad (5)$$

let u_i be their corresponding pseudo-inverse, i.e.

$$u_i(t, z) = \inf\{x \in \mathbb{R} : F_i(t, x) > z\} \quad (\text{for } z \in [0, 1]). \quad (6)$$

Then, system (1) transforms into (cf. Sect. 3)

$$\partial_t u_i(t, z) = m_i \sum_{j=1}^n p_j \int_0^1 W'_{ij}(u_j(t, \xi) - u_i(t, z)) d\xi \quad (i = 1, \dots, n). \quad (7)$$

In terms of system (7), if $\mu \in AC_{\text{loc}}^2([0, \infty); (\mathcal{P}, \mathbf{W}_{\mathcal{P}}))$, one has that $u \in AC_{\text{loc}}^2([0, \infty); L^2([0, 1]; \mathbb{R}^n))$ and for all $t \geq 0$ and all $i \in \{1, \dots, n\}$, $u_i(t, \cdot)$ is a non-decreasing *càdlàg* function on $(0, 1)$. Section 3 is devoted to the analysis of the qualitative behaviour of the solution μ to (1) by means of

investigation of the corresponding solution u to (7). In that part, our main result is a *confinement* property of the solution:

Theorem (Confinement of solutions) *Assume that (W1)–(W5) are satisfied and let an initial datum μ^0 with compact support be given. Then, the solution μ to (1) propagates at finite speed: for each $T > 0$, there exists a constant $K(T, \mu^0) > 0$ depending on T and μ^0 such that*

$$\text{supp } \mu_i(t) \subset [-K(T, \mu^0), K(T, \mu^0)] \quad \forall t \in [0, T] \quad \forall i \in \{1, \dots, n\}.$$

Assume now in addition that the system (1) is irreducible at large distance (see Definition 3.2) and that the following condition is satisfied: there exist $R > 0$ and a matrix $C \in \mathbb{R}^{n \times n}$ such that for each $i, j \in \{1, \dots, n\}$, the map W_{ij} is C_{ij} -(semi)-convex on the interval (R, ∞) and the following holds:

If $n = 1$, then $C > 0$. If $n > 1$, with $\tilde{\eta}_i := \min_{j \neq i} C_{ij} p_j$ for all $i \in \{1, \dots, n\}$,

$$\tilde{\lambda}_0 := \min_{i \in \{1, \dots, n\}} \left[p_i \min(0, m_i C_{ii} - \tilde{\eta}_i) + \frac{1}{2} \sum_{j=1}^n p_j \left(\tilde{\eta}_j + \tilde{\eta}_i \frac{m_i}{m_j} \right) \right] > 0.$$

Then, the solution μ to (1) is uniformly confined: there exists $K(\mu^0) > 0$ such that

$$\text{supp } \mu_i(t) \subset [-K(\mu^0), K(\mu^0)] \quad \forall t \geq 0 \quad \forall i \in \{1, \dots, n\}. \quad (8)$$

Thus, in many cases, mass cannot escape to infinity. In contrast, is it possible to have *concentration* in finite time, i.e. can it occur that absolutely continuous solutions collapse to measures with nonvanishing singular part in finite time? The answer is negative for Lipschitz-continuous W'_{ij} and absolutely continuous initial data with continuous and bounded Lebesgue density (cf. Proposition 3.5).

Section 3.2 is devoted to the study of the long-time behaviour of the solution to (1). We first prove (cf. Theorem 3.6) that if the solution is *a priori* confined to a compact set, the ω -limit set of the system only contains steady states of (1). More specifically, assume that (8) is true and that all W'_{ij} are Lipschitz-continuous on the interval $[-2K, 2K]$. Then,

$$\lim_{t \rightarrow \infty} \left(\frac{d}{dt} \mathcal{W}(\mu(t)) \right) = 0. \quad (9)$$

Moreover, for each sequence $t_k \rightarrow \infty$, there exists a subsequence and a steady state $\bar{\mu} \in \mathcal{P}$ of (1) such that on the subsequence

$$\lim_{k \rightarrow \infty} \mathbf{W}_1(\mu_i(t_k), \bar{\mu}_i) = 0 \quad \forall i \in \{1, \dots, n\}.$$

There, $\mathbf{W}_1$ denotes the L^1 -Wasserstein distance between finite measures. However, this large-time limit $\bar{\mu}$ is not unique since it depends both on the sequence $(t_k)_{k \in \mathbb{N}}$ chosen and the extracted subsequence.

Even if the interaction potential does neither yield uniform geodesic convexity of the energy nor is confining and Lipschitz, we may observe the phenomenon of *collapse*: If the initial datum has compact support and the model

parameters admit $\sum_{j=1}^n \kappa_{ij} p_j > 0$ for all i , the diameter of the support of the solution shrinks exponentially fast over time (cf. Proposition 3.8). Still, the solution does in general not converge to a fixed steady state (see Remark 3.7 for more details). However, in the uniformly geodesically convex regime ($\lambda > 0$ in (4)), we obtain convergence even w.r.t. the stronger topology of the L^∞ -Wasserstein distance $\mathbf{W}_\infty$,

$$\lim_{t \rightarrow \infty} \mathbf{W}_\infty(\mu_i(t), \mu_i^\infty) = 0 \quad \forall i \in \{1, \dots, n\},$$

for initial data with compact support. In contrast to convergence w.r.t. $\mathbf{W}_\mathcal{P}$ (cf. Proposition 2.15), we do not obtain a specific rate of convergence.

Comparison to the scalar case For the scalar equation $n = 1$, an analogous confinement property of the solution to (1) has been derived e.g. in [3, 20] in arbitrary spatial dimension d under relatively general requirements on the regularity of the interaction potential W , assuming that W is (at least in a weak sense) attractive at large distance from the origin. In this work, we investigate the qualitative behaviour of solutions to (1) in one spatial dimension $d = 1$ only, concentrating on possible complications arising from the coupling in the equations (1). We obtain here a multi-component condition guaranteeing that W is confining, which is similar to the conditions from [3, 20], but taking into account that not all self- and cross-interactions have to be attractive. Indeed, we require W to behave like a uniformly convex potential in the sense of our new condition (4). A connatural property is assumed in the scalar case, see [20] in view of McCann's condition [44]. At the present state, it is unclear if the approach from [3, 20] to prove uniform compactness of the support of the solution μ to (1) in more than one spatial dimension can be carried over to the case of systems $n > 1$. In contrast, using the approach with inverse distribution functions for $d = 1$ allows us to give a more complete description of the qualitative behaviour of solutions: exclusion of blow-up in finite time can be shown with the method from [15] also in the case of systems $n > 1$. Moreover, our result on the structure of the ω -limit set of the dynamical system associated to (1), as a generalization of the scalar result from [48], also illustrates the use of energy methods, even if $\mathcal{W}$ is not uniformly convex along generalized geodesics.

1.3. Background

System (1) is a natural generalization of the scalar nonlocal evolution equation

$$\partial_t \mu = \operatorname{div}[m \mu \nabla(W * \mu)], \quad (10)$$

to multiple components. For the corresponding interaction energy functional

$$\mathcal{W}(\mu) = \frac{1}{2} \int_{\mathbb{R}^d} (W * \mu) \, d\mu, \quad (11)$$

McCann provided in his seminal paper [44] a criterion for λ -geodesic convexity with respect to the L^2 -Wasserstein distance (see also [56, Thm. 5.15(c)]). In a nutshell, if W is κ -convex in the Euclidean sense on $\mathbb{R}^d$ for some $\kappa \in \mathbb{R}$, then $\mathcal{W}$ is $\min(0, \kappa)$ -geodesically convex on the space of probability measures

endowed with the L^2 -Wasserstein distance. On the subspace of those measures having fixed weighted first moment E , $\mathcal{W}$ is κ -geodesically convex (i.e. uniform convexity is retained in the metric framework). It was proven by Ambrosio, Gigli and Savaré in [2] that geodesic convexity essentially leads to existence and uniqueness of weak solutions for the associated gradient flow evolution equation and to contractivity of the associated flow map. Geodesic convexity also yields useful error estimates, e.g. for the semi-discrete JKO scheme [38, 47] often used to construct weak solutions to equations with gradient flow structure. An immediate consequence of λ -geodesic convexity of functionals—for strictly positive $\lambda \in \mathbb{R}$ —is existence and uniqueness of minimizers (for recent results without using convexity, see e.g. [16, 18]). For more general genuine systems of equations with gradient flow structure, geodesic convexity has been studied in [58].

Model equations of the form (10) have arisen in the study of population dynamics in many cases (e.g. [10, 11, 14, 21, 28, 39, 40, 43, 45, 51, 52]) often derived as the infinite-particle limit of a individual-based model (e.g. [12, 36, 46]):

- In the parabolic-elliptic Keller–Segel model for chemotaxis in two spatial dimensions, the interaction potential is given by (the negative of) the Newtonian potential, i.e. $W(z) = \frac{1}{2\pi} \log(|z|)$, which is singular at $z = 0$ and attractive.
- Typical mathematical models of swarming processes include so-called *attractive-repulsive* potentials of, for instance, the form $W(z) = -C_a e^{-|z|/l_a} + C_r e^{-|z|/l_r}$, a special case of which is the attractive *Morse potential* $W(z) = -e^{-|z|}$. Also, Gaussian-type attractive-repulsive potentials $W(z) = -C_a e^{-|z|^2/l_a} + C_r e^{-|z|^2/l_r}$ are conceivable.

Nonlocal interaction potentials also appear in several models of physical applications such as models for granular media [4, 24, 25, 27, 42, 53], opinion formation [54] or interactions between particles (e.g. in crystals [50] or fluids [57]) with a broad range of reasonable interaction potentials. One can e.g. consider

- convex and C^1 -regular potentials, e.g. $W(z) = |z|^q$ for $q > 1$.
- non-convex, but regular potentials such as the *double-well potential* $W(z) = |z|^4 - |z|^2$,
- non-convex and singular potentials, e.g. the *Lennard–Jones* potential.

In the case of a radially symmetric potential $W(z) = w(|z|)$, the effect of the interaction potential is reflected by the sign of w' : If w' is positive, the individuals of the population *attract* each other, whereas in the case of negative w' the dynamics are *repulsive*. The force generated by the potential W points towards or away from the origin for positive or negative w' , respectively. Radially symmetric potentials describe interactions only depending on the distance of the particles. With the sum of convolutions appearing in the flux on the r.h.s. of system (1), we take into account that every species generates a—probably long-range—force on every other species.

Naturally, aggregation processes modelled by nonlocal interaction potentials are often combined with diffusive processes yielding (nonlinear) drift-diffusion equations as mathematical models. The question of global existence

of solutions to equations of these forms has been addressed in various publications. Using the theory of gradient flows, global existence of measure-valued solutions was proven in [19, 23], also for non-smooth potentials, in generalization of [24, 25]. Methods from optimal transportation theory were also useful for proving uniqueness, see e.g. [26, 29]. Well-posedness in the measure-valued sense was also studied in [17, 30] for a similar system for two species (see below).

A second field of study is the analysis of the qualitative behaviour of solutions to equations like (10), such as the speed of propagation, finite- and infinite-time blow-up of solutions and possible attractors, also with focus on self-similarity of solutions. It is not surprising that (10) exhibits blow-ups if the potential is sufficiently attractive. The aforementioned properties were investigated e.g. in [3, 5–9, 20, 41, 49]. One specific object of study is equation (10) considered in one spatial dimension. For instance, in [33, 34, 48] by Raoul and Fellner, rewriting in terms of inverse distribution functions allowed for the characterization of the long-time behaviour and the set of possible steady states of (10). One-dimensional models with nonlinear diffusion have been studied e.g. by Burger and Di Francesco in [15].

Genuine systems of the specific form (1) have been studied only in the case of two species as a physical model for two-component mixtures [55], fluids [57] or particle interactions [35]. From a more mathematical point of view, it was analysed by Di Francesco and Fagioli in [30, 31], where the results from [19, 33, 34, 48] were generalized to the case of two components. There, also the case of non-symmetric interaction has been studied.

2. Geodesic convexity and existence of solutions

In this section, we derive a sufficient condition for λ -convexity along generalized geodesics of the interaction energy $\mathcal{W}$ (cf. formula (2)) and conclude existence and uniqueness of solutions to (1). Throughout this part, the assumptions (W1)–(W5) above shall be fulfilled.

We begin with our definition of convexity along generalized geodesics, which is a straightforward generalization of the respective definition in the scalar case (see [2, Ch. 9]) to our vector-valued setting:

Definition 2.1. (*λ -convexity along generalized geodesics*) Given $\lambda \in \mathbb{R}$, we say that a functional $\mathfrak{A} : \mathcal{P} \rightarrow \mathbb{R}_\infty$ is λ -convex along generalized geodesics on $(\mathcal{P}, \mathbf{W}_\mathcal{P})$, if for any triple $\mu^1, \mu^2, \mu^3 \in \mathcal{P}$, there exists a n -vector-valued Borel measure $\boldsymbol{\mu}$ on $\mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R}^d$ such that:

- For all $j \in \{1, \dots, n\}$ and all $k \in \{1, 2, 3\}$: $\mu_j^k = \pi^k \# \boldsymbol{\mu}_j$.
- For $k \in \{2, 3\}$ and all $j \in \{1, \dots, n\}$, the measure $\pi^{(1,k)} \# \boldsymbol{\mu}_j$ is optimal in $\Gamma(\mu_j^1, \mu_j^k)$, i.e. it realizes the minimum in

$$\inf \left\{ \int_{\mathbb{R}^d \times \mathbb{R}^d} |x^1 - x^k|^2 \, d\gamma_j(x^1, x^k) \mid \gamma_j \in \Gamma(\mu_j^1, \mu_j^k) \right\}.$$

- Defining for $s \in [0, 1]$ and all $j \in \{1, \dots, n\}$ the *generalized geodesic* μ_s connecting μ^2 and μ^3 (with *base point* μ^1) by

$$\mu_{s,j} := [(1-s)\pi^2 + s\pi^3]_{\#} \mu_j,$$

one has for all $s \in [0, 1]$:

$$\begin{aligned} \mathfrak{A}(\mu_s) &\leq (1-s)\mathfrak{A}(\mu^2) + s\mathfrak{A}(\mu^3) \\ &\quad - \frac{\lambda}{2}s(1-s) \sum_{j=1}^n \frac{1}{m_j} \int_{\mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R}^d} |x^3 - x^2|^2 d\mu(x^1, x^2, x^3). \end{aligned} \quad (12)$$

We call the energy *uniformly* geodesically convex if it is λ -convex along generalized geodesics, for some $\lambda > 0$.

Notice that inequality (12) is sensible since the generalized geodesics μ_s connecting μ^2 and μ^3 remain in $\mathcal{P}$: by definition, one has for all $s \in (0, 1)$ that

$$\begin{aligned} &\sum_{j=1}^n \frac{1}{m_j} \int_{\mathbb{R}^d} x d\mu_{s,j}(x) \\ &= \sum_{j=1}^n \frac{1}{m_j} \int_{\mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R}^d} ((1-s)x^2 + sx^3) d\mu_j(x^1, x^2, x^3) \\ &= (1-s) \sum_{j=1}^n \frac{1}{m_j} \int_{\mathbb{R}^d} y d\mu_j^2(y) + s \sum_{j=1}^n \frac{1}{m_j} \int_{\mathbb{R}^d} y d\mu_j^3(y) \\ &= E, \end{aligned}$$

since $\mu^2, \mu^3 \in \mathcal{P}$.

Moreover, λ -convexity along generalized geodesics implies λ -convexity along geodesics in the usual sense, that is, for every pair $\mu^0, \mu^1 \in \mathcal{P}$, there exists a constant-speed geodesic curve $(\mu^s)_{s \in [0,1]}$ in $\mathcal{P}$ connecting μ^0 and μ^1 for which one has for all $s \in [0, 1]$:

$$\mathcal{W}(\mu^s) \leq (1-s)\mathcal{W}(\mu^0) + s\mathcal{W}(\mu^1) - \frac{1}{2}s(1-s)\lambda \mathbf{W}_{\mathcal{P}}(\mu^0, \mu^1)^2. \quad (13)$$

For convexity along *generalized* geodesics, an inequality of the form (13) is required for a wider class of curves joining μ^0 and μ^1 .

The following sufficient criterion is useful in verifying convexity along generalized geodesics as it allows to consider *absolutely continuous* measures and transport *maps*.

Theorem 2.2 (Sufficient criterion for convexity along generalized geodesics [2, Prop. 9.2.10]) *Let $\mathfrak{A} : \mathcal{P} \rightarrow \mathbb{R}_{\infty}$ be lower semicontinuous such that for all $\mu \in \mathcal{P}$, there exists a sequence $(\mu^k)_{k \in \mathbb{N}}$ on the subspace $\mathcal{P}^{\text{ac}}$ of absolutely continuous measures in $\mathcal{P}$ with $\lim_{k \rightarrow \infty} \mathbf{W}_{\mathcal{P}}(\mu^k, \mu) = 0$ and $\lim_{k \rightarrow \infty} \mathfrak{A}(\mu^k) = \mathfrak{A}(\mu)$.*

Assume moreover that for each $\mu \in \mathcal{P}^{\text{ac}}$ and $t_1, \dots, t_n, \tilde{t}_1, \dots, \tilde{t}_n : \mathbb{R}^d \rightarrow \mathbb{R}^d$ such that $t_j, \tilde{t}_j \in L^2(\mathbb{R}; d\mu_j)$ for all $j \in \{1, \dots, n\}$, the following estimate

holds along the interpolating curve $(\mu^s)_{s \in [0,1]}$ defined as $\mu_j^s := [(1-s)\tilde{t}_j + st_j] \# \mu_j$ for all $s \in [0,1]$ and $j \in \{1, \dots, n\}$:

$$\begin{aligned} \mathfrak{A}(\mu^s) &\leq (1-s)\mathfrak{A}(\mu^0) + s\mathfrak{A}(\mu^1) \\ &\quad - \frac{\lambda}{2}s(1-s) \sum_{j=1}^n \frac{1}{m_j} \int_{\mathbb{R}^d} |t_j(x) - \tilde{t}_j(x)|^2 d\mu(x) \\ &\text{for all } s \in [0,1]. \end{aligned} \tag{14}$$

Then, $\mathfrak{A}$ is λ -convex along generalized geodesics on $(\mathcal{P}, \mathbf{W}_{\mathcal{P}})$.

2.1. Geodesic convexity of the multi-component interaction energy

We first prove some basic properties of the interaction energy $\mathcal{W}$.

Lemma 2.3. (Proper domain and lower semicontinuity) *The following statements hold:*

- (a) For all $\mu \in \mathcal{P}$, one has $|\mathcal{W}(\mu)| < \infty$.
- (b) $\mathcal{W}$ is continuous on the metric space $(\mathcal{P}, \mathbf{W}_{\mathcal{P}})$.
- (c) Let $\rho \in C_c^\infty(\mathbb{R}^d)$ be defined via

$$\rho(x) := Z \exp\left(\frac{1}{|x|^2 - 1}\right) \mathbf{1}_{\mathbb{B}_1(0)}(x),$$

where $Z > 0$ is such that $\|\rho\|_{L^1} = 1$, and put $\rho_\varepsilon(x) := \varepsilon^{-d} \rho\left(\frac{x}{\varepsilon}\right)$ for $\varepsilon > 0$.

Then, for each $\mu \in \mathcal{P}$, the sequence $(\mu^k)_{k \in \mathbb{N}}$ with $\mu_j^k := \rho_{\frac{1}{k}} * \mu_j$ for $k \in \mathbb{N}$, $j \in \{1, \dots, n\}$, belongs to $\mathcal{P}^{\text{ac}}$ and $\lim_{k \rightarrow \infty} \mathbf{W}_{\mathcal{P}}(\mu^k, \mu) = 0$.

Proof. The key observation for parts (a) and (b) is the sub-quadratic growth of $\mathcal{W}$ as a consequence of condition (W4): there exists a constant $C > 0$ such that for all $\mu \in \mathcal{P}$, one has

$$|\mathcal{W}(\mu)| \leq C \sum_{j=1}^n \int_{\mathbb{R}^d} (1 + |x|^2) d\mu_j(x). \tag{15}$$

Clearly, (a) follows. If $(\mu^k)_{k \in \mathbb{N}}$ is a sequence converging to μ in $(\mathcal{P}, \mathbf{W}_{\mathcal{P}})$, in particular their second moments converge componentwise. Hence, the integrand on the r.h.s. in (15) is uniformly integrable which yields (b) as in [2, Lemma 5.1.7], using the continuity of $\mathcal{W}$. For part (c), we observe for $j \in \{1, \dots, n\}$ that μ_j^k is an absolutely continuous measure on $\mathbb{R}^d$ with Lebesgue density

$$x \mapsto \int_{\mathbb{R}^d} \rho_{\frac{1}{k}}(x - y) d\mu_j(y).$$

Clearly, $\mu_j^k(\mathbb{R}^d) = p_j$. Moreover, the weighted first moment E is unchanged by convolution with $\rho_{\frac{1}{k}}$ since for all $j \in \{1, \dots, n\}$ and all $k \in \mathbb{N}$:

$$\int_{\mathbb{R}^d} x d\mu_j^k(x) = \int_{\mathbb{R}^d} x d\mu_j(x).$$

Indeed, by transformation and Fubini's theorem,

$$\begin{aligned} \int_{\mathbb{R}^d} x \, d\mu_j^k(x) &= \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} (z+y) \rho_{\frac{1}{k}}(z) \, d\mu_j(y) \, dz \\ &= \int_{\mathbb{R}^d} y \, d\mu_j(y) + p_j \int_{\mathbb{R}^d} z \rho_{\frac{1}{k}}(z) \, dz. \end{aligned}$$

The last integral above vanishes since ρ is an even function. Along the same lines, one proves convergence of the second moments:

$$\begin{aligned} \mathbf{m}_2(\mu_j^k) &= \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} |z+y|^2 \rho_{\frac{1}{k}}(z) \, d\mu_j(y) \, dz \\ &= \mathbf{m}_2(\mu_j) + p_j \int_{\mathbb{R}^d} |z|^2 \rho_{\frac{1}{k}}(z) \, dz \\ &\quad + 2 \left(\int_{\mathbb{R}^d} z \rho_{\frac{1}{k}}(z) \, dz \right)^T \left(\int_{\mathbb{R}^d} y \, d\mu_j(y) \right). \end{aligned}$$

Since the last term vanishes again, we see

$$\mathbf{m}_2(\mu_j^k) = \mathbf{m}_2(\mu_j) + \frac{1}{k^2} \int_{\mathbb{R}^d} |x|^2 \rho(x) \, dx \xrightarrow{k \rightarrow \infty} \mathbf{m}_2(\mu_j).$$

It remains to prove narrow convergence of μ_j^k to μ_j . Fix $f : \mathbb{R}^d \rightarrow \mathbb{R}$ continuous and bounded. Using Fubini's theorem again, we get, since ρ is even,

$$\int_{\mathbb{R}^d} f \, d\mu_j^k - \int_{\mathbb{R}^d} f \, d\mu_j = \int_{\mathbb{R}^d} (\rho_{\frac{1}{k}} * f - f) \, d\mu_j.$$

Since f is continuous, $\rho_{\frac{1}{k}} * f$ converges to f pointwise on $\mathbb{R}^d$ (see, for instance, [32, App. C]). Clearly, $\rho_{\frac{1}{k}} * f$ is k -uniformly bounded. The dominated convergence theorem now yields

$$\int_{\mathbb{R}^d} (\rho_{\frac{1}{k}} * f - f) \, d\mu_j \xrightarrow{k \rightarrow \infty} 0,$$

proving the claim. $\square$

Lemma 2.4 (Growth control on the gradient) *There exists a matrix $\overline{C} \in \mathbb{R}^{n \times n}$ such that for all $z \in \mathbb{R}^d$ and all $i, j \in \{1, \dots, n\}$:*

$$|\nabla W_{ij}(z)| \leq \overline{C}_{ij}(|z| + 1). \quad (16)$$

Proof. We give a short proof for the sake of completeness. From (W2) and (W5), it easily follows for all $x, y \in \mathbb{R}^d$ that

$$W_{ij}(y) - W_{ij}(x) - \frac{\kappa_{ij}}{2} |y - x|^2 \geq \nabla W_{ij}(x)^T (y - x).$$

Putting

$$\alpha := \begin{cases} |4\overline{W}_{ij} - \kappa_{ij}|^{-1} & \text{if } 4\overline{W}_{ij} > \kappa_{ij}, \\ 1 & \text{otherwise,} \end{cases} \quad \text{and} \quad y := x + \alpha \nabla W_{ij}(x),$$

we get, using (W4) and Young's inequality:

$$\alpha |\nabla W_{ij}(x)|^2 \leq \overline{W}_{ij}(2 + 3|x|^2) + \frac{1}{2} \alpha^2 (4\overline{W}_{ij} - \kappa_{ij}) |\nabla W(x)|^2.$$

Consequently, in both cases, we have

$$|\nabla W_{ij}(x)| \leq (2\alpha^{-1}\overline{W}_{ij}(2+3|x|^2))^{1/2}$$

which implies an estimate of the form (16) via the elementary estimate $\sqrt{a+b} \leq \sqrt{a} + \sqrt{b}$ holding for $a, b \geq 0$. $\square$

Remark 2.5. (*Invariants*) Along the flow of system (1), the set $\mathcal{P}$ is positively invariant. We give a formal indication of this fact: Let an initial datum $\mu^0 \in \mathcal{P}$ be given. Since (1) is in divergence form, we immediately obtain the conservation of mass:

$$\frac{d}{dt} \int_{\mathbb{R}^d} d\mu_i(t, x) = 0.$$

Furthermore, by formal integration by parts, one has

$$\frac{d}{dt} \sum_{i=1}^n \mathbf{m}_2(\mu_i(t)) = - \sum_{i=1}^n 2m_i \int_{\mathbb{R}^d} x^T \left(\sum_{j=1}^n \nabla W_{ij} * \mu_j(t) \right) (x) d\mu_i(t, x),$$

from which it is possible to derive using the Young and Jensen inequalities and Lemma 2.4 the estimate

$$\frac{d}{dt} \sum_{i=1}^n \mathbf{m}_2(\mu_i(t)) \leq A \sum_{i=1}^n \mathbf{m}_2(\mu_i(t)) + B,$$

for suitable $A, B \in \mathbb{R}$. Gronwall's lemma now yields finiteness of second moments at a fixed time $t \geq 0$. Finally,

$$\frac{d}{dt} \sum_{i=1}^n \frac{1}{m_i} \int_{\mathbb{R}^d} x d\mu_i(t, x) = - \sum_{i=1}^n \sum_{j=1}^n \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \nabla W_{ij}(x-y) d\mu_j(t, y) d\mu_i(t, x).$$

Using assumptions (W1) and (W3) in combination with Fubini's theorem, we observe that the r.h.s. above is in fact equal to 0.

Definition 2.6. (*Irreducible systems*) We call a system of the form (1) *irreducible*, if the graph $G = (V_G, E_G)$ with nodes $V_G = \{1, \dots, n\}$ and edges $E_G = \{(i, j) \in V_G \times V_G : \nabla W_{ij} \not\equiv 0 \text{ on } \mathbb{R}^d\}$ is connected. That is, irreducible systems cannot be split up into independent subsystems.

The main result of this section is concerned with the geodesic convexity of the interaction energy $\mathcal{W}$:

Theorem 2.7. (Criterion for geodesic convexity) *Let $n > 1$ and let (1) be irreducible. Define for $i \in \{1, \dots, n\}$ the quantity $\eta_i := \min_{j \neq i} \kappa_{ij} m_j \in \mathbb{R}$. Then, $\mathcal{W}$ is λ_0 -convex along generalized geodesics on $\mathcal{P}$ w.r.t. $\mathbf{W}_{\mathcal{P}}$ with*

$$\lambda_0 := \min_{i \in \{1, \dots, n\}} \left[p_i \min(0, m_i \kappa_{ii} - \eta_i) + \frac{1}{2} \sum_{j=1}^n p_j \left(\eta_j + \eta_i \frac{m_i}{m_j} \right) \right]. \quad (17)$$

Proof. Thanks to the properties from Lemma 2.3, we are allowed to use Theorem 2.2. Let therefore $\mu \in \mathcal{P}^{\text{ac}}$ and let $t_1, \dots, t_n, \tilde{t}_1, \dots, \tilde{t}_n : \mathbb{R}^d \rightarrow \mathbb{R}^d$ such that $t_j, \tilde{t}_j \in L^2(\mathbb{R}; d\mu_j)$ for all $j \in \{1, \dots, n\}$. With the notation from Theorem 2.2, we have, using the convexity condition (W5) and the definition of the push-forward:

$$\begin{aligned} \mathcal{W}(\mu^s) &= \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} W_{ij}(\tilde{t}_j(x) - \tilde{t}_i(y)) \\ &\quad + s[t_j(x) - t_i(y) - (\tilde{t}_j(x) - \tilde{t}_i(y))] d\mu_j(x) d\mu_i(y) \\ &\leq (1-s)\mathcal{W}(\mu^0) + s\mathcal{W}(\mu^1) - \frac{1}{2}s(1-s) \\ &\quad \times \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \kappa_{ij} |t_j(x) - t_i(y) - (\tilde{t}_j(x) - \tilde{t}_i(y))|^2 d\mu_j(x) d\mu_i(y). \end{aligned}$$

In view of (12), we have to verify that

$$\begin{aligned} &\frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \kappa_{ij} |t_j(x) - t_i(y) - (\tilde{t}_j(x) - \tilde{t}_i(y))|^2 d\mu_j(x) d\mu_i(y) \\ &\geq \lambda_0 \sum_{i=1}^n \frac{1}{m_i} \int_{\mathbb{R}^d} |t_i(x) - \tilde{t}_i(x)|^2 d\mu_i(x). \end{aligned} \quad (18)$$

We first split up the l.h.s. of (18) into its diagonal and off-diagonal part and perform an estimate on the latter introducing the numbers $\eta_i = \min_{j \neq i} \kappa_{ij} m_j$:

$$\begin{aligned} &\frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \kappa_{ij} |t_j(x) - t_i(y) - (\tilde{t}_j(x) - \tilde{t}_i(y))|^2 d\mu_j(x) d\mu_i(y) \\ &\geq \frac{1}{2} \sum_i \sum_{j \neq i} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \frac{\eta_i}{m_j} |(t_j(x) - \tilde{t}_j(x)) - (t_i(y) - \tilde{t}_i(y))|^2 d\mu_j(x) d\mu_i(y) \\ &\quad + \frac{1}{2} \sum_i \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \kappa_{ii} |(t_i(x) - \tilde{t}_i(x)) - (t_i(y) - \tilde{t}_i(y))|^2 d\mu_i(x) d\mu_i(y). \end{aligned}$$

Expanding the squares yields

$$\begin{aligned} &\frac{1}{2} \sum_i \sum_{j \neq i} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \frac{\eta_i}{m_j} |(t_j(x) - \tilde{t}_j(x)) - (t_i(y) - \tilde{t}_i(y))|^2 d\mu_j(x) d\mu_i(y) \\ &\quad + \frac{1}{2} \sum_i \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \kappa_{ii} |(t_i(x) - \tilde{t}_i(x)) - (t_i(y) - \tilde{t}_i(y))|^2 d\mu_i(x) d\mu_i(y) \\ &= \frac{1}{2} \sum_i \sum_{j \neq i} \left(\int_{\mathbb{R}^d} \frac{p_i \eta_i}{m_j} |t_j(x) - \tilde{t}_j(x)|^2 d\mu_j(x) \right. \\ &\quad \left. + \int_{\mathbb{R}^d} \frac{p_j \eta_i}{m_j} |t_i(x) - \tilde{t}_i(x)|^2 d\mu_i(x) \right) \end{aligned}$$

$$\begin{aligned}
& - \sum_i \left(\sum_{j \neq i} \int_{\mathbb{R}^d} \frac{1}{m_j} (t_j(x) - \tilde{t}_j(x)) \, d\mu_j(x) \right)^T \\
& \cdot \left(\int_{\mathbb{R}^d} \eta_i(t_i(x) - \tilde{t}_i(x)) \, d\mu_i(x) \right) \\
& + \sum_i \kappa_{ii} \left(\int_{\mathbb{R}^d} p_i |t_i(x) - \tilde{t}_i(x)|^2 \, d\mu_i(x) \right) \\
& - \sum_i \kappa_{ii} \left| \int_{\mathbb{R}^d} (t_i(x) - \tilde{t}_i(x)) \, d\mu_i(x) \right|^2. \tag{19}
\end{aligned}$$

Now, the special structure of $\mathcal{P}$ comes into play: since the weighted first moment E is fixed on $\mathcal{P}$, one has

$$E = \sum_{j=1}^n \frac{1}{m_j} \int_{\mathbb{R}^d} x \, d(t_j \# \mu_j) = \sum_{j=1}^n \frac{1}{m_j} \int_{\mathbb{R}^d} x \, d(\tilde{t}_j \# \mu_j),$$

and consequently

$$\sum_{j \neq i} \int_{\mathbb{R}^d} \frac{1}{m_j} (t_j(x) - \tilde{t}_j(x)) \, d\mu_j(x) = - \int_{\mathbb{R}^d} \frac{1}{m_i} (t_i(x) - \tilde{t}_i(x)) \, d\mu_i(x).$$

We exploit this fact in order to rewrite the r.h.s. of formula (19) above:

r.h.s.

$$\begin{aligned}
& = \frac{1}{2} \sum_i \sum_j \left(\int_{\mathbb{R}^d} \frac{p_j \eta_i}{m_j} |t_j(x) - \tilde{t}_j(x)|^2 \, d\mu_j(x) \right. \\
& \quad + \int_{\mathbb{R}^d} \frac{p_j \eta_i}{m_j} |t_i(x) - \tilde{t}_i(x)|^2 \, d\mu_i(x) \Big) \\
& \quad - \sum_i \int_{\mathbb{R}^d} \frac{p_i \eta_i}{m_i} |t_i(x) - \tilde{t}_i(x)|^2 \, d\mu_i(x) \\
& \quad - \sum_i \left(- \int_{\mathbb{R}^d} \frac{1}{m_i} (t_i(x) - \tilde{t}_i(x)) \, d\mu_i(x) \right)^T \\
& \quad \cdot \left(\int_{\mathbb{R}^d} \eta_i (t_i(x) - \tilde{t}_i(x)) \, d\mu_i(x) \right) \\
& \quad - \sum_i \kappa_{ii} \left(\int_{\mathbb{R}^d} p_i |t_i(x) - \tilde{t}_i(x)|^2 \, d\mu_i(x) \right) \\
& \quad - \sum_i \kappa_{ii} \left| \int_{\mathbb{R}^d} (t_i(x) - \tilde{t}_i(x)) \, d\mu_i(x) \right|^2 \\
& = \sum_i \left\{ \left| \int_{\mathbb{R}^d} (t_i(x) - \tilde{t}_i(x)) \, d \left(\frac{1}{p_i} \mu_i \right) (x) \right|^2 p_i^2 \left(\frac{\eta_i}{m_i} - \kappa_{ii} \right) \right. \\
& \quad \left. + \int_{\mathbb{R}^d} |t_i(x) - \tilde{t}_i(x)|^2 \, d \left(\frac{1}{p_i} \mu_i \right) (x) p_i^2 \left(\kappa_{ii} - \frac{\eta_i}{m_i} \right) \right\}
\end{aligned}$$

$$\begin{aligned}
& + \frac{1}{m_i} \int_{\mathbb{R}^d} |t_i(x) - \tilde{t}_i(x)|^2 d\mu_i(x) \cdot \frac{1}{2} \sum_j p_j \left(\eta_j + \eta_i \frac{m_i}{m_j} \right) \Big\} \\
& =: \sum_i S_i.
\end{aligned}$$

We analyse each S_i separately.

If $\frac{\eta_i}{m_i} - \kappa_{ii} \geq 0$, the first term in S_i is nonnegative, so

$$\begin{aligned}
S_i & \geq \frac{1}{m_i} \int_{\mathbb{R}^d} |t_i(x) - \tilde{t}_i(x)|^2 d\mu_i(x) \\
& \quad \times \left[p_i(m_i \kappa_{ii} - \eta_i) + \frac{1}{2} \sum_j p_j \left(\eta_j + \eta_i \frac{m_i}{m_j} \right) \right].
\end{aligned}$$

If $\frac{\eta_i}{m_i} - \kappa_{ii} < 0$, the sum of the first two terms in S_i is nonnegative thanks to Jensen's inequality. Hence,

$$S_i \geq \frac{1}{m_i} \int_{\mathbb{R}^d} |t_i(x) - \tilde{t}_i(x)|^2 d\mu_i(x) \cdot \frac{1}{2} \sum_j p_j \left(\eta_j + \eta_i \frac{m_i}{m_j} \right).$$

Defining λ_0 as in (17) clearly leads to (18), completing the proof. $\square$

Remark 2.8. (*Non-irreducible systems*) If system (1) is *not* irreducible, there exists an I -integer partition ($I \in \mathbb{N}$) of $n \in \mathbb{N}$ into $n_1 + n_2 + \dots + n_I = n$ such that (1) decomposes into I independent irreducible subsystems having the same structure as (1), but with n replaced by $n_1, \dots, n_I$, respectively. The modulus of geodesic convexity of the interaction energy $\mathcal{W}$ can now be computed as the minimum of the respective convexity moduli of each subsystem: if $n_k > 1$ for some $k \in \{1, \dots, I\}$, formula (17) applies; if $n_k = 1$, McCann's criterion [44] applies (and yields convexity modulus $m\kappa p$ for the respective m, κ, p of the k^{th} subsystem in our framework).

Why do we need the condition of irreducibility at all? Consider a one-parameter family of admissible interaction potentials $(W^\beta)_{\beta>0}$ inducing irreducible systems (1) for each $\beta \rightarrow 0$ and assume that $W^\beta \rightarrow W$ as $\beta \rightarrow 0$, where the limit potential W is such that the associated system (1) is no longer irreducible. This means that system (1) decouples asymptotically as $\beta \rightarrow 0$. Clearly, by definition of irreducibility, we can permute the components $\mu_1, \dots, \mu_n$ such that W becomes block-diagonal. Hence, one has $\lim_{\beta \rightarrow 0} \eta_i^\beta \leq 0$ for all $i \in \{1, \dots, n\}$, and consequently, $\lim_{\beta \rightarrow 0} \lambda_0^\beta \leq 0$. On the other hand—considering each irreducible subsystem associated to W separately—the modulus of convexity of the limit functional might be strictly positive, e.g. if W is diagonal with positive entries on the diagonal.

Proposition 2.9 (Necessary condition for $\lambda_0 > 0$) *If $\lambda_0 > 0$ in (17), then for all $i \in \{1, \dots, n\}$:*

$$\sum_{j=1}^n \kappa_{ij} p_j > 0.$$

This condition is not sufficient (see the examples from the introduction).

Proof. Fix $i \in \{1, \dots, n\}$. The following holds:

$$\begin{aligned} m_i \sum_j \kappa_{ij} p_j &= p_i m_i \kappa_{ii} + \sum_{j \neq i} m_i \kappa_{ij} p_j \\ &= p_i (m_i \kappa_{ii} - \eta_i) + p_i \eta_i + \frac{1}{2} \sum_{j \neq i} p_j \kappa_{ji} m_i + \frac{1}{2} \sum_{j \neq i} p_j \kappa_{ij} m_j \frac{m_i}{m_j}, \end{aligned}$$

where we used the symmetry of κ . Now, we estimate using the definition of η_i, η_j :

$$\begin{aligned} m_i \sum_j \kappa_{ij} p_j &\geq p_i \min(0, m_i \kappa_{ii} - \eta_i) + \frac{1}{2} p_i \eta_i + \frac{1}{2} \sum_{j \neq i} p_j \eta_j + \frac{1}{2} p_i \eta_i \frac{m_i}{m_i} + \frac{1}{2} \sum_{j \neq i} p_j \eta_i \frac{m_i}{m_j} \\ &= p_i \min(0, m_i \kappa_{ii} - \eta_i) + \frac{1}{2} \sum_j p_j \left(\eta_j + \eta_i \frac{m_i}{m_j} \right) \geq \lambda_0 > 0. \end{aligned}$$

□

Remark 2.10 (*Optimal constants*) In general, the constant λ_0 obtained in (17) is not optimal. In some scenarios, however, it might be: for *purely quadratic* interaction potentials W_{ij} in one spatial dimension $d = 1$, i.e., $W_{ij}(x) = \frac{1}{2} \kappa_{ij} x^2$, system (1) can be solved analytically transforming (1) into the *linear* system (7) for the inverse distribution functions $u_1, \dots, u_n$ from (6), and the quantities $v_i(t) = \int_0^1 u_i(t, z) dz$ for $i \in \{1, \dots, n\}$. The optimal rate in the contractivity estimate (25) is then given as the maximal eigenvalue of the matrix associated to this linear system. Obviously, an explicit formula for this eigenvalue is much more involved than our condition (17) for λ_0 which already incorporates much of the qualitative behaviour one expects.

More specifically, for two components $n = 2$, equal masses $p_1 = p_2 = 1$ and mobilities $m_1 = m_2 = 1$, one obtains the following—almost decoupled—system using the invariance of E :

$$\begin{aligned} \partial_t u_1(t, z) &= -(\kappa_{11} + \kappa_{12})u_1(t, z) + (\kappa_{11} - \kappa_{12})v_1(t) + \kappa_{12}E, \\ \partial_t u_2(t, z) &= -(\kappa_{12} + \kappa_{22})u_2(t, z) + (\kappa_{22} - \kappa_{12})v_2(t) + \kappa_{12}E, \\ \frac{d}{dt} v_i(t) &= -2\kappa_{12}v_i(t) + \kappa_{12}E \quad (\text{for both } i \in \{1, 2\}). \end{aligned}$$

One sees that the maximal eigenvalue of the matrix associated to the linear system above is $-\min\{2\kappa_{12}, \kappa_{11} + \kappa_{12}, \kappa_{22} + \kappa_{12}\}$ which coincides with $-\lambda_0$, recall that in this case

$$\lambda_0 = \min\{\kappa_{11}, \kappa_{12}, \kappa_{22}\} + \kappa_{12}.$$

This shows that the rate in the contractivity estimate (25) cannot be improved beyond λ_0 .

We close this section with an investigation of the *local slope* $|\partial\mathcal{W}|$ of $\mathcal{W}$, defined by (see [2, Ch. 1])

$$|\partial\mathcal{W}|(\mu) := \limsup_{\nu \rightarrow \mu} \frac{\max\{\mathcal{W}(\mu) - \mathcal{W}(\nu), 0\}}{\mathbf{W}_{\mathcal{P}}(\mu, \nu)}.$$

The following statement is a generalization of the respective result [2, Thm. 10.4.11] for the scalar case to the vector-valued setting:

Proposition 2.11 (Estimate on $|\partial\mathcal{W}|$) *Let $\mu \in \mathcal{P}$ and define $w = (w_1, \dots, w_n)$ by*

$$w_i := \sum_{j=1}^n \nabla W_{ij} * \mu_j, \quad (20)$$

for each $i \in \{1, \dots, n\}$. If for all i , one has $w_i \in L^2(\mathbb{R}^d; d\mu_i)$, then

$$|\partial\mathcal{W}|(\mu) \leq \left(\|w_i\|_{L^2(\mathbb{R}^d; d\mu_i)}^2 \right)^{1/2}. \quad (21)$$

Proof. By [2, Ch. 10] (which can easily be extended to our tensorized setting), it suffices to prove that for all $\nu \in \mathcal{P}$ and all $\gamma_1, \dots, \gamma_n$ optimal in $\mathbf{W}_{\mathcal{P}}(\mu, \nu)$, i.e.,

$$\mathbf{W}_{\mathcal{P}}^2(\mu, \nu) = \sum_{i=1}^n \frac{1}{m_i} \int_{\mathbb{R}^d \times \mathbb{R}^d} |x - \hat{x}|^2 d\gamma_i(\hat{x}, x),$$

that

$$\mathcal{W}(\nu) - \mathcal{W}(\mu) \geq \sum_{i=1}^n \int_{\mathbb{R}^d \times \mathbb{R}^d} w_i(\hat{x})^T (x - \hat{x}) d\gamma_i(\hat{x}, x) + \lambda_0 \mathbf{W}_{\mathcal{P}}(\mu, \nu)^2. \quad (22)$$

First, by doubling the variables, we obtain

$$\begin{aligned} \mathcal{W}(\nu) - \mathcal{W}(\mu) &= \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} (W_{ij}(x - y) - W_{ij}(\hat{x} - \hat{y})) \\ &\quad \times \frac{1}{p_i p_j} d\nu_i(x) d\nu_j(y) d\mu_i(\hat{x}) d\mu_j(\hat{y}), \end{aligned}$$

which can be rewritten as

$$\begin{aligned} \mathcal{W}(\nu) - \mathcal{W}(\mu) &= \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \left[W_{ij}((\hat{x} - \hat{y}) + s((x - y) - (\hat{x} - \hat{y}))) \right. \\ &\quad \left. - \frac{1}{2} \kappa_{ij} |(\hat{x} - \hat{y}) + s((x - y) - (\hat{x} - \hat{y}))|^2 \right]_{s=0}^{s=1} \\ &\quad \times \frac{1}{p_i p_j} d\nu_i(x) d\nu_j(y) d\mu_i(\hat{x}) d\mu_j(\hat{y}) \\ &\quad + \sum_{i=1}^n \sum_{j=1}^n \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \frac{\kappa_{ij}}{4 p_i p_j} [|x - y|^2 - |\hat{x} - \hat{y}|^2] \\ &\quad \times d\nu_i(x) d\nu_j(y) d\mu_i(\hat{x}) d\mu_j(\hat{y}). \end{aligned}$$

Now, thanks to convexity, the quantity

$$\frac{1}{\hat{s}} \left[W_{ij}((\hat{x} - \hat{y}) + s((x - y) - (\hat{x} - \hat{y}))) - \frac{1}{2} \kappa_{ij} |(\hat{x} - \hat{y}) + s((x - y) - (\hat{x} - \hat{y}))|^2 \right]_{s=0}^{s=\hat{s}}$$

is nondecreasing w.r.t. $\hat{s} \in (0, 1]$, so the monotone convergence theorem yields as $\hat{s} \downarrow 0$ that

$$\begin{aligned} & \mathcal{W}(\nu) - \mathcal{W}(\mu) \\ & \geq \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} (\nabla W_{ij}(\hat{x} - \hat{y}) - \kappa_{ij}(\hat{x} - \hat{y}))^T ((x - y) - (\hat{x} - \hat{y})) \\ & \quad \times \frac{1}{p_i p_j} d\nu_i(x) d\nu_j(y) d\mu_i(\hat{x}) d\mu_j(\hat{y}) \\ & \quad + \sum_{i=1}^n \sum_{j=1}^n \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \frac{\kappa_{ij}}{4p_i p_j} [|x - y|^2 - |\hat{x} - \hat{y}|^2 - 4(\hat{x} - \hat{y})^T (x - y)] \\ & \quad \times d\nu_i(x) d\nu_j(y) d\mu_i(\hat{x}) d\mu_j(\hat{y}). \end{aligned}$$

The fact that ∇W_{ij} is an odd function yields, interchanging variable names, that

$$\begin{aligned} & \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \nabla W_{ij}(\hat{x} - \hat{y})^T ((x - \hat{x}) - (y - \hat{y})) \\ & \quad \times \frac{1}{p_i p_j} d\nu_i(x) d\nu_j(y) d\mu_i(\hat{x}) d\mu_j(\hat{y}) \\ & = \sum_{i=1}^n \sum_{j=1}^n \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \frac{\nabla W_{ij}(\hat{x} - \hat{y})^T (x - \hat{x})}{p_i p_j} d\nu_i(x) d\nu_j(y) d\mu_i(\hat{x}) d\mu_j(\hat{y}) \\ & = \sum_{i=1}^n \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \frac{1}{p_i} w_i(\hat{x})^T (x - \hat{x}) d\nu_i(x) d\mu_i(\hat{x}) \\ & = \sum_{i=1}^n \int_{\mathbb{R}^d \times \mathbb{R}^d} w_i(\hat{x})^T (x - \hat{x}) d\gamma_i(\hat{x}, x). \end{aligned}$$

Hence, in order to show (22), we have to verify that

$$\begin{aligned} \lambda_0 \mathbf{W}_{\mathcal{P}}(\mu, \nu)^2 & \leq \sum_{i=1}^n \sum_{j=1}^n \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \frac{\kappa_{ij}}{4p_i p_j} [|x - y|^2 - |\hat{x} - \hat{y}|^2 - 4(\hat{x} - \hat{y})^T (x - y)] \\ & \quad \times d\nu_i(x) d\nu_j(y) d\mu_i(\hat{x}) d\mu_j(\hat{y}). \end{aligned} \quad (23)$$

By elementary calculations, the right-hand side in (23) can be rewritten as

r.h.s.

$$= \sum_{i=1}^n \sum_{j=1}^n \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \frac{\kappa_{ij}}{4} |(x - \hat{x}) - (y - \hat{y})|^2 d\gamma_i(\hat{x}, x) d\gamma_j(\hat{y}, y),$$

the structure of which—probably not a surprise—is similar to the left-hand side of formula (18) in the proof of Theorem 2.7. We are now in position to carry out exactly the same method as for the derivation of λ_0 in the proof of Theorem 2.7, and eventually end up with (23). We omit these highly technical but conceptually not novel steps for the sake of brevity. $\square$

Remark 2.12. With even more technical effort, one can show that equality holds in (21) (see [2, Ch. 10] for more details).

2.2. Existence and uniqueness of gradient flow solutions

With the results of Lemma 2.3 and Theorem 2.7 at hand, the following statement follows thanks to [2, Chapter 11]:

Theorem 2.13 (Existence and uniqueness) *Consider (1) endowed with an initial datum $\mu^0 \in \mathcal{P}$. Then, there exists a gradient flow solution $\mu \in AC_{\text{loc}}^2([0, \infty); (\mathcal{P}, \mathbf{W}_{\mathcal{P}}))$ to this initial-value problem: System (1) holds in the sense of distributions and one has $\mu(0) = \mu^0$. Moreover, with λ_0 from (17), the evolution variational estimate holds for almost every $t > 0$ and all $\nu \in \mathcal{P}$:*

$$\frac{1}{2} \frac{d^+}{dt} \mathbf{W}_{\mathcal{P}}(\mu(t), \nu)^2 + \frac{\lambda_0}{2} \mathbf{W}_{\mathcal{P}}(\mu(t), \nu)^2 \leq \mathcal{W}(\nu) - \mathcal{W}(\mu(t)). \quad (24)$$

Given another initial datum $\nu^0 \in \mathcal{P}$ and the respective gradient flow solution $\nu \in AC_{\text{loc}}^2([0, \infty); (\mathcal{P}, \mathbf{W}_{\mathcal{P}}))$, the following contraction estimate holds for all $t \geq 0$:

$$\mathbf{W}_{\mathcal{P}}(\mu(t), \nu(t)) \leq e^{-\lambda_0 t} \mathbf{W}_{\mathcal{P}}(\mu^0, \nu^0), \quad (25)$$

which implies in particular the uniqueness of solutions.

Notice that for Theorem 2.13 to hold for $\lambda_0 > 0$, one explicitly needs λ_0 -convexity along *generalized* geodesics, since the weaker λ_0 -convexity along (non-generalized) geodesics does not suffice to infer existence of the gradient flow: the main reason are the curvature properties of the space $(\mathcal{P}, \mathbf{W}_{\mathcal{P}})$ being as in the scalar Wasserstein setting discussed in [2]. In contrast to that, our derivation of λ_0 in the proof of Theorem 2.7 can be carried out in the same way to obtain the same condition for convexity along non-generalized geodesics. For interaction energy functionals like $\mathcal{W}$, the optimal moduli of convexity along generalized and non-generalized geodesics coincide (see e.g. [2, 44]).

Our next result is concerned with gradient flow solutions associated to steady states of (1):

Proposition 2.14 (Stationary solutions are gradient flow solutions) *Assume that $\mu^0 \in \mathcal{P}$ is a steady state of (1), that is, for all $i \in \{1, \dots, n\}$, one has*

$$\sum_{j=1}^n \nabla W_{ij} * \mu_j^0 = 0 \quad \text{on} \quad \text{supp } \mu_i^0.$$

Then, the curve $\mu(t) = \mu^0$ for all $t \geq 0$ satisfies the evolution variational estimate (24). Thus, the stationary solution $\mu \equiv \mu^0$ to (1) coincides with the unique gradient flow solution in the sense of Theorem 2.13.

Proof. On the one hand, by Proposition 2.11, we have that $|\partial \mathcal{W}|(\mu^0) = 0$. On the other hand, thanks to the λ_0 -convexity of $\mathcal{W}$ along generalized geodesics, the local slope can also be characterized as in [2, Thm. 2.4.9]:

$$|\partial \mathcal{W}|(\mu^0) = \sup_{\nu \neq \mu^0} \max \left\{ 0, \frac{\mathcal{W}(\mu^0) - \mathcal{W}(\nu)}{\mathbf{W}_{\mathcal{P}}(\nu, \mu^0)} + \frac{1}{2} \lambda_0 \mathbf{W}_{\mathcal{P}}(\nu, \mu^0) \right\}.$$

Hence, for all $\nu \in \mathcal{P}$:

$$\mathcal{W}(\mu^0) - \mathcal{W}(\nu) + \frac{1}{2} \lambda_0 \mathbf{W}_{\mathcal{P}}(\nu, \mu^0)^2 \leq 0,$$

which is (24) for the curve $\mu \equiv \mu^0$. $\square$

Proposition 2.15 (The uniformly convex case) *If (17) yields $\lambda_0 > 0$, the measure*

$$\mu^\infty := (p_1, \dots, p_n)^\top \delta_{x^\infty}, \quad \text{with } x^\infty := E \left[\sum_{j=1}^n \frac{p_j}{m_j} \right]^{-1} \in \mathbb{R}^d,$$

is the unique minimizer of $\mathcal{W}$ and the unique stationary state of (1) on $\mathcal{P}$. It is globally asymptotically stable: The solution from Theorem 2.13 converges exponentially fast in $(\mathcal{P}, \mathbf{W}_{\mathcal{P}})$ at rate λ_0 to μ^∞ .

Proof. One easily sees—thanks to the symmetry property $\nabla W_{ij}(0) = 0$ from conditions (W2) and (W3)—that μ^∞ is a steady state of (1). It is unique by uniform geodesic convexity of $\mathcal{W}$: the contraction estimate (25) in combination with Proposition 2.14 necessarily implies that two steady states on $\mathcal{P}$ coincide. On the other hand, uniform geodesic convexity of $\mathcal{W}$ implies existence and uniqueness of a minimizer of $\mathcal{W}$ which is—since $\mathcal{W}$ is nonincreasing along the solution $\mu(t)$ —a steady state of (1). $\square$

As for scalar equations of the form (10), system (1) can be viewed as a continuum limit of a multi-particle system. To this end, we introduce the concept of *particle solutions* as a conclusion to this section.

Remark 2.16. (*Particle solutions*) Assume that the initial datum is discrete, i.e. each component μ_i^0 is a finite linear combination of Dirac measures:

$$\mu_i^0 = \sum_{k=1}^{N_i} p_i^k \delta_{x_i^{0,k}} \quad (i = 1, \dots, n).$$

There, the $N_i \in \mathbb{N}$ *particles* of species i have *mass* $p_i^k > 0$ and are at initial *position* $x_i^{0,k} \in \mathbb{R}^d$, for $k = 1, \dots, N_i$, respectively. Let $N := \sum_{i=1}^n N_i$ and let a family $x = (x_i^k)$ ($k = 1, \dots, N_i$; $i = 1, \dots, n$) of L^2 -absolutely continuous curves $x_i^k \in AC^2([0, \infty); \mathbb{R}^d)$ be given, such that the following initial-value problem for a system of N ordinary differential equations on $\mathbb{R}^d$ is globally solved:

$$\begin{aligned} \frac{d}{dt} x_i^k(t) &= -m_i \sum_{j=1}^n \sum_{l=1}^{N_j} p_j^l \nabla W_{ij}(x_i^k(t) - x_j^l(t)), & x_i^k(0) &= x_i^{0,k}, \\ (k &= 1, \dots, N_i; i = 1, \dots, n). \end{aligned} \quad (26)$$

Then it is easy to verify that the *particle solution*

$$\mu_i(t) = \sum_{k=1}^{N_i} p_i^k \delta_{x_i^k(t)} \quad (i = 1, \dots, n) \quad (27)$$

is the unique gradient flow solution to system (1) with initial datum μ^0 given above. However, it is a non-trivial question if such x exist *globally in time* and are unique, since (W1)–(W5) do not imply global Lipschitz-continuity of the r.h.s. in (26). Nevertheless, (26) admits locally absolutely continuous solutions since this system possesses an underlying (discrete) gradient flow structure: define the finite-dimensional space

$$\mathcal{P}_d := \left\{ x \in \prod_{i=1}^n \prod_{k=1}^{N_i} \mathbb{R}^d \cong \mathbb{R}^{Nd} : p_i = \sum_{k=1}^{N_i} p_i^k \ (i = 1, \dots, n); E = \sum_{i=1}^n \frac{1}{m_i} \sum_{k=1}^{N_i} p_i^k x_i^k \right\},$$

endowed with the (weighted Euclidean) distance

$$\mathbf{d}(x, y) = \left[\sum_{i=1}^n \frac{1}{m_i} \sum_{k=1}^{N_i} p_i^k |x_i^k - y_i^k|^2 \right]^{1/2},$$

and define the *discrete interaction energy* $\mathcal{W}_d$ on $\mathcal{P}_d$ as

$$\mathcal{W}_d(x) := \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \sum_{k=1}^{N_i} \sum_{l=1}^{N_j} p_i^k p_j^l W_{ij}(x_i^k - x_j^l).$$

Applying the same method of proof as for Theorem 2.7 *mutatis mutandis* for the discrete framework, one can show that $\mathcal{W}_d$ is λ_0 -geodesically convex on $(\mathcal{P}_d, \mathbf{d})$ with the same modulus of convexity λ_0 as in the continuous case (17). We can again invoke [2] to obtain existence and uniqueness of a solution curve $x \in AC_{\text{loc}}^2([0, \infty); (\mathcal{P}_d, \mathbf{d}))$ to the particle system (26). Conversely, thanks to the uniqueness of solutions to both (1) and (26), a gradient flow solution μ to (1) of the form (27) can be represented by a solution x to (26).

The particle system (26) might also provide an alternative approach for proving the well-posedness of (1). Approximating the initial datum μ^0 by discrete ones letting the number of particles $N_i \rightarrow \infty$, a suitable existence result on the system (26) of ordinary differential equations and a contractivity estimate like (25) possibly yield also the well-posedness of (1) by passing to the continuum limit. This strategy might also be applicable for more singular

interaction potentials as considered here [13, 19, 22, 37] as well as for systems without gradient structure [30, 31].

3. Qualitative properties in one spatial dimension

In this section, we analyse the qualitative behaviour of the solution from Theorem 2.13 in the general scenario, i.e. the criterion for geodesic convexity may only yield $\lambda_0 \leq 0$. In this case, the contraction estimate (25) does not allow for conclusions on the long-time behaviour of the solution.

From now on, consider (1) in one spatial dimension $d = 1$; and let μ be the solution to (1) with initial datum $\mu^0 \in \mathcal{P}$, as given in Theorem 2.13. First, we rewrite system (1) in terms of the inverse distribution functions $u = (u_1, \dots, u_n)$; recall their definition from (5) and (6).

For all $z \in [0, 1]$, one has $z = F_i(t, u_i(t, z))$. Differentiation w.r.t. t yields

$$\begin{aligned} 0 &= \partial_t F_i(t, u_i(t, z)) + \partial_x F_i(t, u_i(t, z)) \partial_t u_i(t, z) \\ &= \int_{-\infty}^{u_i(t, z)} \frac{1}{p_i} \partial_y \left(\sum_{j=1}^n m_j \mu_j(t, y) (W'_{ij} * \mu_j)(t, y) \right) dy \\ &\quad + \frac{1}{p_i} \mu_i(t, u_i(t, z)) \partial_t u_i(t, z) \\ &= \frac{m_i}{p_i} \sum_{j=1}^n \mu_j(t, u_i(t, z)) \int_{\mathbb{R}} W'_{ij}(u_i(t, z) - y) \mu_j(t, y) dy \\ &\quad + \frac{1}{p_i} \mu_i(t, u_i(t, z)) \partial_t u_i(t, z). \end{aligned}$$

Rearranging yields—with the help of (W3) and the transformation $\xi := F_j(t, y)$:

$$\partial_t u_i(t, z) = m_i \sum_{j=1}^n p_j \int_0^1 W'_{ij}(u_j(t, \xi) - u_i(t, z)) d\xi \quad (i = 1, \dots, n). \quad (28)$$

It is a consequence of Theorem 2.13 that given a gradient flow solution μ to (1), the corresponding curve of pseudo-inverse distribution functions $u \in AC_{\text{loc}}^2([0, \infty); L^2([0, 1]; \mathbb{R}^n))$ solves (28). Furthermore, since $\mu(t) \in \mathcal{P}$ for all $t \geq 0$, $u_i(t, \cdot)$ is a non-decreasing càdlàg function on $(0, 1)$. Conservation of the weighted first moment E over time is reflected in terms of u by the identity

$$E = \sum_{j=1}^n \frac{p_j}{m_j} \int_0^1 u_j(t, z) dz \quad \forall t \geq 0. \quad (29)$$

The concept of inverse distribution functions substantially simplifies the analysis of solutions to (1) since there does not appear any spatial derivative on the right-hand side of (28) anymore. However, this approach can be employed in one spatial dimension $d = 1$ only.

3.1. Speed of propagation and confinement

In this section, we investigate the rate of propagation of the solution to (1) in space over time, given an initial datum with compact support. We first obtain—for arbitrary potentials satisfying (W1)–(W5)—boundedness of the support of $\mu(t)$ for fixed time $t > 0$, and second—under more restrictive requirements on the potential W — t -uniform boundedness of $\text{supp } \mu(t)$.

Proposition 3.1. (Finite speed of propagation) *Let an initial datum μ^0 with compact support and $T > 0$ be given. Then, there exists a constant $K = K(T, \mu^0) > 0$ such that for all $t \in [0, T]$,*

$$\text{supp } \mu(t) \subset [-K, K].$$

Proof. For $t \in [0, T]$ and $i \in \{1, \dots, n\}$, denote

$$\begin{aligned} u_i(t, 1^-) &:= \lim_{\varepsilon \searrow 0} u_i(t, 1 - \varepsilon) \in \mathbb{R} \cup \{+\infty\} \quad \text{and} \\ u_i(t, 0^+) &:= \lim_{\varepsilon \searrow 0} u_i(t, \varepsilon) \in \mathbb{R} \cup \{-\infty\}. \end{aligned}$$

The assertion will follow from finiteness of those limits. Let $\varepsilon > 0$. Then,

$$\partial_t(u_i(t, 1 - \varepsilon)^2) \leq 2|u_i(t, 1 - \varepsilon)| \sum_{j=1}^n m_i p_j \int_0^1 |W'_{ij}(u_j(t, \xi) - u_i(t, 1 - \varepsilon))| d\xi.$$

Lemma 2.4, Hölder's and Young's inequality eventually lead to

$$\begin{aligned} & 2|u_i(t, 1 - \varepsilon)| \sum_{j=1}^n m_i p_j \int_0^1 |W'_{ij}(u_j(t, \xi) - u_i(t, 1 - \varepsilon))| d\xi \\ & \leq 2|u_i(t, 1 - \varepsilon)| \sum_{j=1}^n \bar{C}_{ij} m_i p_j \left(\int_0^1 |u_j(t, \xi)| d\xi + |u_i(t, 1 - \varepsilon)| + 1 \right) \\ & \leq 2 \left[\sum_{j=1}^n \bar{C}_{ij} m_i p_j + 1 \right] u_i(t, 1 - \varepsilon)^2 + \left(\sum_{j=1}^n \bar{C}_{ij} m_i p_j \right)^2 \\ & \quad + 2 \max_j \left(\bar{C}_{ij}^2 m_i^2 m_j p_j \right) \sum_{j=1}^n \int_0^1 \frac{p_j}{m_j} u_j(t, \xi)^2 d\xi. \end{aligned} \tag{30}$$

With the transformation $\xi := F_j(t, x)$, we observe that the sum in the last term on the r.h.s. of (30) can be expressed in terms of the second moments $\mathbf{m}_2(\mu_j(t))$ and of $\mathbf{W}_{\mathcal{P}}(\mu(t), \delta_0 \mathbf{e})$, where $\mathbf{e} = (1, 1, \dots, 1)^T \in \mathbb{R}^n$:

$$\sum_{j=1}^n \int_0^1 \frac{p_j}{m_j} u_j(t, \xi)^2 d\xi = \sum_{j=1}^n \frac{1}{m_j} \mathbf{m}_2(\mu_j(t)) = \mathbf{W}_{\mathcal{P}}^2(\mu(t), \delta_0 \mathbf{e}).$$

Since $\mu \in AC^2([0, T]; (\mathcal{P}, \mathbf{W}_{\mathcal{P}}))$, there exists $\varphi \in L^2([0, T])$ such that

$$\mathbf{W}_{\mathcal{P}}(\mu(t), \mu^0) \leq \int_0^t \varphi(s) ds, \quad \forall t \in [0, T].$$

We obtain

$$\begin{aligned} \sum_{j=1}^n \int_0^1 \frac{p_j}{m_j} u_j(t, \xi)^2 d\xi &\leq 2\mathbf{W}_{\mathcal{D}}^2(\mu(t), \mu^0) + 2\mathbf{W}_{\mathcal{D}}^2(\mu^0, \delta_0 \mathbf{e}) \\ &\leq 2 \left(\int_0^t \varphi(s) ds \right)^2 + 2\mathbf{W}_{\mathcal{D}}^2(\mu^0, \delta_0 \mathbf{e}) \\ &\leq 2T \|\varphi\|_{L^2([0, T])}^2 + 2 \sum_{j=1}^n \frac{1}{m_j} \mathbf{m}_2(\mu_j^0), \end{aligned}$$

which is a constant depending on T and μ^0 . Inserting into (30), we observe

$$\partial_t(u_i(t, 1 - \varepsilon)^2) \leq A u_i(t, 1 - \varepsilon)^2 + B(T, \mu^0),$$

for suitable constants $A, B > 0$. We apply Gronwall's lemma, let $\varepsilon \searrow 0$ and use that—since μ^0 has compact support by assumption—the limit $u_i(0, 1^-)$ exists in $\mathbb{R}$:

$$u_i(t, 1^-)^2 \leq \left[u_i(0, 1^-)^2 + \frac{B}{A} \right] \exp(AT) \quad \forall t \in [0, T].$$

Thus, $u_i(t, 1^-)$ is a finite value, at each $t \in [0, T]$. Along the same lines, it can be shown that

$$u_i(t, 0^+)^2 \leq \left[u_i(0, 0^+)^2 + \frac{B}{A} \right] \exp(AT) \quad \forall t \in [0, T].$$

Since by construction $\text{supp } \mu_i(t) \subset [u_i(t, 0^+), u_i(t, 1^-)]$, the assertion is proven. $\square$

The statement of Proposition 3.1 shows that at fixed $t \geq 0$, the limits

$$u_i(t, 1^-) := \lim_{\varepsilon \searrow 0} u_i(t, 1 - \varepsilon) \quad \text{and} \quad u_i(t, 0^+) := \lim_{\varepsilon \searrow 0} u_i(t, \varepsilon)$$

exist (in $\mathbb{R}$). In order to prove uniform confinement of the solution, we show uniform boundedness of those limits. We first introduce a requirement on the potential by the following

Definition 3.2 (*Confining potentials*) We call an interaction potential W satisfying (W1)–(W5) *confining* if there exists $R > 0$ such that:

- (i) System (1) is *irreducible at large distance*, that is the graph $G' = (V_{G'}, E_{G'})$ with nodes $V_{G'} = \{1, \dots, n\}$ and edges $E_{G'} = \{(i, j) \in V_{G'} \times V_{G'} : W'_{ij} \not\equiv 0 \text{ on } (R, \infty)\}$ is connected.
- (ii) There exists a matrix $C \in \mathbb{R}^{n \times n}$ such that for each $i, j \in \{1, \dots, n\}$, the map W_{ij} is C_{ij} -(semi-)convex on the interval (R, ∞) and the following holds:

If $n = 1$, then $C > 0$. If $n > 1$, with $\tilde{\eta}_i := \min_{j \neq i} C_{ij} p_j$ for all $i \in \{1, \dots, n\}$,

$$\tilde{\lambda}_0 := \min_{i \in \{1, \dots, n\}} \left[p_i \min(0, m_i C_{ii} - \tilde{\eta}_i) + \frac{1}{2} \sum_{j=1}^n p_j \left(\tilde{\eta}_j + \tilde{\eta}_i \frac{m_i}{m_j} \right) \right] > 0. \quad (31)$$

Remark 3.3 (*Geodesic convexity and confinement*) In the scalar case $n = 1$, uniform geodesic convexity of the interaction energy $\mathcal{W}$ is *equivalent* to κ -convexity of W with $\kappa > 0$ [44]. So, the potential is confining. Also for genuine systems, if $\lambda_0 > 0$ in (17), the definition $C := \kappa$ yields $\tilde{\lambda}_0 = \lambda_0 > 0$. Hence, our criterion for uniform geodesic convexity of $\mathcal{W}$ necessarily implies that W is a confining potential. Naturally, if the system is not irreducible at large distance, the independent irreducible subsystems should be considered separately.

Theorem 3.4 (Confinement) *Assume that W is confining and let μ^0 have compact support. Then, there exists a constant $K = K(\mu^0) > 0$ independent of t such that for all $t \geq 0$:*

$$\text{supp } \mu(t) \subset [-K, K]. \quad (32)$$

Proof. We prove the assertion in the case of genuine systems $n > 1$.

Step 1 Estimate on second moments of μ .

Let $\varepsilon > 0$ be sufficiently small such that replacing C_{ij} by $C_{ij}^\varepsilon := C_{ij} - \varepsilon$ in (31) still yields a number $\tilde{\lambda}_0^\varepsilon > 0$, possibly with $\tilde{\lambda}_0^\varepsilon < \tilde{\lambda}_0$. From the C_{ij} -convexity of W_{ij} on (R, ∞) and with the help of Young's inequality, we get for all $z > R$:

$$W_{ij}(z) \geq W_{ij}(R) + W'_{ij}(R)(z - R) + \frac{1}{2}C_{ij}(z - R)^2 \geq \frac{1}{2}C_{ij}^\varepsilon z^2 - D_{ij},$$

for appropriate constants $D_{ij} > 0$. Thanks to (W2)&(W3), enlarging the constants, there exists $D > 0$ such that for all $i, j \in \{1, \dots, n\}$ and for all $z \in \mathbb{R}$:

$$W_{ij}(z) \geq \frac{1}{2}C_{ij}^\varepsilon z^2 - D. \quad (33)$$

We now use boundedness of the energy $\mathcal{W}$ along the (gradient flow) solution to obtain with (33) for all $t \geq 0$:

$$\begin{aligned} 2\mathcal{W}(\mu^0) &\geq 2\mathcal{W}(\mu(t)) \\ &\geq \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \int_{\mathbb{R}} \int_{\mathbb{R}} C_{ij}^\varepsilon (x - y)^2 d\mu_j(x) d\mu_i(y) - D \left(\sum_{j=1}^n p_j \right)^2. \end{aligned}$$

The first term on the r.h.s. has precisely the same structure as the l.h.s. in (18) for $t_i = t_j \equiv 0$ and $\tilde{t}_i = \tilde{t}_j = \text{id}$. Arguing exactly as in the proof of Theorem 2.7, we obtain

$$2\mathcal{W}(\mu^0) \geq \tilde{\lambda}_0^\varepsilon \sum_{j=1}^n \frac{1}{m_j} \mathbf{m}_2(\mu_j(t)) - D \left(\sum_{j=1}^n p_j \right)^2.$$

All in all, we have proven uniform boundedness of the second moments: There exists $C_2 > 0$ such that for all $t \geq 0$ and all $i \in \{1, \dots, n\}$, one has $\mathbf{m}_2(\mu_i(t)) \leq C_2$.

Step 2 L^∞ estimate for u .

We first prove an upper bound. For each $t \geq 0$, we consider those indices $i \in \{1, \dots, n\}$, where

$$u_i(t, 1^-) \geq \max_{j \in \{1, \dots, n\}} u_j(t, 1^-) - R.$$

That is, for all $\xi \in [0, 1)$ and all $j \in \{1, \dots, n\}$: $u_i(t, 1^-) \geq u_j(t, \xi) - R$. We thus have, for each $j \in \{1, \dots, n\}$, a partition of $[0, 1)$ into two sets A_1^j and A_2^j , where

$$\begin{aligned} A_1^j &:= \{\xi \in [0, 1) : u_j(t, \xi) - u_i(t, 1^-) < -R\}, \\ A_2^j &:= \{\xi \in [0, 1) : |u_j(t, \xi) - u_i(t, 1^-)| \leq R\}. \end{aligned}$$

Since W'_{ij} is continuous thanks to (W2), it is bounded on the interval $[-R, R]$. The C_{ij} -convexity of W on $(-\infty, -R)$ yields

$$W'_{ij}(z) - C_{ij}z \leq W'_{ij}(-R) - C_{ij}(-R) \quad \forall z < -R,$$

which can be rewritten as follows using (W3):

$$W'_{ij}(z) \leq C_{ij}z + C_{ij}R - W'_{ij}(R) \quad \forall z < -R.$$

Hence, we obtain

$$\partial_t u_i(t, 1^-) \leq \sum_{j=1}^n \int_{A_1^j} m_i C_{ij} p_j (u_j(t, \xi) - u_i(t, 1^-)) d\xi + C_0,$$

for some constant $C_0 > 0$. Then, with the help of Hölder's inequality,

$$\begin{aligned} & \sum_{j=1}^n \int_{A_1^j} m_i C_{ij} p_j (u_j(t, \xi) - u_i(t, 1^-)) d\xi \\ & \leq m_i \sum_{j=1}^n C_{ij} p_j \left[\int_0^1 (u_j(t, \xi) - u_i(t, 1^-)) d\xi - \int_{A_2^j} (u_j(t, \xi) - u_i(t, 1^-)) d\xi \right] \\ & \leq -m_i \sum_{j=1}^n C_{ij} p_j u_i(t, 1^-) + C' \sum_{j=1}^n \int_0^1 p_j u_j(t, \xi)^2 d\xi + C_1 \\ & = -m_i \sum_{j=1}^n C_{ij} p_j u_i(t, 1^-) + C' \sum_{j=1}^n \mathbf{m}_2(\mu_j(t)) + C_1, \end{aligned}$$

for some constants $C', C_1 > 0$. We now employ step 1 and observe that, as in Proposition 2.9, we have $\sum_{j=1}^n C_{ij} p_j \geq \tilde{\lambda}_0 > 0$:

$$\partial_t u_i(t, 1^-) \leq -m_i \tilde{\lambda}_0 u_i(t, 1^-) + C'',$$

for $C'' > 0$. Gronwall's lemma yields—thanks to $u_i(0, 1^-) < \infty$ —the existence of a constant $K > 0$ such that $\max_{j \in \{1, \dots, n\}} u_j(t, 1^-) \leq K$ for all $t \geq 0$.

In analogy, we now consider those $i \in \{1, \dots, n\}$ such that

$$u_i(t, 0^+) \leq \min_{j \in \{1, \dots, n\}} u_j(t, 0^+) + R,$$

yielding for each $j \in \{1, \dots, n\}$ a partition $[0, 1) = B_1^j \cup B_2^j$ with

$$\begin{aligned} B_1^j &:= \{\xi \in [0, 1) : u_j(t, \xi) - u_i(t, 0^+) > R\}, \\ B_2^j &:= \{\xi \in [0, 1) : |u_j(t, \xi) - u_i(t, 0^+)| \leq R\}. \end{aligned}$$

Similarly as above, using the symmetry property (W3), we get

$$\begin{aligned} & -\partial_t u_i(t, 0^+) \\ & \leq -m_i \sum_{j=1}^n C_{ij} p_j(-u_i(t, 0^+)) \\ & \quad + C' \sum_{j=1}^n \mathbf{m}_2(\mu_j(t)) + C_1 \leq -m_i \tilde{\lambda}_0(-u_i(t, 0^+)) + C'', \end{aligned}$$

allowing us to proceed as before.

Putting the bounds together finishes the proof:

$$\sup_{t \geq 0} \max_{j \in \{1, \dots, n\}} \|u_j(t, \cdot)\|_{L^\infty([0,1])} \leq K.$$

□

We thus know, given a confining potential, that the solution lives on a fixed compact interval. It is now a natural question to ask if, for absolutely continuous initial conditions, partial or total collapse of the support can occur in *finite* time. This question is addressed in the following

Proposition 3.5 (Exclusion of finite-time blow-up) *Let $i \in \{1, \dots, n\}$ be fixed, but arbitrary. Assume that for all $j \in \{1, \dots, n\}$ the maps W'_{ij} are Lipschitz-continuous. Suppose moreover that $\text{supp } \mu_i^0$ is a (possibly unbounded) interval and μ_i^0 is absolutely continuous w.r.t. the Lebesgue measure. Assume that its Lebesgue density is continuous on the interior of $\text{supp } \mu_i^0$ and globally bounded. Then, $\mu_i(t)$ is absolutely continuous for all $t \geq 0$.*

Proof. Our method of proof is an adaptation of the proof of [15, Thm. 2.9] to the situation at hand. We show that for all $t \geq 0$, there exists $\gamma(t) > 0$ such that for all $z \in [0, 1)$ and all $h > 0$ with $z + h < 1$:

$$\frac{1}{h}(u_i(t, z + h) - u_i(t, z)) \geq \gamma(t) > 0. \quad (34)$$

That is, $u_i(t, \cdot)$ is *strictly* increasing at each $t \geq 0$. The assumptions on the initial datum above ensure that (34) is true at $t = 0$ with some $\gamma(0) > 0$. If (34) holds at a given t_0 , the cumulative distribution function $F_i(t_0, \cdot)$ is Lipschitz-continuous, which implies absolute continuity of $\mu_i(t_0)$.

From (28), we get

$$\begin{aligned} & \partial_t(u_i(t, z + h) - u_i(t, z)) \\ & = m_i \sum_{j=1}^n p_j \int_0^1 [W'_{ij}(u_j(t, \xi) - u_i(t, z + h)) - W'_{ij}(u_j(t, \xi) - u_i(t, z))] \, d\xi. \end{aligned}$$

Denote by $L_{ij} > 0$ the Lipschitz constant of W'_{ij} . From the monotonicity $u_i(t, z + h) - u_i(t, z) \geq 0$, it follows that

$$\partial_t(u_i(t, z + h) - u_i(t, z)) \geq -m_i \sum_{j=1}^n L_{ij} p_j(u_i(t, z + h) - u_i(t, z)).$$

We subsequently obtain for $\tilde{C}_i := m_i \sum_{j=1}^n L_{ij} p_j$ that

$$\partial_t[(u_i(t, z+h) - u_i(t, z))e^{\tilde{C}_i t}] \geq 0,$$

and hence

$$\frac{1}{h}(u_i(t, z+h) - u_i(t, z)) \geq \frac{1}{h}e^{-\tilde{C}_i t}(u_i(0, z+h) - u_i(0, z)) \geq e^{-\tilde{C}_i t}\gamma(0) > 0.$$

Letting $\gamma(t) := e^{-\tilde{C}_i t}\gamma(0)$, (34) follows. $\square$

Naturally, the above result does not extend to $t \rightarrow \infty$ since e.g. in the uniformly geodesically convex case, the solution collapses to a Dirac measure in the large-time limit.

3.2. Long-time behaviour

We now analyse the long-time behaviour of the solution to (1) in the non-uniformly convex case.

Theorem 3.6 (Long-time behaviour) *Assume that the solution μ to (1) is uniformly confined, i.e. there exists $K > 0$ such that $\text{supp } \mu_i(t) \subset [-K, K]$ holds for all $t \geq 0$ and all $i \in \{1, \dots, n\}$ as in (32). Moreover, suppose that the maps W'_{ij} are Lipschitz-continuous on the interval $[-2K, 2K]$ for all $i, j \in \{1, \dots, n\}$. Set, for $t \geq 0$, $\mathcal{W}^t := \mathcal{W}(\mu(t))$. The following holds:*

(a) *There exists $\mathcal{W}^\infty \in \mathbb{R}$ such that*

$$\lim_{t \rightarrow \infty} \mathcal{W}^t = \mathcal{W}^\infty, \quad (35)$$

and

$$\lim_{t \rightarrow \infty} \left(\frac{d}{dt} \mathcal{W}^t \right) = 0. \quad (36)$$

(b) *For each sequence $(t_k)_{k \in \mathbb{N}}$ in $(0, \infty)$ with $t_k \rightarrow \infty$ as $k \rightarrow \infty$, there exists a subsequence $(t_{k_l})_{l \in \mathbb{N}}$ and a steady state $\bar{\mu} \in \mathcal{P}$ of (1) such that for all $i \in \{1, \dots, n\}$:*

$$\lim_{l \rightarrow \infty} \mathbf{W}_1(\mu_i(t_{k_l}), \bar{\mu}_i) = 0. \quad (37)$$

Thus, the ω -limit set of the dynamical system associated to (1) can only contain steady states of (1).

Proof. We proceed similarly to the proof of [48, Prop. 1] and observe that along the solution μ , the dissipation of $\mathcal{W}$ reads

$$\begin{aligned} \frac{d}{dt} \mathcal{W}^t &= - \sum_{i=1}^n m_i \int_{\mathbb{R}} \left(\sum_{j=1}^n W'_{ij} * \mu_j(t) \right)^2 d\mu_i(t) \\ &= - \sum_{i=1}^n m_i p_i \int_0^1 \left(\sum_{j=1}^n p_j \int_0^1 W'_{ij}(u_i(t, z) - u_j(t, \xi)) d\xi \right)^2 dz, \end{aligned} \quad (38)$$

which is non-positive. By (32), all u_i are bounded in time and space by the constant K . Since W'_{ij} is Lipschitz-continuous, it is differentiable almost everywhere on $[-2K, 2K]$. So, another differentiation of the dissipation w.r.t. t shows

$$\begin{aligned} \frac{d^2}{dt^2} \mathcal{W}^t &= -2 \sum_{i=1}^n m_i p_i \int_0^1 \left(\sum_{j=1}^n \int_0^1 p_j W'_{ij}(u_i(t, z) - u_j(t, \zeta)) d\zeta \right) \\ &\quad \times \left(\sum_{j=1}^n p_j \int_0^1 W''_{ij}(u_i(t, z) - u_j(t, \xi)) \right. \\ &\quad \times \left[\sum_{k=1}^n m_i p_k \int_0^1 W'_{ik}(u_k(t, \zeta) - u_i(t, z)) d\zeta \right. \\ &\quad \left. \left. - \sum_{k=1}^n m_j p_k \int_0^1 W'_{jk}(u_k(t, \zeta) - u_j(t, \xi)) d\zeta \right] d\xi \right) dz. \end{aligned}$$

By elementary estimates, using in particular that $|W''_{ij}(z)| \leq L_{ij}$ a.e. on $[-2K, 2K]$ by Lipschitz-continuity, we find

$$\sup_{t \geq 0} \left| \frac{d^2}{dt^2} \mathcal{W}^t \right| \leq C_2, \quad (39)$$

for some $C_2 > 0$. Furthermore, it is easy to conclude from (W2) and (32) that

$$\inf_{t \geq 0} \mathcal{W}^t \geq -C_0, \quad (40)$$

for another constant $C_0 > 0$. Putting (38) and (40) together yields the existence of $\mathcal{W}^\infty \in \mathbb{R}$ such that (35) holds. We now use (39) to prove (36):

Define for $t > \frac{2}{C_2} \sqrt{\mathcal{W}^0 - \mathcal{W}^\infty} > 0$ the positive quantity $\tau(t) := \frac{1}{C_2} \sqrt{\mathcal{W}^{t/2} - \mathcal{W}^\infty}$. Since $\mathcal{W}^t$ is nonincreasing, we also have $\tau(t) < \frac{t}{2}$. Moreover,

$$\begin{aligned} \frac{d}{dt} \mathcal{W}^t &= \frac{1}{\tau(t)} (\mathcal{W}^t - \mathcal{W}^\infty) - \frac{1}{\tau(t)} (\mathcal{W}^{\tau(t)} - \mathcal{W}^\infty) \\ &\quad + \frac{1}{\tau(t)} \int_{t-\tau(t)}^t \int_s^t \frac{d^2}{d\sigma^2} \mathcal{W}^\sigma d\sigma ds, \end{aligned}$$

from which with (39) and (35) the desired result (36) follows:

$$\begin{aligned} \left| \frac{d}{dt} \mathcal{W}^t \right| &\leq \frac{2}{\tau(t)} (\mathcal{W}^{t/2} - \mathcal{W}^\infty) + \frac{1}{\tau(t)} C_2 \tau(t)^2 \\ &= (2C_2 + 1) \sqrt{\mathcal{W}^{t/2} - \mathcal{W}^\infty} \xrightarrow{t \rightarrow \infty} 0. \end{aligned}$$

For part (b), let a sequence of time points $t_k \rightarrow \infty$ be given. The family $u_i(t_k, \cdot)$ ($k \in \mathbb{N}$; $i \in \{1, \dots, n\}$) of nondecreasing functions is uniformly bounded in $L^\infty([0, 1])$ (by the constant K)—hence bounded in the space $BV([0, 1])$. Thus, there exist a subsequence $(t_{k_l})_{l \in \mathbb{N}}$ and nondecreasing maps $\bar{u}_1, \dots, \bar{u}_n$ with $\|\bar{u}_i\|_{L^\infty([0, 1])} \leq K$ for all $i \in \{1, \dots, n\}$, such that $u_i(t_{k_l}, \cdot)$ converges to $\bar{u}_i$ in $L^1([0, 1])$ and almost everywhere on $[0, 1]$, as $l \rightarrow \infty$ (for details, see e.g. [1]). The corresponding measure $\bar{\mu}$ belongs to $\mathcal{P}$ thanks to the dominated

convergence theorem. It remains to show that $\bar{\mu}$ is a steady state of system (1). Define

$$\omega := - \sum_{i=1}^n m_i p_i \int_0^1 \left(\sum_{j=1}^n p_j \int_0^1 W'_{ij}(\bar{u}_i(z) - \bar{u}_j(\xi)) d\xi \right)^2 dz.$$

Elementary calculations—using e.g. Lemma 2.4—show

$$\begin{aligned} & \left| \frac{d}{dt} \mathcal{W}^{t_{k_l}} - \omega \right| \\ & \leq C_0 \sum_{i=1}^n \sum_{j=1}^n \int_0^1 \int_0^1 |W'_{ij}(u_i(t_{k_l}, z) - u_j(t_{k_l}, \xi)) - W'_{ij}(\bar{u}_i(z) - \bar{u}_j(\xi))| d\xi dz, \end{aligned}$$

for a suitable constant $C_0 > 0$. The Lipschitz-continuity of the W'_{ij} on $[-2K, 2K]$, the triangle inequality and (37) then imply

$$\begin{aligned} & \left| \frac{d}{dt} \mathcal{W}^{t_{k_l}} - \omega \right| \\ & \leq C_0 \sum_{i=1}^n \sum_{j=1}^n L_{ij} (\|u_i(t_{k_l}, \cdot) - \bar{u}_i\|_{L^1([0,1])} + \|u_j(t_{k_l}, \cdot) - \bar{u}_j\|_{L^1([0,1])}) \xrightarrow{t \rightarrow \infty} 0. \end{aligned}$$

Hence, because of (36), $\omega = 0$. Specifically, this means that for each $i \in \{1, \dots, n\}$ and almost every $x \in \text{supp } \bar{\mu}_i$, the following holds:

$$\sum_{j=1}^n \int_{\mathbb{R}} W'_{ij}(x - y) d\bar{\mu}_j(y) = 0.$$

So, $\bar{\mu}$ is a solution to (1) and the proof is complete. $\square$

Remark 3.7 The result of Theorem 3.6 does neither yield uniqueness of steady states of (1) nor convergence of the entire curve μ to some specific element in $\mathcal{P}$ as $t \rightarrow \infty$. Due to the possible non-strict convexity of $\mathcal{W}$, the ω -limit set might contain more than one element. Still, as it is also the case for a finite-dimensional gradient system, every ω -limit point is an equilibrium. Thus, the solution orbits might in general approach, as $t \rightarrow \infty$, a continuum of steady states on certain level sets of the energy $\mathcal{W}$.

Our result may be viewed as an extension of the well-known *LaSalle invariance principle* from the theory of ordinary differential equations: The existence of the Lyapunov functional $\mathcal{W}$ and certain compactness properties (here, the confinement of the solution) guarantee convergence of the solution orbit to the set with vanishing energy dissipation.

If there only exists the trivial steady state μ^∞ from Proposition 2.15 in the set of those elements from $\mathcal{P}$ with support contained in $[-K, K]$, then Theorem 3.6 implies

$$\lim_{t \rightarrow \infty} \mathbf{W}_1(\mu_i(t), \mu_i^\infty) = 0 \quad \forall i \in \{1, \dots, n\},$$

without obtaining any specific rate of convergence.

If the potential is not confining, convergence may not occur. However, we might observe that the support of each component μ_i collapses to a single (but not necessarily fixed) point as $t \rightarrow \infty$.

Proposition 3.8 (Collapse of the support) *Let $i \in \{1, \dots, n\}$ be fixed, but arbitrary. Assume that the support of μ_i^0 is compact and that $S_i := \sum_{j=1}^n \kappa_{ij} p_j > 0$ holds. Then,*

$$\text{diam supp } \mu_i(t) \leq e^{-m_i S_i t} \text{diam supp } \mu_i^0.$$

That is, the support of μ_i contracts at exponential speed.

Proof. Recall that $\text{diam supp } \mu_i(t) = u_i(t, 1^-) - u_i(t, 0^+)$. We have

$$\begin{aligned} & \partial_t(u_i(t, 1^-) - u_i(t, 0^+)) \\ &= \sum_{j=1}^n m_i p_j \int_0^1 [W'_{ij}(u_j(t, \xi) - u_i(t, 1^-)) - W'_{ij}(u_j(t, \xi) - u_i(t, 0^+))] \, d\xi \\ &\leq \sum_{j=1}^n m_i p_j \kappa_{ij} \int_0^1 [(u_j(t, \xi) - u_i(t, 1^-)) - (u_j(t, \xi) - u_i(t, 0^+))] \, d\xi \\ &= -m_i S_i (u_i(t, 1^-) - u_i(t, 0^+)), \end{aligned}$$

the second-to-last step being a consequence of κ_{ij} -convexity (W5). Applying Gronwall's lemma completes the proof. $\square$

In the regime where Proposition 3.8 is applicable for all $i \in \{1, \dots, n\}$, system (1) behaves asymptotically as $t \rightarrow \infty$ like the particle system (26) in the case of only one (heavy) particle for each component ($N_i = 1$ for all i). Obviously, by Proposition 2.9, the condition $S_i > 0$ above is met in the scenario with uniformly geodesically convex energy. This enables us to improve the convergence result from Sect. 2.2 in one spatial dimension for compactly supported initial data:

Proposition 3.9 (The uniformly convex case in one spatial dimension) *Assume that the criterion for geodesic convexity (17) yields $\lambda_0 > 0$ and suppose that μ^0 has compact support. Then, for each $i \in \{1, \dots, n\}$,*

$$\lim_{t \rightarrow \infty} \mathbf{W}_\infty(\mu_i(t), \mu_i^\infty) = 0,$$

where μ^∞ is the unique steady state from Proposition 2.15.

In view of Proposition 2.15, we obtain convergence w.r.t. the stronger topology of the L^∞ -Wasserstein distance, but lose the exponential rate of convergence.

Proof. Fix $i \in \{1, \dots, n\}$. Since $\lambda_0 > 0$ and $\text{supp } \mu_i^0$ is compact, we know from Proposition 2.15, Theorem 3.4 and Proposition 3.8 that $\|u_i(t, \cdot) - x^\infty\|_{L^2([0,1])} \rightarrow 0$ as $t \rightarrow \infty$, $\|u_i(t, \cdot)\|_{L^\infty([0,1])} \leq K$ for all $t \geq 0$ and $u_i(t, 1^-) - u_i(t, 0^+) \rightarrow 0$ as

$t \rightarrow \infty$. Obviously, if $\lim_{t \rightarrow \infty} u_i(t, 1^-) = x^\infty$ holds, the desired result follows immediately from

$$\|u_i(t, \cdot) - x^\infty\|_{L^\infty([0,1])} = \max(|u_i(t, 1^-) - x^\infty|, |u_i(t, 0^+) - x^\infty|),$$

since then also $\lim_{t \rightarrow \infty} u_i(t, 0^+) = x^\infty$ holds. So, assume that $u_i(t, 1^-)$ does *not* converge to x^∞ as $t \rightarrow \infty$. Then, there exists $\varepsilon > 0$ and a sequence $t_k \rightarrow \infty$ such that

$$|u_i(t_k, 1^-) - x^\infty| \geq \varepsilon \quad \forall k \in \mathbb{N}. \quad (41)$$

Thanks to the observations above, there exist a subsequence $(t_{k_l})_{l \in \mathbb{N}}$ and $\omega \in \mathbb{R}$ such that

$$\lim_{l \rightarrow \infty} (u_i(t_{k_l}, 1^-) - x^\infty) = \omega, \quad \text{and} \quad \lim_{l \rightarrow \infty} u_i(t_{k_l}, z) = x^\infty \quad \text{for a.e. } z \in (0, 1).$$

Immediately, it follows that $\lim_{l \rightarrow \infty} u_i(t_{k_l}, 0^+) = x^\infty + \omega$ and consequently $\omega = 0$ by monotonicity. But $\omega = 0$ is a contradiction to (41). $\square$

Acknowledgements

This research has been supported by the German Research Foundation (DFG), SFB TRR 109. The author thanks Daniel Matthes for helpful discussion and comments.

References

- [1] Ambrosio, L., Fusco, N., Pallara, D.: (2000) Functions of bounded variation and free discontinuity problems. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York (2000)
- [2] Ambrosio, L., Gigli, N., Savaré, G.: Gradient Flows in Metric Spaces and in the Space of Probability Measures, 2nd edn. Lectures in Mathematics ETH Zürich. Birkhäuser Verlag, Basel (2008)
- [3] Balagué, D., Carrillo, J.A., Yao, Y.: Confinement for repulsive-attractive kernels. Discret. Contin. Dyn. Syst. Ser. B **19**(5), 1227–1248 (2014)
- [4] Benedetto, D., Caglioti, E., Pulvirenti, M.: A kinetic equation for granular media. RAIRO Modél. Math. Anal. Numér **31**(5), 615–641 (1997)
- [5] Bertozzi, A.L., Brandman, J.: Finite-time blow-up of L^∞ -weak solutions of an aggregation equation. Commun. Math. Sci. **8**(1), 45–65 (2010)
- [6] Bertozzi, A.L., Carrillo, J.A., Laurent, T.: Blow-up in multidimensional aggregation equations with mildly singular interaction kernels. Nonlinearity **22**(3), 683–710 (2009)
- [7] Bertozzi, A.L., Laurent, T.: Finite-time blow-up of solutions of an aggregation equation in $\mathbf{R}^n$. Comm. Math. Phys **274**(3), 717–735 (2007)

- [8] Bertozzi, A.L., Laurent, T., Rosado, J.: L^p theory for the multidimensional aggregation equation. *Comm. Pure Appl. Math.* **64**(1), 45–83 (2011)
- [9] Biler, P., Karch, G., Laurençot, P.: Blowup of solutions to a diffusive aggregation model. *Nonlinearity* **22**(7), 1559–1568 (2009)
- [10] Blanchet, A., Carlen, E.A., Carrillo, J.A.: Functional inequalities, thick tails and asymptotics for the critical mass Patlak–Keller–Segel model. *J. Funct. Anal.* **262**(5), 2142–2230 (2012)
- [11] Blanchet, A., Dolbeault, J., Perthame, B.: Two-dimensional Keller–Segel model: optimal critical mass and qualitative properties of the solutions. *Electron. J. Differ. Equ.* pages No. 44, pp. 32 (2006) (electronic)
- [12] Bodnar, M., Velázquez, J.J.L.: An integro-differential equation arising as a limit of individual cell-based models. *J. Differ. Equ.* **222**(2), 341–380 (2006)
- [13] Bonaschi, G.A., Carrillo, J.A., Di Francesco, M., Peletier, M.A.: Equivalence of gradient flows and entropy solutions for singular nonlocal interaction equations in 1D. *ESAIM Control Optim. Calc. Var.* **21**(2), 414–441 (2015)
- [14] Burger, M., Capasso, V., Morale, D.: On an aggregation model with long and short range interactions. *Nonlinear Anal. Real World Appl.* **8**(3), 939–958 (2007)
- [15] Burger, M., Di Francesco, M.: Large time behavior of nonlocal aggregation models with nonlinear diffusion. *Netw. Heterog. Media* **3**(4), 749–785 (2008)
- [16] Cañizo, J.A., Carrillo, J.A., Patacchini, F.S.: Existence of compactly supported global minimisers for the interaction energy. *Arch. Ration. Mech. Anal.* **217**(3), 1197–1217 (2015)
- [17] Cañizo, J.A., Carrillo, J.A., Rosado, J.: A well-posedness theory in measures for some kinetic models of collective motion. *Math. Models Methods Appl. Sci.* **21**(3), 515–539 (2011)
- [18] Carrillo, J.A., Castorina, D., Volzone, B.: Ground states for diffusion dominated free energies with logarithmic interaction. *SIAM J. Math. Anal.* **47**(1), 1–25 (2015)
- [19] Carrillo, J.A., Di Francesco, M., Figalli, A., Laurent, T., Slepčev, D.: Global-in-time weak measure solutions and finite-time aggregation for nonlocal interaction equations. *Duke Math. J.* **156**(2), 229–271 (2011)
- [20] Carrillo, J.A., Di Francesco, M., Figalli, A., Laurent, T., Slepčev, D.: Confinement in nonlocal interaction equations. *Nonlinear Anal.* **75**(2), 550–558 (2012)
- [21] Carrillo, J.A., D’Orsogna, M.R., Panferov, V.: Double milling in self-propelled swarms from kinetic theory. *Kinet. Relat. Models* **2**(2), 363–378 (2009)
- [22] Carrillo, J.A., James, F., Lagoutière, F., Vauchelet, N.: The Filippov characteristic flow for the aggregation equation with mildly singular potentials. *J. Differ. Equ.* **260**(1), 304–338 (2016)

- [23] Carrillo, J.A., Lisini, S., Mainini, E.: Gradient flows for non-smooth interaction potentials. *Nonlinear Anal.* **100**, 122–147 (2014)
- [24] Carrillo, J.A., McCann, R.J., Villani, C.: Kinetic equilibration rates for granular media and related equations: entropy dissipation and mass transportation estimates. *Rev. Mat. Iberoam.* **19**(3), 971–1018 (2003)
- [25] Carrillo, J.A., McCann, R.J., Villani, C.: Contractions in the 2-Wasserstein length space and thermalization of granular media. *Arch. Ration. Mech. Anal.* **179**(2), 217–263 (2006)
- [26] Carrillo, J.A., Rosado, J.: Uniqueness of bounded solutions to aggregation equations by optimal transport methods. In: *European Congress of Mathematics*, pp. 3–16. European Mathematical Society, Zürich (2010)
- [27] Carrillo, J.A., Toscani, G.: Wasserstein metric and large-time asymptotics of nonlinear diffusion equations. In: Fergola, P., Capone, F., Gentile, M., Guerriero, G. (eds.) *New Trends in Mathematical Physics*, pp. 234–244. World Scientific Publishing, Hackensack, NJ, (2004)
- [28] Colombo, R.M., Lécureux-Mercier, M.: Nonlocal crowd dynamics models for several populations. *Acta Math. Sci. Ser. B Engl. Ed* **32**(1), 177–196 (2012)
- [29] Crippa, G., Lécureux-Mercier, M.: Existence and uniqueness of measure solutions for a system of continuity equations with non-local flow. *NoDEA Nonlinear Differ. Equ. Appl.* **20**(3), 523–537 (2013)
- [30] Di Francesco, M., Fagioli, S.: Measure solutions for non-local interaction PDEs with two species. *Nonlinearity* **26**(10), 2777–2808 (2013)
- [31] Di Francesco, M., Fagioli, S.: A nonlocal swarm model for predators–prey interactions. *Math. Models Methods Appl. Sci.* **26**(2), 319–355 (2016)
- [32] Evans, L.C.: *Partial Differential Equations*, vol. 19 of Graduate Studies in Mathematics, 2nd edn. American Mathematical Society, Providence (2010)
- [33] Fellner, K., Raoul, G.: Stable stationary states of non-local interaction equations. *Math. Models Methods Appl. Sci.* **20**(12), 2267–2291 (2010)
- [34] Fellner, K., Raoul, G.: Stability of stationary states of non-local equations with singular interaction potentials. *Math. Comput. Modell.* **53**(7–8), 1436–1450 (2011)
- [35] Giacomini, G., Lebowitz, J.L., Marra, R.: Macroscopic evolution of particle systems with short- and long-range interactions. *Nonlinearity* **13**(6), 2143–2162 (2000)
- [36] Golse, F.: The mean-field limit for the dynamics of large particle systems. *Journées “Équations aux Dérivées Partielles”*, pp. Exp. No. IX, 1–47. Univ. Nantes, Nantes (2003)
- [37] James, F., Vauchelet, N.: Equivalence between duality and gradient flow solutions for one-dimensional aggregation equations. *Discret. Contin. Dyn. Syst.* **36**(3), 1355–1382 (2016)

- [38] Jordan, R., Kinderlehrer, D., Otto, F.: The variational formulation of the Fokker–Planck equation. *SIAM J. Math. Anal.* **29**(1), 1–17 (1998)
- [39] Kang, K., Perthame, B., Stevens, A., Velázquez, J.J.L.: An integro-differential equation model for alignment and orientational aggregation. *J. Differ. Equ.* **246**(4), 1387–1421 (2009)
- [40] Kolokolnikov, T., Huang, Y., Pavlovski, M.: Singular patterns for an aggregation model with a confining potential. *Phys. D* **260**, 65–76 (2013)
- [41] Laurent, T.: Local and global existence for an aggregation equation. *Commun. Partial Differ. Equ.* **32**(10–12), 1941–1964 (2007)
- [42] Li, H., Toscani, G.: Long-time asymptotics of kinetic models of granular flows. *Arch. Ration. Mech. Anal.* **172**(3), 407–428 (2004)
- [43] Luckhaus, S., Sugiyama, Y., Velázquez, J.J.L.: Measure valued solutions of the 2D Keller–Segel system. *Arch. Ration. Mech. Anal.* **206**(1), 31–80 (2012)
- [44] McCann, R.J.: A convexity principle for interacting gases. *Adv. Math.* **128**(1), 153–179 (1997)
- [45] Mogilner, A., Edelstein-Keshet, L.: A non-local model for a swarm. *J. Math. Biol.* **38**(6), 534–570 (1999)
- [46] Morale, D., Capasso, V., Oelschläger, K.: An interacting particle system modelling aggregation behavior: from individuals to populations. *J. Math. Biol.* **50**(1), 49–66 (2005)
- [47] Otto, F.: The geometry of dissipative evolution equations: the porous medium equation. *Commun. Partial Differ. Equ.* **26**(1–2), 101–174 (2001)
- [48] Raoul, G.: Nonlocal interaction equations: stationary states and stability analysis. *Differ. Integral Equ.* **25**(5–6), 417–440 (2012)
- [49] Sun, H., Uminsky, D., Bertozzi, A.L.: Stability and clustering of self-similar solutions of aggregation equations. *J. Math. Phys.* **53**(11), 115610 (2012)
- [50] Theil, F.: A proof of crystallization in two dimensions. *Commun. Math. Phys.* **262**(1), 209–236 (2006)
- [51] Topaz, C.M., Bertozzi, A.L.: Swarming patterns in a two-dimensional kinematic model for biological groups. *SIAM J. Appl. Math.* **65**(1), 152–174 (2004)
- [52] Topaz, C.M., Bertozzi, A.L., Lewis, M.A.: A nonlocal continuum model for biological aggregation. *Bull. Math. Biol.* **68**(7), 1601–1623 (2006)
- [53] Toscani, G.: One-dimensional kinetic models of granular flows. *M2NMath. Model. Numer. Anal* **34**(6), 1277–1291 (2000)
- [54] Toscani, G.: Kinetic models of opinion formation. *Commun. Math. Sci.* **4**(3), 481–496 (2006)
- [55] Giessen, A.E. van., Widom, B.: Path dependence of surface-tension scaling in binary mixtures. *Fluid Phase Equilib.* **164**(1), 1–12 (1999)

- [56] Villani, C.: Topics in Optimal Transportation. Graduate Studies in Mathematics, vol. 58. American Mathematical Society, Providence (2003)
- [57] Zhang, J., Kwok, D.Y.: A mean-field free energy lattice Boltzmann model for multicomponent fluids. *Eur. Phys. J. Spec. Top* **171**(1), 45–53 (2009)
- [58] Zinsl, J., Matthes, D.: Transport distances and geodesic convexity for systems of degenerate diffusion equations. *Calc. Var. Partial Differ. Equ.* **54**(4), 3397–3438 (2015)

Jonathan Zinsl
Zentrum für Mathematik
Technische Universität München
85747 Garching, Germany
e-mail: zinsl@ma.tum.de

Received: 17 December 2015.

Accepted: 9 July 2016.