

JORDAN–KRONECKER INVARIANTS OF LIE ALGEBRA REPRESENTATIONS AND DEGREES OF INVARIANT POLYNOMIALS

A. BOLSINOV*

Department
of Mathematical Sciences
Loughborough University
Loughborough, LE11 3TU, UK

and

Faculty
of Mechanics and Mathematics
Moscow State University
Moscow, Russia, 119991
A.Bolsinov@lboro.ac.uk

A. IZOSIMOV**

Department
of Mathematics
University of Arizona
Tucson AZ 85721, USA
izosimov@math.arizona.edu

I. KOZLOV***

Faculty
of Mechanics and Mathematics
Moscow State University
Moscow, Russia, 119991
ikozlov90@gmail.com

Abstract. For an arbitrary representation ρ of a complex finite-dimensional Lie algebra, we construct a collection of numbers that we call the *Jordan–Kronecker invariants* of ρ . Among other interesting properties, these numbers provide lower bounds for degrees of polynomial invariants of ρ . Furthermore, we prove that these lower bounds are exact if and only if the invariants are independent outside of a set of large codimension. Finally, we show that under certain additional assumptions our bounds are exact if and only if the algebra of invariants is freely generated.

1. Introduction

The main idea of this paper has its roots in the theory of bi-Hamiltonian systems where it was discovered that the algebraic structure of a pair of compatible Poisson brackets $\{, \}_0$ and $\{, \}_1$ essentially affects the differential geometry of the pencil $\{, \}_0 + \lambda \{, \}_1$

DOI: 10.1007/s00031-021-09661-0

*Supported by the Russian Science Foundation, project No.17-11-01303.

**Supported by NSF grant DMS-2008021.

***Supported by the Russian Science Foundation, project No.17-11-01303.

Received August 30, 2020. Accepted December 17, 2020.

Published online June 17, 2021.

Corresponding Author: A. Bolsinov, e-mail: A.Bolsinov@lboro.ac.uk

and even the dynamical properties of bi-Hamiltonian systems related to it (see, e.g., [16], [18], [10], [2], [1]). This observation has recently been used in [3] to introduce Jordan–Kronecker invariants for a finite-dimensional Lie algebra \mathfrak{g} , which are directly related to a natural pencil of compatible Poisson brackets on \mathfrak{g}^* . From the algebraic viewpoint, this construction is based on a simple fact that every element $x \in \mathfrak{g}^*$ defines a natural skew-symmetric bilinear form $\mathcal{A}_x(\xi, \eta) = \langle x, [\xi, \eta] \rangle$ on \mathfrak{g} . The Jordan–Kronecker invariant of \mathfrak{g} is, by definition, the algebraic type of the pencil of forms $\mathcal{A}_{a+\lambda b}$ for a generic pair $(a, b) \in \mathfrak{g}^* \times \mathfrak{g}^*$. All possible algebraic types are described by the Jordan–Kronecker theorem on a canonical form of a pencil of skew-symmetric matrices (see, e.g., [15], [4]) and, in each dimension, there are only finitely many of them.

In the present paper, this construction is generalized to arbitrary finite-dimensional representations of finite-dimensional Lie algebras. In particular, we show how the classical theorem on the canonical form of a pair of linear maps can be applied in the study of Lie algebra representations and their invariants. In what follows, we assume that all objects are defined over the field \mathbb{C} of complex numbers, although everything can be generalized to the case of an arbitrary field of characteristic zero (in which case some of the objects we consider are defined over the algebraic closure of the initial field).

Our construction can be outlined as follows. Let $\rho : \mathfrak{g} \rightarrow \mathfrak{gl}(V)$ be a linear representation of a finite-dimensional Lie algebra \mathfrak{g} on a finite-dimensional vector space V . To this representation and an arbitrary element $x \in V$, one can naturally associate an operator $R_x : \mathfrak{g} \rightarrow V$ defined by $R_x(\xi) = \rho(\xi)x$. Consider a pair of such operators R_a, R_b and the pencil $R_a + \lambda R_b = R_{a+\lambda b}$ generated by them. It is well known that such a pencil can be completely characterized by a collection of quite simple numerical invariants (see Section 2 for details). In the present paper we show that many important and interesting properties of the representation $\rho : \mathfrak{g} \rightarrow \mathfrak{gl}(V)$ are related to and can be derived from the invariants of such a pencil $R_{a+\lambda b}$ generated by a generic pair $(a, b) \in V \times V$. In particular, one of our main results gives lower bounds for degrees of invariant polynomials of the representation ρ in terms of the numerical invariants of the associated pair of operators R_a, R_b (Theorem 26). Furthermore, we show that these lower bounds are exact if and only if the invariant polynomials of ρ are independent outside of a subset of codimension ≥ 2 (Theorem 31). Finally, we prove that under certain additional assumptions our bound is exact if and only if the algebra of invariant polynomials of ρ is freely generated (Theorem 36). This generalizes several previously known results (see, in particular, [8], [5], [9]) which relate polynomiality of the algebra of invariants with the *sum* of degrees of the invariants (while our statements concern *individual* invariants). In the last Section 6 we show that our results have quite non-trivial implications already in the case of standard representations of simple Lie groups.

2. Canonical form of a pair of linear maps

In this section, we recall the normal form theorem for a pair of linear maps. We first state the theorem in the matrix form, and then discuss the invariant meaning of the ingredients involved.

Theorem 1 (On the Jordan–Kronecker normal form [4]). *Consider two complex vector spaces U and V . Then for every two linear maps $A, B : U \rightarrow V$ there are bases in U and V in which the matrices of the pencil $\mathcal{P} = \{A + \lambda B\}$ have the following block-diagonal form:*

$$A + \lambda B = \text{diag}(0_{m,n}, A_1 + \lambda B_1, \dots, A_k + \lambda B_k) \quad (1)$$

where $0_{m,n}$ is the zero $m \times n$ -matrix, and each pair of the corresponding blocks A_i and B_i takes one of the following forms:

1. *Jordan block with eigenvalue* $\lambda_0 \in \mathbb{C}$

$$A_i = \begin{pmatrix} \lambda_0 & 1 & & \\ & \lambda_0 & \ddots & \\ & & \ddots & 1 \\ & & & \lambda_0 \end{pmatrix}, \quad B_i = \begin{pmatrix} -1 & & & \\ & -1 & & \\ & & \ddots & \\ & & & -1 \end{pmatrix};$$

2. *Jordan block with eigenvalue* ∞

$$A_i = \begin{pmatrix} 1 & & & \\ & 1 & & \\ & & \ddots & \\ & & & 1 \end{pmatrix}, \quad B_i = \begin{pmatrix} 0 & -1 & & \\ & 0 & \ddots & \\ & & \ddots & -1 \\ & & & 0 \end{pmatrix};$$

3. *Horizontal Kronecker block*

$$A_i = \begin{pmatrix} 0 & 1 & & \\ & \ddots & \ddots & \\ & & 0 & 1 \end{pmatrix}, \quad B_i = \begin{pmatrix} -1 & 0 & & \\ & \ddots & \ddots & \\ & & -1 & 0 \end{pmatrix};$$

4. *Vertical Kronecker block*

$$A_i = \begin{pmatrix} 0 & & & \\ 1 & \ddots & & \\ & \ddots & 0 & \\ & & & 1 \end{pmatrix}, \quad B_i = \begin{pmatrix} -1 & & & \\ 0 & \ddots & & \\ & \ddots & -1 & \\ & & & 0 \end{pmatrix}.$$

The number and types of blocks in decomposition (1) are uniquely defined up to permutation.

Remark 2. The negative signs in the matrices B_i are not important and are introduced to simplify some of the formulas to follow.

Remark 3. In what follows, we interchangeably use the following two notions of a pencil: $\{A + \lambda B\}$ and $\{\alpha A + \beta B\}$. These two notions are equivalent if we allow λ to be infinite (in which case $A + \lambda B = B$) and do not distinguish between operators that are scalar multiples of each other. Under this convention, $(\beta : \alpha)$ in $\{\alpha A + \beta B\}$ are simply homogeneous coordinates of $\lambda \in \bar{\mathbb{C}}$ in $\{A + \lambda B\}$.

It is convenient to regard the zero block $0_{m,n}$ in (1) as a block-diagonal matrix that is composed of m vertical Kronecker blocks of size 1×0 and n horizontal Kronecker blocks of size 0×1 .

Definition 4. The *horizontal indices* h_1, \dots, h_p of the pencil $\mathcal{P} = \{A + \lambda B\}$ are defined to be the horizontal dimensions (widths) of horizontal Kronecker blocks (i.e., each horizontal index is the number of columns in the corresponding horizontal Kronecker block). Similarly, the *vertical indices* v_1, \dots, v_q are the vertical dimensions (heights) of vertical blocks.

In particular, in view of the above interpretation of the $0_{m,n}$ block, the first m horizontal indices and first n vertical indices are equal to 1. We will denote the total number p of horizontal indices by n_h , and the total number q of vertical indices by n_v .

Remark 5. There also exist closely related notions of *minimal row and column indices*. Namely, minimal column indices are equal to the numbers $h_i - 1$, while minimal row indices are equal to $v_i - 1$. However, we find the terms *horizontal and vertical indices* more intuitive and more suitable for our purposes.

Definition 6. The total number of columns in horizontal blocks $h_{\text{tot}} = \sum_{i=1}^{n_h} h_i$ is said to be the *total Kronecker h-index* of the pencil \mathcal{P} . Similarly, the total number of rows in the vertical Kronecker blocks $v_{\text{tot}} = \sum_{i=1}^{n_v} v_i$ is said to be the *total Kronecker v-index* of \mathcal{P} .

We now give an invariant interpretation for eigenvalues of Jordan blocks, as well as for vertical and horizontal indices. We begin with Jordan blocks.

Definition 7. The *rank of the pencil* $\mathcal{P} = \{A + \lambda B\}$ is the number $\text{rk } \mathcal{P} = \max_{\lambda \in \mathbb{C}} \text{rk}(A + \lambda B)$.

Definition 8. The *characteristic polynomial* $\chi(\alpha, \beta)$ of the pencil \mathcal{P} is defined as the greatest common divisor of all the $r \times r$ minors of the matrix $\alpha A + \beta B$, where $r = \text{rk } \mathcal{P}$.

One can show that the polynomial $\chi(\alpha, \beta)$ does not depend on the choice of bases and therefore is an invariant of the pencil. Furthermore, it is easy to see that $\chi(\alpha, \beta)$ is the product of characteristic polynomials of all the Jordan blocks. These polynomials, in turn, are called *elementary divisors* of the pencil and also admit a natural invariant interpretation, see [4] for details.

Proposition 9. *The eigenvalues of Jordan blocks can be characterized as those $\lambda \in \mathbb{C}$ for which the rank of $A + \lambda B$ drops, i.e., $\text{rk}(A + \lambda B) < r = \text{rk } \mathcal{P}$. The infinite eigenvalue appears in the case when $\text{rk } B < r$. In other words, the eigenvalues of Jordan blocks, written as $\lambda = (\beta : \alpha) \in \mathbb{C}\mathbb{P}^1$, are solutions of the characteristic equation $\chi(\alpha, \beta) = 0$. Moreover, the multiplicity of each eigenvalue coincides with the multiplicity of the corresponding root of the characteristic equation. Jordan blocks are absent if and only if the rank of all non-trivial linear combinations $\alpha A + \beta B$ is the same.*

Proof. This is easily verified when A and B are written in the Jordan–Kronecker form (1). The crucial point is that $\text{rk}(A_i + \lambda B_i) = \text{const}$ for any pair (A_i, B_i) of Kronecker blocks, so only Jordan blocks affect the dependence of $\text{rk}(A_i + \lambda B_i)$ on λ . \square

Now we discuss the meaning of vertical and horizontal indices. First of all, the total numbers of horizontal and vertical indices can be defined as the *coranks* of the pencil:

Proposition 10. *Let $\text{rk } \mathcal{P} = \max_{\lambda \in \mathbb{C}} \text{rk}(A + \lambda B)$ be the rank of the pencil $\mathcal{P} = \{A + \lambda B\}$ of operators $A, B : U \rightarrow V$. Then:*

- (1) *The number n_h of horizontal indices (or equivalently, the number of horizontal Kronecker blocks) is equal to $\dim U - \text{rk } \mathcal{P}$.*
- (2) *The number n_v of vertical indices (or equivalently, the number of vertical Kronecker blocks) is equal to $\dim V - \text{rk } \mathcal{P}$.*

In other words, $n_h = \dim \text{Ker}(A + \lambda B)$ and $n_v = \dim \text{Ker}(A + \lambda B)^$ for generic $\lambda \in \mathbb{C}$.*

Proof. Indeed, for generic λ the kernel of $A_i + \lambda B_i$ is one-dimensional if the pair of blocks (A_i, B_i) is horizontal Kronecker, and is trivial otherwise. So, $\dim \text{Ker}(A + \lambda B)$ is equal to the number of horizontal blocks. Similarly, $\dim \text{Ker}(A + \lambda B)^*$ is the number of vertical blocks. \square

We will also need the following relation between the total indices and the degree of the characteristic polynomial:

Proposition 11. *We have*

$$\mathbf{v}_{\text{tot}} + \mathbf{h}_{\text{tot}} = \dim V + \dim U - \text{rk } P - \deg \chi. \tag{2}$$

Proof. The number of columns in matrices A and B is equal to $\dim U$. On the other hand, this number can be computed as the total number of columns in Jordan blocks, which is equal to $\deg \chi$, plus the total number of columns in horizontal Kronecker blocks, which is equal to \mathbf{h}_{tot} , plus the total number of columns in vertical blocks, which is equal to $\mathbf{v}_{\text{tot}} - n_v = \mathbf{v}_{\text{tot}} - \dim V + \text{rk } P$ (see Proposition 10). The result follows. \square

Finally, we characterize the horizontal and vertical indices themselves. We begin with the following preliminary statement.

Proposition 12. *Let A be regular in a pencil $\mathcal{P} = \{A + \lambda B\}$, i.e., $\text{rk } A = \text{rk } \mathcal{P}$. Then for every $u_0 \in \text{Ker } A$ there exists a sequence of vectors $\{u_0, \dots, u_l \in U\}$ such that the expression $u(\lambda) = \sum_{j=0}^l u_j \lambda^j$ is a solution of the equation*

$$(A + \lambda B)u(\lambda) = 0. \tag{3}$$

Similarly, for any $u_0 \in \text{Ker } A^$ there exists a sequence of vectors $\{u_0, \dots, u_l \in V^*\}$ such that the expression $u(\lambda) = \sum_{j=0}^l u_j \lambda^j$ is a solution of the equation*

$$(A + \lambda B)^* u(\lambda) = 0. \tag{4}$$

Proof. We prove only the first statement, as the second one is similar. Notice that since A is regular, its kernel is generated by basis vectors in U corresponding to the leftmost columns in horizontal Kronecker blocks. So, by linearity, it suffices to consider the case when u_0 is one of those basis vectors. In this case, as u_1, u_2, \dots , one takes the remaining basis vectors in U generating the given horizontal block (which, in particular, means that the number l is equal to one of the horizontal indices minus one). \square

Proposition 13. *Let $\mathcal{P} = \{A + \lambda B\}$ and $A \in \mathcal{P}$ be regular. Let also*

$$u_i(\lambda) = \sum_{j=0}^{\deg u_i} u_{ij} \lambda^j, \quad i = 1, \dots, l,$$

where $u_{ij} \in U$, be polynomial solutions of (3) such that the vectors $u_i(0) = u_{i0}$ are linearly independent. Suppose also that $\deg u_1 \leq \dots \leq \deg u_l$. Then, for each $i = 1, \dots, l$ we have

$$\deg u_i \geq \mathbf{h}_i - 1, \tag{5}$$

where $\mathbf{h}_1 \leq \mathbf{h}_2 \leq \dots$ is the ordered sequence of horizontal indices of \mathcal{P} .

Similarly, if u_i 's are polynomial solutions of the dual problem (4), then

$$\deg u_i \geq \mathbf{v}_i - 1, \tag{6}$$

where $\mathbf{v}_1 \leq \mathbf{v}_2 \leq \dots$ are vertical indices of \mathcal{P} .

Note that the proof of Proposition 12 provides a way to construct polynomials $u_i(\lambda)$ for which inequalities (5) and (6) become equalities. This, along with Proposition 13, immediately implies the following:

Corollary 14. *Horizontal indices h_1, \dots, h_p are given by $h_i = r_i + 1$, where r_1, \dots, r_p are the minimal degrees of independent solutions of (3). Similarly, vertical indices v_1, \dots, v_q are given by $v_i = r'_i + 1$, where r'_1, \dots, r'_q are the minimal degrees of independent solutions of the dual problem (4).*

Remark 15. Recall that the numbers $h_i - 1$, $v_i - 1$ are called minimal indices (see Remark 14). Corollary 14 explains in which sense these numbers are minimal.

Proof of Proposition 13. We only consider the horizontal case, as the vertical one is analogous. Begin with the first statement. Let $\tilde{u}_i(\lambda)$, $i = 1, \dots, n_h$, be polynomial solutions of (3) constructed in the proof of Proposition 12, with $\deg \tilde{u}_i = h_i - 1$. For dimension reasons, these solutions form a basis in the kernel of $A + \lambda B$, when the latter is considered as a matrix over the field of formal Laurent series (we consider Laurent series that are finite in the negative direction). Therefore, we have

$$u_i(\lambda) = \sum_{j=1}^{n_h} f_{ij}(\lambda) \tilde{u}_j(\lambda),$$

where f_{ij} 's are formal Laurent series. Furthermore, from the linear independence of coefficients of \tilde{u}_j 's it follows that f_{ij} 's are in fact power series (otherwise the left-hand side would have terms of negative degree in λ). Now assume that (5) does not hold for some $i = k$, and k is the minimal number with this property. Then

$$\deg u_1 \leq \dots \leq \deg u_k < h_k - 1 = \deg \tilde{u}_k \leq \dots \leq \deg \tilde{u}_p,$$

which, along with independence of coefficients of \tilde{u}_j 's, implies $f_{ij} = 0$ for $i \leq k, j \geq k$. But this means that the rank of the matrix f_{ij} is not maximal, which contradicts linear independence of the vectors $u_i(0)$. So, the proposition is proved. \square

3. Jordan–Kronecker invariants of Lie algebra representations

In this section we define the main object of the present paper: Jordan–Kronecker invariants of Lie algebra representations.

Consider a finite-dimensional linear representation $\rho : \mathfrak{g} \rightarrow \mathfrak{gl}(V)$ of a finite-dimensional Lie algebra \mathfrak{g} . To each point $x \in V$, the representation ρ assigns a linear operator $R_x : \mathfrak{g} \rightarrow V$, $R_x(\xi) = \rho(\xi)x \in V$. Since the mapping $x \mapsto R_x$ is in essence equivalent to ρ , many natural algebraic objects related to ρ can be defined in terms of R_x .

Example 16. The *stabilizer* of $x \in V$ can be defined as

$$\text{St}_x = \text{Ker } R_x = \{ \xi \in \mathfrak{g} \mid R_x(\xi) = \rho(\xi)x = 0 \} \subset \mathfrak{g}.$$

Now consider the pencil of such operators generated by a pair of vectors $a, b \in V$. By the *algebraic type* of a pencil $R_a + \lambda R_b = R_{a+\lambda b}$, we will understand the following collection of discrete invariants:

- the number of distinct eigenvalues of Jordan blocks,
- the number and sizes of the Jordan blocks associated with each eigenvalue,
- horizontal and vertical indices.

Proposition 17. *The algebraic type of a pencil $R_a + \lambda R_b$ does not change under replacing a and b with any linearly independent combinations of them $a' = \alpha a + \beta b$ and $b' = \gamma a + \delta b$.*

In other words, the type characterizes two-dimensional subspaces in V or, which is the same, one-dimensional projective subspaces in the projectivization of V .

Proof. It is easy to see that replacing two operators with their independent linear combinations does not change the discrete invariants in the Jordan–Kronecker normal form (only eigenvalues of the Jordan blocks do change). \square

Since the number of different algebraic types is finite, it is easily seen that in the space $V \times V$ there exists a non-empty Zariski open subset of pairs (a, b) for which the algebraic type of the pencil $R_{a+\lambda b}$ will be one and the same (cf. [3, Prop. 1]).

Definition 18. A pair $(a, b) \in V \times V$ from this subspace and the corresponding pencil $R_{a+\lambda b}$ will be called *generic*.

Definition 19. The *Jordan–Kronecker invariant* of ρ is the algebraic type of a generic pencil $R_{a+\lambda b}$.

In particular, horizontal and vertical indices of a generic pencil will be denoted by $h_1(\rho), \dots, h_p(\rho)$ and $v_1(\rho), \dots, v_q(\rho)$ and will be called *horizontal and vertical indices* of the representation ρ .

4. Interpretation of Jordan–Kronecker invariants

In this section we give an interpretation for some of the Jordan–Kronecker invariants. All properties we discuss here are quite elementary but will be useful in the sequel. We begin with the Jordan part.

Definition 20. A point $a \in V$ is called *regular*, if

$$\dim \text{St}_a \leq \dim \text{St}_x \quad \text{for all } x \in V.$$

Those points which are not regular are called *singular*.

The set of singular points will be denoted by $\text{Sing} \subset V$. In terms of R_x we have

$$\text{Sing} = \{y \in V \mid \text{rk } R_y < r = \max_{x \in V} \text{rk } R_x\}.$$

The dimension of the stabilizer of a regular point is a natural characteristic of ρ and we will denote it by $\dim \text{St}_{\text{reg}}$. Though in our paper we never use the action of the Lie group G associated with the Lie algebra \mathfrak{g} , it will be convenient to keep in mind the action and its orbits. We will need, however, not the orbits themselves but their dimensions only. In particular, for the dimension of a regular orbit we will use the notation $\dim \mathcal{O}_{\text{reg}}$. Notice that

$$T_x \mathcal{O}_x = \text{Im } R_x \quad \text{and} \quad \dim \mathcal{O}_x = \text{rk } R_x.$$

Proposition 21.

- (1) *The eigenvalues of Jordan blocks of a pencil $R_{a+\lambda b}$ are those values of $\lambda \in \mathbb{C}$ for which the line $a + \lambda b$ intersects the singular set Sing . In particular, $R_{a+\lambda b}$ has a Jordan block with eigenvalue ∞ if and only if b is singular.*
- (2) *A generic pencil $R_a + \lambda R_b$ has no Jordan blocks if and only if the codimension of the singular set Sing is greater or equal than 2.*

Proof. The first statement is an immediate corollary of Proposition 9. Furthermore, it follows from Proposition 9 that a generic pencil $R_a + \lambda R_b$ has no Jordan blocks if and only if all these operators are of the same rank, i.e., a generic line $a + \lambda b$ does not intersect the singular set Sing . Clearly, the latter condition is fulfilled if and only if $\text{codim } \text{Sing} \geq 2$. \square

Let us discuss the case $\text{codim Sing} = 1$ in more detail. Consider the matrix of the operator R_x and take all of its minors of size $r \times r$, where $r = \dim \mathcal{O}_{\text{reg}}$, that do not vanish identically (such minors certainly exist). We consider them as polynomials $p_1(x), \dots, p_N(x)$ on V . The singular set $\text{Sing} \subset V$ is then given by the system of polynomial equations

$$p_i(x) = 0, \quad i = 1, \dots, N.$$

This set is of codimension one if and only if these polynomials possess a non-trivial greatest common divisor which we denote by \mathfrak{p}_ρ .

Thus, we have $p_i(x) = \mathfrak{p}_\rho(x)h_i(x)$, which implies that the singular set Sing can be represented as the union of two subsets:

$$\text{Sing}_0 = \{x \in V \mid \mathfrak{p}_\rho(x) = 0\} \quad \text{and} \quad \text{Sing}_1 = \{x \in V \mid h_i(x) = 0, \quad i = 1, \dots, N\}.$$

It is easy to see that $\mathfrak{p}_\rho(x)$ is a semi-invariant of the representation ρ . This follows from the fact that the action of G leaves the singular set Sing_0 invariant and therefore may only multiply \mathfrak{p}_ρ by a character of G . We will refer to this polynomial \mathfrak{p}_ρ as the *fundamental semi-invariant* of ρ . The fundamental semi-invariant is closely related to the characteristic polynomial $\chi_{a,b}$ of the pencil $R_{a+\lambda b}$:

Proposition 22. *Let $a, b \in V$ be such that the set $\{\alpha a + \beta b \mid (\alpha, \beta) \neq (0, 0)\}$ of their non-trivial linear combinations does not intersect Sing_1 and is not completely contained in Sing (i.e., contains at least one regular element). Then*

$$\chi_{a,b}(\alpha, \beta) = \mathfrak{p}_\rho(\alpha a + \beta b).$$

Proof. As above, let $p_1(x), \dots, p_N(x)$ be the $r \times r$ minors of the matrix R_x . Then $p_i(x) = \mathfrak{p}_\rho(x)h_i(x)$ for certain polynomials $h_i(x)$. Substituting $x = \alpha a + \beta b$, we get

$$p_i(\alpha a + \beta b) = \mathfrak{p}_\rho(\alpha a + \beta b)h_i(\alpha a + \beta b).$$

Further, notice that since the set $\{\alpha a + \beta b \mid (\alpha, \beta) \neq (0, 0)\}$ contains a regular element, it follows that the rank of the pencil $R_{a+\lambda b}$ is equal to $\dim \mathcal{O}_{\text{reg}}$. Therefore, $\chi_{a,b}$ is, by definition, the greatest common divisor of the expressions $p_i(\alpha a + \beta b)$, where the latter are viewed as polynomials in α and β . So,

$$\chi_{a,b} = \gcd\{\mathfrak{p}_\rho(\alpha a + \beta b)h_i(\alpha a + \beta b)\} = \mathfrak{p}_\rho(\alpha a + \beta b) \cdot \gcd\{h_i(\alpha a + \beta b)\}.$$

For the sake of contradiction, assume the latter factor is non-constant. Then there exist α and β , not simultaneously equal to zero, such that $h_i(\alpha a + \beta b) = 0$ for every i (here we use that the polynomials $h_i(\alpha a + \beta b)$ are homogeneous). But this means that the set $\{\alpha a + \beta b \mid (\alpha, \beta) \neq (0, 0)\}$ intersects Sing_1 , which is not the case. So, $\chi_{a,b} = \mathfrak{p}_\rho(\alpha a + \beta b)$, up to a constant factor. \square

Corollary 23. *The degree of the fundamental semi-invariant \mathfrak{p}_ρ is equal to the sum of the sizes of all Jordan blocks for a generic pencil $R_{a+\lambda b}$.*

Proof. Indeed, by Proposition 22 for generic a, b we get $\deg \mathfrak{p}_\rho = \deg \chi_{a,b}$. But the degree of $\chi_{a,b}$ is exactly the sum of the sizes of Jordan blocks. \square

We now turn to the Kronecker part. Following Section 2, for an arbitrary pencil $R_{a+\lambda b}$ we define the numbers $\mathfrak{v}_{\text{tot}}(a, b)$ and $\mathfrak{h}_{\text{tot}}(a, b)$. These numbers computed for a generic pair (a, b) are invariants of the representation ρ . We denote them $\mathfrak{v}_{\text{tot}}(\rho)$, $\mathfrak{h}_{\text{tot}}(\rho)$ and call the *total Kronecker v-index and h-index* of ρ . Similarly, let $n_h(\rho), n_v(\rho)$ be the numbers of horizontal and vertical indices for a generic pencil $R_{a+\lambda b}$.

Proposition 24.

- (1) The number $n_h(\rho)$ of horizontal indices of ρ is equal to $\dim \text{St}_{\text{reg}}$.
- (2) The number $n_v(\rho)$ of vertical indices of ρ is equal to $\text{codim } \mathcal{O}_{\text{reg}}$.

Proof. This follows from $\text{rk } R_{a+\lambda b} = \dim \mathcal{O}_{a+\lambda b}$ and Proposition 10. \square

Remark 25. Notice that (cf. (2))

$$v_{\text{tot}}(\rho) + h_{\text{tot}}(\rho) = \dim V + \dim \text{St}_{\text{reg}} - \deg \mathfrak{p}_\rho = \dim \mathfrak{g} + \text{codim } \mathcal{O}_{\text{reg}} - \deg \mathfrak{p}_\rho. \quad (7)$$

Similarly, if ρ has an open orbit, i.e., $\text{codim } \mathcal{O}_{\text{reg}} = 0$, then vertical Kronecker blocks are absent and we have $h_{\text{tot}}(\rho) = \dim \mathfrak{g} - \deg \mathfrak{p}_\rho$. If in addition $\text{codim } \text{Sing} \geq 2$, then $h_{\text{tot}}(\rho) = \dim \mathfrak{g}$.

5. Degrees of invariant polynomials and vertical indices

This section contains our main results. The first result gives a bound for degrees of invariant polynomials in terms of vertical indices. In the case of the coadjoint representation it was obtained by A. Vorontsov [17].

Theorem 26 (Lower bounds for degrees of polynomial invariants). *Let $\rho: \mathfrak{g} \rightarrow \mathfrak{gl}(V)$ be a representation of a finite-dimensional Lie algebra \mathfrak{g} on a finite-dimensional vector space V . Assume that f_1, \dots, f_m are homogeneous algebraically independent invariant polynomials of ρ , and $\deg f_1 \leq \dots \leq \deg f_m$. Let also $v_1(\rho) \leq \dots \leq v_q(\rho)$ be the vertical indices of ρ . Then*

$$\deg f_i \geq v_i(\rho) \quad (8)$$

for $i = 1, \dots, m$.

Proof. Let $(a, b) \in V \times V$ be generic. Expanding $f_i(a + \lambda x)$ in powers of λ , we get

$$f_i(a + \lambda x) = f_{i0}(x) + \lambda f_{i1}(x) + \dots + \lambda^{m_i} f_{im_i}(x), \quad m_i = \deg f_i.$$

Furthermore, since f_i is an invariant, we have

$$(R_a + \lambda R_b)^* \sum_{j=0}^{m_i} \lambda^j df_{ij}(b) = 0,$$

and since $f_{i0}(x) = f_i(a)$, the first term in the latter sum is zero, so we can divide the latter equation by λ :

$$(R_a + \lambda R_b)^* \sum_{j=0}^{m_i-1} \lambda^j df_{i,j+1}(b) = 0.$$

Also notice that $df_{i,1}(x) = df_i(a)$ for any $x \in V$, so if we take a such that the differentials of f_i 's at a are independent, then Proposition 13 gives exactly the desired estimate (8). \square

Corollary 27. *Let f_1, f_2, \dots, f_q , $q = \text{codim } \mathcal{O}_{\text{reg}}$, be algebraically independent invariant polynomials of ρ . Then*

$$\sum_{i=1}^q \deg f_i \geq v_{\text{tot}}(\rho).$$

Proof. This is obtained by adding up inequalities (8). \square

Taking into account Remark 25, we also get

Corollary 28. *Let f_1, f_2, \dots, f_q , $q = \text{codim } \mathcal{O}_{\text{reg}}$, be algebraically independent invariant polynomials of ρ . Suppose that the stabilizer of a regular point is trivial, i.e., $\text{St}_{\text{reg}} = \{0\}$. Then*

$$\sum_{i=1}^q \deg f_i \geq \dim V - \deg \rho.$$

Moreover, if in addition $\text{codim Sing} \geq 2$, then

$$\sum_{i=1}^q \deg f_i \geq \dim V. \tag{9}$$

Remark 29. An estimate similar to (9) was obtained by F. Knop and P. Littelmann [7].

Another immediate corollary of Theorem 26 is the following:

Corollary 30. *Suppose that there exist algebraically independent invariant polynomials f_1, f_2, \dots, f_q , $q = \text{codim } \mathcal{O}_{\text{reg}}$, of a representation ρ satisfying the condition*

$$\sum_{i=1}^q \deg f_i = \mathbf{v}_{\text{tot}}(\rho). \tag{10}$$

Then

$$\mathbf{v}_i(\rho) = \deg f_i. \tag{11}$$

Proof. Indeed, (10) is equivalent to

$$\sum_{i=1}^q \deg f_i = \sum_{i=1}^q \mathbf{v}_i(\rho),$$

which, in view of (8), is equivalent to (11). \square

We now investigate in more detail the case when one of the equivalent conditions (10), (11) hold.

Theorem 31 (On the set where the invariants become dependent). *Let $\rho: \mathfrak{g} \rightarrow \mathfrak{gl}(V)$ be a representation of a finite-dimensional Lie algebra \mathfrak{g} on a finite-dimensional vector space V . Assume that f_1, f_2, \dots, f_q are algebraically independent invariant polynomials of ρ and $q = \text{codim } \mathcal{O}_{\text{reg}}$. Let also $\mathbf{v}_1(\rho), \dots, \mathbf{v}_q(\rho)$ be the vertical indices of ρ . Then the following conditions are equivalent:*

- (1) *The degrees of f_i 's are equal to the vertical indices of ρ : $\deg f_i = \mathbf{v}_i(\rho)$.*
- (2) *The sum of the degrees of f_i 's is equal to the total vertical index of ρ : $\sum \deg f_i = \mathbf{v}_{\text{tot}}(\rho)$.*
- (3) *The set where the differentials df_1, \dots, df_q are linearly dependent has codimension ≥ 2 in V .*
- (4) *The set where the differentials df_1, \dots, df_q are linearly dependent is contained in the set Sing_1 , i.e., in the codimension ≥ 2 stratum of the set of singular points of ρ in V .*

The proof is based on the following lemma, which, in particular, gives an interpretation of the total Kronecker indices:

Lemma 32. *Let $r = \dim \mathcal{O}_{\text{reg}}$. For $x \in V$, consider the operator $\Lambda^r R_x: \Lambda^r \mathfrak{g} \rightarrow \Lambda^r V$, where $\Lambda^r R_x(\xi_1 \wedge \cdots \wedge \xi_r) = R_x(\xi_1) \wedge \cdots \wedge R_x(\xi_r)$. Then*

$$\Lambda^r R_x = \mathfrak{p}_\rho(x) \cdot \omega_h(x) \otimes \omega_v(x), \tag{12}$$

where ω_h, ω_v are homogeneous polynomials in x with values in $\Lambda^r \mathfrak{g}^*$, $\Lambda^r V$ respectively. The polynomials ω_h, ω_v are defined uniquely up to constant factors, not divisible by any non-trivial scalar polynomial of x , and are of degrees

$$\begin{aligned} \deg \omega_h &= \mathfrak{h}_{\text{tot}}(\rho) - n_h(\rho) = \mathfrak{h}_{\text{tot}}(\rho) - \dim \text{St}_{\text{reg}}, \\ \deg \omega_v &= \mathfrak{v}_{\text{tot}}(\rho) - n_v(\rho) = \mathfrak{v}_{\text{tot}}(\rho) - \text{codim } \mathcal{O}_{\text{reg}}. \end{aligned} \tag{13}$$

Remark 33. Formulas (13) can be also rewritten as

$$\deg \omega_h = \sum_{i=1}^{n_h(\rho)} (\mathfrak{h}_i(\rho) - 1), \quad \deg \omega_v = \sum_{i=1}^{n_v(\rho)} (\mathfrak{v}_i(\rho) - 1),$$

i.e., the degrees of ω_h, ω_v are given by sums of minimal indices. Also note that two different formulas for each of the degrees in (13) are equivalent due to Proposition 24.

Proof of Lemma 32. The matrix entries of the operator $\Lambda^r R_x$ are $r \times r$ minors of the matrix R_x . The greatest common divisor of those minors is, by definition, the fundamental semi-invariant $\mathfrak{p}_\rho(x)$. So, $\Lambda^r R_x = \mathfrak{p}_\rho(x)S(x)$, where $S(x)$ is a polynomial in x with values in $\text{Hom}(\Lambda^r \mathfrak{g}, \Lambda^r V)$. Further, note that the rank of R_x for regular x is $r = \dim \mathcal{O}_{\text{reg}}$, so $\dim \text{Im } R_x = r$, and hence the space $\text{Im } \Lambda^r R_x = \Lambda^r \text{Im } R_x$ is generically one-dimensional. Therefore, the image of $S(x)$ is generically one-dimensional too. So, if we regard $S(x)$ as a matrix over the field of rational functions in x , its image has dimension 1, which means that it is decomposable:

$$S(x) = \omega_h(x) \otimes \omega_v(x),$$

where ω_h, ω_v are rational functions in x with values in $\Lambda^r \mathfrak{g}^*$ and $\Lambda^r V$ respectively. Furthermore, multiplying, if necessary, ω_h by a scalar rational function of x and dividing ω_v by the same function, we can arrange that ω_h is polynomial, and that the greatest common divisor of its coefficients is equal to 1. But then ω_v must be polynomial too, because the product $\omega_h \otimes \omega_v = S$ is polynomial. Hence, existence of factorization (12) is proved.

To prove that ω_h, ω_v are not divisible by any non-trivial scalar polynomial, recall that \mathfrak{p}_ρ is the greatest common divisor of the matrix entries of $\Lambda^r R_x$. Therefore, the greatest common divisor of the matrix entries of $S = \frac{1}{\mathfrak{p}_\rho} \Lambda^r R_x$ is 1, and hence the same is true for its factors ω_h, ω_v , as desired.

To prove uniqueness, we use that the representation of a rank 1 operator as a tensor product is unique up to multiplying the first factor by an element of the base field, and dividing the second factor by the same element. Therefore, if we have another representation

$$S(x) = \omega'_h(x) \otimes \omega'_v(x),$$

then there exists a scalar rational function $\mu(x)$, such that

$$\omega'_h(x) = \mu(x)\omega_h(x), \quad \omega'_v(x) = \frac{1}{\mu(x)}\omega_v(x).$$

If, moreover, ω'_h, ω'_v are polynomials, then it follows that ω_h is divisible by the denominator of μ , while ω_v is divisible by the numerator of μ (we assume that μ is written

in the reduced form). But we know that ω_h, ω_v do not have non-trivial polynomial factors. Therefore, $\mu(x)$ must be constant, which proves that factorization (12) is unique. This also shows that the polynomials ω_h, ω_v are homogeneous, because if they were not, then the expression $\omega_h(\lambda x) \otimes \omega_v(\lambda x)$, divided by a suitable constant, would provide another factorization of $S(x)$.

Now it remains to compute the degrees of ω_h, ω_v . To that end, we restrict (12) to a 2-plane $x = \alpha a + \beta b$, where $a, b \in V \times V$ are generic. This gives

$$\Lambda^r R_{\alpha a + \beta b} = \chi_{a,b}(\alpha, \beta) \cdot \omega_h(\alpha a + \beta b) \otimes \omega_v(\alpha a + \beta b), \tag{14}$$

where we used Proposition 22 to rewrite the first factor in the right-hand side. Now, consider the Jordan–Kronecker normal form of the pencil $R_{\alpha a + \beta b}$. Take all columns of all vertical Kronecker blocks of non-zero width (these can be naturally viewed as vectors in V). In addition to that, take basis vectors in V corresponding to rows of all Jordan blocks and all horizontal Kronecker blocks of non-zero height. Taken together, all these vectors form a basis in $\text{Im } R_{\alpha a + \beta b}$ for generic α, β . Thus, their wedge product, which we denote by $f(\alpha, \beta)$, is a non-trivial polynomial of α, β valued in $\text{Im } \Lambda^r R_{\alpha a + \beta b}$. At the same time, by (14) the latter space is generated by $\omega_v(\alpha a + \beta b)$, so we must have

$$f(\alpha, \beta) = \mu(\alpha, \beta) \omega_v(\alpha a + \beta b),$$

where μ is a rational function. Note also that since $\omega_v(x)$ is not divisible by a non-trivial scalar polynomial, it follows that its vanishing set in $\mathbb{P}V$ has codimension at least 2. A generic projective line $\alpha a + \beta b$ does not intersect this set, which means that $\omega_v(\alpha a + \beta b)$ does not vanish at all (unless $\alpha = \beta = 0$), and hence $\mu(\alpha, \beta)$ is actually a polynomial function. This gives

$$\deg \omega_v(\alpha a + \beta b) \leq \deg f(\alpha, \beta) = \mathbf{v}_{\text{tot}}(\rho) - n_v(\rho),$$

and thus

$$\deg \omega_v(x) \leq \mathbf{v}_{\text{tot}}(\rho) - n_v(\rho), \tag{15}$$

where we used that $\deg f = \mathbf{v}_{\text{tot}}(\rho) - n_v(\rho)$ by construction. Furthermore, an analogous argument for the dual pencil gives

$$\deg \omega_h(x) \leq \mathbf{h}_{\text{tot}}(\rho) - n_h(\rho). \tag{16}$$

At the same time

$$\deg \omega_v + \deg \omega_h = \deg \Lambda^r R_x - \deg \mathbf{p}_\rho = \dim \mathcal{O}_{\text{reg}} - \deg \mathbf{p}_\rho,$$

while the sum of right-hand sides in (15) and (16) is

$$\begin{aligned} \mathbf{v}_{\text{tot}}(\rho) + \mathbf{h}_{\text{tot}}(\rho) - n_v(\rho) - n_h(\rho) &= \mathbf{v}_{\text{tot}}(\rho) + \mathbf{h}_{\text{tot}}(\rho) - \dim \text{St}_{\text{reg}} - \text{codim } \mathcal{O}_{\text{reg}} \\ &= \mathbf{v}_{\text{tot}}(\rho) + \mathbf{h}_{\text{tot}}(\rho) - \dim \text{St}_{\text{reg}} + \dim \mathcal{O}_{\text{reg}} - \dim V \\ &= \dim \mathcal{O}_{\text{reg}} - \deg \mathbf{p}_\rho, \end{aligned}$$

where the first equality uses Proposition 24, while the last one uses (7). So, the sum of left-hand sides in (15) and (16) is equal to the sum of right-hand sides, and thus both inequalities must be equalities, providing the desired formulas for $\deg \omega_v, \deg \omega_h$. \square

Proof of Theorem 31. Equivalence of Conditions 1 and 2 follows from Corollary 30. Also, implication $4 \Rightarrow 3$ is obvious. So, it suffices to prove implications $2 \Rightarrow 4$ and $3 \Rightarrow 2$. We begin with $2 \Rightarrow 4$. Let $\Omega \in \Lambda^{\dim V} V$ be a constant non-zero top-degree multi-vector on V (i.e., a top-degree form on V^*). Let also $r = \dim \mathcal{O}_{\text{reg}}$. Define an r -vector ω on V by

$$\omega(x)(\xi_1, \dots, \xi_r) = \Omega(df_1(x), \dots, df_q(x), \xi_1, \dots, \xi_r).$$

In other words, ω is a multi-vector dual to the form $df_1 \wedge \dots \wedge df_q$. Note that for generic x the differentials $df_1(x), \dots, df_q(x)$ span the annihilator of the tangent space $T_x \mathcal{O}_x$ to the orbit of ρ . Therefore, $\omega \in \Lambda^r T_x \mathcal{O}_x$. But the latter space is generically one-dimensional and generated by the form $\omega_v(x)$ given by (12). Therefore, we must have

$$\omega(x) = \mu(x)\omega_v(x), \tag{17}$$

where $\mu(x)$ is a rational function. But the form ω_v is not divisible by any scalar polynomial (see Lemma 32), so $\mu(x)$ is in fact a polynomial. Furthermore, we have

$$\deg \omega = \sum \deg df_i = \sum (\deg f_i - 1) = \left(\sum \deg f_i \right) - \text{codim } \mathcal{O}_{\text{reg}}. \tag{18}$$

So, assuming Condition 2 of the theorem, i.e., $\sum \deg f_i = \mathbf{v}_{\text{tot}}(\rho)$, we get

$$\deg \omega = \mathbf{v}_{\text{tot}}(\rho) - \text{codim } \mathcal{O}_{\text{reg}}.$$

But the latter number is equal to $\deg \omega_v$ by Lemma 32, so we must conclude that $\mu(x)$ in (17) is a constant function, and zeros of $\omega(x)$ are the same as zeros of $\omega_v(x)$. Furthermore, zeros of ω are exactly those points where df_1, \dots, df_q become linearly dependent, while zeros of ω_v are contained in the set of zeros of $\omega_h(x) \otimes \omega_v(x) = \frac{1}{p_\rho} \Lambda^r R_x$. But the latter set is exactly Sing_1 , which proves the implication $2 \Rightarrow 4$.

We now prove $3 \Rightarrow 2$. Condition 3 says that df_1, \dots, df_q are linearly dependent on a set of codimension ≥ 2 , which is equivalent to saying that the zero set of $\omega(x)$ has codimension ≥ 2 . But this is only possible if $\mu(x)$ in (17) is a constant function, in which case we have $\deg \omega = \deg \omega_v$. In view of (18), this gives

$$\sum \deg f_i = \deg \omega_v + \text{codim } \mathcal{O}_{\text{reg}} = \mathbf{v}_{\text{tot}}(\rho),$$

as desired. \square

As can be seen from the proof, in general the set Sing_1 consists of two components: the zeros of ω_v (which, under the conditions of Theorem 31, coincide with the set where the invariants become dependent), and the zeros of ω_h . This immediately gives the following:

Corollary 34. *Let \mathfrak{g} be a finite-dimensional Lie algebra, and let $\rho: \mathfrak{g} \rightarrow \mathfrak{gl}(V)$ be either its coadjoint representation, or any finite-dimensional representation such that the stabilizer of a generic element in V is trivial. Assume that $q = \text{codim } \mathcal{O}_{\text{reg}}$ and f_1, f_2, \dots, f_q are algebraically independent invariant polynomials of ρ whose degrees are equal to the vertical indices: $\deg f_i = \mathbf{v}_i(\rho)$ (equivalently, $\sum \deg f_i = \mathbf{v}_{\text{tot}}(\rho)$). Then their differentials df_1, df_2, \dots, df_q are linearly dependent exactly on the set Sing_1 .*

Remark 35. In the case of the coadjoint representation, this is equivalent to a result of D.Panyushev (Theorem 1.2. in [11]), proved in the case $\text{codim } \text{Sing} \geq 2$.

Proof of Corollary 34. Indeed, for the coadjoint representation due to skew-symmetry of R_x we have $\omega_h = \omega_v$, while in the trivial stabilizer case we have that ω_h is of degree 0 and hence constant. In both cases, the zero set of $\omega_h \otimes \omega_v$ is the same as the zero set of ω_v , hence the result. \square

Theorem 36 (On polynomiality of the algebra of invariants). *Let $\rho: \mathfrak{g} \rightarrow \mathfrak{gl}(V)$ be a representation of a finite-dimensional Lie algebra \mathfrak{g} on a finite-dimensional vector space V . Assume that f_1, f_2, \dots, f_q are algebraically independent invariant polynomials of ρ and $q = \text{codim } \mathcal{O}_{\text{reg}}$. Let also $v_1(\rho), \dots, v_q(\rho)$ be the vertical indices of ρ . Then we have the following:*

- (1) *If the degrees of f_i 's are equal to the vertical indices: $\deg f_i = v_i(\rho)$ (equivalently, $\sum \deg f_i = v_{\text{tot}}(\rho)$), then the algebra $\mathbb{C}[V]^{\mathfrak{g}}$ of polynomial invariants of ρ is freely generated by f_1, f_2, \dots, f_q (i.e., it is a polynomial algebra).*
- (2) *Conversely, if the algebra $\mathbb{C}[V]^{\mathfrak{g}}$ of polynomial invariants of ρ is freely generated by f_1, f_2, \dots, f_q , and, in addition, ρ has no proper semi-invariants (i.e., any semi-invariant is an invariant), then the degrees of f_i 's are equal to the vertical indices.*

Remark 37. The condition on semi-invariants holds, for example, when \mathfrak{g} is perfect, i.e., coincides with its derived subalgebra. This condition cannot be omitted, as shown by the following example. Consider a 3-dimensional Lie algebra with relations $[z, x] = x, [z, y] = -2y$. Then the algebra of invariants of its coadjoint representation is freely generated by a degree 3 function x^2y , while the only vertical index is equal to 2 (and, as predicted by Theorem 31, the gradient of the invariant vanishes on a hypersurface). Theorem 36 does not apply in this case because x and y are proper semi-invariants.

Remark 38. A similar result is obtained in [5] (see Corollary 5.5): if the algebra of polynomial invariants is freely generated by f_1, f_2, \dots, f_q , and, in addition, a certain semi-invariant is an invariant, then $\sum \deg f_i$ is equal to the degree of a certain form. A novel feature of our work is that we, first, compute the degree of the latter form (and hence show that the number $\sum \deg f_i$ is equal to the total vertical index), and second, provide formulas for the degrees $\deg f_i$ themselves, and not just for their sum.

Proof of Theorem 36. Assume that the degrees of f_i 's are equal to the vertical indices. Take any polynomial invariant $f \in \mathbb{C}[V]^{\mathfrak{g}}$ of ρ . Then, since $\text{tr.deg. } \mathbb{C}[V]^{\mathfrak{g}} = q$, it follows that f is algebraically dependent with f_1, f_2, \dots, f_q . Given also that f_1, f_2, \dots, f_q are independent outside of a subset of codimension ≥ 2 (Theorem 31), it follows from Theorem 1.1 of [12] that f is a polynomial function of f_1, f_2, \dots, f_q . So, the polynomials f_1, f_2, \dots, f_q generate $\mathbb{C}[V]^{\mathfrak{g}}$, and since those polynomials are independent, the algebra is freely generated, as desired.

Conversely, assume that $\mathbb{C}[V]^{\mathfrak{g}}$ is freely generated by f_1, f_2, \dots, f_q . Assume, for the sake of contradiction, that df_1, df_2, \dots, df_q are linearly dependent on a subset $S \subset V$ of codimension 1. Then the union of codimension 1 irreducible components of S is given by a single polynomial equation $h_f(x) = 0$. Since f_1, f_2, \dots, f_q are invariants, it follows that h_f is a semi-invariant. So, under the assumptions of the theorem, h_f must be an invariant, and hence a scalar by Proposition 5.2 of [5]. Therefore, f_1, f_2, \dots, f_q are actually dependent on a subset of codimension ≥ 2 , and the result follows by Theorem 31. \square

Corollary 39. *Let $\rho: \mathfrak{g} \rightarrow \mathfrak{gl}(V)$ be a representation of a finite-dimensional Lie algebra \mathfrak{g} on a finite-dimensional vector space V . Assume that ρ admits no proper semi-invariants, and that the algebra $\mathbb{C}[V]^{\mathfrak{g}}$ of polynomial invariants of ρ is freely generated by f_1, f_2, \dots, f_q . Then*

$$\sum_{i=1}^q \deg f_i \leq \dim V.$$

Proof. Indeed, in these assumptions we have

$$\sum_{i=1}^q \deg f_i = v_{\text{tot}}(\rho) \leq \dim V. \quad \square$$

Remark 40. When \mathfrak{g} is simple, this corollary is a particular case of Popov’s conjecture [14, p. 350] proved in [6].

6. JK invariants for standard representations of simple Lie algebras

In this section we consider classical simple Lie algebras, namely $\mathfrak{sl}(n)$, $\mathfrak{so}(n)$, and $\mathfrak{sp}(n)$. For each of them we calculate the JK invariants for the sums of their standard representations. In each case we give the answer first and then show how it correlates with the general results from Sections 4 and 5, most importantly, Theorems 26, 31 and 36 about vertical indices and the algebra of invariants.

In all cases we consider a matrix subalgebra $\mathfrak{g} \subset \mathfrak{gl}(n)$ and the sum of its m standard representations

$$\rho^{\oplus m} : \mathfrak{g} \rightarrow \mathfrak{gl}(V^{\oplus m}).$$

Fix a basis of V , so that we could identify all the linear maps with matrices. Then an element of $V^{\oplus m}$ is given by a matrix $X \in \text{Mat}_{n \times m}$ and it defines a linear mapping

$$\begin{aligned} R_X : \mathfrak{g} &\rightarrow \text{Mat}_{n \times m}, \\ Y &\rightarrow YX. \end{aligned}$$

Lemma 41. *Consider the sum of m standard representations for any of the Lie algebras $\mathfrak{sl}(n)$, $\mathfrak{so}(n)$, or $\mathfrak{sp}(n)$. Define $n \times m$ matrices X and A as follows:*

- If $m < n$, then

$$X = \begin{pmatrix} 0 \\ I_m \end{pmatrix}, \quad A = \begin{pmatrix} -I_m \\ 0 \end{pmatrix}.$$

- If $n = m$, then

$$X = \begin{pmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{pmatrix}, \quad A = \begin{pmatrix} -1 & & \\ & \ddots & \\ & & -1 \end{pmatrix},$$

where λ_i ’s are distinct numbers.

- If $m > n$, then

$$X = (0 \quad I_n), \quad A = (-I_n \quad 0).$$

Then the pencil $R_{X+\lambda A}$ is generic.

Proof. We need to show that there exists an open dense subset $U \subset \text{Mat}_{n \times m} \times \text{Mat}_{n \times m}$ such that for any pair $(\tilde{X}, \tilde{A}) \in U$, the pencil $R_{\tilde{X}+\lambda\tilde{A}}$ has the same algebraic type as the pencil $R_{X+\lambda A}$. What is well-known is that there exists an open dense subset $U \subset \text{Mat}_{n \times m} \times \text{Mat}_{n \times m}$ such that, for any pair $(\tilde{X}, \tilde{A}) \in U$, its orbit under the natural left-right action of $\text{GL}(n) \times \text{GL}(m)$ on $\text{Mat}_{n \times m}$ contains a pair (X, A) of the above form, see, e.g., [13, p. 119]. In other words, any generic pair of operators $\mathbb{R}^m \rightarrow \mathbb{R}^n$ can be written in the above form (X, A) in a suitable basis. So, to show that the pencil $R_{X+\lambda A}$ is generic, it suffices to prove that for any $C \in \text{GL}(n)$, $D \in \text{GL}(m)$ the JK decompositions of the pencils $R_X + \lambda R_A$ and $R_{CXD} + \lambda R_{CAD}$ coincide. The latter is equivalent to the existence of invertible linear mappings $\varphi : \mathfrak{g} \rightarrow \mathfrak{g}$ and $\psi : \text{Mat}_{n \times m} \rightarrow \text{Mat}_{n \times m}$ (where φ does not have to be a Lie algebra automorphism) such that the following diagram commutes

$$\begin{array}{ccc}
 \mathfrak{g} & \xrightarrow{R_{CXD} + \lambda R_{CAD}} & \text{Mat}_{n \times m} \\
 \varphi \downarrow & & \uparrow \psi \\
 \mathfrak{g} & \xrightarrow{R_X + \lambda R_A} & \text{Mat}_{n \times m}
 \end{array}$$

For $\mathfrak{sl}(n)$, a suitable choice of φ, ψ is

$$\varphi(Y) = C^{-1}YC, \quad \psi(Z) = CZD.$$

Similarly for $\mathfrak{so}(n)$ and $\mathfrak{sp}(n)$ we take

$$\varphi(Y) = C^*YC, \quad \psi(Z) = (C^*)^{-1}ZD,$$

where $C^* = C^\top$ for $\mathfrak{so}(n)$ and $C^* = \Omega^{-1}C^\top\Omega$ for $\mathfrak{sp}(n)$, with Ω being the matrix of the symplectic form given by formula (19) below. In all cases we see that the automorphisms φ, ψ intertwine the pencils $R_X + \lambda R_A$ and $R_{CXD} + \lambda R_{CAD}$, so those pencils have the same JK type, as desired. \square

6.1. Standard representations of $\mathfrak{sl}(n)$

Proposition 42. *Let ρ be the sum of m standard representations of $\mathfrak{sl}(n)$.*

- (1) *Assume that $m < n$. Let q be the integral part of the quotient $m / (n - m)$, and r be the remainder. Then the JK invariants of ρ consist of $n(n - m) - 1$ horizontal indices*

$$\underbrace{q + 1, \dots, q + 1}_{n(n-m-r)-(q+1)}, \quad \underbrace{q + 2, \dots, q + 2}_{nr+q}.$$

- (2) *Assume that $m = n$. Then the JK invariants of ρ consist of one vertical index $v_1 = n$ and n distinct eigenvalues with $n - 1$ Jordan 1×1 blocks corresponding to each eigenvalue.*
- (3) *Assume that $m > n$. Let q be the integral part of the quotient $n / (m - n)$, and r be the remainder. Then the JK invariants of ρ consist of the following $n(m - n) + 1$ vertical indices:*

- *If $r \neq 0$, then the vertical indices are*

$$\underbrace{q + 1, \dots, q + 1}_{n(m-n-r)+(q+2)}, \quad \underbrace{q + 2, \dots, q + 2}_{nr-(q+1)}.$$

- *If $r = 0$, then the vertical indices are*

$$\underbrace{q, \dots, q}_{q+1}, \quad \underbrace{q + 1, \dots, q + 1}_{n(m-n)-q}.$$

Proof. We need to compute the Jordan–Kronecker normal form for explicitly given operators R_X, R_A , where X, A are defined in Lemma 41. This is a tedious, yet direct, calculation. \square

Now, let us demonstrate how Theorem 36 works in this case.

Proposition 43. *Let $\rho: \mathfrak{sl}(n) \rightarrow \mathfrak{gl}(\text{Mat}_{n \times m})$ be the sum of m standard representations of $\mathfrak{sl}(n)$, and let $X \in \text{Mat}_{n \times m}$. Then:*

- (1) *If $m < n$, the algebra of invariants of ρ is trivial.*
- (2) *If $m = n$, the algebra of invariants is freely generated by the polynomial $\det X$.*
- (3) *If $m = n + 1$, the algebra of invariants is freely generated by $n \times n$ minors of X .*
- (4) *If $m > n + 1$, the algebra of invariants is not freely generated.*

Remark 44. Of course, these results are well known. For $m > n$, the algebra of invariants of ρ can be identified with the homogeneous coordinate ring of the Grassmannian $\text{Gr}(n, m)$. If $m = n + 1$, then the Grassmannian $\text{Gr}(n, m)$ is the projective space \mathbb{P}^n , whose homogeneous coordinate ring is the ring of polynomials in $n + 1$ variables, hence freely generated. For $m > n + 1$, the homogeneous coordinate ring of the Grassmannian $\text{Gr}(n, m)$ is generated by Plücker coordinates, subject to Plücker relations.

Proof of Proposition 43. There are clearly no invariants in the $m < n$ case, because in this case the action of the group $\text{SL}(n)$ on m -tuples of vectors in \mathbb{R}^n has an open orbit. (One could also say that in this case there are no vertical indices and hence no invariants by Theorem 26.) As for the case $m = n$, in this case we have one vertical index equal to n , and a polynomial invariant $\det X$ of degree n . So, by Theorem 36, the algebra of invariants is freely generated by $\det X$ (which is obvious in this case, because the transcendence degree of the algebra of invariants is equal to 1, so it must be freely generated).

Further, when $m = n + 1$, the vertical indices are n, \dots, n ($n + 1$ times), which again coincides with the degrees of the $n \times n$ minors of X . So, these minors generate the algebra of invariants by Theorem 36 (they are clearly independent, as one can find a matrix X with any prescribed values of these minors).

Finally, consider the case $m > n + 1$. Observe that in this case there may also be no invariants of degree $k < n$. Indeed, assume that f is such an invariant. Then there exist k indices $1 \leq i_1 < \dots < i_k \leq n$ such that f has non-trivial restriction to the subspace of matrices whose all columns, except for the ones with indices i_1, \dots, i_k , vanish. But this restriction must be an invariant of the representation $\mathfrak{sl}(n) \rightarrow \mathfrak{gl}(\text{Mat}_{n \times k})$, and hence trivial, which is a contradiction. So, any non-trivial invariant of ρ has degree at least n . On the other hand, it is easy to see from Proposition 42 that at least one of the vertical indices is less than n . So, by the second part of Theorem 36, the algebra of invariants is not freely generated, as desired (this statement applies in our case, because $\mathfrak{sl}(n)$ is simple and thus none of its representations admit proper semi-invariants). \square

6.2. Standard representations of $\mathfrak{so}(n)$ and $\mathfrak{sp}(n)$

Let us now consider the Lie algebras $\mathfrak{so}(n)$, which is the Lie algebra of skew-symmetric matrices, and $\mathfrak{sp}(n)$. For $\mathfrak{sp}(n)$ we always assume that n is even $n = 2k$ and we denote by $\mathfrak{sp}(n)$ what would usually be denoted by $\mathfrak{sp}(2k)$, that is the space of $n \times n$ matrices X given by the equation

$$X^\top \Omega + \Omega X = 0, \quad \Omega = \begin{pmatrix} 0 & I_{n/2} \\ -I_{n/2} & 0 \end{pmatrix}. \tag{19}$$

In other words, the elements of $\mathfrak{sp}(n)$ have the form $X = \Omega^{-1}S$, where S is symmetric. Note that for both $\mathfrak{so}(n)$ and $\mathfrak{sp}(n)$ the index n is the dimension of the underlying space. Since these Lie algebras correspond to skew-symmetric and symmetric matrices of the same size, most formulas for them differ by some choices of signs. For this reason it is convenient to write the formulas in terms of the number ε , which is equal to $+1$ for $\mathfrak{sp}(n)$ and -1 for $\mathfrak{so}(n)$.

Proposition 45. *Let ρ be the sum of m standard representations of $\mathfrak{so}(n)$ or $\mathfrak{sp}(n)$.*

- (1) *Assume that $m < n$. Let q be the integral part of the quotient $m / (n - m)$, and r be the remainder. Then the JK invariants of ρ are*
 - $(n - m)(n - m + \varepsilon)/2$ horizontal indices:

$$\underbrace{2q + 1, \dots, 2q + 1}_{(n-m-r)(n-m-r+\varepsilon)/2}, \quad \underbrace{2q + 2, \dots, 2q + 2}_{(n-m-r)r}, \quad \underbrace{2q + 3, \dots, 2q + 3}_{r(r+\varepsilon)/2},$$

- $m(m - \varepsilon)/2$ vertical indices $\mathbf{v}_i = 2$.

- (2) *Assume that $m = n$. Then, in the $\mathfrak{so}(n)$ case, the JK invariants of ρ consist of $n(n + 1)/2$ vertical indices:*

$$\underbrace{1, \dots, 1}_n, \quad \underbrace{2, \dots, 2}_{n(n-1)/2},$$

while in the $\mathfrak{sp}(n)$ case the JK invariants are $n(n - 1)/2$ vertical indices $\mathbf{v}_i = 2$ and n Jordan 1×1 blocks with different eigenvalues.

- (3) *Assume that $m > n$. Then the JK invariants of ρ are $n(m - n) + n(n - \varepsilon)/2$ vertical indices:*

$$\underbrace{1, \dots, 1}_{(m-n-\varepsilon)n}, \quad \underbrace{2, \dots, 2}_{n(n+\varepsilon)/2}.$$

The proof of this result is analogous to that of Proposition 42. Now, we use this to study the algebra of invariants. As in the $\mathfrak{sl}(n)$ case, the structure of the algebra of invariants for sums of standard representations of $\mathfrak{so}(n)$ and $\mathfrak{sp}(n)$ is well known, and we discuss it here for the sole purpose of demonstrating the power of Theorem 36.

Proposition 46. *Let $\rho: \mathfrak{so}(n) \rightarrow \mathfrak{gl}(\text{Mat}_{n \times m})$ be the sum of m standard representations of $\mathfrak{so}(n)$, and let $X \in \text{Mat}_{n \times m}$. Then:*

- (1) *For $m < n$, the algebra of invariants of ρ is freely generated by $m(m + 1)/2$ pairwise inner products of columns of X .*
- (2) *For $m \geq n$, the algebra of invariants of ρ is not freely generated.*

Proof of Proposition 46. The first statement is a direct consequence of Theorem 36, since all pairwise inner products of columns of X have degree 2 and hence coincide with vertical indices. To prove the second statement, observe that the same restriction argument as we used in the proof of Proposition 43 shows that there can be no invariants of degree 1. On the other hand, some of the vertical indices are equal to 1, so the algebra of invariants is not freely generated by Theorem 36. \square

Remark 47. For $m \geq n$, the algebra of invariants of ρ is known to be generated by pairwise inner products of columns of X , supplemented by $n \times n$ minors of X . It is easy to see that pairwise inner products alone do not generate the algebra of invariants, as any minor of X can be expressed as the square root of the Gram determinant of its columns, which is a non-polynomial function in terms of the pairwise inner products.

The next statement is proved similarly to Proposition 46.

Proposition 48. *Let $\rho: \mathfrak{sp}(n) \rightarrow \mathfrak{gl}(\text{Mat}_{n \times m})$ be the sum of m standard representations of $\mathfrak{sp}(n)$, and let $X \in \text{Mat}_{n \times m}$. Then:*

- (1) *For $m \leq n + 1$, the algebra of invariants of ρ is freely generated by $m(m - 1)/2$ pairwise symplectic products of columns of X .*
- (2) *For $m > n + 1$, the algebra of invariants of ρ is not freely generated.*

Remark 49. In contrast to the $\mathfrak{so}(n)$ case, the minors of X in the symplectic case can be expressed in terms of pairwise symplectic products. Namely, any minor is equal to the Pfaffian of the symplectic Gram matrix of its columns. So, the algebra of invariants of ρ is generated just by the symplectic products. The relations between these symplectic products for $m > n + 1$ are given by Plücker relations between minors of X .

References

- [1] A. V. Bolsinov, A. M. Izosimov, *Singularities of bi-Hamiltonian systems*, Comm. Math. Physics **331** (2014), no. 2, 507–543.
- [2] A. V. Bolsinov, A. A. Oshemkov, *Bi-Hamiltonian structures and singularities of integrable systems*, Regular and Chaotic Dynamics **14** (2009), no. 4–5, 325–348.
- [3] A. V. Bolsinov, P. Zhang, *Jordan–Kronecker invariants of finite-dimensional Lie algebras*, Transform. Groups **21** (2016), no. 1, 51–86.
- [4] F. Gantmacher, *Theory of Matrices*, Vols. 1, 2, AMS Chelsea Publishing, New York, 1959.
- [5] A. Joseph, D. Shafrir, *Polynomiality of invariants, unimodularity and adapted pairs*, Transform. Groups **15** (2010), no. 4, 851–882.
- [6] F. Knop, *Über die Glattheit von Quotientenabbildungen*, Manuscr. Math. **56** (1986), no. 4, 419–427.
- [7] F. Knop, P. Littelmann, *Der Grad erzeugender Funktionen von Invariantenringen*, Math. Z. **196** (1987), no. 2, 221–229.
- [8] A. I. Ooms, *The Poisson center and polynomial, maximal Poisson commutative subalgebras, especially for nilpotent Lie algebras of dimension at most seven*, J. Algebra **365** (2012), 83–113.
- [9] A. I. Ooms, M. Van den Bergh, *A degree inequality for Lie algebras with a regular Poisson semi-center*, J. Algebra **323** (2010), 305–322.
- [10] A. Panasyuk, *Veronese webs for bi-Hamiltonian structures of higher rank*, Banach Center Publications **51** (2000), 251–261.
- [11] D. Panyushev, *On the coadjoint representation of \mathbb{Z}_2 -contractions of reductive Lie algebras*, Adv. Math. **213** (2007), 380–404.
- [12] D. Panyushev, A. Premet, O. Yakimova, *On symmetric invariants of centralisers in reductive Lie algebras*, J. Algebra **313** (2007), no. 1, 343–391.
- [13] A. Pokrzywa, *On perturbations and the equivalence orbit of a matrix pencil*, Linear Algebra Appl. **82** (1986), 99–121.
- [14] В. Л. Попов, *Теорема конечности для представлений со свободной алгеброй инвариантов*, Изв. АН СССР. Сер. матем. **46** (1982), вып. 2, 347–370. Engl. transl.: V. L. Popov, *A finiteness theorem for representations with a free algebra of invariants*, Math USSR-Izv. **20** (1983), no. 2, 333–354.
- [15] R. C. Thompson, *Pencils of complex and real symmetric and skew matrices*, Linear Algebra Appl. **147** (1991), 323–371.
- [16] F.-J. Turiel, *Classification locale simultanée de deux formes symplectiques compatibles*, Manuscr. Math. **82** (1994), no. 3–4, 349–362.
- [17] А. С. Воронцов, *Кронеckerovy индексы алгебры Ли и оценка степеней инвариантов*, Вестник Моск. Унив. Сер. I Мат. Мех. (2011), вып. 1, 26–30. Engl.

transl.: A. S. Vorontsov, *Kronecker indices of the Lie algebra and an estimate for the degrees of invariants*, Moscow Univer. Math. Bulletin **66** (2011), no. 1, 25–29.

- [18] I. Zakharevich, *Kronecker webs, bihamiltonian structures, and the method of argument translation*, Transform. Groups **6** (2001), no. 3, 267–300.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.