Correction to: Electrophysiological and behavioral responses *Dendroctonus frontalis* and *D. terebrans* (Coleoptera: Curculionidae) to resin odors of host pines (*Pinus* spp.)

Holly L. Munro¹ · Kamal J. K. Gandhi¹ · Brittany F. Barnes¹ · Cristian R. Montes¹ · John T. Nowak³ · William P. Shepherd² · Caterina Villari¹ · Brian T. Sullivan²

Published online: 14 October 2020
© Springer Nature Switzerland AG 2020

Correction to: Chemoecology (2020) 30:215–231
<https://doi.org/10.1007/s00049-020-00311-7>

In Table 2 (Page 220) the units given for release rates of the *endo*- and *exo*-brevicomin flexlures should be mg rather than µg (i.e., both had a release rate of 0.12 mg/day).

The original article can be found online at <https://doi.org/10.1007/s00049-020-00311-7>.

✉ Holly L. Munro
hmunro@uga.edu

¹ D.B. Warnell School of Forestry and Natural Resources,
University of Georgia, 180 East Green Street, Athens,
GA 30602, USA

² USDA Forest Service, Southern Research Station, 2500
Shreveport Highway, Pineville, LA 71360, USA

³ USDA Forest Service, Forest Health Protection, 200
W.T. Weaver Blvd, Asheville, NC 28804, USA