CORRECTION

Correction to: Left-sided breast cancer and risks of secondary lung cancer and ischemic heart disease—Effects of modern radiotherapy techniques

Stefanie Corradini¹ · Hendrik Ballhausen¹ · Helmut Weingandt¹ · Philipp Freisleder¹ · Stephan Schönecker¹ · Maximilian Niyazi¹ · Cristoforo Simonetto² · Markus Eidemüller² · Ute Ganswindt^{1,3} · Claus Belka¹

Published online: 2 November 2017
© Springer-Verlag GmbH Deutschland 2017

Correction to:
Strahlenther Onkol 2017
<https://doi.org/10.1007/s00066-017-1213-y>

Unfortunately, during copy editing, the titles of Fig. 2a and 2b were removed.

The correct Fig. 2a and 2b are shown below. The original article has been corrected accordingly.

The online version of the original article can be found at <https://doi.org/10.1007/s00066-017-1213-y>

✉ Dr. med. Stefanie Corradini
stefanie.corradini@med.uni-muenchen.de

¹ Department of Radiation Oncology, University Hospital, LMU Munich, Marchioninistraße 15, 81377 Munich, Germany

² Institute of Radiation Protection, Helmholtz Zentrum München, Neuherberg, Germany

³ Department of Radiation Oncology, Medical University, Innsbruck, Austria

Fig. 2 The 10-year excess absolute risk (EAR) for major coronary events and secondary lung cancer for medium-risk and high-risk profiles based on OED_{linear} calculations for different treatment modalities and representative treatment ages **a** age 50 and **b** age 70. *3D-CRT* three-dimensional conformal radiotherapy, *VMAT* volumetric-modulated arc radiotherapy, *DIBH* deep-inspiration breath-hold, *FB* free-breathing