

Literatur

- Alexander CL, Shankland GS, Carman W, Williams C (2011) Introduction of a dermatophyte polymerase chain reaction assay to the diagnostic mycology service in Scotland. *Br J Dermatol* 164:966–972
- Badgwell Doherty C, Doherty SD, Rosen T (2010) Thermotherapy in dermatologic infections. *J Am Acad Dermatol* 62:909–927
- Barchiesi F, Arzeni D, Camiletti V et al (2001) In vitro activity of posaconazole against clinical isolates of dermatophytes. *J Clin Microbiol* 39:4208–4209
- Bergmans AM, Ent M van der, Klaassen A et al (2010) Evaluation of a single-tube real-time PCR for detection and identification of 11 dermatophyte species in clinical material. *Clin Microbiol Infect* 16:704–710
- Brasch J, Beck-Jendroscheck V, Gläser R (2011) Fast and sensitive detection of *Trichophyton rubrum* in superficial tinea and onychomycosis by use of a direct PCR assay. *Mycoses* 54:e313–317
- Brasch J, Köppl G (2009) Persisting onychomycosis caused by *Fusarium solani* in an immunocompetent patient. *Mycoses* 52:285–286
- Brillowska-Dabrowska A, Nielsen SS, Nielsen HV, Arendrup MC (2010) Optimized 5-hour multiplex PCR test for the detection of tinea unguium: performance in a routine PCR laboratory. *Med Mycol* 48:828–831
- Budihardja D, Freund V, Maysner P (2010) Widespread erosive tinea corporis by *Arthroderma benhamiae* in a renal transplant recipient: case report. *Mycoses* 53:530–532
- Bueno JG, Martinez C, Zapata B et al (2010) In vitro activity of fluconazole, itraconazole, voriconazole and terbinafine against fungi causing onychomycosis. *Clin Exp Dermatol* 35:658–663
- Cafarchia C, Gasser RB, Figueredo LA et al (2011) Advances in the identification of *Malassezia*. *Mol Cell Probes* 25:1–7
- Carrillo-Muñoz AJ, Giusiano G, Guarro J et al (2007) In vitro activity of voriconazole against dermatophytes, *Scopulariopsis brevicaulis* and other opportunistic fungi as agents of onychomycosis. *Int J Antimicrob Agents* 30:157–161
- Drouot S, Mignon B, Fratti M et al (2009) Pets as the main source of two zoonotic species of the *Trichophyton mentagrophytes* complex in Switzerland, *Arthroderma vanbreuseghemii* and *Arthroderma benhamiae*. *Vet Dermatol* 20:13–18
- Erhard E, Hipler UC, Burmester A et al (2008) Identification of dermatophyte species causing onychomycosis and tinea pedis by MALDI-TOF mass spectrometry. *Exp Dermatol* 17:356–361
- Jackson CJ, Mochizuki T, Barton RC (2006) PCR fingerprinting of *Trichophyton mentagrophytes* var. *interdigitale* using polymorphic subrepeat loci in the rDNA nontranscribed spacer. *J Med Microbiol* 55:1349–1355
- Kamiya A, Kikuchi A, Tomita Y, Kanbe T (2004) PCR and PCR-RFLP techniques targeting the DNA topoisomerase II gene for rapid clinical diagnosis of the etiologic agent of dermatophytosis. *J Dermatol Sci* 34:35–48
- Kanbe T (2008) Molecular approaches in the diagnosis of dermatophytosis. *Mycopathologia* 166:307–317
- Kano R, Okabayashi K, Nakamura Y et al (2000) Differences among chitin synthase I gene sequences in *Trichophyton rubrum* and *T. violaceum*. *Med Mycol* 38:47–50
- Kondori N, Abrahamsson AL, Ataollahy N, Wennerås C (2010) Comparison of a new commercial test, Dermatophyte-PCR kit, with conventional methods for rapid detection and identification of *Trichophyton rubrum* in nail specimens. *Med Mycol* 48:1005–1008
- Krishna G, Beresford E, Ma L, Vickery D et al (2010) Skin concentrations and pharmacokinetics of posaconazole after oral administration. *Antimicrob Agents Chemother* 54:1807–1810
- Lyon JP, Moreira LM, Moraes PC de et al (2011) Photodynamic therapy for pathogenic fungi. *Mycoses* 54:e265–271
- Pryce TM, Palladino S, Kay ID, Coombs GW (2003) Rapid identification of fungi by sequencing the ITS1 and ITS2 regions using an automated capillary electrophoresis system. *Med Mycol* 41:369–381
- Qiao J, Li R, Ding Y, Fang H (2010) Photodynamic therapy in the treatment of superficial mycoses: an evidence-based evaluation. *Mycopathologia* 170:339–343
- Saunte DM, Simmel F, Frimodt-Moller N et al (2007) In vivo efficacy and pharmacokinetics of voriconazole in an animal model of dermatophytosis. *Antimicrob Agents Chemother* 51:317–321
- Savin C, Huck S, Roland C et al (2007) Multicenter evaluation of a commercial PCR-enzyme-linked immunosorbent assay diagnostic kit (Onychodiag) for diagnosis of dermatophytic onychomycosis. *J Clin Microbiol* 45:1205–1210
- Scheinfeld N (2007) A review of the new antifungals: posaconazole, micafungin, and anidulafungin. *J Drugs Dermatol* 6:1249–1251
- Seyfarth F, Wiegand C, Erhard M et al (2011) Identification of yeast isolated from dermatological patients by MALDI-TOF mass spectrometry. *Mycoses* 2012 Feb 13. doi: 10.1111/j.1439-0507.2012.02171.x. [Epub ahead of print]
- Skiada A, Petrikos G (2009) Cutaneous zygomycosis. *Clin Microbiol Infect* 15(Suppl 5):41–45
- Steensels D, Verhaegen J, Lagrou K (2011) Matrix-assisted laser desorption ionization-time of flight mass spectrometry for the identification of bacteria and yeasts in a clinical microbiological laboratory: a review. *Acta Clin Belg* 66:267–273
- Theel ES, Hall L, Mandrekar J, Wengenack NL (2011) Dermatophyte identification using matrix-assisted laser desorption ionization-time of flight mass spectrometry. *J Clin Microbiol* 49:4067–4071
- Wilsmann-Theis D, Sareika F, Bieber T et al (2011) New reasons for histopathological nail-clipping examination in the diagnosis of onychomycosis. *J Eur Acad Dermatol Venereol* 25:235–237 *Ecum quia voluta nullorem sant dendand elentem ea coreritat.*

Erratum

Hautarzt 2012 · 63:395–395
DOI 10.1007/s00105-012-2391-2
© Springer-Verlag 2012

H. Stanisz · C. Pfoehler · T. Vogt · C.S.L. Müller

Klinik für Dermatologie, Allergologie und Venerologie, Homburg/Saar

Erratum zu: Disseminierte infiltrierte juckende Läsionen am Körper

Hautarzt 63:230–232
<http://dx.doi.org/10.1007/s00105-012-2338-7>

In diesem Beitrag wurden ein falscher englischer Titel und ein fehlerhaftes Abstract angegeben. Die korrekten Daten finden Sie im Folgenden – wir bitten um Beachtung.

Disseminated infiltrated pruritic papules

Abstract

The diagnosis of scabies is clinically not always simple without the detection of mites. Histology can be difficult, particularly without clinico-pathologic correlation. Even after adequate antiscabies treatment the dermatological symptoms and complaints can persist and necessitate an intensive follow-up treatment. The case example described here underlines that in consideration of the multifaceted clinical appearance and the sometimes difficult detection of mites, scabies should always be considered as a differential diagnosis in cases of persistent pruritus.

Keywords

Scabies · Mites · Detection · Itching dermatitis · Pruritus

Korrespondenzadresse

H. Stanisz
Klinik für Dermatologie, Allergologie und Venerologie
Kirrberger Str., 66421 Homburg/Saar
hedwig.stanisiz@uks.eu