Correction to: Galectin-3 and soluble RAGE as new biomarkers of post-infarction cardiac remodeling

Alfredo Redondo¹ · Beatriz Paradelo-Dobarro^{2,3} · Isabel Moscoso^{2,3} · María Moure-Álvarez¹ · María Cebro-Márquez² · José Ramón González-Juanatey^{1,2,3} · Javier García-Seara^{1,2,3} · Ezequiel Álvarez^{3,4}

Published online: 13 March 2021

© Springer-Verlag GmbH Germany, part of Springer Nature 2021

Correction to: Journal of Molecular Medicine

<https://doi.org/10.1007/s00109-021-02054-6>

The original publication contains an error. Correct figure 1 is presented in this paper.

The original article has been corrected.

The online version of the original article can be found at <https://doi.org/10.1007/s00109-021-02054-6>

✉ Ezequiel Álvarez
ezequiel.alvarez.castro@gmail.com

¹ Servicio de Cardiología y Unidad de Hemodinámica, Complejo Hospitalario Universitario de Santiago de Compostela (CHUS), SERGAS, Travesía da Choupana s/n, Santiago de Compostela, 15706 A Coruña, Spain

² Instituto de Investigación Sanitaria de Santiago de Compostela (IDIS), Complejo Hospitalario Universitario de Santiago de Compostela (CHUS), SERGAS, Travesía da Choupana s/n, Santiago de Compostela, 15706 A Coruña, Spain

³ CIBERCV, Madrid, Spain

⁴ Laboratorio No. 6. Edif. Consultas Externas (Planta-2), Instituto de Investigación Sanitaria de Santiago de Compostela (IDIS), Complejo Hospitalario Universitario de Santiago de Compostela (CHUS), SERGAS, Travesía da Choupana s/n, Santiago de Compostela, 15706 A Coruña, Spain

956 Fig. 1 Linear correlation between plasma levels during 48 h after STEMI of galectin-3 (a) and sRAGE (b) with the percentage of left ventricle ejection fraction (LVEF). The straight lines show total adjustment with their confidence intervals (curved lines). r_s = Spearman's correlation coefficient

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.