

## **Erratum to: The antennal sensilla of *Melipona quadrifasciata* (Hymenoptera: Apidae: Meliponini): a study of different sexes and castes**

**Samira Veiga Ravaiano · Rúdo de Paiva Ferreira ·  
Lucio Antonio de Oliveira Campos ·  
Gustavo Ferreira Martins**

Published online: 2 July 2014  
© Springer-Verlag Berlin Heidelberg 2014

**Erratum to: Naturwissenschaften**  
**DOI 10.1007/s00114-014-1184-0**

The original publication of this article has incorrect presentation of Table 2. Correct table is shown on the following page.

---

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00114-014-1184-0>.

---

S. V. Ravaiano · G. F. Martins  
Departamento de Entomologia, Universidade Federal de Viçosa  
(UFV), 36570-900 Viçosa, Minas Gerais, Brazil

R. d. P. Ferreira · L. A. d. O. Campos · G. F. Martins (✉)  
Departamento de Biologia Geral, Universidade Federal de Viçosa  
(UFV), 36570-900 Viçosa, Minas Gerais, Brazil  
e-mail: [gmartins@ufv.br](mailto:gmartins@ufv.br)

**Table 2** Sizes of sensilla in the antenna of *M. quadrifasciata*. Different letters in the same column indicate significant differences (ANOVA,  $p < 0.05$ ). QLM- queen-like male; ST- sensillum trichodeum; SCa- sensillum chaeticum; SP- sensillum placodeum; SB- sensillum basiconicum; SCo- sensillum coeloconicum; SA- sensillum ampullaceum and SCp- sensillum campaniformium

Class of individual	Sizes of sensilla ( $\mu\text{m}$ )												
	ST I	ST II	ST III	ST IV	ST V	ST VI	SCa I	SCa II	SP	SB	SC	SA	SCp
worker	15.14 ( $\pm 1.38$ ) <sup>a</sup>	13.75 ( $\pm 1.13$ ) <sup>a</sup>	26.75 ( $\pm 2.20$ ) <sup>a</sup>	17.40 ( $\pm 2.21$ ) <sup>a</sup>	19.63 ( $\pm 1.25$ ) <sup>b</sup>	20.81 ( $\pm 2.18$ ) <sup>b</sup>	17.10 ( $\pm 3.09$ ) <sup>a</sup>	-	15.29 ( $\pm 1.42$ ) <sup>a</sup>	8.93 ( $\pm 1.14$ ) <sup>a</sup>	1.72 ( $\pm 0.40$ )	0.52 ( $\pm 0.15$ )	4.88 ( $\pm 0.12$ )
queen	13.78 ( $\pm 1.39$ ) <sup>b</sup>	15.04 ( $\pm 2.48$ ) <sup>a</sup>	26.67 ( $\pm 3.79$ ) <sup>a</sup>	18.04 ( $\pm 2.10$ ) <sup>b</sup>	30.08 ( $\pm 4.73$ ) <sup>a</sup>	32.42 ( $\pm 4.17$ ) <sup>a</sup>	18.69 ( $\pm 2.10$ ) <sup>a</sup>	-	13.64 ( $\pm 1.01$ ) <sup>b</sup>	9.23 ( $\pm 2.02$ ) <sup>a</sup>	2.51 ( $\pm 0.08$ )	0.73 ( $\pm 0.17$ )	4.59 ( $\pm 0.29$ )
n male	6.80 ( $\pm 0.69$ ) <sup>c</sup>	7.82 ( $\pm 0.70$ ) <sup>b</sup>	13.77 ( $\pm 0.80$ ) <sup>b</sup>	18.51 ( $\pm 2.46$ ) <sup>b</sup>	19.26 ( $\pm 2.22$ ) <sup>b</sup>	21.62 ( $\pm 2.17$ ) <sup>b</sup>	-	8.43 ( $\pm 1.35$ ) <sup>a</sup>	8.09 ( $\pm 0.64$ ) <sup>c</sup>	-	-	-	-
male 2n	6.10 ( $\pm 0.93$ ) <sup>c</sup>	7.35 ( $\pm 1.01$ ) <sup>b</sup>	13.94 ( $\pm 0.85$ ) <sup>b</sup>	12.74 ( $\pm 2.13$ ) <sup>c</sup>	19.07 ( $\pm 2.61$ ) <sup>b</sup>	20.09 ( $\pm 1.57$ ) <sup>b</sup>	-	8.55 ( $\pm 1.20$ ) <sup>a</sup>	8.63 ( $\pm 1.16$ ) <sup>c</sup>	-	-	-	3.26 ( $\pm 0.09$ )
QLM	8.26 ( $\pm 0.51$ ) <sup>d</sup>	8.81 ( $\pm 0.68$ ) <sup>b</sup>	16.54 ( $\pm 1.80$ ) <sup>b</sup>	15.50 ( $\pm 2.75$ ) <sup>a</sup>	19.09 ( $\pm 1.87$ ) <sup>b</sup>	27.00 ( $\pm 8.46$ ) <sup>b</sup>	-	9.06 ( $\pm 1.00$ ) <sup>a</sup>	9.31 ( $\pm 0.95$ ) <sup>c</sup>	-	-	-	4.06 ( $\pm 1.23$ )