

- parallel trial of amantadine, pemoline, and placebo. *Neurology* 45(11):1956–1961
78. Weinschenker BG, Penman M, Bass B, Ebers GC, Rice GP (1992) A double-blind, randomized, crossover trial of pemoline in fatigue associated with multiple sclerosis. *Neurology* 42(8):1468–1471
 79. Cohen RA, Fisher M (1989) Amantadine treatment of fatigue associated with multiple sclerosis. *Arch Neurol* 46(6):676–680
 80. Rosenberg GA, Appenzeller O (1988) Amantadine, fatigue, and multiple sclerosis. *Arch Neurol* 45(10):1104–1106
 81. The Canadian MS Research Group (1987) A randomized controlled trial of amantadine in fatigue associated with multiple sclerosis. *Can J Neurol Sci* 14(3):273–278
 82. Mücke M, Mochamat, Cuhls H, Peuckmann-Post V, Minton O, Stone P, Radbruch L (2015) Pharmacological treatments for fatigue associated with palliative care. In: *The Cochrane Collaboration (Hrsg) Cochrane Database of Systematic Reviews*. John Wiley & Sons, Chichester
 83. Pucci E, Branäs P, D'Amico R, Giuliani G, Solari A, Taus C (2007) Amantadine for fatigue in multiple sclerosis. *Cochrane Database Syst Rev* 2007:CD002818
 84. Tejani AM, Wasdell M, Spiwak R, Rowell G, Nathwani S (2010) Carnitine for fatigue in multiple sclerosis. In: *The Cochrane Collaboration (Hrsg) Cochrane Database of Systematic Reviews*. John Wiley & Sons, Chichester
 85. Peuckmann-Post V, Elsner F, Krumm N, Trottenberg P, Radbruch L (2010) Pharmacological treatments for fatigue associated with palliative care. In: *The Cochrane Collaboration (Hrsg) Cochrane Database of Systematic Reviews*. John Wiley & Sons, Chichester
 86. Grossman P, Kappos L, Gensicke H, D'Souza M, Mohr DC, Penner IK, Steiner C (2010) MS quality of life, depression, and fatigue improve after mindfulness training: A randomized trial. *Neurology* 75(13):1141–1149
 87. Moss-Morris R, McCrone P, Yardley L, Kessel K van, Wills G, Dennison L (2012) A pilot randomised controlled trial of an Internet-based cognitive behavioural therapy self-management programme (MS Invigor8) for multiple sclerosis fatigue. *Behav Res Ther* 50(6):415–421
 88. Kessel K van, Moss-Morris R, Willoughby E, Chalder T, Johnson MH, Robinson E (2008) A randomized controlled trial of cognitive behavior therapy for multiple sclerosis fatigue. *Psychosom Med* 70(2):205–213
 89. Rietberg MB, Brooks D, Uitdehaag BM, Kwakkel G (2005) Exercise therapy for multiple sclerosis. In: *The Cochrane Collaboration (Hrsg) Cochrane Database of Systematic Reviews*. John Wiley & Sons, Chichester
 90. Hebert JR, Corboy JR, Manago MM, Schenkman M (2011) Effects of vestibular rehabilitation on multiple sclerosis-related fatigue and upright postural control: A randomized controlled trial. *Phys Ther* 91(8):1166–1183
 91. Braley TJ, Segal BM, Chervin RD (2015) Hypnotic use and fatigue in multiple sclerosis. *Sleep Med* 16(1):131–137
 92. Veauthier C (2015) Hypnotic use and multiple sclerosis related fatigue: a forgotten confounder. *Sleep Med* 16(3):319
 93. Veauthier C, Paul F (2012) Fatigue in multiple sclerosis: which patient should be referred to a sleep specialist? *Mult Scler* 18(2):248–249

Nervenarzt 2016 · 87:1321
 DOI 10.1007/s00115-016-0233-7
 Online publiziert: 17. November 2016
 © Springer-Verlag Berlin Heidelberg 2016



W. Heide

Neurologische Klinik, Allgemeines Krankenhaus Celle, Celle, Deutschland

Erratum zu: „Non-heart-beating donors“ sind nicht geeignet

Erratum zu

Nervenarzt (2016)

DOI: 10.1007/s00115-015-0048-y

In der Korrespondenzadresse wurden versehentlich die akademischen Titel des Autors nicht genannt.

Korrekt ist: Prof. Dr. W. Heide.

Wir bitten diesen Fehler zu entschuldigen.

Korrespondenzadresse

Prof. Dr. W. Heide

Neurologische Klinik, Allgemeines

Krankenhaus Celle

Siemensplatz 4, 29223 Celle, Deutschland

wolfgang.heide@akh-celle.de

Die Onlineversion des Originalbeitrages finden Sie unter <http://dx.doi.org/10.1007/s00115-015-0048-y>.