Correction to: Genome-wide mapping of adult plant stripe rust resistance in wheat cultivar Toni

Xinli Zhou¹ · Tian Hu¹ · Xin Li¹ · Ma YU¹ · Yuanyuan Li¹ · Suizhuang Yang¹ · Keping Huang¹ · Dejun Han² · Zhensheng Kang²

Published online: 30 May 2019

© Springer-Verlag GmbH Germany, part of Springer Nature 2019

Correction to: Theoretical and Applied Genetics

<https://doi.org/10.1007/s00122-019-03308-1>

Unfortunately, Figs. 2 and 3 were interchanged in the results section. The figures should swap positions, whereas the legends should stay in the given order.

The correct Figs. 2 and 3 are given below:

The original article can be found online at <https://doi.org/10.1007/s00122-019-03308-1>.

✉ Suizhuang Yang
yangszh@126.com

Zhensheng Kang
kangzs@nwsuaf.edu.cn

¹ Wheat Research Institute, School of Life Sciences and Engineering, Southwest University of Science and Technology, Mianyang 621010, Sichuan, People's Republic of China

² State Key Laboratory of Crop Stress Biology in Arid Areas and College of Plant Protection, Northwest A&F University, Yangling 712100, Shaanxi, China

Fig. 2 Frequency distributions of mean infection types (IT) (A) and disease severities (DS) for 171 F_7 RILs from cross Mingxian169 × Toni grown at Yangling and Tianshui in 2016–2017. Arrows indicate the values of the parental lines

Fig. 3 Circos diagram showing LOD values on genetic maps for 21 wheat chromosomes. Track **a** chromosomes and genetic scale (cM). Track **b** stripe rust resistance QTL detected by ICIM in the M169/Toni RIL population across four environments based on mean infection type. Track **c** stripe rust resistance QTL detected by ICIM in the M169/Toni RIL population across four environments based on mean disease severity (MDS) (track **c**)