



Correction to: Dissection of two quantitative trait loci with pleiotropic effects on plant height and spike length linked in coupling phase on the short arm of chromosome 2D of common wheat (*Triticum aestivum* L.)

Lingling Chai^{1,2} · Zhaoyan Chen^{1,2} · Ruolin Bian^{1,2} · Huijie Zhai³ · Xuejiao Cheng^{1,2} · Huiru Peng^{1,2} · Yingyin Yao^{1,2} · Zhaorong Hu^{1,2} · Mingming Xin^{1,2} · Weilong Guo^{1,2} · Qixin Sun^{1,2} · Aiju Zhao⁴ · Zhongfu Ni^{1,2}

Published online: 24 September 2019
© Springer-Verlag GmbH Germany, part of Springer Nature 2019

Correction to:

Theoretical and Applied Genetics (2018) 131:2621–2637
<https://doi.org/10.1007/s00122-018-3177-4> and
Theoretical and Applied Genetics (2019) 132:1815–1831
<https://doi.org/10.1007/s00122-019-03318-z>

This article was unintentionally published twice in this journal. The following should be considered the version of record and used for citation purposes: Lingling Chai, Zhaoyan Chen, Ruolin Bian, Huijie Zhai, Xuejiao Cheng, Huiru Peng, Yingyin Yao, Zhaorong Hu, Mingming Xin, Weilong Guo, Qixin Sun, Aiju Zhao, Zhongfu Ni, Dissection of two quantitative trait loci with pleiotropic effects on plant height and spike length linked in coupling phase on the short arm of chromosome 2D of common wheat (*Triticum aestivum* L.), Theoretical and Applied Genetics, Volume 132,

Issue 6, 2019, pages 1815–1831, <https://doi.org/10.1007/s00122-019-03318-z>.

The duplicate Lingling Chai, Zhaoyan Chen, Ruolin Bian, Huijie Zhai, Xuejiao Cheng, Huiru Peng, Yingyin Yao, Zhaorong Hu, Mingming Xin, Weilong Guo, Qixin Sun, Aiju Zhao, Zhongfu Ni, Dissection of two quantitative trait loci with pleiotropic effects on plant height and spike length linked in coupling phase on the short arm of chromosome 2D of common wheat (*Triticum aestivum* L.), Theoretical and Applied Genetics, Volume 131, Issue 12, 2018, pages 2621–2637, <https://doi.org/10.1007/s00122-018-3177-4> is to be ignored.

Springer Nature apologizes to the readers of the journal for not detecting the duplication during the publication process.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s00122-019-03318-z>.

✉ Aiju Zhao
zhaoaiju81@126.com

✉ Zhongfu Ni
nizf@cau.edu.cn

- ¹ State Key Laboratory for Agrobiotechnology/Key Laboratory of Crop Heterosis and Utilization, the Ministry of Education/Key Laboratory of Crop Genetic Improvement, Beijing Municipality, China Agricultural University, Beijing 100193, China
- ² National Plant Gene Research Centre, Beijing 100193, China
- ³ College of Agronomy, Henan Agricultural University, Zhengzhou 450002, China
- ⁴ Institute of Cereal and Oil Crops, Hebei Academy of Agriculture/Forestry Sciences, Hebei Crop Genetic Breeding Laboratory, Shijiazhuang 050035, China