



Correction to: A recessive gene *Cmpmr2F* confers powdery mildew resistance in melon (*Cucumis melo* L.)

Taifeng Zhang^{1,2} · Haonan Cui^{3,4} · Feishi Luan^{1,2} · Hongyu Liu^{1,2} · Zhuo Ding^{3,4} · Sikandar Amanullah^{1,2} · Manlin Zhang^{1,2} · Tingting Ma^{1,2} · Peng Gao^{1,2}

Published online: 7 April 2023
© Springer-Verlag GmbH Germany, part of Springer Nature 2023

Correction to:

Theoretical and Applied Genetics (2023) 136:4
<https://doi.org/10.1007/s00122-023-04269-2>

The references 30, 32, 33 and 59 have been incorrectly published in the original publication. The complete correct references are given below.

Lebeda A, Kříštková E, Sedláková B et al (2008) New concept for determination and denomination of pathotypes and races of cucurbit powdery mildew. In: Pitrat M (ed) Cucurbitaceae 2008. Proceedings of the IXth EUCARPIA Meeting on Genetics and Breeding of Cucurbitaceae. INRA, Avignon (France), pp 125–134 (30th reference)

Lebeda A, Sedláková B (2006) Identification and survey of cucurbit powdery mildew races in Czech populations. In:

Holmes G J (ed) Proceedings of Cucurbitaceae 2006. Universal Press, Raleigh, North Carolina, USA, pp 444–452 (32th reference)

Lebeda A, Sedláková B (2010) Chapter 19—Screening for resistance to cucurbit powdery mildews (*Golovinomyces cichoracearum*, *Podosphaera xanthii*). In: Spencer M M, Lebeda A (eds) Mass Screening Techniques for Selecting Crops Resistant to Disease. International Atomic Energy Agency (IAEA), Vienna, Austria (ISBN 978-92-0-105110-3), pp 295–307 (33th reference)

Sedláková B, Lebeda A, Gryczová K et al (2014) Virulence structure (pathotypes, races) of cucurbit powdery mildew populations in the Czech Republic in the years 2010–2012. In: Havey M, Weng Y, Day B, Grumet R (eds) Cucurbitaceae 2014 Proceedings. Bay Harbor, Michigan, USA, October 12–16, 2014, Michigan State University and University of Wisconsin-Madison, USA. American Society

The original article can be found online at <https://doi.org/10.1007/s00122-023-04269-2>.

✉ Peng Gao
gaopeng_neau@163.com

Taifeng Zhang
ztf18104603353@163.com

Haonan Cui
chn4051@hevttc.edu.cn

Feishi Luan
luanfeishi@neau.edu.cn

Hongyu Liu
hyliu@neau.edu.cn

Zhuo Ding
1025132245@qq.com

Sikandar Amanullah
sikandaraman@yahoo.com

Manlin Zhang
zml15231228878@163.com

Tingting Ma
2430567867@qq.com

¹ College of Horticulture and Landscape Architecture, Northeast Agricultural University, Harbin 150036, Heilongjiang, China

² Key Laboratory of Biology and Genetic Improvement of Horticulture Crops (Northeast Region), Ministry of Agriculture and Rural Affairs, Northeast Agricultural University, Harbin 150036, Heilongjiang, China

³ College of Horticulture Science and Technology, Hebei Normal University of Science and Technology, Qinhuangdao 066004, China

⁴ Hebei Key Laboratory of Horticultural Germplasm Excavation and Innovative Utilization, Qinhuangdao 066004, China

for Horticultural Science, Alexandria, VA, USA (ISBN 978-0-9830932-2-0), pp 28–31 (59th reference)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article has been corrected.