

Erratum

Common genetic polymorphisms in the promoter of resistin gene are major determinants of plasma resistin concentrations in humans

Y. M. Cho^{1, 2} · B.-S. Youn³ · S. S. Chung² · K. W. Kim² · H. K. Lee¹ · K.-Y. Yu³ · H. J. Park³ · H. D. Shin⁴ · K. S. Park^{1, 2}

¹ Department of Internal Medicine, Seoul National University College of Medicine, Seoul, Korea

² Genome Research Center for Diabetes and Endocrine Disease, Clinical Research Institute, Seoul National University Hospital, Seoul, Korea

³ KOMED Institute for Life Science, Graduate School of Biotechnology, Korea University, Seoul, Korea

⁴ Department of Genetic Epidemiology, SNP Genetics, Seoul, Korea

Diabetologia (2004) 47:559–565

Figure 1a. The correct positions of the arrows for Band 1 and Band 2 are shown below:

Published online: 25 June 2004

© Springer-Verlag 2004

K. S. Park (✉)
Department of Internal Medicine,
Seoul National University College of Medicine,
28 Yongon-Dong Chongno-Gu, Seoul, 110–744 Korea
E-mail: kspark@snu.ac.kr

The online version of the original article can be found at
<http://dx.doi.org/10.1007/s00125-004-1319-x>



a g.-420C>G**b** g.-537A>C

Fig. 1a. Binding of nuclear proteins to -420 and -537 regions of the resistin gene. The probe, -433 to -406 bp of the resistin gene containing -420G, was labelled as described in Materials and Methods. Lane 1: probe without nuclear proteins; lane 2:

nuclear proteins (5 μ g) of 3T3-L1; lanes 3-7: nuclear proteins (10 μ g) of 3T3-L1; lane 8: nuclear proteins (5 μ g) of THP-1; lanes 9-13: nuclear proteins (10 μ g) of THP-1. Competitors were added to the probe at 200- or 500-fold molar excesses