

K. A. McAuley · C. M. Hopkins · K. J. Smith ·
R. T. McLay · S. M. Williams · R. W. Taylor ·
J. I. Mann

Comparison of high-fat and high-protein diets with a high-carbohydrate diet in insulin-resistant obese women

Published online: 16 April 2005
© Springer-Verlag 2005

Diabetologia (2005) 48:8–16

Unfortunately there was a mistake in the Abstract of this paper. The third sentence in the 'Results' section should have read:

When compared with the HC diet, the HF and HP diets were shown to produce significantly ($p < 0.01$) greater reductions in several parameters, including weight loss (HF -2.8 kg, HP -2.7 kg), waist circumference (HF -3.5 cm, HP -2.7 cm) and triglycerides (HF -0.30 mmol/l, HP -0.22 mmol/l).

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00125-004-1603-4>

K. A. McAuley (✉) · J. I. Mann
Edgar National Centre for Diabetes Research,
Medical and Surgical Sciences,
University of Otago,
PO Box 56, Dunedin, New Zealand
e-mail: kirsten.mcauley@stonebow.otago.ac.nz
Tel.: +64-3-4798153
Fax: +64-3-4747641

K. A. McAuley · C. M. Hopkins · K. J. Smith · R. T. McLay ·
R. W. Taylor · J. I. Mann
Department of Human Nutrition,
University of Otago,
Dunedin, New Zealand

S. M. Williams
Department of Preventive and Social Medicine,
University of Otago,
Dunedin, New Zealand