

Erratum to: The proteomic signature of insulin-resistant human skeletal muscle reveals increased glycolytic and decreased mitochondrial enzymes

J. Giebelstein · G. Poschmann · K. Højlund ·
W. Schechinger · J. W. Dietrich · K. Levin ·
H. Beck-Nielsen · K. Podwojski · K. Stühler ·
H. E. Meyer · H. H. Klein

Published online: 29 April 2012
© Springer-Verlag 2012

Erratum to: Diabetologia
DOI 10.1007/s00125-012-2456-x

Unfortunately, “negatively” and “positively” were inadvertently reversed in the legend to Fig. 6. The third sentence of the legend should read: A complete dark red or dark blue label indicates proteins where combined spot volumes (combination of all spots where the respective protein was identified) were significantly positively (red) or negatively (blue) correlated with GDR ($p < 0.05$).

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00125-012-2456-x>.

J. Giebelstein · W. Schechinger · J. W. Dietrich · H. H. Klein (✉)
Medizinische Klinik I, Berufsgenossenschaftliches,
Universitätsklinikum Bergmannsheil,
Klinikum der Ruhr Universität Bochum,
Bürkle-de-la-Camp-Platz 1,
44789 Bochum, Germany
e-mail: harald.klein@rub.de

G. Poschmann · K. Podwojski · K. Stühler · H. E. Meyer
Medizinisches Proteom-Center, Ruhr Universität Bochum,
Bochum, Germany

K. Højlund · K. Levin · H. Beck-Nielsen
Diabetes Research Center, Department of Endocrinology,
Odense University Hospital,
Odense, Denmark

W. Schechinger
Bioavid Diagnostics,
Dieburg, Germany

K. Podwojski
Department of Statistical Methods in Genetics and Chemometrics,
Technical University Dortmund,
Dortmund, Germany

K. Stühler
Molecular Proteomics Laboratory,
Heinrich Heine Universität Düsseldorf,
Düsseldorf, Germany