

Erratum to: Prevalence of diabetes and impaired fasting glucose in the Chinese adult population: International Collaborative Study of Cardiovascular Disease in Asia (InterASIA)

D. Gu · K. Reynolds · X. Duan · X. Xin · J. Chen ·
X. Wu · J. Mo · P. K. Whelton · J. He ·
for the InterASIA Collaborative Group

Published online: 16 August 2012
© Springer-Verlag 2012

Erratum to: Diabetologia
DOI 10.1007/s00125-003-1167-8

Age-specific numbers were inadvertently switched between the ‘Undiagnosed diabetes’ and ‘Impaired fasting glucose’ columns in Table 3. The corrected section of the table is reproduced here.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00125-003-1167-8>.

D. Gu · X. Duan · X. Wu
Cardiovascular Institute and Fuwai Hospital of Chinese Academy
of Medical Sciences, Peking Union Medical College,
Beijing, China

K. Reynolds · X. Xin · P. K. Whelton · J. He (✉)
Department of Epidemiology,
Tulane University School of Public Health and Tropical Medicine,
1430 Tulane Avenue SL18,
New Orleans, LA 70112, USA
e-mail: jhe@tulane.edu

J. Chen · P. K. Whelton · J. He
Tulane University School of Medicine,
New Orleans, LA, USA

J. Mo
Pfizer Inc,
New York, NY, USA

Table 3 Prevalence of undiagnosed diabetes and impaired fasting glucose^a in the adult population aged 35–74 Years in China, 2000–2001

Age, yrs	Men		Women		Total	
	Undiagnosed Diabetes, % (SE)	Impaired Fasting Glucose, % (SE)	Undiagnosed Diabetes, % (SE)	Impaired Fasting Glucose, % (SE)	Undiagnosed Diabetes, % (SE)	Impaired Fasting Glucose, % (SE)
North						
35–44	3.2 (0.5)	8.7 (1.0)	3.7 (0.6)	5.1 (0.8)	3.4 (0.4)	7.0 (0.7)
45–54	3.3 (0.7)	7.6 (1.1)	5.5 (0.9)	6.6 (1.0)	4.4 (0.6)	7.1 (0.7)
55–64	5.8 (1.2)	8.8 (1.4)	5.6 (1.1)	11.2 (1.6)	5.7 (0.8)	10.0 (1.1)
65–74	3.9 (1.3)	12.2 (2.4)	10.0 (1.8)	9.1 (2.0)	6.9 (1.1)	10.7 (1.6)
35–74	3.8 (0.4)	8.8 (0.6)	5.3 (0.5)	7.1 (0.6)	4.6 (0.3)	8.0 (0.4)
Age-Standardized ^b	4.3 (0.4)	9.1 (0.6)	5.7 (0.4)	6.8 (0.5)	5.0 (0.3)	7.9 (0.4)