

Erratum to: Body iron stores and risk of type 2 diabetes: results from the European Prospective Investigation into Cancer and Nutrition (EPIC)-Potsdam study

J. Montonen · H. Boeing · A. Steffen · R. Lehmann ·
A. Fritsche · H.-G. Joost · M. B. Schulze · T. Pischon

Published online: 4 September 2012
© Springer-Verlag 2012

Erratum to: Diabetologia
DOI 10.1007/s00125-012-2633-y

Instances of ‘mg/l’ for ferritin values should instead have read ‘µg/dl’. (To convert values in µg/dl into the SI unit of pmol/l, multiply by 22.47.)

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00125-012-2633-y>.

J. Montonen · H. Boeing · A. Steffen (✉) · T. Pischon
Department of Epidemiology, German Institute of Human
Nutrition Potsdam-Rehbruecke,
Arthur-Scheunert-Allee 114-116,
14558 Nuthetal, Germany
e-mail: annika.steffen@dife.de

R. Lehmann
Central Laboratory/Clinical Chemistry and Pathobiochemistry,
University Hospital Tübingen,
Tübingen, Germany

R. Lehmann · A. Fritsche
Paul Langerhans Institute Tübingen (Inst. for Diabetes Research
and Metabolic Diseases of the Helmholtz Centre Munich
at the University of Tübingen),
Tübingen, Germany

A. Fritsche
Department of Internal Medicine IV, University of Tübingen,
Tübingen, Germany

H.-G. Joost
Department of Pharmacology, German Institute of Human
Nutrition Potsdam-Rehbruecke,
Nuthetal, Germany

M. B. Schulze
Department of Molecular Epidemiology,
German Institute of Human Nutrition Potsdam-Rehbruecke,
Nuthetal, Germany

T. Pischon
Molecular Epidemiology Group,
Max Delbrück Center for Molecular Medicine (MDC),
Berlin-Buch, Germany