

# A diabetic milieu promotes OCT4 and NANOG production in human visceral-derived adipose stem cells

P. Dentelli · C. Barale · G. Togliatto · A. Trombetta ·  
C. Olgasi · M. Gili · C. Riganti · M. Toppino ·  
M. F. Brizzi

Received: 18 July 2012 / Accepted: 30 August 2012 / Published online: 12 October 2012  
© Springer-Verlag Berlin Heidelberg 2012

## Abstract

**Aims/hypothesis** Successful outcomes have been obtained by exploiting adipose-derived stem cells (ASCs) in regenerative medicine. NADPH oxidase (NOX)-generated reactive oxygen species (ROS) are known to control stem cell self-renewal. Several high glucose (HG)-mediated effects depend on NOX-generated ROS. In this study, we investigated whether, and how mechanistically, HG concentrations control ASC fate in patients with diabetes.

**Methods** ASCs from the visceral adipose tissue of non-diabetic (N-ASCs) and diabetic participants (D-ASCs), identified by surface markers, were counted and evaluated for ROS generation and stem cell properties. Their ability to release soluble factors was assessed by BioPlex analysis. To reproduce an in vitro diabetic glucose milieu, N-ASCs were cultured in HG (25 mmol/l) or normal glucose (NG) concentration (5 mmol/l), as control. ASC pluripotency was assessed by in vitro study. The p47<sup>phox</sup> NOX subunit, AKT and octamer-binding transcription factor 4 (OCT4; also known as POU5F1) were knocked down by small-interfering RNA

technology. Stem-cell features were evaluated by sphere cluster formation.

**Results** The ASC number was higher in diabetic patients than in non-diabetic controls. Production of OCT4 and NANOG, stem-cell-specific transcription factors, was upregulated in D-ASCs compared with N-ASCs. Moreover, we found that ROS production and AKT activation drove D-ASC, but not N-ASC, secretion. When N-ASCs were cultured in vitro in the presence of HG, they also expressed OCT4/NANOG and formed spheres. By knock-down of the p47<sup>phox</sup> NOX subunit, AKT and OCT4 we demonstrated that NOX-generated ROS and their downstream signals are crucial for HG-mediated ASC de-differentiation and proinflammatory cytokine production.

**Conclusions/interpretation** We herein provide a rationale for exploiting D-ASCs in regenerative medicine and/or exploiting HG preconditioning to increase ASCs ex vivo.

**Keywords** Adipose-derived stem cells · Cytokines · High glucose · Visceral adipose tissue

**Electronic supplementary material** The online version of this article (doi:10.1007/s00125-012-2734-7) contains peer-reviewed but unedited supplementary material, which is available to authorised users.

P. Dentelli and C. Barale contributed equally to this study.

P. Dentelli · C. Barale · G. Togliatto · A. Trombetta · C. Olgasi ·  
M. Gili · M. F. Brizzi (✉)  
Department of Medical Sciences, University of Torino,  
Corso Dogliotti 14,  
10126 Torino, Italy  
e-mail: mariafelice.brizzi@unito.it

C. Riganti  
Department of Oncology, University of Torino,  
Torino, Italy

M. Toppino  
Department of Surgical Sciences, University of Torino,  
Torino, Italy

## Abbreviations

ASC	Adipose-derived stem cell
D-ASCs	Adipose-derived stem cells from diabetic participants
DCF-DA	5-(and-6)-Carboxy-2',7'-dichlorofluorescein diacetate
HG	High glucose
HM	High mannitol
MSC	Mesenchymal stem cell
N-ASCs	Adipose-derived stem cells from non-diabetic participants
NG	Normal glucose
NOX	NADPH oxidase
OCT4	Octamer-binding transcription factor 4
PCNA	Proliferating cell nuclear antigen
PI3K	Phosphoinositide 3-kinase

qRT-PCR	Quantitative real-time PCR
ROS	Reactive oxygen species
siRNA	Small interfering RNA

## Introduction

Human adult mesenchymal stem cells (MSCs), owing to their useful properties, have become of increasing interest to clinicians [1–4]. Indeed, their ability to differentiate into several mesenchymal lineages has spurred clinicians to exploit these cells in tissue engineering and in regenerative medicine to repair damaged organs [1–4]. Different tissues have been discovered as sources of adult stem cells and recently adipose tissue has been considered a novel promising, abundant and accessible resource of adipose-derived stem cells (ASCs) [5–7]. Successful outcomes have been reported from a number of clinical trials exploiting ASCs, mainly in regeneration or repair of failing organs [8], although a number of relevant issues remain to be clarified. In particular, whether the therapeutic effect of ASCs depends merely on their engraftment and differentiation is still debated [9]. Moreover, very few data are as yet available on the molecular mechanisms that regulate their self-renewal and differentiation, both in physiological and pathological settings [10].

The stem cell microenvironment, the ‘stem cell niche’, in adipose tissue is crucial for preserving ASC self-renewal, denoting ‘stemness’ potential and controlling differentiation [11]. To date, data have been provided on the negative effects of obesity on the adipogenic niche [10]. However, how mechanistically abnormal metabolic conditions act on niches to control ASC fate is still poorly understood. Likewise, whether and how abnormal metabolic conditions might alter the ability of ASCs to release soluble mediators, a mechanism required for their therapeutic potential [12], is still unknown.

Stem cell pluripotency and differentiation are strictly controlled by a coordinated network of transcription factors [13]. Among them, octamer-binding transcription factor 4 (OCT4) and NANOG, the levels of which are strictly controlled by OCT4 and SRY-box 2 (SOX2) [14], have been long recognised as crucial transcriptional regulators of stem cell self-renewal during embryogenesis [15]. More recently, it has been shown that both OCT4 and NANOG are also produced by adult undifferentiated MSCs and can be recovered from bone marrow, human cord blood and umbilical cord matrix [15–19].

The ability to reciprocally de-differentiate, re-differentiate or trans-differentiate in response to a specific stimulus is defined as cell plasticity [20–22]. *OCT4* overexpression has been shown to induce de-differentiation of ASCs into a more immature status by activating the AKT/phosphoinositide 3-kinase (PI3K) and extracellular signal-related kinase (ERK1/

2) signalling pathways [23]. As in developmental processes, in adult tissues several factors are engaged to modulate and maintain the balance between self-renewal and differentiation [21, 22]. Indeed, it has been reported that ASCs possess their own multipotency to reprogramme into more primitive stem cells [23]. However, how this process takes place mechanistically is still controversial.

NADPH oxidase (NOX)-mediated reactive oxygen species (ROS) production [24] has been reported to be involved in the control of stem and progenitor cell fate [25], as well as in neural cell reprogramming [26]. Likewise, ROS generation in response to hypoxia seems to control ASC proliferation [27].

In the diabetic setting, hyperglycaemia, via NOX enzymatic activity, upholds the production of ROS, which in turn acts as a second messenger [28]. Collectively, this might suggest that high glucose (HG) levels, a common feature in diabetes, might also regulate the fate or functional behaviour of stem cells in visceral adipose tissue by activating downstream signals. The present study aimed to investigate whether HG might dictate the balance between self-renewal and differentiation of ASCs in the ‘niche’ and, if so, how HG acts. In particular, this study aimed to investigate the potential clinical application of ASCs in diabetic patients.

## Methods

**Patients and controls** A total of 25 type 2 diabetic patients and 15 non-diabetic participants who underwent abdominal surgery (gallbladder removal, in situ colorectal cancer) in our clinic were included. Non-diabetic participants were used as controls. Biometric data from the participants are reported in the electronic supplementary material (ESM) Table 1. Ethics approval was obtained from the Institutional Review Board of S. Giovanni Battista Hospital, Turin, Italy. Informed consent was obtained according to the Helsinki Declaration. We also declare that for the present study, we had no direct contact with the participants.

**Isolation of human visceral-derived ASCs** Human visceral adipose tissue samples were recovered from 20 type 2 diabetic patients (D-ASCs) and 10 non-diabetic controls (N-ASCs) and processed as detailed in the **ESM Methods**. To select the ASC subpopulation from the stromal vascular fraction, cells obtained from an additional five patients and five controls were subjected to CD31 and CD45 negative selection by magnetic cell sorting and characterised as detailed in the **ESM Methods** [29]. Alternatively, visceral N-ASCs were maintained in normal glucose (NG) DMEM (5 mmol/l D-glucose), HG DMEM (25 mmol/l D-glucose) or high mannitol (HM) DMEM (19 mmol/l D-mannitol, as osmotic control) and processed. Cell viability was evaluated at the end of each experiment using Trypan Blue or MTT assay.

**ASC proliferation** Cell proliferation was assayed at different time intervals by direct cell count or FACS analysis (FACS-Calibur flow cytometer; BD Biosciences, San Jose, CA, USA) using the proliferating cell nuclear antigen (PCNA) antibody [30]. Alternatively, cell proliferation was performed at day 16 after 48 h apocynin treatment or p47<sup>phox</sup> silencing on HG-cultured N-ASCs [31].

**Detection of apoptosis** Apoptosis was evaluated as indicated by a Cell Death Detection ELISA<sup>PLUS</sup> kit (Roche, Mannheim, Germany), following the manufacturer's instructions.

**Senescence assay** Senescence was evaluated by measuring the acidic  $\beta$ -galactosidase activity of N-ASCs that were differently cultured, as previously described [32].

**ASC characterisation** Freshly isolated ASCs, before or after magnetic cell-sorting selection, were characterised by FACS analysis.

**Endothelial and adipogenic ASC differentiation** Cell multipotency was assessed under appropriate culture conditions.

**RNA isolation and quantitative real-time PCR** Total RNA was isolated as previously described [31] from freshly or cultured ASCs, differently treated, and from 'spheroids', as indicated. *IL6*, *IL8*, *TNF $\alpha$*  (also known as *TNFA*) and *IL1 $\beta$*  (also known as *IL1B*) mRNA quantification was performed by quantitative real-time PCR (qRT-PCR).

**Western blot** Cells were lysed and 50  $\mu$ g of protein was subjected to SDS-PAGE, and processed as described [33].

**Endogenous p47<sup>phox</sup>, OCT4 and AKT silencing by small interfering RNA** To transiently knock down p47<sup>phox</sup>, OCT4 or AKT duplex small-interfering RNA (siRNA) (Qiagen, Valencia, CA, USA) was used.

**Sphere cluster (spheroid) formation** ASC de-differentiation status was evaluated by a sphere cluster formation and clonogenic assay. Sphere cluster formation was evaluated by three different operators counting 10 fields at  $\times 20$  magnification and reported as number  $\pm$  SD of spheres per field.

**ROS detection** Cellular ROS level detection on ASCs, cultured and treated as indicated, was performed by using DCF-DA (5-[and-6]-carboxy-2',7'-dichlorofluorescein diacetate, 0.5  $\mu$ mol/l final concentration) (Molecular Probes, Invitrogen, Carlsbad, CA, USA) assay.

**Mitochondria isolation and measurement of complex I–III activity** Cells incubated in the absence or presence of rotenone (50  $\mu$ mol/l for 18 h) were lysed and treated, as detailed

in the **ESM Methods**, to recover cytoplasmic mitochondria and measure complex I–III activity.

**Cytokine detection** Conditioned media from 14 days of NG or HG cultures were collected and processed to detect the levels of cytokines by a BioPlex Array Reader (Bio-Rad Laboratories, Hercules, CA, USA).


**Statistical analysis** Data are representative of at least three independent experiments, performed in triplicate, unless otherwise indicated. Densitometric analysis was used to calculate the differences in the fold induction of protein levels and normalised, with a statistically significant difference between experimental and control values indicated by  $p < 0.05$ . Western blot panels and relative densitometric histograms reported as 'relative units' in the figures are representative of the results obtained by experiments performed in triplicate ( $n = 3$ ,  $n = 4$  or  $n = 5$ ). The significance of differences between experimental and control values (indicated by  $p < 0.05$ ) was calculated using ANOVA with Newman–Keuls and Tukey multicomparison tests. ANOVA multicomparison testing was also used.

All of the methods described above are fully detailed in the **ESM Methods**. Reagents and antibodies are also reported in the **ESM Methods**.

## Results

**Characterisation of ASCs derived from diabetic patients** To investigate whether environmental cues such as hyperglycaemia can control the fate of visceral-derived ASCs, D-ASCs were compared with N-ASCs. ASC availability was first evaluated. An average (mean  $\pm$  SEM) of  $9.16 \times 10^4 \pm 9.2 \times 10^4$  and  $5.95 \times 10^4 \pm 7.5 \times 10^4$  ASCs per gram of visceral adipose tissue were isolated from patients and controls, respectively (Fig. 1a). However, as such a population also contains the stromal vascular fraction, to attain purified ASCs, cells recovered from additional patients and controls (five of each) were subjected to CD31 and CD45 negative selection by magnetic cell sorting [29] and counted. An average (mean  $\pm$  SEM) of  $4.58 \times 10^4 \pm 3.2 \times 10^4$  and  $1.9 \times 10^4 \pm 2.8 \times 10^4$  ASCs per gram of visceral adipose tissue were isolated from patients and controls, respectively (ESM Fig. 1a). Cell viability was evaluated by Trypan Blue staining and by MTT assay (data not shown).

Freshly isolated or purified ASCs were subsequently characterised for their expression profile. N-ASCs and D-ASCs, before or after magnetic cell sorting, expressed similar levels of mesenchymal cell markers, such as CD29 and CD44. By contrast, levels of the stemness markers CD105 and CD90 were higher in ASCs from diabetic patients compared with those from non-diabetic controls (Fig. 1b and ESM Fig. 1b).



**Fig. 1** Characterisation of ASCs derived from diabetic patients and non-diabetic controls. **(a)** The number of visceral ASCs (mean  $\pm$  SEM) per gram of fat tissue at isolation. Data refer to type 2 diabetic patients ( $n=20$  samples) or non-diabetic controls ( $n=10$ ).  $*p<0.05$ , D-ASCs vs N-ASCs. **(b)** Representative FACS analysis of visceral-derived D-ASCs and N-ASCs soon after isolation. Isotype controls were included (black peaks). The percentage of expression  $\pm$  SD of CD44, CD29, CD90 and CD105 markers was obtained by analysing 20 D-ASC and


10N-ASC samples (for N-ASCs: CD44,  $96\pm 2\%$ ; CD29,  $96\pm 2\%$ ; CD90,  $18\pm 1\%$ ; CD105,  $48\pm 3\%$ ; for D-ASCs: CD44,  $95\pm 3\%$ ; CD29,  $93\pm 1\%$ ; CD90,  $62\pm 2\%$ ; CD105,  $62\pm 4\%$ ). **(c)** Cell extracts from N-ASCs and D-ASCs were analysed by western blot for OCT4 (white bars) and NANOG (black bars) content by densitometry. Protein levels were normalised to tubulin content. The results are representative of three different experiments, each performed in triplicate ( $n=3$ ).  $***p<0.001$  vs equivalent group in N-ASCs

ASCs maintain their multipotency, as demonstrated by their ability to differentiate into mature endothelial cells and adipocytes [5, 34], as shown by the levels of specific differentiating markers (ESM Fig. 2). The finding that freshly recovered ASCs subjected to magnetic cell sorting did not produce KDR (kinase insert domain receptor, also known as vascular endothelial growth factor receptor 2 [VEGFR2]) (ESM Fig. 2b) further supports the conclusion that ASC purification was successfully performed. The observation that CD90 levels were higher in D-ASCs than in N-ASCs led us to investigate the content of OCT4 and NANOG, two crucial stemness regulators [13–15]. We found that OCT4 and NANOG production was upregulated in D-ASCs in comparison with N-ASCs (Fig. 1c). The result was further confirmed on the CD31<sup>-</sup>/CD45<sup>-</sup> ASC subpopulation (ESM Fig. 1c), suggesting that the number of de-differentiated ASCs is increased in diabetic patients.

**HG induces NOX-mediated ROS production** To investigate the involvement of ROS production in mediating ASC expansion in the diabetic condition, DCF-DA fluorescence assay was performed. As shown in Fig. 2a, ROS levels were

higher in freshly recovered D-ASCs than in N-ASCs. To further confirm these results, DCF-DA-fluorescence was also performed on the CD31<sup>-</sup>/CD45<sup>-</sup> ASC population. Again, ROS generation was higher in D-ASCs than in N-ASCs (ESM Fig. 1d). As hyperglycaemia is a common feature of diabetes, we hypothesised that ASC de-differentiation status could be reproduced in vitro by culturing N-ASCs in HG conditions. A total of 14 days of culture are required to select a CD31<sup>-</sup>/CD45<sup>-</sup> ASC population (ESM Fig. 3a) [6]. Thus, the impact of HG in mediating ROS generation was evaluated on freshly recovered ASCs cultured in HG-containing media for 14 days. The results reported in Fig. 2b, obtained by N-ASCs cultured with HG or NG, demonstrated that ROS production increased in response to HG, but not to NG. Hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>)-treated cells were used as controls. To investigate the involvement of NOX in ROS generation, HG or NG cultures were subjected to apocynin treatment. Apocynin treatment was able to prevent HG-generated ROS (Fig. 2c). As apocynin is known to act as a radical scavenger [35] and the assembly of p47<sup>phox</sup> and p67<sup>phox</sup> subunits is required for NOX enzymatic activity [36], N-ASCs endogenously depleted of p47<sup>phox</sup> (ESM Fig. 4a) were evaluated for ROS





**Fig. 2** NOX-mediated ROS production controls ASC proliferation. **(a)** DCF-DA assay was performed soon after isolation on ASCs freshly recovered from the visceral adipose tissue of type 2 diabetic patients (D-ASCs; black line) or non-diabetic controls (N-ASCs; grey line), to evaluate intracellular ROS levels. The intracellular DCF fluorescence levels, generated by the presence of ROS, varied linearly with increasing concentrations of oxidants and for this reason were used as a marker to quantify intracellular ROS production. The result is representative of three different experiments, each performed in triplicate ( $n=3$ ). **(b)** DCF-DA assay was performed on N-ASCs cultured for 14 days in the presence of NG (dark-grey line) or HG (black line), to evaluate ROS generation.  $H_2O_2$ -treated cells (light-grey line) were used as a positive control for ROS generation. HG-cultured N-ASCs were also treated with apocynin (light-grey line) or depleted of  $p47^{phox}$  by siRNA (grey line) and analysed for ROS generation. HG, black line.

generation. As shown in Fig. 2c, knock-down of  $p47^{phox}$  prevented ROS production in HG-cultured N-ASCs. To exclude the possibility that mitochondria could contribute to ROS production, the mitochondrial complex I–III inhibitor rotenone [37] was added and ROS generation was evaluated. As shown in ESM Fig. 3b, HG-dependent ROS production was not affected by rotenone treatment. These data were confirmed by measuring complex I–III activity (ESM Fig. 3c). Cell viability was evaluated by MTT assay (data not shown).


*NOX-generated ROS are crucial for HG-mediated ASC proliferation* To evaluate whether increased ROS levels translate into increased cell proliferation, NG- or HG-cultured

N-ASCs were analysed for their proliferative capability. Data reported in Fig. 2d, obtained by directly counting cells or analysing the abundance of a DNA-polymerase- $\delta$  auxiliary protein such as PCNA (ESM Fig. 5a), revealed the rate of cell proliferation was higher in HG-cultured N-ASCs than in those that were cultured with NG. HM (19 mmol/l), used as osmotic control, had no effect (Fig. 2d and ESM Fig. 5a). MTT assay as well as the cell death detection ELISA assay, used as an index of DNA fragmentation (ESM Fig. 6), revealed that non-toxic ROS levels were generated in response to HG treatment. To investigate the role of ROS in HG-mediated N-ASC proliferation, HG-cultured N-ASCs were treated with apocynin for 48 h and evaluated for their ability to undergo cell

proliferation. As reported in Fig. 2e and ESM Fig. 5b, the addition of apocynin led to a reduction of N-ASC proliferation. The inhibitory effect exerted by apocynin was further confirmed by analysing the content of cyclin D1 (a member of the cyclin protein family that drives the G1/S phase transition) and p21<sup>waf</sup> (a cyclin-dependent kinase inhibitor acting as a negative regulator of cell cycle progression at G1) (Fig. 2f). Similar results were obtained in N-ASCs silenced for p47<sup>phox</sup> (Fig. 2e, f and ESM Fig. 5b). The finding that, when depleted of their signalling machinery, ASCs underwent cell senescence could provide an explanation for the reduced numbers of cells detected in these experimental conditions (ESM Fig. 5e). Conversely, both apocynin treatment and p47<sup>phox</sup> depletion did not affect N-ASC proliferation when cultured in the presence of NG (ESM Fig. 5c and 5d).


*HG drives OCT4 and NANOG expression by NOX-mediated ROS generation* FACS analysis was performed to

characterise HG- and NG-cultured N-ASCs. The data reported in Fig. 3a demonstrate that while N-ASCs cultured with HG for 14 days still contained CD90, those cultured with NG retained only mesenchymal cell markers (CD105<sup>+</sup>, CD29<sup>+</sup> and CD44<sup>+</sup>). Similarly to D-ASCs, HG-cultured N-ASCs produced OCT4 and NANOG (Fig. 3b). Undifferentiated adult stem cells, unlike committed cells, grow in suspension and form sphere clusters, in an anchorage-independent fashion [38]. Consistent with OCT4/NANOG content, only HG-cultured N-ASCs were able to form ‘spheroids’ (Fig. 3c). To investigate the role of ROS in this process, HG-cultured N-ASCs were subjected to apocynin or depleted of p47<sup>phox</sup> and evaluated. Data reported in Fig. 4a demonstrate that NOX-mediated ROS generation is essential for spheroid formation. Moreover, the finding that both approaches were able to decrease OCT4/NANOG content (Fig. 4b) further sustains the conclusion that ROS generation is crucial to promote ASC de-differentiation.



**Fig. 3** HG induces OCT4 and NANOG production, as well as sphere cluster formation. **(a)** Representative FACS analysis of N-ASCs cultured for 14 days in NG or HG medium. Isotype controls were included (black peaks). The percentage of expression $\pm$ SD of CD44, CD29, CD90 and CD105 markers was obtained by analysing 10 different samples (NG-cultured N-ASCs: CD44, 96 $\pm$ 2; CD29, 91 $\pm$ 2; CD90, 19 $\pm$ 2; CD105, 30 $\pm$ 2; HG-cultured N-ASCs: CD44, 95 $\pm$ 3; CD29, 94 $\pm$ 1; CD90, 79 $\pm$ 3; CD105, 76 $\pm$ 3). **(b)** Cell extracts from N-ASCs culture for 14 days with NG or HG were analysed by western blot for OCT4 and NANOG (grey bars) content by densitometry. Protein levels were normalised to tubulin content. \*\*\* $p$ <0.001. **(c)** Sphere cluster formation was evaluated 72 h after seeding N-ASCs cultured for 14 days in NG or HG in semi-solid medium, as described in the ESM Methods.

Black bar, P1 HG; grey bar, P1 NG; white bar, P2 HG. \*\* $p$ <0.01, P1 NG vs P1 HG. The clonogenic potential of HG-cultured N-ASCs was investigated by plating single cell suspension derived from P1 spheres for an additional 48 h (P2 HG). †† $p$ <0.01, P1 NG vs P2 HG. Sphere cluster formation was evaluated by three different operators (each experiment was performed in triplicate), counting 10 fields at  $\times 20$  magnification and reporting as number $\pm$ SD of spheres per field in the histogram. Scale bars indicate 50  $\mu$ m in P1 NG and P1 HG ( $\times 20$  magnification) and 20  $\mu$ m in P2 HG ( $\times 40$  magnification). The results are representative of four individual experiments performed in triplicate ( $n=4$ ). P1 corresponds to sphere cluster formation; P2 corresponds to sphere cluster formation of P1-derived cells




**Fig. 4** NOX-mediated ROS generation in response to HG is crucial for both sphere cluster formation and OCT4/NANOG production. **(a)** N-ASCs, subjected to the indicated treatments, were plated in semi-solid medium, as described in the *ESM Methods*, to assess sphere cluster formation. The histogram reports the number $\pm$ SD of spheres per field, counting 10 fields at  $\times 20$  magnification. Black bar, HG; dark-grey bar, HG+apocynin; light-grey bar, HG+p47<sup>phox</sup> siRNA; white bar, HG+OCT4 siRNA. Scale bars indicate 50  $\mu$ m. **\*\*** $p < 0.01$ . The results are representative of four different experiments performed in triplicate ( $n = 4$ ). **(b)** HG-cultured N-ASCs in the presence or absence of apocynin or depleted of p47<sup>phox</sup> were lysed and subjected to SDS-PAGE. White bars, OCT4; black bars, NANOG. **\*\*** $p < 0.01$ . The filters were immunoblotted with anti-OCT4, anti-NANOG and anti-tubulin antibodies. **(c)** Cell extracts from HG-cultured N-ASCs, depleted or not of OCT4, were analysed by western blot for cyclin D1 (white bars) and p21<sup>waf</sup> (black bars) content. Protein levels were normalised to  $\beta$ -actin content. **\*\*\*** $p < 0.001$ , HG+OCT4 siRNA-treated cells vs HG for cyclin D1 and p21<sup>waf</sup> content. The results are representative of three different experiments performed in triplicate ( $n = 3$ )

**OCT4 is crucial for HG-mediated ASC de-differentiation** ASCs overexpressing OCT4 have been shown to undergo cell reprogramming [23]. Thus, the involvement of OCT4 in controlling HG-mediated ASC de-differentiation was further investigated. As reported in Fig. 4a, knock-down of OCT4 (*ESM Fig. 4b*) prevented HG-induced sphere cluster formation. Consistently, decreased expression of cyclin D1 and increased expression of p21<sup>waf</sup> was detected (*Fig. 4c*) in OCT4-silenced cells.

AKT activation is crucial for OCT4 and NANOG expression and for spheroid formation in response to HG. AKT is crucial for OCT4-mediated de-differentiation [23], thus, the role of AKT in HG-mediated ASC reprogramming was investigated. As shown in Fig. 5a, AKT production could be detected in HG-cultured N-ASCs, but not in NG-cultured cells. To gain further insight into the role of AKT in regulating this process, functional studies were performed in AKT-silenced N-ASCs (*ESM Fig. 4c*). Data in Fig. 5b, c show that, as in apocynin-treated and p47<sup>phox</sup>-silenced N-ASCs, OCT4/NANOG content was reduced and spheroid formation prevented in N-ASCs endogenously depleted of AKT. Again, the content of cyclin D1 and p21<sup>waf</sup> validated these data (*Fig. 5d*). To further confirm the role of ROS in controlling AKT activation, N-ASCs were cultured in the presence of apocynin or depleted of p47<sup>phox</sup>. As reported in Fig. 5a in both culture conditions we failed to detect AKT activation. The finding that ROS generation was still detectable in AKT-depleted cells (*Fig. 5e*) indicates that ROS generation occurs upstream to AKT.

**ROS production and AKT activation are required for IL-6 and IL-8 production by spheroids** The role of ROS in mediating ASC paracrine action has been reported [39]. Therefore, the contribution of ROS in mediating HG-induced soluble factor secretion was investigated. BioPlex analysis (on supernatant fractions) and qRT-PCR (on spheroids) were performed. As reported in Fig. 6, the amount of IL-6 and IL-8 produced by N-ASCs cultured in the presence of HG (*Fig. 6a*) and spheroids (*Fig. 6c,e*) was higher than in NG-cultured cells. Next, N-ASC cytokine production was analysed in p47<sup>phox</sup> and AKT-silenced cells. Data shown in Fig. 6 clearly demonstrate that NOX-generated ROS and AKT activation are required for IL-6 and IL-8 production in response to HG only, indicating that different mechanisms account for IL-6 and IL-8 production in NG-cultured N-ASCs (*ESM Fig. 7a*). Moreover, the finding that preventing spheroid formation by silencing p47<sup>phox</sup> and AKT led to reduced cytokine production sustains the possibility that N-ASC paracrine function is enhanced by their de-differentiation status. Consistent with this possibility, we found that the release of IL-6 and IL-8 was significantly reduced when HG-containing media were removed and replaced with NG (*Fig. 6g*). Finally, we failed to detect TNF $\alpha$  and IL-1 $\beta$  in our samples (*Fig. 6b,d,f*).

**ROS generation and AKT activation are required for D-ASC cytokine production** An increased level of proinflammatory cytokines is a common feature in diabetes [40]. Thus, we investigated whether D-ASC de-differentiation status might contribute to this proinflammatory profile. Soon after isolation, both freshly recovered and CD31<sup>-</sup>/CD45<sup>-</sup> ASCs were evaluated for their ability to produce IL-6 and



**Fig. 5** ROS-mediated AKT activation is involved in sphere cluster formation and OCT4/NANOG production in response to HG. **(a)** N-ASCs cultured in different experimental conditions, as indicated, were lysed and analysed by western blot for pAKT (black bars) and normalised to total AKT content. **(b)** Cell lysates from HG-cultured N-ASCs endogenously depleted of AKT were subjected to SDS-PAGE. The filters were immunoblotted with anti-OCT4 and anti-NANOG antibodies. Protein levels were normalised to tubulin content. White bars, OCT4; black bars, NANOG. **(c)** Sphere cluster formation assay was performed by seeding HG-cultured N-ASCs untreated or silenced for


AKT in semi-solid medium. Histogram reports the number $\pm$ SD of spheres per field, counting 10 fields at  $\times 20$  magnification. Scale bars indicate 50  $\mu$ m.  $**p < 0.01$ , HG+AKT siRNA (black bar) vs HG (white bar). **(d)** N-ASCs cultured and treated as indicated above were analysed by western blot for cyclin D1 (white bars), p21<sup>waf</sup> (black bars) and  $\beta$ -actin content. **(e)** DCF-DA assay was performed on HG-cultured N-ASCs, silenced or not for AKT, to evaluate cellular ROS levels. The results are representative of three different experiments, each performed in triplicate ( $n=3$ ).  $***p < 0.001$

IL-8 by qRT-PCR. As reported in Fig. 7 and ESM Fig. 7b, IL-6 and IL-8 production was higher in D-ASCs than in N-ASCs in both conditions. To gain further insight into the mechanisms regulating proinflammatory cytokine production and to avoid long-term culture conditions, apocynin and the specific AKT inhibitor TCL1<sub>10–24</sub> were used instead of siRNA technology. Indeed, by applying these approaches, we demonstrated that ROS generation and AKT activation are essential for IL-6 and IL-8 production (Fig. 7 and ESM Fig. 7b). Again, we failed to detect either TNF $\alpha$  or IL-1 $\beta$ .

## Discussion

White adipose tissue, owing to the easy access by minimally invasive methods, is currently considered an alternative and valuable source of ASCs for tissue regeneration [5, 7, 41]. However, no data have been provided on how mechanistically the balance between self-renewal and differentiation in the ‘niche’ is regulated in pathological conditions. Likewise, studies evaluating the potential impact of ASCs in clinical settings where tissue regeneration is required are still unavailable.






**Fig. 6** ROS generation and AKT activation are required for IL-6 and IL-8 production. **(a, b)** BioPlex analysis was performed, according to the manufacturer’s instructions, to detect IL-6, IL-8 **(a)** and TNFα and IL-1β **(b)** levels in sixfold-concentrated supernatant fractions, recovered from the indicated cultures. White bars, NG; light-grey bars, HG; grey bars, HG+p47<sup>phox</sup> siRNA; dark-grey bars, HG+AKT siRNA; black bars, positive control. IL-6 or IL-8: \*\*\**p*<0.001, HG-cultured N-ASCs vs NG-cultured N-ASCs, †††*p*<0.001, HG+p47<sup>phox</sup> siRNA and HG+AKT siRNA vs HG. TNFα and IL-1β: \*\*\**p*<0.001, experimental vs control value. **(c–f)** *IL6* **(c)**, *IL1β* **(d)**, *IL8* **(e)** and *TNFα* **(f)** mRNA quantification was performed by qRT-PCR on N-ASCs pre-treated as indicated and cultured in semi-solid medium (see sphere cluster formation assay in the **ESM Methods**). Fold expression of the indicated cytokines, normalised to *GAPDH*, is representative of ten samples. \*\*\**p*<0.001, HG-cultured N-ASCs vs NG-cultured N-ASCs; ††*p*<0.01, HG+p47<sup>phox</sup> siRNA and HG+AKT siRNA vs HG for *IL6*; \*\**p*<0.01, HG-cultured N-ASCs vs NG-cultured N-ASCs; †*p*<0.05, HG+p47<sup>phox</sup> siRNA and HG+AKT siRNA vs HG for *IL8*; \*\*\**p*<0.001, experimental vs control value for *TNFα* and *IL1β*. **(g)** BioPlex analysis was also performed to detect IL-6 and IL-8 levels on sixfold concentrated supernatant fractions, recovered from 14 days’ NG (dark-grey bars) and HG (white grey bars) cultures. Alternatively, HG medium was replaced for 48 h with NG medium (light-grey bars) and then collected to detect IL-6 and IL-8 (\*\*\**p*<0.001, NG and HG replaced with NG vs HG-cultured N-ASCs). Black bars, positive control. In **(a–d)**, supernatant fractions or mRNA obtained from activated macrophages were used as positive controls (+). The results in **a** and **d** are representative of three different experiments, each performed in triplicate (*n*=3)

knock-down, used to extend our knowledge on D-ASC stemness behaviour, further supports the role of OCT4 in promoting ASC de-differentiation. This suggests that the increased number of ASCs recovered from diabetic patients may be the result of HG concentrations in the adipose ‘stem cell niche’.

The ‘stem cell niche’ is a specialised microenvironment containing stem cells [42]. As in developmental processes, in adult tissues, the balance between self-renewal and differentiation of stem cells is strictly controlled by environmental cues [13–18, 21, 22]. In this study, we have demonstrated that hyperglycaemic conditions induce and maintain ASC pluripotency. Pluripotency factors, including stemness genes, provide fundamental mechanisms underlying the properties of stem cells [22]. OCT4, SOX2 and NANOG are known regulators of genes encoding molecules involved in the control of stem cell behaviours, including proliferation and differentiation [18, 21, 22]. We provide evidence that in D-ASCs, *OCT4* and *NANOG* expression is upregulated compared with N-ASCs. Consistently, we found that HG-induced *OCT4* and *NANOG* expression is associated with an increased rate of cell proliferation and with their de-differentiation status. siRNA-mediated gene



**Fig. 7** *IL6* and *IL8* expression by CD31<sup>-</sup>/CD45<sup>-</sup> selected ASCs. *IL6*, *IL8*, *TNFα* and *IL1β* mRNA quantification was performed by qRT-PCR on CD31<sup>-</sup>/CD45<sup>-</sup> N-ASCs (grey bars) and D-ASCs (striped bars). CD31<sup>-</sup>/CD45<sup>-</sup> D-ASCs were also treated for 6 h with apocynin (dotted bars) or with the AKT inhibitor TCL1<sub>10-24</sub> (black bars). Fold expression of the indicated cytokines, normalised to *GAPDH*, is representative of five samples tested. \*\*\**p*<0.001, D-ASCs vs N-ASCs; †††*p*<0.001, D-ASCs+apocynin and D-ASCs+TCL1<sub>10-24</sub> vs D-ASCs for *IL6*; \*\*\**p*<0.01, N-ASCs, D-ASCs+apocynin and D-ASCs+TCL1<sub>10-24</sub> vs D-ASCs for *IL8*; \*\*\**p*<0.001, experimental vs control value for *TNFα* and *IL1β*. mRNA from activated macrophages was used as positive control (white bars, +). The results are representative of five different experiments, each performed in triplicate (*n*=5)

Although cellular accumulation of ROS has been considered a crucial driver of cell death in different pathological settings [43], evidence has been provided that the reduction–oxidation balance plays a far more complex role in stem cell biology [26, 27, 44]. Our data demonstrated that HG challenge induces ‘non-toxic’ levels of ROS in ASCs. ROS generation mainly relies on NOX enzymatic activity [28], which in turn depends on its regulatory subunits, p47<sup>phox</sup> and p67<sup>phox</sup> [28]. The finding that p47<sup>phox</sup> knock-down prevented both ROS generation and ASC proliferation indicates that NOX-mediated ROS generation in response to HG acts as second messenger to drive ASC proliferation. Moreover, the observation that interfering with the mitochondrial electron transport system did not affect ROS production suggests that ROS generation in our experimental conditions mainly depends on NOX activity. We could not directly demonstrate the involvement of NOX in mediating ROS production in freshly isolated ASCs (to obtain p47<sup>phox</sup> endogenous depletion would require long-term culture conditions). However, the percentage of HbA<sub>1c</sub> detected in our patients leads us to speculate that, as in vitro, in D-ASC NOX might be crucial in driving ROS generation, eventually leading to the increased de-differentiation status.

To increase stem cell proliferation and self-renewal potential, physiological levels of ROS are required [25, 26, 45]. Consistently, data provided by our study (demonstrating that by silencing p47<sup>phox</sup>, ASCs failed to form spheroids and did not express *OCT4*) indicate that NOX-generated ROS are also crucial for HG-driven ASC de-differentiation. Moreover, the possibility that a decrease in ROS cellular levels has a negative impact on ASC self-renewal is supported by the finding that NG-cultured N-ASCs were unable to form spheroids.

Beside phagocytes, other cell types use NOX-generated ROS as a second messenger [46]. The activation of the MAPK and PI3K/AKT signalling pathways has been reported to be the result of NOX-generated ROS [27]. Le Belle et al have demonstrated that in neural stem cell populations, self-renewal is associated with the ROS-mediated PI3K/AKT signalling pathway [26]. In addition, it has been shown that ASC reprogramming induced by *OCT4* over-expression engages the PI3K/AKT and JAK/STAT3 pathways [23]. By applying siRNA technology to knock-down NOX and AKT, we have demonstrated that AKT activation is also essential for HG-mediated *OCT4*/*NANOG* production and for spheroid formation. Moreover, data provided by knock-down AKT demonstrate that AKT is downstream to ROS and upstream to *OCT4*. Again, further support for the biological relevance of the AKT signalling pathway in driving HG-mediated ASC de-differentiation is provided by data obtained from NG-cultured N-ASCs.

ASCs, besides possessing the ability to regenerate damaged tissue, also secrete soluble mediators [12, 39, 47]. A great deal of interest in adipose tissue has been generated by

the finding that ASCs, by means of their paracrine function, influence the inflammatory response [39, 47]. We demonstrate that microenvironmental cues, by inducing ASC de-differentiation, influence ASC functional behaviour, resulting in increased proinflammatory cytokine production. This suggests that ASCs, exposed to hyperglycaemic conditions via NOX-generated ROS and AKT activation, might actively participate in dysfunctional proinflammatory cytokine production. Epidemiological studies have described a clear link between the C-reactive protein-mediated inflammatory response, its inducer IL-6 and an increased cardiovascular risk profile [48]. Interventions aimed at ameliorating the metabolic profile lead to a reduction in proinflammatory protein plasma levels, including that of IL-6 [49]. Consistently, our results show that restoring glucose concentrations leads to decreased IL-6 and IL-8 production. Therefore, as particular attention has been dedicated to the detrimental consequences of hyperglycaemia, our data suggest that targeting hyperglycaemia could also represent a rational approach to interfering with the ASC dysfunctional cytokine profile.

Adipose tissue transplantation has been proposed as a potential tool to provide favourable metabolic effects in diabetes [3]. Data provided herein open new perspectives for the clinical application of visceral ASCs in diabetic patients. The potential clinical impact is supported by the following observations. First, the number of ASCs recovered from the visceral adipose tissue of diabetic patients is increased compared with that of non-diabetic controls. Second, ASCs obtained from these patients have high content of *OCT4*/*NANOG*, and thus possess a wider therapeutic potential. Third, visceral adipose tissue could be an ideal source of autologous ASCs obtainable with minimally invasive procedures. Finally, consistent with data provided by omentectomy in humans [50], the removal of visceral adipose tissue, besides providing favourable metabolic effects, might also offer an additional advantage such as the possibility of exploiting ASCs to repair failing organs. However, we cannot exclude that as in hematopoietic systems, where a high ROS level is associated with a high proliferation rate and premature exhaustion of the self-renewal potential [25], HG concentrations in the visceral adipose tissue of diabetic patients could lead to a rapid exhaustion of ASC stemness capability, eventually translating in an accelerated impairment of the regenerative potential. Further studies are required to resolve this issue.

Data have indicated that multipotent stem cells can be recovered from the adipose tissue of adults [8]. However, how metabolic conditions can modulate the balance between self-renewal and differentiation in the ‘niche’ is still unclear. In this study, we provide evidence that environmental cues impact ASC biological properties and significantly affect their stemness and secretion profile. Moreover, these results

provide the rationale to develop therapeutic strategies exploiting adipose tissue as an easily accessible source of MSCs to repair failing organs in diabetic patients.

**Acknowledgements** We thank Prof. D. Ghigo (Department of Oncology, University of Torino, Torino, Italy) for his helpful advice.

**Funding** This work was supported by grants from the Italian Association for Cancer Research (AIRC), Ricerca Finalizzata Regione Piemonte and MIUR (Ministero dell'Università e Ricerca Scientifica, cofinanziamento MURST) to MFB.

**Duality of interest** The authors declare that there is no duality of interest associated with this manuscript.

**Contribution statement** PD, CB, GT and AT contributed to the study design, analysis and interpretation of data and to the critical revision of the manuscript. CO, MG and CR contributed to the analysis and interpretation of data, as well as to the revision of the manuscript. MT contributed to the acquisition of data and critical revision of the manuscript. MFB contributed to the study conception and design, and in writing the manuscript. All authors approved the version to be published.

## References

- Schaffler A, Buchler C (2007) Concise review: adipose tissue-derived stromal cells—basic and clinical implications for novel cell-based therapies. *Stem Cells* 25:818–827
- Le Blanc K, Ringdén O (2007) Immunomodulation by mesenchymal stem cells and clinical experience. *J Intern Med* 262:509–525
- Tran TT, Kahn CR (2010) Transplantation of adipose tissue and stem cells: role in metabolism and disease. *Nat Rev Endocrinol* 6:195–213
- Rubina K, Kalinina N, Efimenko A et al (2009) Adipose stromal cells stimulate angiogenesis via promoting progenitor cell differentiation, secretion of angiogenic factors, and enhancing vessel maturation. *Tissue Eng Part A* 15:2039–2050
- Zuk PA, Zhu M, Ashjian P et al (2002) Human adipose tissue is a source of multipotent stem cells. *Mol Biol Cell* 13:4279–4295
- Baglioni S, Francalanci M, Squecco R et al (2009) Characterization of human adult stem-cell populations isolated from visceral and subcutaneous adipose tissue. *FASEB J* 23:3494–3505
- Gimble JM, Katz JA, Bunnell BA (2007) Adipose derived stem cells for regenerative medicine. *Circ Res* 100:1249–1260
- Locke M, Feisst V, Dunbar PR (2011) Concise review: human adipose-derived stem cells: separating promise from clinical need. *Stem Cells* 29:404–411
- Yarak S, Okamoto OK (2010) Human adipose-derived stem cells: current challenges and clinical perspectives. *An Bras Dermatol* 85:647–656
- Roldan M, Macias-Gonzalez M, Garcia R, Tinahones FJ, Martin M (2011) Obesity short-circuits stemness gene network in human adipose multipotent stem cells. *FASEB J* 25:4111–4126
- Schraufstatter IU, Discipio RG, Khaldoyanidi S (2011) Mesenchymal stem cells and their microenvironment. *Front Biosci* 17:2271–2288
- Horwitz EM, Dominici M (2008) How do mesenchymal stromal cells exert their therapeutic benefit? *Cytotherapy* 10:771–774
- Ng HH, Surani MA (2011) The transcriptional and signalling networks of pluripotency. *Nat Cell Biol* 13:490–496
- Kuroda T, Tada M, Kubota H et al (2005) Octamer and Sox elements are required for transcriptional cis regulation of NANOG gene expression. *Mol Cell Biol* 25:2475–2485
- Loh YH, Wu Q, Chew JL et al (2006) The OCT4 and NANOG transcription network regulates pluripotency in mouse embryonic stem cells. *Nat Genet* 38:431–440
- Silva J, Chambers I, Pollard S, Smith A (2006) NANOG promotes transfer of pluripotency after cell fusion. *Nature* 441:997–1001
- Lengner CJ, Welstead GG, Jaenisch R (2008) The pluripotency regulator OCT4: a role in somatic stem cells? *Cell Cycle* 7:725–728
- Park IH, Zhao R, West JA et al (2008) Reprogramming of human somatic cells to pluripotency with defined factors. *Nature* 451:141–146
- Santourlidis S, Wernet P, Ghanjati F et al (2011) Unrestricted somatic stem cells (USSC) from human umbilical cord blood display uncommitted epigenetic signatures of the major stem cell pluripotency genes. *Stem Cell Res* 6:60–69
- Freberg CT, Dahl JA, Timoskainen S, Collas P (2007) Epigenetic reprogramming of OCT4 and NANOG regulatory regions by embryonal carcinoma cell extract. *Mol Biol Cell* 18:1543–1553
- Goodell MA (2003) Stem-cell “plasticity”: befuddled by the muddle. *Curr Opin Hematol* 10:208–213
- Sanges D, Cosma MP (2010) Reprogramming cell fate to pluripotency: the decision-making signalling pathways. *Int J Dev Biol* 54:1575–1587
- Kim JH, Jee MK, Lee SY et al (2009) Regulation of adipose tissue stromal cells behaviors by endogenous OCT4 expression control. *PLoS One* 4:e7166
- Lambeth JD, Krause KH, Clark RA (2008) NOX enzymes as novel targets for drug development. *Semin Immunopathol* 30:339–363
- Jang YY, Sharkis SJ (2007) A low level of reactive oxygen species selects for primitive hematopoietic stem cells that may reside in the low-oxygenic niche. *Blood* 110:3056–3063
- Le Belle JE, Orozco NM, Paucar AA et al (2011) Proliferative neural stem cells have high endogenous ROS levels that regulate self-renewal and neurogenesis in a PI3K/AKT-dependant manner. *Cell Stem Cell* 8:59–71
- Kim JH, Park SH, Park SG, Choi JS, Xia Y, Sung JH (2011) The pivotal role of reactive oxygen species generation in the hypoxia-induced stimulation of adipose-derived stem cells. *Stem Cells Dev* 20:1753–1761
- Takechi T, Yabe-Nishimura (2008) NOX enzymes and diabetic complications. *Semin Immunopathol C* 30:301–314
- Zhang P, Moudgill N, Hager E et al (2011) Endothelial differentiation of adipose-derived stem cells from elderly patients with cardiovascular disease. *Stem Cells Dev* 20:977–988
- Dentelli P, Rosso A, Olgasi C, Camussi G, Brizzi MF (2011) IL-3 is a novel target to interfere with tumor vasculature. *Oncogene* 30:4930–4940
- Togliatto G, Trombetta A, Dentelli P, Rosso A, Brizzi MF (2011) MIR221/MIR222-driven post-transcriptional regulation of P27KIP1 and P57KIP2 is crucial for high-glucose- and AGE-mediated vascular cell damage. *Diabetologia* 54:1930–1940
- Togliatto G, Trombetta A, Dentelli P et al (2010) Unacylated ghrelin rescues endothelial progenitor cell function in individuals with type 2 diabetes. *Diabetes* 59:1016–1025
- Dentelli P, Trombetta A, Togliatto G et al (2009) Formation of STAT5/PPARgamma transcriptional complex modulates angiogenic cell bioavailability in diabetes. *Arterioscler Thromb Vasc Biol* 29:114–120
- Liu TM, Martina M, Huttmacher DW, Hui JH, Lee EH, Lim B (2007) Identification of common pathways mediating

- differentiation of bone marrow- and adipose tissue-derived human mesenchymal stem cells into three mesenchymal lineages. *Stem Cells* 25:750–760
35. Heumüller S, Sven W, Barbosa-Sicard E et al (2008) Apocynin is not an inhibitor of vascular NADPH oxidases but an antioxidant. *Hypertension* 51:211–217
  36. Babior BM (2004) NADPH oxidase. *Curr Opin Immunol* 16:42–47
  37. Brand MD, Affourtit C, Esteves TC et al (2004) Mitochondrial superoxide: production, biological effects, and activation of uncoupling proteins. *Free Radic Biol Med* 37:755–767
  38. Reynolds BA, Weiss S (1996) Clonal and population analyses demonstrate that an EGF-responsive mammalian embryonic CNS precursor is a stem cell. *Dev Biol* 175:1–13
  39. Carrière A, Ebrahimian TG, Dehez S et al (2009) Preconditioning by mitochondrial reactive oxygen species improves the proangiogenic potential of adipose-derived cells-based therapy. *Arterioscler Thromb Vasc Biol* 29:1093–1099
  40. Calle MC, Fernandez ML (2012) Inflammation and type 2 diabetes. *Diabetes Metab* 38:183–191
  41. Maumus M, Peyrafitte JA, D'Angelo R et al (2011) Native human adipose stromal cells: localization, morphology and phenotype. *Int J Obes (Lond)* 35:1141–1153
  42. Li L, Xie T (2005) Stem cell niche: structure and function. *Annu Rev Cell Dev Biol* 21:605–631
  43. Simon HU, Haj-Yehia A, Levi-Schaffer F (2000) Role of reactive oxygen species (ROS) in apoptosis induction. *Apoptosis* 5:415–418
  44. Park SG, Kim JH, Xia Y, Sung JH (2011) Generation of reactive oxygen species in adipose-derived stem cells: friend or foe? *Expert Opin Ther Targets* 15:1297–1306
  45. Noble M, Pröschel C, Mayer-Pröschel M (2011) Oxidative-reductionist approaches to stem and progenitor cell function. *Cell Stem Cell* 8:1–2
  46. Garrido AM, Griendling KK (2009) NADPH oxidases and angiotensin II receptor signaling. *Mol Cell Endocrinol* 302:148–158
  47. Waki H, Tontonoz P (2007) Endocrine functions of adipose tissue. *Annu Rev Pathol* 2:31–56
  48. Pradhan AD, Manson JE, Rifai N et al (2001) C-reactive protein, interleukin 6, and risk of developing type 2 diabetes mellitus. *JAMA* 286:327–334
  49. Fisher G, Hyatt TC, Hunter GR et al (2012) Markers of inflammation and fat distribution following weight loss in African-American and white women. *Obesity (Silver Spring)* 20:715–720
  50. Thörne A, Lönnqvist F, Apelman J, Hellers G, Arner P (2002) A pilot study of long-term effects of a novel obesity treatment: omentectomy in connection with adjustable gastric banding. *Int J Obes Relat Metab Disord* 26:193–199