Sick of sitting

James A. Levine¹

Received: 25 November 2014 / Accepted: 12 March 2015 / Published online: 24 May 2015
© Springer-Verlag Berlin Heidelberg 2015

Abstract Sitting too much kills. Epidemiological, physiological and molecular data suggest that sedentary lifestyle can explain, in part, how modernity is associated with obesity, more than 30 chronic diseases and conditions and high healthcare costs. Excessive sitting—sitting disease—is not innate to the human condition. People were designed to be bipedal and, before the industrial revolution, people moved substantially more throughout the day than they do presently. It is encouraging that solutions exist to reverse sitting disease. Work environments, schools, communities and cities can be re-imagined and re-invented as walking spaces, and people thereby offered more active, happier, healthier and more productive lives.

Keywords Chronic disease · Diabetes · Non-exercise activity thermogenesis · Obesity · Physical activity · Review

Abbreviations

NEAT Non-exercise activity thermogenesis

In the beginning, we were bipedal

There is debate regarding the evolutionary steps that resulted in bipedalism [1]; suffice it to say, the human evolved over hundreds of thousands of years to be upright, two-legged, walking beings [2–6]. Over time, people evolved to explore by foot, to manually invent tools and weapons and to think while upright and responding to environmental cues,

perceived threats and calculated opportunities [7]. Thousands of years ago, living was dynamic. Compare chasing a bison over a cliff to choosing a meat package at the supermarket and contrast hand-chipping a flint for a spear versus engineering a cyber attack. Prior to the industrial revolution 200 years ago, 90% of the world's population lived in agricultural communities where shelter, nutrition and reproduction all required physical exertion. Data from agricultural communities suggest that, prior to the industrial revolution, people sat for 300 min per day and lived actively [8]. From 1760 onward, the industrial revolution precipitated urbanisation; it was the predominant demographic shift into modern history [9–11]. Now more than half the world's population live in cities and urbanisation continues to grow worldwide [12]. In industrialising countries, 1908 saw the introduction of factories that used conveyor belts, and in the 1940s, modern chair-based offices were developed. In both cases, the environments and furnishings were designed to promote productivity and limit movement by having people sit. Walking around factories or offices was perceived as wasted time. Fast-forward to the present day, and office workers can sit for up to 15 h in a single day! [13].

People are designed to work and socialise while on their legs and to sit in order to rest; the default position for people is to be up and moving. Is it a surprise that modern people who default to sitting (e.g. 'take a seat') experience negative physical, medical and psychological consequences? Do modern environments, however, give us any other choice except to sit?

Modern environments foster sitting

Environments foster sedentariness in many ways and at multiple societal levels. On an individual level, a person is confronted with environmental cues to sit throughout the day, e.g. drive-through banks [14]. Modern homes contain many tools of convenience, each one decreasing non-

✉ James A. Levine
levine.james@mayo.edu

¹ Mayo Clinic, 13400 East Shea Blvd, Scottsdale, AZ 85259, USA

exercise activity thermogenesis (NEAT) [15]. Workplaces [16, 17] and schools [18–20] are major environmental drivers of NEAT. Communities encourage inhabitants to be sedentary. For instance, when data for 3,139 US counties were reviewed, encompassing the majority of the US population, sedentariness correlated with poverty, as did obesity and diabetes [21]. No personal choices, behaviour or genetics can explain this association because people do not choose to live in poverty. Other factors that may explain why people living in poverty-dense areas are likely to be sedentary include crime, poor housing conditions, poor health and lack of access to activity-promoting opportunities.

Cityscapes and sitting

More than half the world's population live in cities, with the fastest growth occurring in some of the world's poorest countries [22]. Active leg-based transportation is common in a number of European cities, including Copenhagen and Amsterdam. Rain or shine, Amsterdam is swamped with bicycles [23]. In the USA, there is variability in how cityscapes encourage or inhibit walking. New York City, San Francisco, Boston, Philadelphia and Miami are among the most walkable [24, 25], suggesting that weather is not a key determinant. Los Angeles, on the other hand, exemplifies a city that inhibits walking because of poor air quality and a roadway design that imposes the use of motor transport [26, 27]. Los Angeles has the most congested streets in the USA. The average Los Angeles driver experiences 72 h of traffic delays per year. Overall, US drivers waste 3 billion gallons of gas per year and 6 billion hours sitting in traffic [28]. Cityscapes that promote sedentariness are unhealthy in other ways, too.

Public policy [29] and prevailing social ambience can impact sitting [30–32]. In many societies, gender- and age-defined roles impact sedentariness [32–34]. Stereotypes about obesity can perpetuate low NEAT, promote bullying in children and permit discrimination [35–38]. Religion, too, can factor into promoting sedentariness [39].

Overall, the environment influences sitting in multiple ways. The precise magnitude of environmental influences on NEAT and/or sitting is poorly quantified because it is difficult to conduct societal-wide sedentariness research, gather valid population data and account appropriately for the vast array of variables as wide-ranging as religion, sociology, wealth and weather.

The impact of sitting on the human body. The physiology of sitting too much

Studies illustrate the physiological implications of sedentariness. Sixteen lean, healthy volunteers were overfed

1,000 kcal/day (4.18 MJ/day) above their weight maintenance needs for 2 months. Those volunteers who increased their NEAT the most gained the least amount of body fat. Some volunteers increased their NEAT by >700 kcal/day (>2.93 MJ/day) [40, 41]. Conversely, those people who did not increase their NEAT—those who stayed seated—were the most prone to gain body fat.

Underwear interlaced with sensors that captured body postures and movement was used to examine how a person can increase their NEAT by >700 kcal/day (>2.93 MJ/day). People who increase habitual walking (not purposeful exercise) can 'burn-off' the excesses of consumption. Conversely, people who overeat and stay seated gain the most body fat [41].

Human energetics can explain how NEAT can vary so greatly depending on how a person lives. Basal energy expenditure is the energy a person expends at complete rest. It increases with greater body size [42]. A seated person working at a computer screen generally expends 5% above basal energy expenditure—10–20 kcal/h (0.04–0.08 MJ/h) [42]. Within minutes of standing and walking, a person's energy expenditure doubles. An average person moving at 1–2 mph (1.6–3.2 kph) expends 100–250 kcal/h (0.42–1 MJ/h) above basal energy expenditure [42]. The standard workday of three hypothetical office workers are compared in Fig. 1. The figure illustrates that, during work hours alone, one worker can expend 1,000 kcal/day (4.2 MJ/day) more than another worker, even when both people have similar desk-based jobs. This accounting does not even consider the energy expenditure of leisure, which also varies greatly from sedentary screen time (100 kcal/evening) (0.42 MJ/evening) to active leisure (500–1,000 kcal/evening) (2.1–4.2 MJ/evening). It is not surprising, therefore, that when a group of people with similar desk-based jobs is studied, those who sit more have greater body fat than those who walk more [43, 44]. Interestingly, people who are most active in their leisure time are also most active at work [43]. Overall, people with obesity tend to sit for 2.5 h per day more than lean people with similar professions, economic statuses and home environments [44]. People with obesity tend to sit; walkers tend to be leaner.

There are other physiological factors that influence NEAT, including age, weather, concomitant illness, medications and sleep [45–48]. Sleep deprivation, for example, is associated with obesity [49–51] because, when people are deprived of sleep, their cumulative daily NEAT does not increase, despite their being awake for longer. Energy intake, however, does increase [52]. In sleep deprivation, positive energy balance and weight gain occur through the disassociation of energy intake from NEAT. It is interesting that a key neurochemical involved in wakefulness, orexin, is also a direct mediator of NEAT [53–55] and is integrated into the feeding circuits [56].

Fig. 1 Hypothetical workplace energy expenditures for (a) The chair-locked worker who sits most of the day; (b) the NEAThusiast who converts half of their six, 0.5-h meetings to walking meetings, parks a little farther away from the office, and takes 10 min ‘active breaks’ during the work day and a brief stroll at lunch time; and (c) the NEAThlete who conducts all work meetings while walking, has a chair-less desk, takes a 0.5 h walk at lunchtime and cycles to work

The molecular biology of sitting too much

Minor, short bursts of NEAT break up sitting and activate multiple molecular mechanisms. For example, in muscle, NEAT bursts improve insulin handling [57] via mitochondrial factors, lipolysis and insulin-receptor action [58–62]. The aforementioned associations between sitting, adiposity, insulin sensitivity and metabolic dysfunction can explain the association between excess sitting and cancer, in part [63–65]. Excess sitting also disrupts sex hormone function, insulin-like growth factors, inflammation and vitamin handling [66]. Prolonged sitting adversely impacts bone growth factors [67, 68] and, coupled with decreased weight bearing, sitting is associated with lower bone mineral density and osteoporosis [69, 70]. Being seated alters the activation patterns of multiple weight-bearing muscles and, therefore, excessive desk use is associated with adverse back curvature, back pain [71–74] and upper extremity problems such as carpal tunnel syndrome [75, 76]. Movement is neurologically linked to brain centres outside the motor control centres [53, 77–85]. Perhaps this assembly can explain the relationship between sedentariness, low mood, depression and dulled intellectual function [86, 87]. Furthermore, a person’s predisposition to sitting is partially predicated by genetic factors, although how these factors interface with environmental cues is not understood [14, 88–90].

En masse, the physiological evidence suggests that high levels of NEAT occur in the lean population, whereas low NEAT/sedentariness is linked to obesity. Sitting is ultimately a behaviour with environmental, sociological and biological cues. The organisational hub is in the brain.

The chair-locked brain

When people overeat, NEAT increases. This suggests that circuits exist to integrate food intake and NEAT [91]. For

example, if a person rakes leaves all day, they become hungry. It is unlikely that the integration of NEAT and food intake is limited to humans, because when wolves, horses and dogs run a lot, they eat a lot, too [92].

While overeating increases NEAT, starvation decreases NEAT. Keys and colleagues showed that prisoners decreased their spontaneous movements when starved [93, 94]. The link between starvation and NEAT occurs across species and genera [95, 96]. All species that have been studied show similar responses to starvation: first, there is a short-term increase in NEAT (ascribed to foraging), but once starvation is prolonged, NEAT decreases—even in fish [97]. Excess food and undernutrition evoke heightened and lowered NEAT, respectively, implying that a central connection exists between these two variables.

NEAT and energy intake connect in the hypothalamus. Several neuromodulators influence the paraventricular nucleus of the hypothalamus to regulate NEAT, such as orexin A and neuromedin U [54, 56, 80, 98, 99]. Orexin A is especially interesting, as it is an arousal protein associated with wakefulness—the period of the day when people move. When orexin A is injected into the paraventricular nucleus of the hypothalamus, NEAT increases in a dose-dependent fashion. The central effect of orexin A is blunted in rats inbred for obesity [53, 80]. Furthermore, orexin A is directly integrated with neuromodulators of appetite [100], muscle efficiency [53] and adipokines [100, 101]. The complete circuit for these interactions has not been fully mapped. However, these effects, and others like it, could explain why some people with a tendency to develop obesity are predisposed to sitting and eating in excess of energy needs. Conversely, people with heightened responsiveness to NEAT signals move more, have higher NEAT, maintain less body fat and resist obesity.

NEAT is influenced by other sensory inputs. For instance, how does a person who rakes leaves for several hours know that they are tired and sense the need to take a sitting break? What is the hierarchical interplay between behaviour and

neurochemistry whereby a person decides to conduct a professional meeting while walking instead of sitting? In both of these examples, there are likely to be multiple sensory inputs from the musculature, joints, tendons, nerves, metabolic systems and multiple brain regions. The response systems are likely to be hierarchical and complex, whereby NEAT activation is concomitantly affected by behaviour, environment and social cues. Bench-based biology bolsters the evidence that NEAT is regulated principally in the hypothalamus. However, field-based observations suggest that the integration of NEAT with life is complex and functions across the domains of biology, behaviour, environment and sociology; therefore, across many neuroanatomic domains as well.

NEAT changes in response to sudden environmental stimuli, e.g. jumping out of the way of a bus, and with short- and long-term illness, e.g. influenza and congestive heart failure [102]. NEAT also changes over the long term, e.g. with changes in food availability and with the seasons. NEAT changes across a person's lifespan—compare the frenetic energy of a young child [103, 104] to the deliberate gait of an elderly person [45, 105]. The age-decline in NEAT has been linked to changes in telomere length [90]. Historically, most people in agricultural communities moved throughout their day, and now most people sit for most of their day, so it is self-evident that movers can become sitters. If complex neurological systems regulate NEAT, what is the mechanism for NEAT adaptation? The central nervous system—both anatomically and biochemically—contains adaptive mechanisms that enable the brain to change its structure and function. Some of these neuroplasticity factors have been linked to NEAT and, by inference, to sitting [84, 106]. Several human experiments suggest that sitters can become movers [107, 108], and so it is plausible that central neuroplasticity factors influence human NEAT. Perhaps because the brain adapts more rapidly in the young, adaptation to NEAT-promoting environments occurs more readily in the young than in the elderly. Which epigenetic factors are important in NEAT adaptation remain unknown [106].

Excessive sitting occurs in most people in modern society because of sociological, behavioural and biological cues. Excessive sitting is a concern because it is harmful. The illnesses associated with sedentariness are discussed next.

Sitting sickness

Sedentariness and low NEAT may be important factors in obesity [109], but it is less obvious as to why sedentariness is causally associated with diabetes and 33 other chronic diseases and conditions [90]. To illustrate the impact of sitting, in one experiment, healthy volunteers were provided with three meals and encouraged to remain sedentary thereafter. The

same participants, on another occasion, were provided with similar meals and asked after they had eaten to walk for 15 min at 1.5 mph (2.4 kph) [110]. Continuous glucose monitoring showed that the short walks halved postprandial glycaemic excursion, regardless of when the meal was consumed. In this instance, and in several other experiments [111, 112], the data underscore how prolonged sitting increases insulin resistance and that breaking up sitting time can improve glucose handling [111–113]. These data provide a physiological rationale as to why sitting is associated with type 2 diabetes and gestational diabetes [114–116].

Thirty-five chronic diseases and conditions are associated with sedentariness [90], including frailty in the elderly [117–120], weight regain after therapeutic weight loss [121, 122], hypertension [123], osteoporosis [69, 124–126], malignancies such as breast [23] and prostate [127] cancer, cardiovascular disease [102, 128], male erectile dysfunction [129], depression [130], and back and musculoskeletal pain [75, 131]. Improved physical activity also helps with addictions to alcohol, opiates and cigarettes [132–135]. Chronic diseases and conditions associated with sedentary behaviour [90] impact approximately 70% of patients [136] and the majority of US healthcare costs [137], and so the fiscal consequences of a sedentary lifestyle are enormous.

Conclusion

Excessive sitting is a common pathway that contributes to numerous chronic diseases such as obesity, type 2 diabetes, cardiovascular disease and multiple types of cancer. Physiological and molecular mechanisms can explain these associations. The good news is that people can adapt; if work places, schools and cityscapes could be designed to promote activity, the new default posture could be up and moving. Upwardly mobile people could become healthier.

Sitting science and solutions to sedentariness are in their infancy. Scientific enquiry is important to understand the causality and consequence of sitting and to validate approaches to reverse sedentariness. But if the curse of lethal sitting is to be reversed, a societal-wide approach is needed.

Sedentariness is a transdisciplinary problem that hurts many people and therefore requires a transdisciplinary response. Entrepreneurs, policy makers, health professionals, teachers, industrialists, lawyers, business leaders, scientists, healthcare providers and other key leaders will need to get up, leave the comfort of their specialties, and achieve change in coalition. Individuals too must stand united. Ultimately, individuals are consumers, control school and company boards, work, attend educational centres, buy stock and decide elections. The challenge of sedentariness is as societal as it is personal: *Homo sedentarius*, arise.

Duality of interest The author declares that there is no duality of interest associated with this manuscript. However, JAL declares that he consults to Kersh Health Inc., Gruve Inc, SP Health Inc. and Gentag Inc. He does not receive fees from these organisations but has the potential to earn royalties. The author consults, without fee, to Apple Computer.

Contribution statement The author was the sole contributor to this paper.

References

- Morgan E (1993) Bipedalism. *Nutr Health* 9:193–203
- Pontzer H, Rolian C, Rightmire GP et al (2010) Locomotor anatomy and biomechanics of the Dmanisi hominins. *J Hum Evol* 58:492–504
- Barker G, Barton H, Bird M et al (2007) The ‘human revolution’ in lowland tropical Southeast Asia: the antiquity and behavior of anatomically modern humans at Niah Cave (Sarawak, Borneo). *J Hum Evol* 52:243–261
- Bennett MR, Harris JW, Richmond BG et al (2009) Early hominin foot morphology based on 1.5-million-year-old footprints from Ileret, Kenya. *Science* (New York, NY) 323:1197–1201
- Richards MP, Pettitt PB, Trinkaus E, Smith FH, Paunovic M, Karavanic I (2000) Neanderthal diet at Vindija and Neanderthal predation: the evidence from stable isotopes. *Proc Natl Acad Sci U S A* 97:7663–7666
- Carroll RL (1997) Patterns and processes of vertebrate evolution. Cambridge University Press, Cambridge
- Heitmann BL, Westerterp KR, Loos RJ et al (2012) Obesity: lessons from evolution and the environment. *Obes Rev* 10:910–922
- Levine JA, McCrady SK, Boyne S, Smith J, Cargill K, Forrester T (2011) Non-exercise physical activity in agricultural and urban people. *Urban Stud* 48:2417–2427
- Hibbs DA Jr, Olsson O (2004) Geography, biogeography, and why some countries are rich and others are poor. *Proc Natl Acad Sci U S A* 101:3715–3720
- Perdue WC, Stone LA, Gostin LO (2003) The built environment and its relationship to the public’s health: the legal framework. *Am J Public Health* 93:1390–1394
- Doyle R (2006) Not so revolutionary. Recent advances are no third industrial revolution. *Sci Am* 295:42
- Capon AG (2007) The way we live in our cities. *Med J Aust* 187:658–661
- Matthews CE, Chen KY, Freedson PS et al (2008) Amount of time spent in sedentary behaviors in the United States, 2003–2004. *Am J Epidemiol* 167:875–881
- Robinson S, Yardy K, Carter V (2012) A narrative literature review of the development of obesity in infancy and childhood. *J Child Health Care* 16:339–354
- Lanningham-Foster L, Nysse LJ, Levine JA (2003) Labor saved, calories lost: the energetic impact of domestic labor-saving devices. *Obes Res* 11:1178–1181
- Smith MJ, Conway FT, Karsh BT (1999) Occupational stress in human computer interaction. *Ind Health* 37:157–173
- Thomas L, Williams M (2006) Promoting physical activity in the workplace: using pedometers to increase daily activity levels. *Health Promot J Austr* 17:97–102
- Fernandes M, Sturm R (2010) Facility provision in elementary schools: correlates with physical education, recess, and obesity. *Prev Med* 50(Suppl 1):S30–S35
- Fernandes MM, Sturm R (2011) The role of school physical activity programs in child body mass trajectory. *J Phys Act Health* 8:174–181
- Mendoza JA, Levinger DD, Johnston BD (2009) Pilot evaluation of a walking school bus program in a low-income, urban community. *BMC Public Health* 9:122
- Levine JA (2011) Poverty and obesity in the U.S. *Diabetes* 60:2667–2668
- UN-Habitat (2005) State of the world’s cities 2004/2005—globalization and urban culture. Earthscan, London
- Schnohr P, Gronbaek M, Petersen L, Hein HO, Sorensen TI (2005) Physical activity in leisure-time and risk of cancer: 14-year follow-up of 28,000 Danish men and women. *Scand J Public Health* 33:244–249
- Walk Score. Cities and neighborhoods. Available from www.walkscore.com/cities-and-neighborhoods/. Accessed 20 Nov 2014
- Larouche R (2014) Assessing the health-related outcomes and correlates of active transportation in children and youth. *Appl Physiol Nutr Metab* 39:403
- Hsu YW, Belcher BR, Ventura EE et al (2011) Physical activity, sedentary behavior, and the metabolic syndrome in minority youth. *Med Sci Sports Exerc* 43:2307–2313
- Wu J, Jiang C, Houston D, Baker D, Delfino R (2011) Automated time activity classification based on global positioning system (GPS) tracking data. *Environ Health: Glob Access Sci Source* 10:101
- Gorzalany J (2013) The world’s most traffic-congested cities. Available from www.forbes.com/sites/jimgorzalany/2013/04/25/the-worlds-most-traffic-congested-cities/. Accessed 20 Nov 2014
- Thompson JL, Allen P, Cunningham-Sabo L, Yazzie DA, Curtis M, Davis SM (2002) Environmental, policy, and cultural factors related to physical activity in sedentary American Indian women. *Women Health* 36:59–74
- Levine JA, Koeppe GA (2011) Federal health-care reform: opportunities for obesity prevention. *Obesity* (Silver Spring) 19:897–899
- Hill JO, Peters JC (1998) Environmental contributions to the obesity epidemic. *Science* (New York, NY) 280:1371–1374
- Levine JA, Weisell R, Chevassus S, Martinez CD, Burlingame B, Coward WA (2001) The work burden of women. *Science* (New York, NY) 294:812
- Levine JA, Heet J, Burlingame B (2006) Aging on the job. *Sci Aging Knowl Environ* 28:pe16
- Levine JA, Weisell R, Chevassus S, Martinez CD, Burlingame B (2002) Looking at child labor. *Science* (New York, NY) 296:1025–1026
- Farhat T, Haynie D, Summersett-Ringgold F, Brooks-Russell A, Iannotti RJ (2015) Weight perceptions, misperceptions, and dating violence victimization among U.S. adolescents. *J Interpers Violence* 30:1511–1532
- Husarewycz MN, El-Gabalawy R, Logsetty S, Sareen J (2014) The association between number and type of traumatic life experiences and physical conditions in a nationally representative sample. *Gen Hosp Psychiatry* 36:26–32
- Jansen PW, Verlinden M, Domisse-van Berkel A et al (2014) Teacher and peer reports of overweight and bullying among young primary school children. *Pediatrics* 134:473–480
- Giel KE, Thiel A, Teufel M, Mayer J, Zipfel S (2010) Weight bias in work settings—a qualitative review. *Obes Facts* 3:33–40
- Persson G, Mahmud AJ, Hansson EE, Strandberg EL (2014) Somali women’s view of physical activity—a focus group study. *BMC Women’s Health* 14:129
- Levine JA, Eberhardt NL, Jensen MD (1999) Role of nonexercise activity thermogenesis in resistance to fat gain in humans. *Science* (New York, NY) 283:212–214
- Levine JA, McCrady SK, Lanningham-Foster LM, Kane PH, Foster RC, Manohar CU (2008) The role of free-living daily walking in human weight gain and obesity. *Diabetes* 57:548–554

42. Levine JA, Schleusner SJ, Jensen MD (2000) Energy expenditure of nonexercise activity. *Am J Clin Nutr* 72:1451–1454
43. McCrady SK, Levine JA (2009) Sedentariness at work: how much do we really sit? *Obesity (Silver Spring)* 17:2103–2105
44. Levine JA, Lanningham-Foster LM, McCrady SK et al (2005) Interindividual variation in posture allocation: possible role in human obesity. *Science (New York, NY)* 307:584–586
45. Harris AM, Lanningham-Foster LM, McCrady SK, Levine JA (2007) Nonexercise movement in elderly compared with young people. *Am J Physiol Endocrinol Metab* 292:E1207–E1212
46. Levine J, Melanson EL, Westerterp KR, Hill JO (2001) Measurement of the components of nonexercise activity thermogenesis. *Am J Physiol Endocrinol Metab* 281:E670–E675
47. Morgan MY, Levine JA (1988) Alcohol and nutrition. *Proc Nutr Soc* 47:85–98
48. Thomas DM, Ciesla A, Levine JA, Stevens JG, Martin CK (2009) A mathematical model of weight change with adaptation. *Math Biosci Eng* 6:873–887
49. Evans BD, Rogers AE (1994) 24-hour sleep/wake patterns in healthy elderly persons. *Appl Nurs Res* 7:75–83
50. Shapiro CM (1982) Energy expenditure and restorative sleep. *Biol Psychol* 15:229–239
51. Fraser G, Trinder J, Colrain IM, Montgomery I (1989) Effect of sleep and circadian cycle on sleep period energy expenditure. *J Appl Physiol* 66:830–836
52. Calvin AD, Carter RE, Adachi T et al (2013) Effects of experimental sleep restriction on caloric intake and activity energy expenditure. *Chest* 144:79–86
53. Novak CM, Escande C, Burghardt PR et al (2010) Spontaneous activity, economy of activity, and resistance to diet-induced obesity in rats bred for high intrinsic aerobic capacity. *Horm Behav* 58:355–367
54. Novak CM, Kotz CM, Levine JA (2006) Central orexin sensitivity, physical activity, and obesity in diet-induced obese and diet-resistant rats. *Am J Physiol Endocrinol Metab* 290:E396–E403
55. Kiwaki K, Kotz CM, Wang C, Lanningham-Foster L, Levine JA (2004) Orexin A (hypocretin 1) injected into hypothalamic paraventricular nucleus and spontaneous physical activity in rats. *Am J Physiol Endocrinol Metab* 286:E551–E559
56. Kotz CM, Teske JA, Levine JA, Wang C (2002) Feeding and activity induced by orexin A in the lateral hypothalamus in rats. *Regul Pept* 104:27–32
57. Solomon TP, Thyfault JP (2013) Type 2 diabetes sits in a chair. *Diab Obes Metab* 15:987–992
58. Deb S, Martin RJ (1975) Effects of exercise and of food restriction on the development of spontaneous obesity in rats. *J Nutr* 105:543–549
59. Mayer EJ, Burchfiel CM, Eckel RH, Marshall JA, Haskell WL, Hamman RF (1991) The role of insulin and body fat in associations of physical activity with lipids and lipoproteins in a biethnic population. The San Luis Valley diabetes study. *Arterioscler Thromb* 11:973–984
60. Short KR, Vittone JL, Bigelow ML et al (2003) Impact of aerobic exercise training on age-related changes in insulin sensitivity and muscle oxidative capacity. *Diabetes* 52:1888–1896
61. Ravussin E, Bogardus C (1992) A brief overview of human energy metabolism and its relationship to essential obesity. *Am J Clin Nutr* 55:242S–245S
62. Jensen MD, Sarr MG, Dumesic DA, Southorn PA, Levine JA (2003) Regional uptake of meal fatty acids in humans. *Am J Physiol Endocrinol Metab* 285:E1282–E1288
63. Klein WM, Bloch M, Hesse BW et al (2014) Behavioral research in cancer prevention and control: a look to the future. *Am J Prev Med* 46:303–311
64. Milton K, Macniven R, Bauman A (2014) Review of the epidemiological evidence for physical activity and health from low- and middle-income countries. *Glob Public Health* 9:369–381
65. Biswas A, Oh PI, Faulkner GE et al (2015) Sedentary time and its association with risk for disease incidence, mortality, and hospitalization in adults: a systematic review and meta-analysis. *Ann Intern Med* 162:123–132
66. Lynch BM (2010) Sedentary behavior and cancer: a systematic review of the literature and proposed biological mechanisms. *Cancer Epidemiol Biomarkers Prev* 19:2691–2709
67. Wiktor-Jedrzejczak W, Gordon S (1996) Cytokine regulation of the macrophage (M ϕ) system studied using the colony stimulating factor-1-deficient *op/op* mouse. *Physiol Rev* 76:927–947
68. Stanley ER, Berg KL, Einstein DB et al (1997) Biology and action of colony-stimulating factor-1. *Mol Reprod Dev* 46:4–10
69. Kolbe-Alexander TL, Charlton KE, Lambert EV (2004) Lifetime physical activity and determinants of estimated bone mineral density using calcaneal ultrasound in older South African adults. *J Nutr Health Aging* 8:521–530
70. DiPietro L (2001) Physical activity in aging: changes in patterns and their relationship to health and function. *J Gerontol A Biol Sci Med Sci* 56(Spec No 2):13–22
71. Beach TA, Parkinson RJ, Stothart JP, Callaghan JP (2005) Effects of prolonged sitting on the passive flexion stiffness of the in vivo lumbar spine. *Spine J* 5:145–154
72. Callaghan JP, McGill SM (2001) Low back joint loading and kinematics during standing and unsupported sitting. *Ergonomics* 44:280–294
73. Frymoyer JW (1992) Predicting disability from low back pain. *Clinical orthopaedics and related research*: 101–109
74. Morl F, Bradl I (2013) Lumbar posture and muscular activity while sitting during office work. *J Electromyogr Kinesiol* 23:362–368
75. Jordan JL, Holden MA, Mason EE, Foster NE (2010) Interventions to improve adherence to exercise for chronic musculoskeletal pain in adults. *Cochrane Database Syst Rev*, Issue 1, Art. no.: CD005956
76. Zejda JE, Bugajska J, Kowalska M et al (2009) Upper extremities, neck and back symptoms in office employees working at computer stations. *Med Pr* 60:359–367 [article in Polish]
77. Goldberg E, Bougakov D (2005) Neuropsychologic assessment of frontal lobe dysfunction. *Psychiatr Clin North Am* 28(567–580): 578–579
78. Dehaene S, Changeux JP (2000) Reward-dependent learning in neuronal networks for planning and decision making. *Prog Brain Res* 126:217–229
79. Kreier F, Kap YS, Mettenleiter TC et al (2006) Tracing from fat tissue, liver, and pancreas: a neuroanatomical framework for the role of the brain in type 2 diabetes. *Endocrinology* 147:1140–1147
80. Novak CM, Zhang M, Levine JA (2006) Neuromedin U in the paraventricular and arcuate hypothalamic nuclei increases non-exercise activity thermogenesis. *J Neuroendocrinol* 18:594–601
81. Jean A (2001) Brain stem control of swallowing: neuronal network and cellular mechanisms. *Physiol Rev* 81:929–969
82. Falgairolle M, Ceccato JC, Seze M, Herbin M, Cazalets JR (2013) Metachronal propagation of motor activity. *Front Biosci (Landmark Ed)* 18:820–837
83. Pisotta I, Molinari M (2014) Cerebellar contribution to feedforward control of locomotion. *Front Hum Neurosci* 8:475
84. Luque NR, Garrido JA, Carrillo RR, D'Angelo E, Ros E (2014) Fast convergence of learning requires plasticity between inferior olive and deep cerebellar nuclei in a manipulation task: a closed-loop robotic simulation. *Front Comput Neurosci* 8:97
85. Deliagina TG, Beloozerova IN, Orlovsky GN, Zelenin PV (2014) Contribution of supraspinal systems to generation of automatic postural responses. *Front Integr Neurosci* 8:76

86. Critchley CR, Hardie EA, Moore SM (2012) Examining the psychological pathways to behavior change in a group-based lifestyle program to prevent type 2 diabetes. *Diabetes Care* 35:699–705
87. Sund AM, Larsson B, Wichstrom L (2011) Role of physical and sedentary activities in the development of depressive symptoms in early adolescence. *Soc Psychiatry Psychiatr Epidemiol* 46:431–441
88. Bouchard C, Tremblay A (1990) Genetic effects in human energy expenditure components. *Int J Obes* 14(Suppl 1):49–55, discussion 55–48
89. Blum K, Oscar-Berman M, Bowirrat A et al (2012) Neuropsychiatric genetics of happiness, friendships, and politics: hypothesizing homophily (“birds of a feather flock together”) as a function of reward gene polymorphisms. *J Genet Syndr Gene Ther* 3:1000112
90. Thyfault JP, Du M, Kraus WE, Levine JA, Booth FW (2014) Physiology of sedentary behavior and its relationship to health outcomes. *Med Sci Sports Exerc*. doi:10.1249/MSS.0000000000000518
91. Mellecker RR, Lanningham-Foster L, Levine JA, McManus AM (2010) Energy intake during activity enhanced video game play. *Appetite* 55:343–347
92. Schutz Y (1997) On problems of calculating energy expenditure and substrate utilization from respiratory exchange data. *Z Ernahrungswiss* 36:255–262
93. Keys A, Brozek J, Henschel A, Mickelson O, Taylor HL (1950) The biology of human starvation. University of Minnesota Press, Minneapolis
94. Keys A, Anderson JT, Brozek J (1955) Weight gain from simple overeating. I. Character of the tissue gained. *Metabolism* 4:427–432
95. Sims EA, Horton ES (1968) Endocrine and metabolic adaptation to obesity and starvation. *Am J Clin Nutr* 21:1455–1470
96. Scrimshaw NS (1987) The phenomenon of famine. *Annu Rev Nutr* 7:1–21
97. Novak CM, Jiang X, Wang C, Teske JA, Kotz CM, Levine JA (2005) Caloric restriction and physical activity in zebrafish (*Danio rerio*). *Neurosci Lett* 383:99–104
98. Kotz C, Nixon J, Butterick T, Perez-Leighton C, Teske J, Billington C (2012) Brain orexin promotes obesity resistance. *Ann N Y Acad Sci* 1264:72–86
99. Teske JA, Levine AS, Kuskowski M, Levine JA, Kotz CM (2006) Elevated hypothalamic orexin signaling, sensitivity to orexin A and spontaneous physical activity in obesity resistant rats. *Am J Physiol Regul Integr Comp Physiol* 291:R889–R899
100. Trayhurn P, Bing C (2006) Appetite and energy balance signals from adipocytes. *Philos Trans R Soc Lond B Biol Sci* 361:1237–1249
101. Trayhurn P, Bing C, Wood IS (2006) Adipose tissue and adipokines—energy regulation from the human perspective. *J Nutr* 136:1935S–1939S
102. Toth MJ, Gottlieb SS, Goran MI, Fisher ML, Poehlman ET (1997) Daily energy expenditure in free-living heart failure patients. *Am J Physiol* 272:E469–E475
103. Lanningham-Foster L, Foster RC, McCrady SK et al (2008) Changing the school environment to increase physical activity in children. *Obesity (Silver Spring)* 16:1849–1853
104. Mellecker RR, McManus AM, Lanningham-Foster LM, Levine JA (2009) The feasibility of ambulatory screen time in children. *Int J Pediatr Obes* 4:106–111
105. Starling RD, Toth MJ, Carpenter WH, Matthews DE, Poehlman ET (1998) Energy requirements and physical activity in free-living older women and men: a doubly labeled water study. *J Appl Physiol* 85:1063–1069
106. Reynolds RF, Bronstein AM (2003) The broken escalator phenomenon. Aftereffect of walking onto a moving platform. *Exp Brain Res* 151:301–308
107. Koepp GA, Manohar CU, McCrady-Spitzer SK et al (2013) Treadmill desks: a 1-year prospective trial. *Obesity* 2013:705–711
108. Koepp GA, Manohar CU, McCrady-Spitzer SK, Levine JA (2011) Scalable office-based health care. *Health Serv Manage Res* 24:69–74
109. Ravussin E (2005) A NEAT way to control weight? *Science* 307:530–531
110. Manohar C, Levine JA, Nandy DK et al (2012) The effect of walking on postprandial glycemic excursion in patients with type 1 diabetes and healthy people. *Diabetes Care* 35:2493–2499
111. Dunstan DW, Kingwell BA, Larsen R et al (2012) Breaking up prolonged sitting reduces postprandial glucose and insulin responses. *Diabetes Care* 35:976–983
112. Hamilton MT, Healy GN, Dunstan DW, Zderic TW, Owen N (2008) Too little exercise and too much sitting: inactivity physiology and the need for new recommendations on sedentary behavior. *Curr Cardiovasc Risk Rep* 2:292–298
113. Basu R, Basu A, Johnson CM, Schwenk WF, Rizza RA (2004) Insulin dose-response curves for stimulation of splanchnic glucose uptake and suppression of endogenous glucose production differ in nondiabetic humans and are abnormal in people with type 2 diabetes. *Diabetes* 53:2042–2050
114. Fysekidis M, Cosson E, Banu I, Duteil R, Cyrille C, Valensi P (2014) Increased glycemic variability and decrease of the postprandial glucose contribution to HbA_{1c} in obese subjects across the glycemic continuum from normal glycemia to first time diagnosed diabetes. *Metabolism* 63:1553–1561
115. Ruifrok AE, Althuisen E, Oostdam N, van Mechelen W, Mol BW (2014) The relationship of objectively measured physical activity and sedentary behaviour with gestational weight gain and birth weight. *J Pregnancy* 2014:567379
116. Knowler WC, Barrett-Connor E, Fowler SE et al (2002) Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med* 346:393–403
117. Guimaraes FC, Amorim PR, Dos Reis FF et al (2014) Physical activity and better medication compliance improve mini-mental state examination scores in the elderly. *Dement Geriatr Cogn Disord* 39: 25–31
118. Sardinha LB, Santos DA, Silva AM, Baptista F, Owen N (2015) Breaking-up sedentary time is associated with physical function in older adults. *J Gerontol A: Biol Med Sci* 70:119–124
119. Vaz Fragoso CA, Miller ME, Fielding RA et al (2014) Sleep–wake disturbances in sedentary community-dwelling elderly adults with functional limitations. *J Am Geriatr Soc* 62:1064–1072
120. Pahor M, Guralnik JM, Ambrosius WT et al (2014) Effect of structured physical activity on prevention of major mobility disability in older adults: the LIFE study randomized clinical trial. *JAMA* 311:2387–2396
121. Bennett EM (1991) Weight-loss practices of overweight adults. *Am J Clin Nutr* 53:1519S–1521S
122. Sarlio-Lahteenkorva S, Rissanen A (1998) Weight loss maintenance: determinants of long-term success. *Eat Weight Disord* 3: 131–135
123. Vuori IM (2001) Health benefits of physical activity with special reference to interaction with diet. *Public Health Nutr* 4:517–528
124. Kenny AM, McGee D, Joseph C, Covault J, Abreu C, Raisz LG (2005) Lack of association between androgen receptor polymorphisms and bone mineral density or physical function in older men. *Endocr Res* 31:285–293
125. Nguyen TV, Sambrook PN, Eisman JA (1998) Bone loss, physical activity, and weight change in elderly women: the Dubbo Osteoporosis Epidemiology Study. *J Bone Miner Res* 13:1458–1467

126. Fenton M (2005) Battling America's epidemic of physical inactivity: building more walkable, livable communities. *J Nutr Educ Behav* 37(Suppl 2):S115–S120
127. Patel AV, Rodriguez C, Jacobs EJ, Solomon L, Thun MJ, Calle EE (2005) Recreational physical activity and risk of prostate cancer in a large cohort of U.S. men. *Cancer Epidemiol Biomarkers Prev* 14:275–279
128. Dunstan DW, Thorp AA, Healy GN (2011) Prolonged sitting: is it a distinct coronary heart disease risk factor? *Curr Opin Cardiol* 5: 412–419
129. Leoni LA, Fukushima AR, Rocha LY, Maifrino LB, Rodrigues B (2014) Physical activity on endothelial and erectile dysfunction: a literature review. *Aging Male* 17:125–130
130. Wadden TA, Steen SN, Wingate BJ, Foster GD (1996) Psychosocial consequences of weight reduction: how much weight loss is enough? *Am J Clin Nutr* 63:461S–465S
131. Proper KI, Heymans MW, Paw MJ et al (2006) Promoting physical activity with people in different places—a Dutch perspective. *J Sci Med Sport* 9:371–377
132. Unterwald EM, Ho A, Rubenfeld JM, Kreek MJ (1994) Time course of the development of behavioral sensitization and dopamine receptor up-regulation during binge cocaine administration. *J Pharmacol Exp Ther* 270:1387–1396
133. Greenberg JL, Lewis SE, Dodd DK (1999) Overlapping addictions and self-esteem among college men and women. *Addict Behav* 24:565–571
134. Ussher M, Cropley M, Playle S, Mohidin R, West R (2009) Effect of isometric exercise and body scanning on cigarette cravings and withdrawal symptoms. *Addiction* 104:1251–1257
135. De Maeyer J, Vanderplasschen W, Lammertyn J, van Nieuwenhuizen C, Sabbe B, Broekaert E (2011) Current quality of life and its determinants among opiate-dependent individuals five years after starting methadone treatment. *Qual Life Res* 20:139–150
136. Temple NJ, Burkitt DP (1993) Towards a new system of health: the challenge of Western disease. *J Community Health* 18:37–47
137. Popkin BM, Kim S, Rusev ER, Du S, Zizza C (2006) Measuring the full economic costs of diet, physical activity and obesity-related chronic diseases. *Obes Rev* 7:271–293