ERRATUM

Erratum to: Bedtime ingestion of hypertension medications reduces the risk of new-onset type 2 diabetes: a randomised controlled trial

Ramón C. Hermida¹ · Diana E. Ayala¹ · Artemio Mojón¹ · José R. Fernández¹

Published online: 6 November 2015
© Springer-Verlag Berlin Heidelberg 2015

Erratum to: Diabetologia
DOI 10.1007/s00125-015-3749-7

New-onset diabetes was not a pre-specified endpoint in the published trial design [1]. However, the authors note that the MAPEC trial was designed to investigate, as major outcomes, total cardiovascular events, major vascular events, new-onset diabetes and new-onset chronic kidney disease.

Reference

1. Hermida RC, for the MAPEC Study Investigators (2007) Ambulatory blood pressure monitoring in the prediction of cardiovascular events and effects of chronotherapy: rationale and design of the MAPEC Study. *Chronobiology International* 24:749–775

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00125-015-3749-7>.

✉ Ramón C. Hermida
rhermida@uvigo.es

¹ Bioengineering and Chronobiology Laboratories, E.I. Telecomunicación, University of Vigo, Campus Universitario, Vigo, Pontevedra 36310, Spain