



## Correction to: Bone marrow pericyte dysfunction in individuals with type 2 diabetes

Giuseppe Mangialardi<sup>1</sup> · David Ferland-McCollough<sup>1</sup> · Davide Maselli<sup>1,2,3</sup> · Marianna Santopaolo<sup>1</sup> · Andrea Cordaro<sup>1</sup> · Gaia Spinetti<sup>2</sup> · Maria Sambataro<sup>4</sup> · Niall Sullivan<sup>5</sup> · Ashley Blom<sup>5</sup> · Paolo Madeddu<sup>1</sup>

Published online: 21 May 2019  
© Springer-Verlag GmbH Germany, part of Springer Nature 2019

### Correction to: Diabetologia

<https://doi.org/10.1007/s00125-019-4865-6>

Unfortunately, three errors were made in the conversion of HbA<sub>1c</sub> to per cent values. The corrections are shown in bold in the text and table below:

### Inclusion and exclusion criteria

Type 2 diabetes was diagnosed according to the American Diabetes Association guidelines and was defined as follows: (1) patient/referring doctor reports a previous diagnosis of diabetes; (2) HbA<sub>1c</sub> >48 mmol/mol (**6.5%**) and (3) off insulin for at least 12 months after diagnosis.

**Table 1** Characteristics of participants

Characteristic	Without diabetes	With diabetes
Participants, <i>n</i>	25	14
Age, years	65 ± 2 (39–84)	67 ± 3 (41–79)
Men/women, <i>n</i>	11/14	7/7
BMI, kg/m <sup>2</sup>	29 ± 1 (21–42)	33 ± 2 (24–44)
HbA <sub>1c</sub> , mmol/mol	40 ± 1	60 ± 4
HbA <sub>1c</sub> , %	<b>5.8 ± 0.1</b>	<b>7.6 ± 0.4</b>

Data are mean±SD (range) or mean±SD, unless stated otherwise

The online version of the original article can be found at <https://doi.org/10.1007/s00125-019-4865-6>

✉ Paolo Madeddu  
mdprm@bristol.ac.uk

<sup>1</sup> Bristol Heart Institute, University of Bristol, Bristol Royal Infirmary, Level 7, Upper Maudlin Street, Bristol BS2 8HW, UK

<sup>2</sup> IRCCS Multimedica, Milan, Italy

<sup>3</sup> Department of Biochemistry, University of Sassari, Sassari, Italy

<sup>4</sup> Department of Specialized Medicines, Endocrine, Metabolic and Nutrition Diseases Unit, Santa Maria of Ca' Foncello Hospital, Treviso, Italy

<sup>5</sup> Musculoskeletal Research Unit, School of Clinical Sciences, University of Bristol, Bristol, UK