



Correction to: Altered mitochondrial bioenergetics and ultrastructure in the skeletal muscle of young adults with type 1 diabetes

Cynthia M. F. Monaco¹ · Meghan C. Hughes² · Sofia V. Ramos² · Nina E. Varah¹ · Christian Lamberz³ · Fasih A. Rahman⁴ · Chris McGlory⁵ · Mark A. Tarnopolsky⁶ · Matthew P. Krause⁴ · Robert Laham² · Thomas J. Hawke¹ · Christopher G. R. Perry²

Published online: 28 January 2020
© Springer-Verlag GmbH Germany, part of Springer Nature 2020

Correction to: Diabetologia


<https://doi.org/10.1007/s00125-018-4602-6>

In Fig. 1e the rate of mitochondrial H₂O₂ emission was incorrectly shown as being per second rather than per minute. The correct unit is pmol min⁻¹ [mg wet wt]⁻¹ and the figure part is corrected here.

The online version of the original article can be found at <https://doi.org/10.1007/s00125-018-4602-6>

✉ Thomas J. Hawke
hawke@mcmaster.ca

- ¹ Department of Pathology and Molecular Medicine, McMaster University, 4N65 Health Sciences Centre, 1200 Main Street West, Hamilton, ON L8N 3Z5, Canada
- ² School of Kinesiology and Health Sciences, Muscle Health Research Centre, York University, Toronto, ON, Canada
- ³ German Center for Neurodegenerative Diseases (DZNE), Bonn, Germany
- ⁴ Department of Kinesiology, University of Windsor, Windsor, ON, Canada
- ⁵ Department of Kinesiology, McMaster University, Hamilton, ON, Canada
- ⁶ Department of Pediatrics, McMaster University, Hamilton, ON, Canada



Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.