Correction to: Pregnant women with gestational diabetes and with well controlled glucose levels have decreased concentrations of individual fatty acids in maternal and cord serum

Henar Ortega-Senovilla¹ · Ute Schaefer-Graf² · Emilio Herrera¹

Published online: 1 April 2020

© Springer-Verlag GmbH Germany, part of Springer Nature 2020

Correction to: Diabetologia

<https://doi.org/10.1007/s00125-019-05054-x>

The authors regret that the incorrect figure was shown in Fig. 3d. The corrected figure is reproduced here.

The online version of the original article can be found at <https://doi.org/10.1007/s00125-019-05054-x>

✉ Henar Ortega-Senovilla
henar@ceu.es

¹ Department of Chemistry and Biochemistry, Facultad de Farmacia, Universidad San Pablo-CEU, Ctra. Boadilla del Monte km 5,3, E-28668 Madrid, Spain

² Department of Obstetrics and Gynecology, St Joseph's Hospital Center for Diabetes in Pregnancy, Berlin, Germany

Fig. 3 Correlation between maternal and cord serum of AA and DHA concentrations in control ($n = 90$) (a, c) and GDM ($n = 84$) (b, d) samples. The regression line, 95% CI lines and prediction limit lines are shown, together with the Pearson correlation coefficient. All values were $\log_{10}$ -transformed for statistical comparisons

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.