Association of diabetes-related autoantibodies with the incidence of asthma, eczema and allergic rhinitis in the TRIGR randomised clinical trial

Jeffrey P. Krischer¹ · David Cuthbertson¹ · Marisa Couluris² · Mikael Knip^{3,4} · Suvi M. Virtanen^{5,6,7,8}

Received: 3 December 2019 / Accepted: 15 April 2020 / Published online: 17 June 2020
© Springer-Verlag GmbH Germany, part of Springer Nature 2020

Abstract

Aims/hypothesis This paper presents the relationship between islet autoantibodies, precursors of type 1 diabetes, and the development of persistent asthma, allergic rhinitis and atopic eczema.

Methods A total of 2159 newborns who had a first-degree relative with type 1 diabetes and selected HLA genotypes were followed until the youngest participant reached 10 years of age. Islet cell antibodies (ICA) were detected using indirect immunofluorescence. Autoantibodies to insulin (IAA), GAD (GADA), the tyrosine phosphatase-related insulinoma-associated 2 molecule (IA-2A) and zinc transporter 8 (ZnT8A) were quantified with the use of specific radiobinding assays. As an ancillary study, the incidence of asthma, allergic rhinitis and eczema was assessed in 1106 of these children using the International Study of Asthma and Allergies in Childhood (ISAAC) core questionnaire when the children were 9–11 years old. HRs with 95% CIs were calculated to depict the incidence of these diseases following seroconversion to autoantibody positivity.

Results The cumulative incidence of atopic eczema, allergic rhinitis and persistent asthma were 22%, 9% and 7.5%, respectively, by 9–11 years of age. The occurrence of diabetes-related autoantibodies showed a protective association with subsequently reported incidence of asthma and eczema. The incidence of rhinitis was not significantly related to the occurrence of IAA or GADA (statistical power was limited), but demonstrated the same inverse relationship as did the other diseases with ICA or when multiple autoantibodies first appeared together.

Conclusions/interpretation The findings add evidence to the relationships between these atopic diseases and diabetes-related autoimmunity and also suggest that, for eczema, the interaction depends upon which autoantibody appeared first.

Trial registration [ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT00179777) NCT00179777

Keywords Asthma · Autoimmunity · Eczema · Rhinitis · Type 1 diabetes

Abbreviations

GADA	GAD autoantibodies
IA-2A	Tyrosine phosphatase-related insulinoma-associated 2 molecule autoantibodies
IAA	Insulin autoantibodies
ICA	Islet cell antibodies
ISAAC	International Study of Asthma and Allergies in Childhood

TRIGR	Trial to Reduce Insulin Dependent Diabetes in the Genetically at Risk
ZnT8A	Zinc transporter 8 antibodies

Introduction

The prevalence rates of asthma and other allergic diseases in children share many of the same characteristics as that of type 1 diabetes and its associated autoimmunity. Both have been increasing in prevalence during the last six decades, particularly in industrialised countries, although the increase may have levelled off in some [1–4].

Allergic diseases are caused by over-reactivity of the human immune system against foreign proteins, leading to

✉ Jeffrey P. Krischer
jeffrey.krischer@epi.usf.edu

Extended author information available on the last page of the article

Research in context

What is already known about this subject?

- Prevalence rates of type 1 diabetes and its associated autoimmunity, as well as asthma and allergic diseases in children, have been increasing during the last six decades

What is the key question?

- What was the relationship between the incidence of diabetes-related autoantibodies and the development of persistent asthma, allergic rhinitis and atopic eczema in the TRIGR study?

What are the new findings?

- Occurrence of diabetes-related autoantibodies showed a protective association with subsequently reported incidence of asthma and eczema
- Incidence of rhinitis was not significantly related to the occurrence of the autoantibodies IAA or GADA, but demonstrated the same inverse relationship as did the other diseases with ICA or when multiple autoantibodies first appeared together

How might this impact on clinical practice in the foreseeable future?

- These findings suggest an interaction between these atopic diseases and diabetes-related autoimmunity

asthma, allergic rhinitis, conjunctivitis, atopic eczema and food allergies. Parental heredity is a strong risk factor for the development of both asthma and atopic eczema: if one of the parents has the allergic disease the risk is increased at least twofold; if both, the increase is three- to fourfold [5, 6]. However, the rapid increase in the occurrence of these diseases over time indicates that environmental factors are also crucial in the disease aetiology. The same picture is found in diabetes-related autoimmunity and type 1 diabetes. A family history of type 1 diabetes increases diabetes risk and there is substantial evidence of the role of environmental exposures [7].

The immune abnormalities that lead to allergic diseases are initiated in infancy and possibly even during the fetal period. Diabetes-related autoimmunity has also been shown to occur in infancy for those presenting with insulin autoantibodies (IAA) as their first-appearing autoantibody, but at a later age for those presenting with GAD autoantibodies (GADA) first [8, 9]. Thus, early months of development may be critical for the maturation of the immune system as well as complex immune regulatory networks that result in functional disturbances, autoimmunity and allergic disorders.

The nutritional factors identified in epidemiological studies as possibly affecting the development of islet cell autoimmunity and allergic diseases and asthma include breastfeeding, age at introduction of new foods in infancy, dietary fatty acids, dietary antioxidants, vitamin D, foods that affect gut microbiota and obesity [10, 11]. There is contrasting evidence of whether prolonged feeding with a hydrolysed infant formula compared with a formula based on intact cow's milk reduces infant and childhood allergies or diabetes-related autoimmunity [12–14].

This paper explores the relationship between the incidence of asthma, rhinitis and eczema and the incidence of diabetes-related autoantibodies in a subset of the children enrolled in the Trial to Reduce Insulin Dependent Diabetes in the Genetically at Risk (TRIGR) study ([ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT00179777) Identifier: NCT00179777). The TRIGR infant feeding trial was designed to examine whether weaning to an extensively hydrolysed infant formula, as compared with formula based on cow's milk, delays or prevents the development of islet cell autoimmunity and/or type 1 diabetes in children at increased genetic risk of type 1 diabetes. TRIGR was a double-blind, randomised controlled trial of a dietary intervention (hydrolysed vs standard infant formula) that enrolled 2159 newborn first-degree relatives of patients with type 1 diabetes from 77 clinical sites in 15 countries [15–17]. The intervention period concluded in August 2007. This cross-sectional asthma and allergy ancillary study was performed between 2013 and 2016, when the children were 9–11 years old.

Methods

Participants Newborn infants who had a first-degree relative with type 1 diabetes and defined HLA genotypes were recruited into TRIGR between May 2002 and January 2007, and followed until diagnosis of type 1 diabetes or until the youngest participant reached 10 years of age in 2017. Randomisation of the infants who met the inclusion criteria took place before birth or immediately after birth. Altogether, 2159 newborn infants (1142 boys [52.9%]) with an eligible HLA genotype (41.9% of the genotyped infants) were randomised to the TRIGR intervention study. Randomisation

resulted in balanced groups with 1081 infants in the casein hydrolysate group and 1078 in the cow's milk group. In total, 516 infants (23.9%) carried the high-risk HLA genotype, 953 (44.1%) carried moderate-risk genotypes, 668 (30.9%) carried mild-risk genotypes and 22 (1.0%) carried the rare mild-risk genotype (Table 1). The first-degree relative with type 1 diabetes was the mother in 1055 infants (48.9%), the father in 723 (33.5%) and a sibling in 308 (14.3%), whereas 73 participants (3.4%) had multiple affected relatives. The median follow-up time for the diagnosis of diabetes was 11.5 years (range 3 months to 14.8 years; mean 11.0 years).

Written, informed consent was obtained from the family before enrolment into the TRIGR study. The study was approved by the ethics committees of all participating centres.

Dietary intervention Infants were randomly assigned weaning to either the intervention or control formula, produced specifically for this study. The intervention formula was an extensively hydrolysed casein-based formula, while the control formula was composed of 80% intact cow's milk protein and 20% hydrolysed milk protein, and formulated so that the taste and smell would be indistinguishable from the intervention formula. Study formulas were prepared and coded with the use of four colours by Mead Johnson Nutrition and were blinded to all investigators except the Data Management Unit (DMU). Newborn infants requiring supplemental feeding before randomisation (e.g., infants born at night or on weekends) received banked breast milk or Nutramigen, an extensively hydrolysed casein-based formula.

Breastfeeding was practiced at the discretion of the participating mothers, and maternal diets were unmodified. Breastfeeding was encouraged and exceeded national averages in both groups [18]. The dietary intervention period lasted until the infant was at least 6 months of age, and if by that time the child had not received the study formula for at least 60 days, study formula feeding was continued until 60 days of study formula exposure was reached, but not beyond 8 months of age. Parents were asked not to feed the children any commercial or other baby foods containing bovine protein during the intervention period. Adherence to the protocol was monitored by means of regular family nutrition interviews (at the ages of 0.5, 1, 2, 3, 4, 5, 6, 7 and 8 months) and by the analysis of cow's milk antibodies in serum samples. Detailed study design and methods have been previously published [16–18]. Online informed consent for this ancillary study was obtained for all study participants from a parent or primary caretaker.

HLA genotyping Cord blood or a heel stick blood sample collected on filter paper shortly after birth was immediately sent to the Turku (Europe and Australia) or Pittsburgh (North America) laboratory for HLA genotyping. HLA genotyping for the selected *DQB1* and *DQA1* alleles was performed using

sequence-specific oligonucleotide hybridisation, with quality control between the two laboratories carefully maintained. The following genotypes were regarded as eligible: (1) *HLA DQB1*02/DQB1*03:02* [high-risk]; (2) *HLA DQB1*03:02/x* (x not *DQB1*02*, *DQB1*03:01* or *DQB1*06:02*) [moderate risk]; (3) *HLA DQA1*05-DQB1*02/y* (y not *DQA1*02:01-DQB1*02*, *DQB1*03:01*, *DQB1*06:02* or *DQB1*06:03*) [mild risk]; and (4) *HLA DQA1*03-DQB1*02/y* (y not *DQA1*02:01-DQB1*02*, *DQB1*03:01*, *DQB1*06:02* or *DQB1*06:03*) [rare mild risk].

Diabetes-related autoantibodies Blood samples for autoantibody analyses were obtained (after application of analgesic cream) at the follow-up visits when the children were 3, 6, 9, 12, 18 and 24 months of age, and annually thereafter up to the age of 10–14 years. Serum samples were stored at -70°C until they could be analysed. Islet cell antibodies (ICA) were detected with the use of indirect immunofluorescence, while the four biochemical autoantibodies (IAA, GADA, autoantibodies to the tyrosine phosphatase-related insulinoma-associated 2 molecule [IA-2A] and autoantibodies to zinc transporter 8 [ZnT8A]) were quantified with the use of specific radiobinding assays [19]. Each autoantibody except for ZnT8A was analysed from all samples available. ZnT8A was initially quantified from the samples taken at the ages of 2, 3, 4, 6, 8, 10 and 12 years. If the 2 year sample tested positive, the samples obtained at the ages of 3, 6, 8, 9, 12 and 18 months were also analysed to assess the age at seroconversion. Similarly, any later samples between a negative and a positive sample were analysed.

Study endpoints The incidence of asthma, allergic rhinitis and eczema was assessed using the standardised and validated International Study of Asthma and Allergies in Childhood (ISAAC) core questionnaire on the child's history presented online to the parents of the participating children [20]. The online assessment included additional questions related to the child's use of asthma medications (if applicable), whether the child was ever diagnosed by a physician with certain diseases and food allergies, exposure to possible allergens and other factors related to asthma and/or allergies. Persistent asthma was defined as doctor-diagnosed asthma plus either any wheezing symptom during the preceding 12 months or use of asthma medication during the preceding 12 months. Age at onset of asthma was determined by the question: 'At what age did the symptom of asthma start or at what age did the child first wheeze?' Wheezing was defined as any of the following during the past 12 months: wheezy sound in respiration; wheezy sound during respiration in association with physical activity; difficulties in respiration in the morning on waking up; or wheezy respiration without having sniffles or respiratory infection. Allergic rhinitis was defined as doctor-diagnosed hay fever or allergic rhinoconjunctivitis

Table 1 Demographic characteristics of the whole TRIGR cohort and the present study participants

Variable	All TRIGR participants				All TRIGR participants who completed the ISAAC questionnaire: exposure to formula at 3 months			
	All participants <i>n</i> = 2159	Participants who did not complete ISAAC questionnaire <i>n</i> = 1053	Participants who completed ISAAC questionnaire <i>n</i> = 1106	<i>p</i> value	Exclusively breastfed <i>n</i> = 268	Hydrolysed formula <i>n</i> = 451	Cow's milk formula <i>n</i> = 350	No cow's milk or hydrolysed formula <i>n</i> = 37
ICA ⁺	764 (35.5)	359 (34.4)	405 (36.6)	0.27	91 (34.0)	171 (37.9)	131 (37.4)	12 (32.4)
IAA ⁺	341 (15.9)	186 (17.8)	155 (14.0)	0.02	42 (15.7)	63 (14.0)	44 (12.6)	6 (16.2)
GADA ⁺	369 (17.2)	194 (18.6)	175 (15.8)	0.09	45 (16.8)	71 (15.7)	55 (15.7)	4 (10.8)
IA2A ⁺	213 (9.9)	124 (11.9)	89 (8.1)	<0.01	26 (9.7)	36 (8.0)	27 (7.7)	0 (0.0)
ZnT8A ⁺	184 (8.6)	109 (10.4)	75 (6.8)	<0.01	18 (6.7)	36 (8.0)	19 (5.4)	2 (5.4)
Any Ab ⁺	959 (44.6)	445 (42.6)	514 (46.5)	0.07	122 (45.5)	213 (47.2)	165 (47.1)	14 (37.8)
Multi-Ab ⁺	298 (13.9)	177 (16.9)	121 (10.9)	<0.01	31 (11.6)	51 (11.3)	36 (10.3)	3 (8.1)
First positive autoantibody								
ICA	503 (23.3)	206 (19.6)	297 (26.9)	<0.01	64 (23.9)	125 (27.7)	98 (28.0)	10 (27.0)
IAA	149 (6.9)	72 (6.8)	77 (7.0)		19 (7.1)	34 (7.5)	20 (5.7)	4 (10.8)
GADA	157 (7.3)	77 (7.3)	80 (7.2)		21 (7.8)	29 (6.4)	30 (8.6)	0 (0.0)
IA2A	18 (0.8)	5 (0.5)	13 (1.2)		6 (2.2)	5 (1.1)	2 (0.6)	0 (0.0)
ZnT8A	22 (1.0)	11 (1.0)	11 (1.0)		2 (0.8)	6 (1.3)	2 (0.6)	1 (2.7)
Multiple AB	127 (5.9)	83 (7.9)	44 (4.0)		12 (4.5)	18 (4.0)	14 (4.0)	0 (0.0)
AB-	1183 (54.8)	599 (56.9)	584 (52.8)		144 (53.7)	234 (51.9)	184 (52.6)	22 (59.5)
HLA group								
DQB1*0302/x	953 (44.1)	441 (41.9)	512 (46.0)	0.06	133 (49.6)	207 (45.9)	155 (44.3)	17 (46.0)
DQB1*0302/DQB1*02	516 (23.9)	276 (26.2)	240 (21.7)		58 (21.6)	96 (21.3)	81 (23.1)	5 (13.5)
DQA1*05-DQB1*02/y	668 (30.9)	324 (30.8)	344 (31.1)		75 (28.0)	144 (31.9)	110 (31.4)	15 (40.5)
DQA1*03-DQB1*02/y	22 (1.0)	12 (1.1)	10 (0.9)		2 (0.8)	4 (0.9)	4 (1.1)	0 (0.0)
Sex								
Female	1021 (47.3)	518 (49.2)	503 (45.5)	0.08	129 (48.1)	195 (43.2)	161 (46.0)	18 (48.7)
Male	1138 (52.7)	535 (50.8)	603 (54.5)		139 (51.9)	256 (56.8)	189 (54.0)	19 (51.4)
Region								
Australia	101 (4.7)	61 (5.8)	40 (3.6)	<0.01	12 (4.5)	15 (3.3)	13 (3.7)	0 (0.0)
Europe	1135 (52.6)	447 (42.5)	688 (62.2)		191 (71.3)	246 (54.5)	221 (63.1)	30 (81.1)
Canada	528 (24.5)	282 (26.8)	246 (22.2)		36 (13.4)	131 (29.1)	73 (20.9)	6 (16.2)
USA	395 (18.3)	263 (25.0)	132 (11.9)		29 (10.8)	59 (13.1)	43 (12.3)	1 (2.7)
Mode of delivery								
Caesarean section	943 (43.7)	472 (44.8)	471 (42.6)	0.29	77 (28.7)	222 (49.2)	161 (46.0)	11 (29.7)
Vaginal delivery	1216 (56.3)	581 (55.2)	635 (57.4)		191 (71.3)	229 (50.8)	189 (54.0)	26 (70.3)
Number of older siblings								
0	2135 (98.9)	1045 (99.2)	1090 (98.6)	0.19	266 (99.3)	441 (97.8)	347 (99.1)	36 (97.3)
1	22 (1.0)	8 (0.8)	14 (1.3)		1 (0.4)	10 (2.2)	2 (0.6)	1 (2.7)
2	2 (0.1)	0 (0.0)	2 (0.2)		1 (0.4)	0 (0.0)	1 (0.3)	0 (0.0)
Potatoes/vegetables introduced by 6 months of age								
Yes	1465 (67.9)	669 (63.5)	796 (72.0)	<0.01	168 (62.7)	323 (71.6)	282 (80.6)	23 (62.2)

Table 1 (continued)

Variable	All TRIGR participants				All TRIGR participants who completed the ISAAC questionnaire: exposure to formula at 3 months			
	All participants <i>n</i> = 2159	Participants who did not complete ISAAC questionnaire <i>n</i> = 1053	Participants who completed ISAAC questionnaire <i>n</i> = 1106	<i>p</i> value	Exclusively breastfed <i>n</i> = 268	Hydrolysed formula <i>n</i> = 451	Cow's milk formula <i>n</i> = 350	No cow's milk or hydrolysed formula <i>n</i> = 37
Fruits/fruit juices introduced by 6 months of age	1414 (65.5)	656 (62.3)	758 (68.5)	<0.01	149 (55.6)	317 (70.3)	272 (77.7)	20 (54.1)
Cereals introduced by 6 months of age	1528 (70.8)	733 (69.6)	797 (72.1)	0.21	147 (54.9)	353 (78.3)	275 (78.6)	22 (59.5)
Number of foods introduced by 6 months				<0.01				
0	186 (8.6)	99 (9.4)	87 (7.9)		58 (21.6)	22 (4.9)	0 (0.0)	7 (18.9)
1	250 (11.6)	137 (13.0)	113 (10.2)		38 (14.2)	45 (10.0)	25 (7.1)	5 (13.5)
2	249 (11.5)	124 (11.8)	125 (11.3)		38 (14.2)	51 (11.3)	31 (8.9)	5 (13.5)
3	348 (16.1)	173 (16.4)	175 (15.8)		43 (16.0)	80 (17.7)	47 (13.4)	5 (13.5)
4	379 (17.6)	198 (18.8)	181 (16.4)		39 (14.6)	77 (17.1)	62 (17.7)	3 (8.1)
5	387 (17.9)	185 (17.6)	202 (18.3)		31 (11.6)	97 (21.5)	77 (22.0)	7 (18.9)
6	223 (10.3)	89 (8.5)	134 (12.1)		16 (6.0)	49 (10.9)	67 (19.1)	2 (5.4)
7	120 (5.6)	40 (3.8)	80 (7.2)		14 (5.2)	25 (5.5)	38 (10.9)	3 (8.1)
8	17 (0.8)	8 (0.8)	9 (0.8)		1 (0.4)	5 (1.1)	3 (0.9)	0 (0.0)
Family history of asthma								
Neither parent	–	–	829 (76.6)	–	209 (80.1)	337 (76.4)	255 (74.3)	28 (75.7)
1 parent	–	–	234 (21.6)	–	51 (19.5)	96 (21.8)	80 (23.3)	7 (18.9)
2 parents	–	–	19 (1.8)	–	1 (0.4)	8 (1.8)	8 (2.3)	2 (5.4)
Family history of rhinitis								
Neither parent	–	–	619 (57.5)	–	141 (54.7)	259 (59.0)	200 (58.3)	19 (51.4)
1 parent	–	–	361 (33.5)	–	93 (36.1)	143 (32.6)	112 (32.7)	13 (35.1)
2 parents	–	–	97 (9.0)	–	24 (9.3)	37 (8.4)	31 (9.0)	5 (13.5)
Family history of eczema								
Neither parent	–	–	763 (70.7)	–	182 (69.7)	298 (68.0)	256 (74.6)	27 (73.0)
1 parent	–	–	267 (24.8)	–	63 (24.1)	123 (28.1)	73 (21.3)	8 (21.6)
2 parents	–	–	49 (4.5)	–	16 (6.1)	17 (3.9)	14 (4.1)	2 (5.4)
Mother smoked during pregnancy								
Yes	–	–	69 (6.2)	–	8 (3.0)	36 (8.0)	24 (6.9)	1 (2.7)
Smoking exposure during 1st year								
Yes	–	–	239 (21.6)	–	43 (16.0)	100 (22.2)	87 (24.9)	9 (24.3)
Day/child care exposure during 1st year								
Yes	–	–	122 (11.0)	–	19 (7.1)	55 (12.2)	44 (12.6)	4 (10.8)
Eczema diagnosed within first 6 months of life								
Yes	–	–	9 (0.8)	–	4 (1.5)	3 (0.7)	2 (0.6)	0 (0.0)

Data are *n* (%)

Differences in calculated percentages are due to adjustments for missing data for specific variables

during the preceding 12 months. Age at onset of allergic rhinitis was based on the age of the child at the onset of symptoms including sneezing, nasal congestion or rhinitis other than with respiratory infections, accompanied by itching of the eyes and tearing. Atopic eczema was defined as ever having doctor-diagnosed atopic eczema. Age of onset of atopic eczema was based on the question: ‘At what age did the child first have atopic eczema?’ All online questionnaires, including ISAAC, were available in local languages.

Statistical analyses The cumulative incidence of onset of each endpoint from birth was estimated using a modified Kaplan–Meier disease-free survival function. The differences between groups in the cumulative incidence functions, and the associated hazard functions, were tested using the logrank test according to the actual diet consumed. The relative risk of disease onset between groups was estimated with the Cox proportional hazards model using time-dependent covariates. Multivariate analyses included characteristics that were found to be statistically significantly associated with autoantibody incidence from the univariate analyses. The autoantibody-negative group was defined to be the referent population for comparing disease incidence by first-appearing autoantibody. When comparing characteristics between the extensively hydrolysed formula and cow’s milk formula groups, the *t* test was applied for normally distributed variables and the non-parametric Mann–Whitney *U* test for skewed variables.

Although the overall study sample size was substantial, the statistical power for testing the effect of covariates on each disease was limited due to the number of children having had that diagnosis. The TRIGR study had >80% power to detect HRs of 0.5 or less in autoantibody incidence according to the use of hydrolysed formula. The power to detect differences in the incidence of atopic diseases, which occurred with lower frequency, conditioned upon autoantibody occurrence was much less. The detectable HRs at 80% power ranged from 0.1 to 0.3 for asthma, allergic rhinitis and eczema.

Data were analysed using the Statistical Analysis System software (version 9.4; SAS Institute, Cary, NC, USA). Two-tailed *p* values less than 0.05 were considered to be statistically significant. No adjustment in type 1 error was made for multiple comparisons except in the context of the multiple Cox regression model.

The TRIGR study and this ancillary study was approved by local institutional review or ethics boards and was monitored by an External Evaluation Committee formed by the National Institutes of Health.

Results

Of the total study population, 1106 (51.2%) completed the ISAAC questionnaire and were included in this analysis.

Altogether, 268 children in this asthma and allergy study population were exclusively breastfed at 3 months of age, 451 received hydrolysed formula, 350 received cow’s milk formula and the remaining 37 were not exclusively breastfed but did not receive either of the formulas. There were statistically significant differences between the enrolled total TRIGR cohort and those who completed the ISAAC questionnaire according to their demographic characteristics. A higher proportion of the ISAAC responders were enrolled from TRIGR European countries as compared with North America or Australia (62% vs 43%, $p < 0.01$), had an increased number of foods introduced by 6 months of age (20% vs 13%, $p < 0.01$) (notably potatoes/vegetables [72% vs 64%] and fruits/fruit juices [69% vs 62%]) and were exclusively breastfed for the first 3 months of life (24% vs 18%, $p < 0.01$). With regard to other demographics measured, there were no statistically significant differences between those who responded to the ISAAC questionnaire and those who did not.

Comparing feeding practices during the first 3 months of life for those who completed the ISAAC questionnaire, among those exclusively breastfed or that did not receive hydrolysed or cow’s milk-containing formulas, a higher percentage of respondents were from sites in Europe as compared with the remaining TRIGR sites (72.5% vs 58.3%, $p < 0.001$) or were delivered vaginally (71.1% vs 52.2%, $p < 0.001$), and a lower percentage had diets that included potatoes/vegetables (62.6% vs 75.5%), fruits/fruit juices (55.4% vs 73.5%) and cereals (55.4% vs 78.4%) by 6 months of age (all $p < 0.0001$). Also, respondents who were exclusively breastfed or did not receive hydrolysed or cow’s milk-containing formulas were less likely to come from families where mothers smoked during pregnancy (3.0% vs 7.5%, $p = 0.005$) or whose children lived in a household with an active smoker during the first year of life (17.0% vs 23.3%, $p = 0.023$), as compared with the other feeding groups (Table 1).

The cumulative incidence rates for eczema, allergic rhinitis and asthma were 22% ($N = 251$ [95% CI 18.5%, 25.7%]), 9% ($N = 112$ [95% CI 6.8%, 11.2%]) and 7.5% ($N = 89$ [95% CI 3.8, 11.2%]), respectively, by 9–11 years of age (Fig. 1). Among those who remained autoantibody negative ($N = 599$), the cumulative incidence rates of eczema, allergic rhinitis and asthma were 33% (95% CI 29.3%, 36.9%), 14% (95% CI 10.8%, 17.8%) and 12.4% (95% CI 9.7%, 15.1%) by 9–11 years of age. These rates are lower than anticipated from the ISAAC study conducted in these same countries, but not by much when considering those who remained autoantibody negative through the follow-up period. Among the TRIGR countries, the prevalence of asthma in children between 6 and 14 years of age as defined by the ISAAC study was highest in Australia (26–37% for asthma). For the USA, the prevalence was 17–20%, and in Canada it was 16–19%. The prevalence was lowest among European countries

Fig. 1 Cumulative incidence of persistent asthma, atopic eczema and allergic rhinitis

participating in TRIGR, ranging from the lowest estimates in Estonia (4–5%) to the highest in Italy (11–15%). The prevalence rates of current wheezing according to ISAAC in children of 6–14 years old were: Australia 20–31%, USA 19–22%, Canada 14–15%, Estonia 9–10% and Italy 8–11%. The prevalence rates of allergic sensitisation to any allergen across the TRIGR countries were in the range of 35–40% [19–23].

Disease risk was significantly increased ($p < 0.01$) for each of the three diseases with a family history of that disease. Also, there was a significantly higher risk for asthma (HR 81.02, $p < 0.01$) if eczema was diagnosed 6 months earlier or within the first 6 months of age (HR 2.37, $p = 0.002$). Excluding IA-2A and ZnT8A due to small numbers ($N = 8$ for each), the development of islet cell autoimmunity was significantly associated with a reduced subsequent risk of asthma or eczema, no matter which autoantibody was present, with HRs ranging from 0.13 to 0.18 (Table 2, Fig. 2). The presence of ICA ($N = 233$, HR 0.41, $p < 0.01$) significantly reduced the risk of allergic rhinitis, but neither IAA ($N = 39$) nor GADA ($N = 48$) had a significant impact on allergic rhinitis (HR 0.63, $p = 0.14$ and HR 0.57, $p = 0.09$, respectively). ICA, as the first-appearing autoantibody ($N = 233$), significantly reduced the subsequent risk of asthma (HR 0.22, $p < 0.01$), rhinitis (HR 0.40, $p < 0.01$) and eczema (HR 0.27, $p < 0.01$). IAA or GADA as the first-appearing autoantibody was followed by a significant risk reduction only for eczema (HR 0.29, $p = 0.01$ and HR 0.33, $p = 0.02$, respectively). The only other significant finding was the lower risk (HR 0.55, $p = 0.01$) of asthma for study children from Europe as compared with children from the USA.

Multivariable analyses of these risk factors resulted in consistent statistically significant reduced risk when ICA was the first-appearing autoantibody ($p < 0.01$; HR 0.24, HR

0.40 and HR 0.27 for asthma, allergic rhinitis and eczema, respectively). When multiple autoantibodies appeared first ($N = 169$), the reduced risk remained significant for asthma (HR 0.06, $p < 0.01$) and allergic rhinitis (HR 0.38, $p < 0.01$), but the risk reduction was no longer significant for eczema ($p < 0.12$). For eczema, IAA and GADA, as first-appearing autoantibodies, were each associated with a statistically significant risk reduction (HR 0.30, $p = 0.02$ and HR 0.34, $p = 0.02$, respectively). The appearance of these autoantibodies individually did not result in a statistically significant impact on risk for asthma or allergic rhinitis. Family history (either one or both parents) conveyed a significantly higher risk across all three diagnoses, as did a family history of eczema for asthma, similar to what was seen in the univariate analyses (Table 3).

Discussion

The lower reported incidence of persistent asthma, allergic rhinitis and eczema in the TRIGR countries could be explained, in part, by the higher incidence of diabetes-related autoantibodies in the TRIGR study population and the inverse relationship of their incidence found in this study. This finding is consistent with other studies in North America [24] and Europe [25] that report a lower prevalence of asthma in children with type 1 diabetes, as compared with the general population; however, some findings have differed, especially the USA [26]. Hence, the relationship between type 1 diabetes and atopic disease remains controversial. That the TRIGR study cohort was not population based, but selectively included children at higher genetic risk for diabetes among families with a first-degree relative with type 1 diabetes, could be considered a limitation on the findings reported herein. However, this paper focuses on the relationship between

Table 2 Univariate HRs (95% CIs) and statistical significance of factors associated with the incidence of asthma, rhinitis and eczema

Variable	Asthma N= 87				Rhinitis N= 112				Eczema N= 251										
	N	n	p value	HR	95% CI	N	n	p value	HR	95% CI	N	n	p value	HR	95% CI				
First positive autoantibody	ICA ⁺	375	8	<0.01	0.18	0.09	0.38	361	21	<0.01	0.41	0.26	0.66	335	27	<0.01	0.24	0.16	0.36
	IAA ⁺	141	1	0.01	0.08	0.01	0.55	141	10	0.14	0.63	0.32	1.18	122	9	<0.01	0.27	0.14	0.53
	GADA ⁺	163	2	<0.01	0.13	0.03	0.54	152	10	0.09	0.57	0.30	1.09	142	10	<0.01	0.26	0.14	0.48
	IA2A ⁺	82	2	0.08	0.28	0.07	1.15	77	6	0.39	0.70	0.31	1.59	71	6	<0.01	0.33	0.15	0.73
	ZnT8A ⁺	63	0	0.98	0.00	0.00	0.00	60	5	0.52	0.75	0.31	1.83	57	4	<0.01	0.27	0.10	0.73
	Any Ab ⁺	481	10	<0.01	0.16	0.08	0.30	466	31	<0.01	0.45	0.30	0.68	428	38	<0.01	0.24	0.17	0.34
	Multi-Ab ⁺	109	1	0.02	0.10	0.01	0.73	105	7	0.17	0.58	0.27	1.25	94	6	<0.01	0.24	0.11	0.54
	ICA	233	7	<0.01	0.22	0.10	0.47	225	14	<0.01	0.40	0.23	0.71	211	22	<0.01	0.27	0.18	0.43
	IAA	39	0	0.98	0.00	0.00	0.00	44	5	0.60	0.78	0.32	1.93	38	4	0.01	0.28	0.11	0.76
	GADA	48	1	0.07	0.16	0.02	1.13	44	2	0.10	0.31	0.08	1.25	41	5	0.01	0.33	0.14	0.80
HLA group	IA2A	8	1	0.91	0.90	0.13	6.45	8	1	0.84	0.82	0.11	5.88	6	1	0.42	0.44	0.06	3.17
	ZnT8A	8	0	0.99	0.00	0.00	0.00	7	1	0.92	0.91	0.13	6.53	8	0	0.97	0.00	0.00	0.00
	Multi-Ab	169	1	<0.01	0.04	0.01	0.31	160	9	<0.01	0.37	0.19	0.74	148	6	<0.01	0.11	0.05	0.24
	Ab ⁺ (reference)	599	77	–	–	–	–	563	80	–	–	–	–	654	213	–	–	–	–
	DQB1*0302/x (reference)	511	35	–	–	–	–	483	53	–	–	–	–	512	128	–	–	–	–
	DQB1*0302/DQB1*02	239	22	0.25	1.37	0.80	2.33	229	26	0.78	1.07	0.67	1.71	240	51	0.30	0.84	0.61	1.17
	DQA1*05-DQB1*02/y	344	29	0.40	1.23	0.75	2.02	329	32	0.57	0.88	0.57	1.37	344	69	0.11	0.79	0.60	1.06
	DQA1*03-DQB1*02/y	10	1	0.72	1.45	0.20	10.57	10	1	0.92	0.91	0.13	6.55	10	3	0.73	1.23	0.39	3.85
	Female	501	37	0.55	0.88	0.58	1.34	473	46	0.37	0.84	0.58	1.23	503	125	0.14	1.21	0.94	1.54
	Australia	40	4	0.49	0.69	0.23	2.02	39	4	0.44	0.65	0.22	1.93	40	8	0.58	0.80	0.37	1.74
Sex	Europe	686	42	<0.01	0.41	0.24	0.70	660	68	0.09	0.64	0.38	1.07	688	148	0.48	0.87	0.60	1.28
	Canada	246	22	0.10	0.60	0.32	1.10	234	22	0.07	0.57	0.30	1.05	246	63	0.89	1.03	0.67	1.58
	USA (reference)	132	19	–	–	–	–	118	18	–	–	–	–	132	32	–	–	–	–
	Caesarean section	470	35	0.63	0.90	0.59	1.38	448	56	0.10	1.36	0.94	1.97	471	108	0.86	1.02	0.80	1.31
	3 month formula exposure	267	16	–	–	–	–	257	31	–	–	–	–	268	63	–	–	–	–
	Exclusively breastfed (reference)	450	43	0.10	1.61	0.91	2.86	428	45	0.54	0.87	0.55	1.37	451	99	0.56	0.91	0.66	1.25
	Hydrolysed formula	350	27	0.41	1.30	0.70	2.40	330	34	0.45	0.84	0.52	1.37	350	80	0.81	0.96	0.69	1.34
	Cow's milk formula	37	1	0.42	0.44	0.06	3.30	36	2	0.26	0.44	0.11	1.84	37	9	0.91	1.04	0.52	2.09
	No cow's milk or hydrolysed formula	828	39	–	–	–	–	588	34	–	–	–	–	763	141	–	–	–	–
	Family history of disease	234	42	<0.01	4.14	2.68	6.41	348	57	<0.01	3.00	1.96	4.58	267	90	<0.01	1.96	1.51	2.56
Mother smoked during pregnancy	1 parent	19	5	<0.01	6.17	2.43	15.65	90	21	<0.01	4.56	2.65	7.86	49	20	<0.01	2.45	1.53	3.93
	2 parents	69	5	0.81	0.89	0.36	2.21	66	7	0.98	0.99	0.46	2.13	69	14	0.64	0.88	0.51	1.51
	Yes																		

Table 2 (continued)

Variable	Asthma N = 87			Rhinitis N = 112			Eczema N = 251			
	N	n	p value	HR	95% CI	N	n	p value	HR	95% CI
Smoking exposure during 1st year	239	15	0.30	0.75	0.43 1.30	227	24	0.95	0.99	0.63 1.55
Day/child care exposure during 1st year	122	9	0.83	0.93	0.47 1.85	117	13	0.89	1.04	0.59 1.86
Eczema diagnosed within first 6 months of life	102	9	<0.01	2.37	1.38 4.08	—	—	—	—	—

The number of individuals with a given characteristic (*N*) and the number diagnosed (*n*) for each disease is listed

Fig. 2 Cumulative incidence by prior autoantibody status for asthma (a), eczema (b) and allergic rhinitis (c)

autoimmunity and asthma, allergic rhinitis and eczema, not the prevalence of these disorders. The selection bias would be pertinent only if one thought that the relationship was different among those who developed diabetes-related autoimmunity and these atopic diseases in other, lower risk, HLA groups. There are no reports in the literature to support this contention.

The occurrence of diabetes-related autoantibodies showed a protective association with subsequent development of persistent asthma and atopic eczema. The incidence of allergic rhinitis was not significantly related to the occurrence of IAA or GADA (statistical power was limited), but demonstrated the same inverse relationship, as did the other diseases, with ICA or when multiple autoantibodies first appeared together. Hence, when considering putative confounders and all of the first-appearing endpoints simultaneously, ICA and multiple autoantibodies were associated with persistent asthma and allergic rhinitis, and ICA, IAA and GADA with atopic eczema. The lack of statistical significance in the comparison of effect of the occurrence of IAA or GADA with allergic rhinitis must be interpreted with some caution as the study has limited statistical power, as noted above.

These findings extend the observation of an inverse relationship between type 1 diabetes and later development of asthma, showing the same directional effect but suggesting that the protective effect extends to the first appearance of autoantibodies [27]. As far as we are aware, this has not been shown before. An earlier study has observed a direct association between wheezing during the first year of life and presence of GADA and IA-2A autoimmunity at the age of 2.5 years [28]. The present study did not observe sufficient numbers of cases of seroconversion to autoantibody positivity after a diagnosis of asthma or allergic rhinitis to test the question of whether earlier diagnosis of these diseases increased the incidence of autoimmunity. In this population, autoantibodies most often occurred in early childhood, preceding disease diagnosis, except for eczema, which was diagnosed at a very early age. In this study, islet autoantibodies were rarely seen before 1 year of age. In other studies [8], when they are seen, IAA is most often the first-appearing autoantibody. Also, this study did not observe a difference in disease diagnosis according to the sex of the child. The random allocation to different early feeding patterns also supports the lack of an observed

Table 3 Multivariate HRs (95% CIs) and statistical significance of factors affecting the incidence of asthma, rhinitis and eczema

Disease	<i>p</i> value	HR	95% CI	
Asthma (<i>n</i> = 87)				
First positive autoantibody				
ICA	<0.01	0.21	0.10	0.46
IAA	0.98	0.00	0.00	0.00
GADA	0.05	0.14	0.02	1.03
IA2A	0.82	0.79	0.11	5.96
ZnT8A	0.99	0.00	0.00	0.00
Multi-Ab	<0.01	0.05	0.01	0.37
Ab [−] (reference)		−	−	−
Region				
Australia	0.29	0.55	0.18	1.66
Europe	0.02	0.53	0.31	0.92
Canada	0.26	0.70	0.38	1.30
USA (reference)		−	−	−
Family history of disease				
Neither parent (reference)		−	−	−
1 parent	<0.01	3.82	2.46	5.92
2 parents	<0.01	6.80	2.55	18.14
Eczema diagnosed within first 6 months of life				
Yes	<0.01	2.23	1.29	3.86
Rhinitis (<i>n</i> = 112)				
First positive autoantibody				
ICA	<0.01	0.40	0.23	0.71
IAA	0.68	0.83	0.34	2.04
GADA	0.06	0.26	0.06	1.05
IA2A	0.89	0.87	0.12	6.28
ZnT8A	0.84	0.82	0.11	5.90
Multi-Ab	<0.01	0.38	0.19	0.75
Ab [−] (reference)		−	−	−
Family history of disease				
Neither parent (reference)		−	−	−
1 parent	<0.01	2.98	1.95	4.56
2 parents	<0.01	4.70	2.72	8.10
Eczema (<i>n</i> = 251)				
First positive autoantibody				
ICA	<0.01	0.27	0.18	0.43
IAA	0.01	0.29	0.11	0.77
GADA	0.02	0.34	0.14	0.83
IA2A	0.45	0.47	0.07	3.36
ZnT8A	0.97	0.00	0.00	0.00
Multi-Ab	<0.01	0.11	0.27	0.18
Ab [−] (reference)		−	−	−
Family history of disease				
Neither parent (reference)		−	−	−
1 parent	<0.01	1.79	1.38	2.34
2 parents	<0.01	2.14	1.34	3.43

relationship between atopic disease incidence and early exposure to hydrolysed formula compared with cow's milk-based formula. Despite the inclusion of a series of potential exposures that could be related to disease incidence, only family history and the prior development of diabetes-related autoimmunity were significant.

The findings of the present study add evidence to the relationships between atopic diseases and diabetes-related autoimmunity, and also suggest that the interaction depends upon which autoantibody appeared first for eczema but not for asthma or rhinitis.

Acknowledgements A special acknowledgement to the TRIGR families for their participation in this wonderful study and to the dedication of the TRIGR study coordinators who contributed to the amazing success of TRIGR and this ancillary study.

Data availability The data used in the analysis reported in this paper are available from the TRIGR Data Management Unit upon request to the lead author.

Funding This work was supported by the Eunice Kennedy Shriver National Institute of Child Health & Human Development (NICHD) and the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK), the National Institutes of Health (NIH) (grant numbers HD040364, HD042444 and HD051997), the Canadian Institutes of Health Research, JDRF and the Commission of the European Communities (specific RTD programme 'Quality of Life and Management of Living Resources', contract number QLK1-2002-00372 'Diabetes Prevention'). Other funding came from the EFSD/JDRF/Novo Nordisk Focused Research Grant, the Academy of Finland (Centre of Excellence in Molecular Systems Immunology and Physiology Research 2012-2017, Decision No. 250114), the Dutch Diabetes Research Foundation and the Finnish Diabetes Research Foundation. Mead Johnson Nutrition provided the blinded colour-coded study formulas.

Authors' relationships and activities The authors declare that there are no relationships or activities that might bias, or be perceived to bias, their work.

Contribution statement All authors attest to meeting ICMJE uniform requirements for authorship by making substantial contributions to the conception and design of this paper; acquisition, analysis and interpretation of the data; drafting or revising the article for intellectual content; and giving final approval of the published version. JPK designed the study, proposed the analysis, interpreted the findings and wrote the manuscript. DC performed the analysis and contributed to the manuscript. SMV, MK and MC designed the study and reviewed/edited the manuscript. JPK and DC are the guarantors of this work and, as such, had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis. The TRIGR study funder was not involved in the design of the study; the collection, analysis and interpretation of data; writing of the report; or the decision to submit the report for publication.

References

1. Seaton A, Godden DJ, Brown K (1994) Increase in asthma: a more toxic environment or a more susceptible population? *Thorax* 49:171–174

2. Becker AB (2000) Is primary prevention of asthma possible? *Pulmonol* 30:63–72
3. Devereux G (2006) The increase in the prevalence of asthma and allergy: food for thought. *Nat Rev Immunol* 6:869–874
4. Zöllner IK, Weiland SK, Piechotowski I et al (2005) No increase in the prevalence of asthma, allergies, and atopic sensitization among children in Germany: 1992–2001. *Thorax* 60:545–548
5. Illi S, von Mutius E, Lau S et al (2006) Perennial allergen sensitization early in life and chronic asthma in children: a birth cohort study. *Lancet* 368(9537):763–770
6. Gough H, Grabenhenrich L, Reich A et al (2015) Allergic multimorbidity of asthma, rhinitis and eczema over 20 years in the German birth cohort MAS. *Pediatr Allergy Immunol* 26(5):431–437
7. Steck AK, Vehik K, Bonifacio E et al (2015) Predictors of progression from the appearance of islet autoantibodies to early childhood diabetes: The Environmental Determinants of Diabetes in the Young (TEDDY). *Diabetes Care* 38(5):808–813
8. Ilonen J, Hammaï A, Laine A-P et al (2013) Patterns of autoantibody appearance and genetic associations reveal heterogeneous pathways of β -cell destruction. *Diabetes* 62(10):3636–3640
9. Krischer JP, Lynch KF, Lemmark Å et al (2017) Genetic and environmental interactions modify the risk of diabetes-related autoimmunity by 6 years of age: The TEDDY Study. *Diabetes Care* 40(9):1194–1202
10. Wegienka G, Zoratti E, Johnson CC (2015) The role of the early-life environment in the development of allergic disease. *Immunol Allergy Clin N Am* 35(1):1–17
11. Virtanen SM (2016) Dietary factors in the development of type 1 diabetes. *Pediatr Diabetes* 17:S49–S55
12. Osborn DA, Sinn JKH, Jones LJ (2018) Infant formulas containing hydrolysed protein for prevention of allergic disease. *Cochrane Database Syst Rev* 10:CD003664
13. Knip M, Åkerblom HK, Becker D et al (2014) Hydrolyzed infant formula and early β -cell autoimmunity: a randomized clinical trial. *JAMA* 311(22):2279–2287
14. Greer FR, Sicherer SH, Burks AW, Committee on nutrition, section on allergy and immunology (2019) The effects of early nutritional interventions on the development of atopic disease in infants and children: the role of maternal dietary restriction, breastfeeding, hydrolyzed formulas, and timing of introduction of allergenic complementary foods. *Pediatrics* 143(4):e20190281
15. The TRIGR Study Group (2007) Study design of the Trial to Reduce IDDM in the Genetically at Risk (TRIGR). *Pediatr Diabetes* 8(3):117–137
16. Åkerblom HK, Knip M, Becker D et al (2007) The TRIGR Study Group. The TRIGR trial: testing the potential link between weaning diet and type 1 diabetes. *Immun Endocr Metab Agents Med Chem* 7(3): 251–263
17. The TRIGR Study Group (2011) The Trial to Reduce IDDM in the Genetically at Risk (TRIGR) study: recruitment, intervention and follow-up. *Diabetologia* 54:627–633
18. Sorkio S, Cuthbertson D, Bärlund S et al (2010) Breastfeeding patterns of mothers with type 1 diabetes: results from an infant feeding trial. *Diabetes/Metab Res Rev* 26(3):206–211
19. Knip M, Virtanen SM, Seppä K et al (2010) Dietary intervention in infancy and later signs of beta-cell autoimmunity. *N Engl J Med* 363(20):1900–1908
20. ISAAC (2012) ISAAC Phase Three Data. Available from <http://isaac.auckland.ac.nz/phases/phasethree/results/results.php>. Accessed 1 Apr 2012
21. Prescott SL, Tang ML (2005) The Australasian Society of Clinical Immunology and Allergy position statement: summary of allergy prevention in children. *Med J Aust* 182(9):464–467
22. Flöistrup H, Swartz J, Bergström A et al (2006) Allergic disease and sensitization in Steiner school children. *J Allergy Clin Immunol* 117(1):59–66
23. von Hertzen L, Haahtela T (2005) Signs of reversing trends in prevalence of asthma. *Allergy* 60(3):283–292
24. Cardwell CR, Shields MD, Carson DJ, Patterson CC (2003) A meta-analysis of the association between childhood type 1 diabetes and atopic disease. *Diabetes Care* 26(9):2568–2574. <https://doi.org/10.2337/diacare.26.9.2568>
25. Hörtenhuber T, Kiess W, Fröhlich-Reiterer E et al (2018) Asthma in children and adolescents with type 1 diabetes in Germany and Austria: Frequency and metabolic control. *Pediatr Diabetes* 19(4): 727–732. <https://doi.org/10.1111/pedi.12618>
26. Black MH, Anderson A, Bell RA et al (2011) Prevalence of asthma and its association with glycemic control among youth with diabetes. *Pediatrics* 128(4):e839–e847. <https://doi.org/10.1542/peds.2010-3636>
27. Metsälä J, Lundqvist A, Virta LJ et al (2018) The association between asthma and type 1 diabetes: a paediatric case-cohort study in Finland, years 1981–2009. *Int J Epidemiol* 47(2):409–416
28. Wahlberg J, Vaarala O, Ludvigsson J, for the ABIS Study Group (2011) Asthma and allergic symptoms and type 1 diabetes-related autoantibodies in 2.5-yr-old children. *Pediatr Diabetes* 12(7):604–610

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Affiliations

Jeffrey P. Krischer¹ · David Cuthbertson¹ · Marisa Couluris² · Mikael Knip^{3,4} · Suvi M. Virtanen^{5,6,7,8}

¹ Health Informatics Institute, Morsani College of Medicine, University of South Florida, 3650 Spectrum Boulevard, Suite 100, Tampa, FL 33612, USA

² Department of Pediatrics, Morsani College of Medicine, University of South Florida, Tampa, FL, USA

³ Children's Hospital, Pediatric Research Center, University of Helsinki and Helsinki University Hospital, Helsinki, Finland

⁴ Research Program for Clinical and Molecular Metabolism, Faculty of Medicine, University of Helsinki, Helsinki, Finland

⁵ Unit of Public Health Promotion, National Institute for Health and Welfare, Helsinki, Finland

⁶ Faculty of Social Sciences/Health Sciences, Tampere University, Tampere, Finland

⁷ Research, Development and Innovation Center, Tampere University Hospital, Tampere, Finland

⁸ Tampere Center for Child Health Research, Tampere University Hospital, Tampere, Finland