Correction to: Acute hyperglycaemia does not have a consistent adverse effect on exercise performance in recreationally active young people with type 1 diabetes: a randomised crossover in-clinic study

Karen M. Rothacker^{1,2} • Sam Armstrong^{1,2} • Grant J. Smith² • Nat Benjanuvatra³ • Brendan Lay³ • Peter Adolfsson^{4,5} • Timothy W. Jones^{1,2,6} • Paul A. Fournier³ • Elizabeth A. Davis^{1,2,6}

Published online: 11 June 2021

© Crown 2021

Correction to: Diabetologia.

<https://doi.org/10.1007/s00125-021-05465-9>

Unfortunately, one of the test conditions (Hyper 5) was incorrectly stated as ‘hyperglycaemia (17 mmol/l) with mU [5 m² BSA]^{−1} min^{−1} insulin’ instead of ‘hyperglycaemia (17 mmol/l) with 5 mU [m² BSA]^{−1} min^{−1} insulin’ in the Abstract. The original article has been corrected.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The online version of the original article can be found at <https://doi.org/10.1007/s00125-021-05465-9>

✉ Karen M. Rothacker
karen.rothacker@health.wa.gov.au

- ¹ Department of Endocrinology and Diabetes, Perth Children's Hospital, Nedlands, Western Australia, Australia
- ² The Centre for Child Health Research, Telethon Kids Institute, The University of Western Australia, Subiaco, Western Australia, Australia
- ³ School of Human Sciences, Division of Sport Science, Exercise and Health, The University of Western Australia, Crawley, Western Australia, Australia
- ⁴ Department of Pediatrics, The Hospital of Halland, Kungälv, Sweden
- ⁵ Institute of Clinical Sciences, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden
- ⁶ School of Paediatrics and Child Health, The University of Western Australia, Crawley, Western Australia, Australia