



Correction to: Progression likelihood score identifies substages of presymptomatic type 1 diabetes in childhood public health screening

Andreas Weiss^{1,2} · Jose Zapardiel-Gonzalo^{1,2} · Franziska Voss¹ · Manja Jolink¹ · Joanna Stock¹ · Florian Haupt^{1,2,3} · Kerstin Kick⁴ · Tiziana Welzhofer⁴ · Anja Heublein¹ · Christiane Winkler^{1,2,3} · Peter Achenbach^{1,2,3,4} · Anette-Gabriele Ziegler^{1,2,3,4} · Ezio Bonifacio^{2,5,6} · for the Fr1da-study group

Published online: 4 October 2022

© Springer-Verlag GmbH Germany, part of Springer Nature 2022

Correction to: Diabetologia

<https://doi.org/10.1007/s00125-022-05780-9>

In the Conclusions/interpretation section of the Abstract, the following, potentially misleading, text has been changed:

Original wording: ‘Public health screening for islet autoantibodies found 0.027% of children to have undiagnosed clinical type 1 diabetes and 0.038% to have undiagnosed presymptomatic type 1 diabetes. Of the latter group, our novel likelihood ratio score predicted approximately 50% to be at risk of developing clinical type 1 diabetes within 2 years.’

Modified wording: ‘Public health screening for islet autoantibodies found 0.027% of children to have undiagnosed clinical type 1 diabetes and 0.038% to have undiagnosed presymptomatic stage 2 or stage 1b type 1 diabetes, with 50% risk to develop clinical type 1 diabetes within 2 years.’

The original article has been corrected.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The online version of the original article can be found at <https://doi.org/10.1007/s00125-022-05780-9>

✉ Anette-Gabriele Ziegler
anette-g.ziegler@helmholtz-muenchen.de

¹ Institute of Diabetes Research, Helmholtz Munich, German Research Center for Environmental Health, Munich, Germany

² German Center for Diabetes Research (DZD), Munich, Germany

³ Forschergruppe Diabetes e.V. at Helmholtz Zentrum München, Munich, Germany

⁴ Technical University Munich, School of Medicine, Forschergruppe Diabetes at Klinikum rechts der Isar, Munich, Germany

⁵ Center for Regenerative Therapies Dresden, Faculty of Medicine, Technische Universität Dresden, Dresden, Germany

⁶ Paul Langerhans Institute Dresden of Helmholtz Centre Munich at University Clinic Carl Gustav Carus of TU Dresden, Faculty of Medicine, Dresden, Germany