

Erratum to: Evidence of sex differences in the relationship between current tobacco use and past-year serious psychological distress: 2005–2008 National Survey on Drug Use and Health

Nicholas Peiper · Brad Rodu

Published online: 1 April 2014
© Springer-Verlag Berlin Heidelberg 2014

Erratum to: Soc Psychiatry Psychiatr Epidemiol (2013) 48:1261–1271
DOI 10.1007/s00127-012-0644-0

In the original publication of this article, there was an error in the reporting of the weighted prevalence of past-year psychiatric morbidity among females in Table 4 and a formatting issue related to the adjusted odds ratios and 95 % confidence intervals for past-year serious psychological distress among male smokers in Table 5. The corrected versions of Tables 4 and 5 are shown below.

Table 4 Weighted prevalence of past-year psychiatric morbidity among adults age 18+ years, National Survey on Drug Use and Health 2005–2008

Psychiatric morbidity	Never user % (SE)	Lifetime user % (SE)	Current tobacco use ^a		
			Cigarettes % (SE)	Smokeless % (SE)	Dual % (SE)
All adults					
SPD	9.1 (0.2)	8.2 (0.2)	18.1 (0.3)	7.0 (0.6)	14.4 (1.0)
MDE	5.7 (0.2)	6.0 (0.2)	11.4 (0.3)	4.3 (0.6)	9.1 (0.9)
AD	3.4 (0.2)	4.0 (0.2)	7.7 (0.2)	2.8 (0.5)	5.8 (0.7)
SUD	3.2 (0.1)	8.2 (0.2)	18.3 (0.3)	14.5 (0.9)	36.3 (1.5)
Males					
SPD	7.3 (0.3)	5.9 (0.2)	12.8 (0.4)	6.8 (0.6)	13.3 (1.0)
MDE	3.7 (0.3)	4.1 (0.2)	7.7 (0.4)	4.4 (0.6)	8.5 (0.9)
AD	1.8 (0.2)	2.6 (0.2)	4.3 (0.2)	2.8 (0.5)	5.2 (0.7)
SUD	5.0 (0.3)	9.5 (0.3)	22.2 (0.5)	14.9 (0.9)	36.5 (1.5)
Females					
SPD	9.8 (0.3)	11.7 (0.4)	24.0 (0.5)	9.5 (2.5)	39.7 (6.7)
MDE	6.5 (0.2)	9.0 (0.3)	15.4 (0.4)	3.2 (1.5)	22.5 (6.3)
AD	4.1 (0.2)	6.1 (0.3)	11.5 (0.4)	3.3 (1.8)	17.3 (5.1)
SUD	2.4 (0.1)	6.1 (0.3)	14.0 (0.4)	8.9 (2.3)	33.0 (5.2)

SE standard error, SPD serious psychological distress, MDE major depressive episode, AD any anxiety disorder, SUD substance use disorder

^a All values expressed as weighted column percentage (SE)

The online version of the original article can be found under doi:10.1007/s00127-012-0644-0.

N. Peiper (✉)
Department of Epidemiology and Population Health,
School of Public Health and Information Sciences,
University of Louisville, 485 East Gray Street,
Louisville, KY 40202, USA
e-mail: npeip01@louisville.edu

B. Rodu
Department of Medicine, University of Louisville,
Louisville, KY, USA

Table 5 Adjusted prevalence odds of past-year serious psychological distress and psychiatric morbidity among adults age 18+ years, National Survey on Drug Use and Health 2005–2008

Psychiatric morbidity	Never user OR (95 % CI)	Lifetime user OR (95 % CI)	Current tobacco use ^a		
			Cigarettes OR (95 % CI)	Smokeless OR (95 % CI)	Dual OR (95 % CI)
All adults ^a					
SPD [†]	1.00	0.99 (0.91–1.08)	1.45 (1.33–1.59)	0.86 (0.67–1.08)	1.05 (0.84–1.32)
MDE	1.00	1.18 (1.08–1.30)	1.37 (1.23–1.52)	0.90 (0.66–1.22)	1.25 (0.96–1.62)
AD	1.00	1.25 (1.09–1.43)	1.74 (1.52–2.01)	1.19 (0.82–1.73)	1.77 (1.27–2.47)
Males ^b					
SPD	1.00	0.82 (0.72–0.93)	1.14 (0.99–1.31)	0.71 (0.55–0.91)	0.86 (0.65–1.14)
MDE	1.00	1.05 (0.89–1.26)	1.21 (0.99–1.48)	0.88 (0.66–1.19)	1.15 (0.82–1.63)
AD	1.00	1.26 (0.97–1.64)	1.53 (1.17–2.00)	1.21 (0.79–1.87)	1.73 (1.13–2.65)
Females ^c					
SPD	1.00	1.05 (0.93–1.19)	1.60 (1.45–1.77)	0.95 (0.51–1.77)	2.24 (1.10–4.58)
MDE	1.00	1.23 (1.10–1.38)	1.43 (1.26–1.63)	0.38 (0.13–1.10)	1.68 (0.71–3.98)
AD	1.00	1.22 (1.04–1.43)	1.83 (1.56–2.14)	0.99 (0.35–2.81)	2.76 (1.09–7.00)

SPD serious psychological distress, MDE major depressive episode, AD any anxiety disorder, OR prevalence odds ratio, 95 % CI 95 % confidence interval

[†] *P* value for sex interaction less than 0.05

^a Adjusted models control for age, sex, race/ethnicity, education, total family income, marital status, population density, survey year, and past-year diabetes, cardiovascular disease, respiratory disease, psychiatric comorbidity, and substance use disorders

^b Male models include age, race/ethnicity, education, total family income, marital status, population density, survey year, and past-year diabetes, cardiovascular disease, respiratory disease, psychiatric comorbidity, and substance use disorders

^c Female models include all covariates from male models plus past-year pregnancy