Incidence of major depression diagnoses in the Canadian Armed Forces: longitudinal analysis of clinical and health administrative data

François L. Thériault^{1,2} · R. A. Hawes¹ · B. G. Garber^{1,2} · F. Momoli^{2,3,5} · W. Gardner^{2,3} · M. A. Zamorski^{1,4} · I. Colman²

Received: 18 January 2019 / Accepted: 20 August 2019 / Published online: 26 September 2019
© The Author(s) 2019

Abstract

Purpose Major depression is a leading cause of morbidity in military populations. However, due to a lack of longitudinal data, little is known about the rate at which military personnel experience the onset of new episodes of major depression. We used a new source of clinical and administrative data to estimate the incidence of major depression diagnoses in Canadian Armed Forces (CAF) personnel, and to compare incidence rates between demographic and occupational factors.

Methods We extracted all data recorded in the electronic medical records of CAF Regular Force personnel, at every primary care and mental health clinical encounter since 2016. Using a 12-month lookback period, we linked data over time, and identified all patients with incident diagnoses of major depression. We then linked clinical data to CAF administrative records, and estimated incidence rates. We used multivariate Poisson regression to compare adjusted incidence rates between demographic and occupational factors.

Results From January to December 2017, CAF Regular Force personnel were diagnosed with major depression at a rate of 29.2 new cases per 1000 person-years at risk. Female sex, age 30 years and older, and non-officer ranks were associated with significantly higher incidence rates.

Conclusions We completed the largest study to date on diagnoses of major depression in the Canadian military, and have provided the first estimates of incidence rates in CAF personnel. Our results can inform future mental health resource allocation, and ongoing major depression prevention efforts within the Canadian Armed Forces and other military organizations.

Keywords Depression · Military · Epidemiology · Risk · ICD code · DSM code

Introduction

Major depression is a recurring disorder that disproportionately affects military personnel. In Australia, Canada, the United Kingdom, and the United States, the prevalence of major depression in military personnel is at least twice that

in comparable civilian populations [1–4]. Depressed soldiers are at increased risk of suicidal behaviors [5] poor job performance and absenteeism [6], and early separation from the military [7]. Because of its high prevalence and deleterious impact, major depression is a public health concern for military organizations. Epidemiological data are, therefore, needed to inform depression prevention strategies, and optimize the allocation of military mental health resources.

Until now, cross-sectional surveys have been the main source of epidemiological data on major depression in military personnel. These surveys have provided valuable information on the prevalence of major depression across different military organizations [1, 3, 8]. They have also shed new light on the occupational factors associated with major depression in military personnel [9] and on stigma perceptions [10] and care-seeking behaviors [11] related to this condition. However, cross-sectional surveys cannot provide information on the rate of major depression incidence.

✉ François L. Thériault
francois.theriault3@forces.gc.ca

¹ Canadian Forces Health Services Group, Department of National Defence, Ottawa, Canada

² School of Epidemiology and Public Health, University of Ottawa, Ottawa, Canada

³ Children's Hospital of Eastern Ontario Research Institute, Ottawa, Canada

⁴ Department of Family Medicine, University of Ottawa, Ottawa, Canada

⁵ Ottawa Hospital Research Institute, Ottawa, Canada

the professional authority to assign formal diagnoses (e.g., triage nurses, social workers, etc.) must record at least one symptom to complete their notes (Fig. 1). These symptoms are selected from a drop-down list of standardized clinical impressions.

External specialists do not have access to CAF electronic medical records. But to be reimbursed for services provided to externally referred patients, external specialists must send a paper copy of their clinical notes to the local CAF clinic. These notes are then scanned into the CAF electronic medical records of externally referred patients. A CAF health-care provider—usually a nurse or social worker by training—periodically reviews the clinical progress of externally referred patients by completing an electronic note. External Provider Review notes are part of the CAF electronic medical record, and must contain at least one symptom code (Fig. 1).

Case definition

We sought to identify all personnel with an incident diagnosis of major depression, from January to December 2017. To do so, we extracted all diagnostic and symptom codes entered in CAF electronic medical records between January 2016 and June 2018.

We considered that personnel with at least one diagnostic code for major depressive episode (F32.x) or major depressive disorder (F33.x) in their electronic medical records were patients with major depression, irrespective of co-morbid psychiatric conditions. For all patients with major depression, we identified all clinical encounters related to major depression care as either: (1) clinical visits during which a diagnostic code for major depression was recorded, or (2) clinical visits that occurred within 6 months of a major depression diagnosis, and during which a code for “depressive symptoms” was recorded. Consistent with other studies [14–16], incident diagnoses were defined as the first clinical encounter for major depression in a 12-month period. With data available from January 2016 to June 2018, and with the case definition presented above, we were able to report on incident diagnoses from January to December 2017.

Patients with major depression who are exclusively assessed and treated by external specialists may never generate a depression code in CAF electronic medical records (Fig. 1). We, therefore, reviewed the scanned clinical notes of all externally referred patients with a “depressive symptom” code recorded on an External Provider Review note between January 2016 and June 2018 ($n = 444$); we, thus, identified an additional 116 patients who had been diagnosed with major depression over this time period. For these 116 patients, the incidence date was defined as the first encounter with the external specialist.

Demographic data

CAF administrative databases were used to determine the sex, age, rank, military unit, military trade, and duration of enlistment, for all CAF Regular Force personnel, from January to December 2017. Age was categorized into four groups: (a) 18–29 years, (b) 30–39 years, (c) 40–49 years, and (d) 50–60 years. Ranks were dichotomized as Non-Commissioned Members (NCM) and Officers.

Occupational data

Military units were linked to one of the six commands: (a) the Canadian Army, (b) the Royal Canadian Air Force, (c) the Royal Canadian Navy, (d) Military Personnel Command, (e) Joint and Special Forces, and (f) other central agencies. Military Personnel Command includes Joint Personnel Support Units (JPSU) that are administrative cells where ill or injured personnel from any command may be posted, if a medical condition precludes them from performing regular duties for a period of at least 6 months. JPSU personnel are expected to have much higher morbidity than other members of Military Personnel Command; we, therefore, reported on their incidence rates separately.

Military trades were linked to one of the ten occupational groups, as outlined on the CAF recruitment website (www.forces.ca): (a) technicians, (b) combat specialists, (c) administration and support, (d) engineers, (e) telecommunications, (f) air and ship crew, (g) health care, (h) sensor and radar, (i) public protection, including military police and firefighters, and (g) other.

Statistical analyses

For the estimation of incidence rates, we excluded all Regular Force personnel with any clinical encounter for major depression from January to December 2016. We then computed population-level counts of person-years at risk of an incident major depression diagnosis in remaining Regular Force personnel, from January to December 2017. We calculated the incidence of new major depression diagnoses per person-years at risk. We treated demographic categories (i.e., age group, rank category, command, and occupational group) as time-varying covariates, and used multivariate Poisson regression models to compare incidence rates across patient characteristics, adjusting for sex, age, and rank. We entered the log of person-years at risk as an offset in all multivariate Poisson analyses.

All analyses were performed using Stata, version 14.0 [17].

Secondary analysis: assessment of data source agreement

As recommended in current guidelines [13], we also sought to assess the validity of our case definition. Data source agreement is an accepted method for assessing the quality of case definitions in health administrative data research [18], and a large cross-sectional survey recently estimated that 6.0% of Regular Force personnel with major depression used mental health services in the previous year [11]. In an effort to assess the quality of data used in the current study, we identified all individuals who were serving in the Regular Force on the last day of our study period (i.e., December 31st 2017), and used our case definition to determine whether they had any clinical encounters for major depression at any point from January to December 2017. We qualitatively compared the sex and age distribution of patients with past-year clinical encounters for major depression to the results of the aforementioned survey [11].

Ethical considerations

The study was approved by the CAF Director of Force Health Protection, the CAF Director of Mental Health, and the University of Ottawa Office of Research Ethics and Integrity.

Results

A total of 72,054 individuals served in the CAF Regular Force at some point in 2017. We excluded 3638 (5.1%) of these individuals from our study because they had at least one diagnosis of major depression recorded in the 12 months preceding the study start date. The remaining 68,416 individuals contributed 61,997 service-years at risk of an incident major depression diagnosis during the study period; most of this person-time was contributed by males (86.0%) under the age of 40 years (69.5%) (Table 1).

During the study period, 1812 incident diagnoses of major depression were observed. Regular Force personnel were diagnosed at a rate of 29.2 new cases per 1000 person-years at risk (95% CI 27.9, 30.6). Female sex, age 30 years and older, and non-officer ranks were associated with higher incidence rates. After adjusting for demographic differences, technicians, and air and ship crewmembers, had significantly lower incidence rates than combat specialists. Adjusted rates were also lower in the Air Force than in the Army. Individuals posted to Joint Personnel Support Units, who were already ill or injured,

Table 1 Person-years at risk of an incident major depression diagnosis, CAF Regular Force personnel, January–December 2017

	Person-years	(%)
Sex		
Female	8698	(14.0)
Male	53,299	(86.0)
Age (years)		
18–29	21,708	(35.0)
30–39	21,400	(34.5)
40–49	12,877	(20.8)
50–60	6012	(9.7)
Rank		
Non-commissioned members	46,337	(74.7)
Officers	15,660	(25.3)
Command		
Canadian Army	21,952	(35.4)
Royal Canadian Air Force	12,779	(20.6)
Military Personnel Command	8288	(13.4)
Royal Canadian Navy	8036	(13.0)
Joint and Special Forces	3270	(5.3)
Joint Personnel Support Units	907	(1.5)
Other ^a	6766	(10.9)
Military Trade		
Technicians	13,907	(22.4)
Combat Specialists	11,835	(19.1)
Administration and Support	10,731	(17.3)
Engineers	6146	(9.9)
Telecommunications	4742	(7.6)
Air and Ship Crew	4711	(7.6)
Health Care	2812	(4.5)
Sensor and Radar	2463	(4.0)
Public Protection	1876	(3.0)
Other	2774	(4.5)

^aIncludes personnel posted to central agencies, including the offices of the Minister and Associate Minister of National Defence, and of the Chief and Vice-Chief of the Defence Staff

were nearly three times more likely to have an incident diagnosis of major depression, compared to Army personnel of the same sex, age, and rank (Table 2).

For our secondary analysis, we identified all 66,605 individuals who were serving in the Regular Force on the last day of the 12-month study period. Using the case definition developed for our study, we estimated that 5.6% (95% CI 5.4%, 5.8%) of these individuals had a clinical encounter for major depression diagnosis at some point in the previous 12 months. Among these patients, 24.5% (95% CI 23.1, 25.9) were females, and 18.5% (95% CI 17.3%, 19.8%) were under the age of 30 years (Table 3). These results can be compared to those of a recent cross-sectional survey (see discussion below).

Table 2 Incidence of new major depression diagnoses in CAF Regular Force personnel, January–December 2017

	Rate per 1000 p–y	(95% CI)	aIRR ^b	(95% CI)	<i>p</i> value
Sex					
Females	44.2	(39.7, 48.6)	Ref		
Males	26.8	(25.4, 28.2)	0.61	(0.54, 0.68)	< 0.001
Age (years)					
18–29	20.7	(18.8, 22.6)	Ref		
30–39	31.5	(29.1, 33.9)	1.51	(1.34, 1.70)	< 0.001
40–49	39.2	(35.8, 42.6)	1.92	(1.69, 2.18)	< 0.001
50–60	30.4	(26.0, 34.8)	1.56	(1.32, 1.86)	< 0.001
Rank					
Non-commissioned members	32.5	(30.8, 34.1)	Ref		
Officers	19.6	(17.4, 21.8)	0.57	(0.50, 0.64)	< 0.001
Command					
Canadian Army	27.5	(25.3, 29.7)	Ref		
Royal Canadian Air Force	24.4	(21.7, 27.1)	0.85	(0.74, 0.98)	0.025
Military Personnel Command	30.8	(27.0, 34.5)	1.14	(0.98, 1.33)	0.087
Royal Canadian Navy	28.7	(25.0, 32.5)	0.99	(0.85, 1.15)	0.899
Joint and Special Forces	31.5	(25.4, 37.6)	1.07	(0.87, 1.32)	0.522
Joint Personnel Support Units	97.0	(76.7, 117.3)	2.90	(2.31, 3.64)	< 0.001
Other ^a	32.4	(28.1, 36.7)	1.12	(0.96, 1.32)	0.157
Military trade					
Combat Specialists	27.4	(24.4, 30.4)	Ref		
Technicians	26.1	(23.4, 28.8)	0.80	(0.69, 0.93)	0.004
Administration and Support	41.1	(37.3, 44.9)	1.12	(0.96, 1.31)	0.151
Engineers	22.1	(18.4, 25.8)	0.89	(0.73, 1.10)	0.286
Telecommunications	33.7	(28.5, 39.0)	1.06	(0.88, 1.28)	0.539
Air and Ship Crew	14.9	(11.4, 18.3)	0.66	(0.51, 0.87)	0.003
Health Care	35.2	(28.3, 42.1)	1.06	(0.84, 1.35)	0.614
Sensor and Radar	30.4	(23.6, 37.3)	0.97	(0.76, 1.25)	0.827
Public Protection	33.0	(24.8, 41.3)	1.00	(0.76, 1.31)	0.997
Other	29.6	(23.2, 36.0)	0.95	(0.74, 1.21)	0.652

^aIncludes personnel posted to central agencies, including the offices of the Minister and Associate Minister of National Defence, and of the Chief and Vice-Chief of the Defence Staff

^bIncidence rate ratio, adjusted for sex, age group, and rank category

Discussion

Table 3 Characteristics of CAF Regular Force personnel with incident or prevalent major depression diagnoses in the previous 12 months, as of December 2017

	Percent	(95% CI)
Sex		
Female	24.5	(23.1, 25.9)
Male	75.5	(74.1, 76.9)
Age (years)		
18–29	18.5	(17.3, 19.8)
30–39	38.6	(37.0, 40.2)
40–49	31.1	(29.7, 32.6)
50–60	11.8	(10.8, 12.9)

This paper presents the most comprehensive examination to date of major depression incidence in serving CAF Regular Force personnel. We found that major depression amongst CAF personnel was diagnosed at a rate of 29.2 new cases per 1000 person-years of service, and that this rate varied considerably across demographic and organizational characteristics.

The overall incidence rate reported in this study is somewhat higher than that observed in the US Armed Forces, where 24.2 new cases of depressive disorders are diagnosed per 1000 person-years [19]. Several factors may account for this difference. First, CAF personnel suffering from major depression may be more likely to seek care

than their American counterparts, and may, therefore, be more likely to receive a formal diagnosis; previous studies have found that 75% of depressed Canadian soldiers access mental health care in a 12-month period [11], compared to less than 50% in the US military [20–22]. This apparent difference in care-seeking propensity may be partly explained by the substantial difference in age composition between the two militaries; on average, CAF personnel are older than their American counterparts [23, 24], and older age is associated with higher mental health service use in depressed patients [25]. Second, our case-finding algorithm was likely more inclusive than that used for the US military study. In fact, US Armed Forces researchers only considered major depression diagnoses if they were recorded in the first or second diagnostic position during medical encounters [19]. In contrast, we used any major depression diagnosis to identify incident cases in our study, even if these diagnoses were recorded along with several co-morbid conditions.

We found that female sex, age 30 years and older, and non-officer ranks were associated with higher incidence of major depression diagnoses. These findings largely mirror those of previous studies. Indeed, a large body of research has already shown that, compared to males, females are more likely to suffer from major depression [26], and are more likely to seek care when experiencing depressive symptoms [25]. It is, therefore, not surprising to find that females were more likely to be diagnosed with major depression, compared to their male counterparts. Similarly, non-officer rank [27] and older age [25] have also been associated with greater likelihood of seeking mental healthcare, when needed.

Until now, researchers have depended on data from either population surveys (e.g., [28]) or manual reviews of medical charts (e.g., [29]) to investigate mental health care utilization in CAF personnel, and have lacked the statistical power required to compare occupational groups with the level of granularity reported in our study. With our large sample size, we found that combat specialists had a greater incidence of major depression than technicians, and air and ship crewmembers. However, incidence rates in combat specialists were not significantly different than in numerous other non-combat trades. This finding has important implications. Although combat experiences are associated with a substantial increase in depression risk [30], they only account for a small proportion of the major depression burden observed in Canada's military [31]. Similarly, most American soldiers who attempt suicide have never deployed [5]. Since the early 2000s, the CAF have improved access to mental health care, implemented mental health training programs across the organization, and provided post-deployment mental health screening to all personnel returning from an overseas deployment, irrespective of their specific occupations [32].

Our findings support this holistic strategy, in the context of major depression prevention.

After excluding Joint Personnel Support Units (JPSU, i.e., service centers where seriously ill or injured personnel are posted if a medical condition precludes them from performing regular military duties for a period of at least 6 months) from Military Personnel Command, we found no large difference in the incidence rate of major depression diagnoses between the six CAF commands. Of note, JPSU personnel were nearly three times more likely to be diagnosed with major depression compared to any other personnel, even after using a 12-month lookback period to exclude prevalent cases. This elevated incidence rate is, therefore, not a consequence of personnel with prevalent major depression diagnoses being transferred to a JPSU. Rather, our finding suggests that individuals are at particularly high risk of being newly diagnosed with major depression after being transferred to a JPSU. This could be explained by several factors. First, individuals who are posted to a JPSU without a prevalent major depression diagnosis must have another physical or mental health condition. These pre-existing conditions could be strong risk factors for major depression onset [33]. Second, individuals who are separating from the military on medical grounds are often posted to a JPSU, and may be more willing to disclose their depressive symptoms. Fear of negative career repercussions is commonly reported as a barrier to mental health care among military personnel [34], but that barrier is unlikely to impact individuals already in the process of releasing from military service. Third, the process of transitioning out of an operational unit and into a JPSU may represent a stressful life event for some individuals which may, therefore, increase their risk of major depression onset [35]. In any case, our findings suggest that military personnel posted to administrative units such as JPSUs may benefit from counseling and other enhanced mental health prevention efforts.

When used to assess past-year prevalence of clinical encounters related to major depression, our case definition produced results that were highly consistent with previous research on CAF personnel. Indeed, a large population-based survey conducted in 2013 found that 6.0% of all CAF Regular Force personnel had sought care for major depression in the previous 12 months [11]; this estimate is remarkably close to the 5.6% past-year prevalence of diagnosed major depression reported herein. The estimated age and sex distribution of diagnosed patients was also similar between studies. Such a high level of data source agreement supports the quality of our data, and the validity of the case definitions used in our study.

Our study had noteworthy limitations. First, not all depressed individuals seek care. In fact, a recent study found that only 75% of CAF Regular Force personnel who met diagnostic criteria for major depression at any point

in the previous year had sought professional mental health care over the same time period [11]. While this treatment gap is much smaller than in other populations [36, 37], 25% of depressed personnel remain undiagnosed. Clinical data can only be used to identify major depression in care-seeking individuals. Our study, therefore, misclassified a substantial number of depressed patients, and underestimated the true incidence of major depression in the CAF Regular Force. While our results cannot be used to measure major depression morbidity in the Canadian military, they nonetheless reflect current mental health service utilization rates for this condition.

Second, we relied on diagnostic codes extracted from electronic medical records to identify depressed patients. Previous authors have argued that diagnostic codes can be inaccurate; psychiatrists may incorrectly code their patients' diagnoses, and may also code only one of the several co-morbid conditions [38]. Depressed patients may, therefore, not generate a diagnostic code for major depression every time they encounter a health professional. We mitigated this risk of misclassification bias by creating a case-finding algorithm, and linking data elements over time. Although we did not validate our case-finding algorithm against a gold-standard measure of major depression, our estimates of past-year treatment prevalence were highly consistent with those obtained from a 2013 population-based survey [11], as discussed above. As already noted, data source agreement is an accepted method for assessing the quality of case-finding algorithms using routinely collected data sources [18].

Despite the limitations listed above, we completed the largest study to date on major depression diagnoses in the Canadian military, and provided the first estimates of incident rates in CAF personnel. Our results have highlighted that the rates of major depression diagnoses vary substantially across sex, age, and rank categories, but are very similar between combat and non-combat trades. Our findings make an important contribution to military population health surveillance, and may be used by military organizations to better prepare for their future mental healthcare needs. For example, our estimated incidence rates could be used in predictive models to forecast the expected number of new major depression cases in a military population, given the age, sex, and occupational profile of its personnel. Furthermore, our study clearly demonstrated the feasibility of using electronic medical records as a powerful tool to study the incidence of mental disorder diagnoses in military populations.

Acknowledgements The authors wish to thank the dedicated healthcare providers, business analysts, and programmers without whom data used for this study could not have been extracted from CAF electronic medical records. This research was undertaken, in part, thanks to funding from the Canada Research Chairs program for IC, and from the Surgeon General's Health Research Program for FLT.

Compliance with ethical standards

Conflict of interest On behalf of all authors, the corresponding author states that there is no conflict of interest.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

1. McFarlane A, Sriapada R, Ganoczy D, et al (2011) Mental health in the Australian Defence Force: 2010 ADF mental health and wellbeing study: Full report. Canberra
2. Rusu C, Zamorski MA, Boulos D, Garber BG (2016) Prevalence comparison of past-year mental disorders and suicidal behaviours in the Canadian Armed Forces and the Canadian General Population. *Can J Psychiatry* 61:46S–55S. <https://doi.org/10.1177/0706743716628856>
3. Goodwin L, Wessely S, Hotopf M et al (2015) Are common mental disorders more prevalent in the UK serving military compared to the general working population? *Psychol Med* 45:1881–1891. <https://doi.org/10.1017/S0033291714002980>
4. Gadermann AM, Engel CC, Naifeh JA et al (2012) Prevalence of DSM-IV major depression among U.S. military personnel: meta-analysis and simulation. *Mil Med* 177:47–59
5. Ursano RJ, Kessler RC, Stein MB et al (2016) Risk factors, methods, and timing of suicide attempts among US Army Soldiers. *JAMA psychiatry* 73:741–749. <https://doi.org/10.1001/jamapsychiatry.2016.0600>
6. Pflanz SE, Ogle AD (2006) Job stress, depression, work performance, and perceptions of supervisors in military personnel. *Mil Med* 171:861–865. <https://doi.org/10.7205/MILMED.171.9.861>
7. Littman AJ, Jacobson IG, Boyko EJ et al (2013) Weight change following US military service. *Int J Obes* 37:244–253. <https://doi.org/10.1038/ijo.2012.46>
8. Zamorski MA, Bennett RE, Rusu C et al (2016) Prevalence of past-year mental disorders in the Canadian Armed Forces, 2002–2013. *Can J Psychiatry* 61:26S–35S. <https://doi.org/10.1177/0706743716628854>
9. Nelson C, Cyr KS, Corbett B et al (2011) Predictors of post-traumatic stress disorder, depression, and suicidal ideation among Canadian Forces personnel in a National Canadian Military Health Survey. *J Psychiatr Res* 45:1483–1488. <https://doi.org/10.1016/j.jpsychires.2011.06.014>
10. Weeks M, Zamorski MA, Rusu C, Colman I (2017) Mental illness-related stigma in Canadian military and civilian populations: a comparison using population health survey data. *Psychiatr Serv* 68:710–716. <https://doi.org/10.1176/appi.ps.201600398>
11. Thériault FL, Garber BG, Momoli F et al (2019) Mental Health Service utilization in depressed Canadian armed forces personnel. *Can J Psychiatry* 64:59–67. <https://doi.org/10.1177/0706743718787792>
12. Tannenbaum C, Lexchin J, Tamblyn R, Romans S (2009) Indicators for measuring mental health: towards better surveillance. *Healthc Policy* 5:e177–e186
13. Benchimol EI, Smeeth L, Guttman A et al (2015) The REporting of studies Conducted using Observational Routinely-collected

- health Data (RECORD) Statement. PLoS Med. <https://doi.org/10.1371/JOURNAL.PMED.1001885>
14. Spettell CM, Wall TC, Allison J et al (2003) Identifying physician-recognized depression from administrative data: consequences for quality measurement. *Health Serv Res* 38:1081–1102
 15. Yoon G, Petrakis IL, Rosenheck RA (2015) Correlates of major depressive disorder with and without comorbid alcohol use disorder nationally in the veterans health administration. *Am J Addict* 24:419–426. <https://doi.org/10.1111/ajad.12219>
 16. Shrestha A, Cornum BGR, Vie LL et al (2018) Protective effects of psychological strengths against psychiatric disorders among soldiers. *Mil Med* 183:386–395. <https://doi.org/10.1093/milmed/usx189>
 17. StataCorp (2015) Stata Statistical Software: Release 14
 18. Weiskopf NG, Weng C (2013) Methods and dimensions of electronic health record data quality assessment: enabling reuse for clinical research. *J Am Med Inform Assoc* 20:144–151. <https://doi.org/10.1136/amiajnl-2011-000681>
 19. Stahlman S, Oetting AA (2018) Mental health disorders and mental health problems, active component, U.S. Armed Forces, 2007–2016. *MSMR* 25:2–11
 20. Hoge CW, Castro CA, Messer SC et al (2004) Combat duty in Iraq and Afghanistan, mental health problems, and barriers to care. *N Engl J Med* 351:13–22. <https://doi.org/10.1056/NEJMoa040603>
 21. Welsh MM, Federinko SP, Burnett DG et al (2014) Deployment-related depression screening, 2001–2008: comparing clinical versus research surveys. *Am J Prev Med* 47:531–540. <https://doi.org/10.1016/j.amepre.2014.07.036>
 22. Morgan JK, Hourani L, Lane ME, Tueller S (2016) Help-seeking behaviors among active-duty military personnel: utilization of Chaplains and Other Mental Health Service Providers. *J Health Care Chaplain* 22:102–117. <https://doi.org/10.1080/08854726.2016.1171598>
 23. Office of the Deputy Assistant Secretary of Defense for Military Community and Family Policy (2015) 2015 Demographics Profile of the Military Community
 24. Thériault FL, Gabler K, Naicker K (2016) Health and lifestyle information survey of Canadian Armed Forces Personnel 2013/2014—Regular Forces Report. Department of National Defence, Ottawa, Canada
 25. Twomey CD, Baldwin DS, Hopfe M, Cieza A (2015) A systematic review of the predictors of health service utilisation by adults with mental disorders in the UK. *BMJ Open* 5:e007575. <https://doi.org/10.1136/bmjopen-2015-007575>
 26. Ferrari AJ, Somerville AJ, Baxter AJ et al (2013) Global variation in the prevalence and incidence of major depressive disorder: a systematic review of the epidemiological literature. *Psychol Med* 43:471–481. <https://doi.org/10.1017/S0033291712001511>
 27. Hines LA, Goodwin L, Jones M et al (2014) Factors affecting help seeking for mental health problems after deployment to Iraq and Afghanistan. *Psychiatr Serv* 65:98–105. <https://doi.org/10.1176/appi.ps.004972012>
 28. Fikretoglu D, Liu A, Pedlar D, Brunet A (2010) Patterns and predictors of treatment delay for mental disorders in a nationally representative, active Canadian military sample. *Med Care* 48:10–17. <https://doi.org/10.1097/MLR.0b013e3181bd4bf9>
 29. Boulos D, Zamorski MA (2016) Delay to mental healthcare in a cohort of Canadian Armed Forces personnel with deployment-related mental disorders, 2002–2011: a retrospective cohort study. *BMJ Open* 6:e012384. <https://doi.org/10.1136/bmjopen-2016-012384>
 30. Bonde JP, Utzon-Frank N, Bertelsen M et al (2016) Risk of depressive disorder following disasters and military deployment: systematic review with meta-analysis. *Br J Psychiatry* 208:330–336. <https://doi.org/10.1192/bjp.bp.114.157859>
 31. Boulos D, Zamorski MA (2016) Contribution of the mission in Afghanistan to the burden of past-year mental disorders in Canadian Armed Forces Personnel, 2013. *Can J Psychiatry* 61:64S–76S. <https://doi.org/10.1177/0706743716628857>
 32. Zamorski MA, Bennett RE, Boulos D et al (2016) The 2013 Canadian Forces Mental Health Survey. *Can J Psychiatry* 61:10S–25S. <https://doi.org/10.1177/0706743716632731>
 33. Katon WJ (2011) Epidemiology and treatment of depression in patients with chronic medical illness. *Dialogues Clin Neurosci* 13:7–23
 34. Hom MA, Stanley IH, Schneider ME, Joiner TE (2017) A systematic review of help-seeking and mental health service utilization among military service members. *Clin Psychol Rev* 53:59–78. <https://doi.org/10.1016/j.cpr.2017.01.008>
 35. Kendler KS, Karkowski LM, Prescott CA (1999) Causal relationship between stressful life events and the onset of major depression. *Am J Psychiatry* 156:837–841. <https://doi.org/10.1176/ajp.156.6.837>
 36. Mojtabai R, Olfson M (2006) Treatment Seeking for Depression in Canada and the United States. *Psychiatr Serv* 57:631–639. <https://doi.org/10.1176/ps.2006.57.5.631>
 37. Quartana PJ, Wilk JE, Thomas JL et al (2014) Trends in mental health services utilization and stigma in US soldiers from 2002 to 2011. *Am J Public Health* 104:1671–1679. <https://doi.org/10.2105/AJPH.2014.301971>
 38. Crumlish N, Kelly BD (2009) How psychiatrists think. *Adv Psychiatr Treat* 15:72–79. <https://doi.org/10.1192/apt.bp.107.005298>